

Small sawtooth regimes in JET plasmas

M. F. F. Nave¹, S. Coda³, R. Galvão¹, J. Graves³, R. Koslowski⁶, M. Mantsinen⁴, F. Nabais¹, S. Sharapov², P. de Vries², R. Buttery², M. de Baar⁷, C. Challis², J. Ferreira¹, C. Giroud², M. Mayoral², S. D. Pinches⁵, M. Stamp², and JET-EFDA contributors*

¹Associação EURATOM/IST, Centro de Fusão Nuclear, Lisbon, Portugal

²Euratom/UKAEA Fusion Association, Culham Science Centre, Abingdon, UK

³CRPP, Association EURATOM-Confederation Suisse, EPFL, Lausanne, Switzerland

⁴Helsinki University of Technology, Association Euratom-Tekes, Finland

⁵Max-Planck-Institut für Plasmaphysik, IPP-EURATOM Assoziation, Garching, Germany

⁶Association EURATOM-FZ-Juelich, Institut fuer Plasmaphysik, Juelich, Germany


⁷FOM Inst. Voor Plasmafysica "Rijnhuizen", Netherlands

*see appendix of the paper by J. Pamela "Overview of recent JET results", Proc. IAEA conference on Fusion Energy, Sorrento, 2000



I – Introduction – Increased sawtooth stabilization leading to long sawtooth periods is a known observation of auxiliary heated plasmas. However, small sawtooth regimes characterised by sawtooth periods, $\tau_{ST}=20-100$ ms, that are shorter than observed in Ohmic regimes have been obtained at JET with ion-cyclotron radio frequency (ICRF) heating [1] as well as with neutral beam injection (NBI) heating [2]. At JET, decreasing the sawtooth period is known to delay the onset of NTMs. Results of two sawtooth studies from recent JET experimental campaigns are reported here.

ICRF heating is usually associated with long sawtooth periods and large crashes referred to as "monster" sawtooth ($\tau_{ST} > 400$ ms). Sawtooth stabilization is understood as increased stability of the internal kink mode from ICRF driven fast ions [3]. In contrast, JET plasmas with high fast ion energy content, obtained by applying ICRF heating to a low-density target plasma, show a very unstable sawtooth regime. As the density is increased an abrupt change from small sawteeth to monster sawteeth is observed [1]. The transition occurs for densities in the range $1.5-2 \times 10^{19} \text{ m}^{-3}$.

Another regime of small sawtooth is observed with NBI heating in counter injection and the toroidal field, B_T , reversed [2]. With counter-NBI the observed plasma rotation in the plasma core is reduced when compared with the usual B_T , co-rotating plasmas. During a recent reversed B_T experimental campaign an experiment was performed to investigate rotation effects on sawtooth stability and in particular, to assess the internal kink stability models [4] that predict increased instability in regimes of low shear rotation.




co-NBI where the observed sawtooth period increases with power, with counter-NBI the sawtooth period decreased up to a minimum at 4 MW (figure 2). A similar trend was observed with core toroidal plasma rotation. With counter-NBI the observed plasma rotation in the core of the plasma (obtained from charge exchange measurements) is reduced when compared with co-rotating plasmas. With co-NBI the sawtooth period increased with plasma rotation. With counter-NBI the sawtooth period reduced to a minimum for plasma rotation at ~ 2 kHz (figure 3), corresponding to the minimum sawtooth period obtained as a function of P_{NBI} .



With reversed B_T , the toroidal rotation is in opposition to characteristic frequencies ω_{hi} and $\langle \omega_{hi} \rangle$. Plasma rotation in the core of the plasma is reduced when compared with co-NBI. A

candidate mechanism consistent with the main features of the observations is the dependence of the internal kink stability on sheared toroidal rotation [4]. Kinetic effects from thermal ions in a regime where the mode frequency $\omega \sim \omega_{*i} \sim \omega_{di}$, are expected to be destabilizing.

III –Sawteeth observations with ICRH in low density plasmas - Sawtooth behaviour was studied in plasmas with a high fast-ion energy content obtained by applying ICRF heating to low density discharges, with $B_T=2.7$ T, $I_p= 2.5$ MA (In the usual B_T configuration). ICRF powers of 3-6 MW were applied using a minority Hydrogen heating in Deuterium plasmas. Different ICRF toroidal phases and resonant positions were considered. Here results for on-axis heating, dipole phasing, are shown. In discharges with ICRF heating only the plasma density was controlled by gas puffing (figure 4). In others the density increased when a small amount of NBI power ($P_{NBI}=1-1.5$ MW) was added (figure 5). In both cases, an abrupt transition from small sawtooth to monster sawtooth, was observed when the density increased.



Small sawtooth, with periods ≤ 100 ms are observed for central densities $< 2 \times 10^{19} \text{ m}^{-3}$. The density threshold is in the range $1.5-2.0 \times 10^{19} \text{ m}^{-3}$ depending on heating details. For similar discharges, the critical density was lower when a small amount of NBI was added (figure 6). The fast ion stabilizing influence on the internal kink mode is predicted to reduce for large ICRF heating powers [3]. Modelling with the CASTOR-K code confirms that as the fast ion temperature increases ($T_{hot} \propto n_e^{-1}$) and the fast particle orbits become wider the fast ion stabilization effect decreases [5]. In this case kinetic effects of trapped thermal ions and




Figure.6 - Central density (top) and central electron temperature (bottom) versus sawtooth period, for similar pulses with ICRF only (open) and with added NBI

diamagnetic effects on the ideal MHD growth rate, not included in [5], may play a role in determining the sawtooth stability.

The $n=1$ sawtooth precursor and postcursor oscillations observed inside the sawtooth inversion radius during the small sawtooth regime indicate low shear rotation with $f_{\text{MHD}}=100\text{-}500$ Hz. Low shear rotation is destabilizing [3]. The observed mode frequency is $\omega < \omega_{*1}$. If NBI is applied, the mode frequency increases rapidly to values 1-2 kHz, making $\omega \geq \omega_{*1}$. Charge exchange measurements show that the rotation profiles

become peaked. The increased stability with rotation might explain the lower density threshold for transition into “monster” sawteeth that is observed when NBI is applied.

IV – Conclusions – In the usual JET machine operation, with co-NBI the sawtooth period increases with P_{NBI} . In contrast with counter-NBI the sawtooth period decreases up to a minimum at $P_{\text{NBI}} = 4\text{MW}$. Sawtooth periods shorter than those for Ohmic heating are obtained. A similar trend is observed with plasma rotation. Kinetic effects from trapped thermal ions can be destabilising for small sheared counter rotation [4].

With ICRF heating the sawtooth period depends sensitively on the density. A critical density ($n_e < 1.5\text{-}2 \times 10^{19} \text{m}^{-3}$) separates the small sawtooth from the usual “monster” sawtooth. Modelling shows that for low densities the stabilizing kinetic effects from fast ions is reduced, as the orbits of the ICRF driven fast ions increases. Recent experiments indicated the possible relevance of diamagnetic and shear rotation effects on the sawtooth stability in this regime. The abrupt change in sawtooth behaviour is not yet understood.

Acknowledgements

This work was performed under the European Fusion Development Agreement. It received financial support from Fundação para a Ciencia e Technology (FCT), Portugal, from the United Kingdom Engineering and Physical Sciences Research Council and from EURATOM.

References

- [1] M. Mantsinen et al, Nuclear Fusion 42 (2000) 1291
- [2] A. Edwards et al, Proc. 19th EPS Conf. On Contr. Fusion and Plasma Phys. (Innsbruck 29 June- 3 July 1992) vol. 16 C Part I, p. 379
- [3] F. Porcelli et al, Phys. Fluids, B 4 (1992) 10
- [4] J. Graves et al, Plasma Phys. Control Fusion 42 (2000) 1049
- [5] F. Nabais et al, submitted to Plasma Phys. Control Fusion (2004)