

PAPER • OPEN ACCESS

# Test of the electric charge conservation law with Borexino detector

To cite this article: A Vishneva *et al* 2016 *J. Phys.: Conf. Ser.* **675** 012025

View the [article online](#) for updates and enhancements.

## Related content

- [Test of the electron stability with the Borexino detector](#)  
A Vishneva, M Agostini, K Altenmüller et al.
- [Recent results from Borexino](#)  
G Testera, M Agostini, K Altenmüller et al.
- [Overview and accomplishments of the Borexino experiment](#)  
G Ranucci, M Agostini, S Appel et al.

## Recent citations

- [Test of the electron stability with the Borexino detector](#)  
A Vishneva *et al*

# Test of the electric charge conservation law with Borexino detector

A Vishneva<sup>1,29,\*</sup>, M Agostini<sup>2</sup>, S Appel<sup>2</sup>, G Bellini<sup>3</sup>, J Benziger<sup>4</sup>,  
 D Bick<sup>5</sup>, G Bonfini<sup>6</sup>, D Bravo<sup>7</sup>, B Caccianiga<sup>3</sup>, F Calaprice<sup>8</sup>,  
 A Caminata<sup>9</sup>, P Cavalcante<sup>6</sup>, A Chepurinov<sup>10</sup>, D D'Angelo<sup>3</sup>,  
 S Davini<sup>11</sup>, A Derbin<sup>12</sup>, L Di Noto<sup>9</sup>, I Drachnev<sup>11</sup>, A Etenko<sup>13</sup>,  
 K Fomenko<sup>1</sup>, D Franco<sup>14</sup>, F Gabriele<sup>6</sup>, C Galbiati<sup>8</sup>, C Ghiano<sup>9</sup>,  
 M Giammarchi<sup>3</sup>, M Goeger-Neff<sup>2</sup>, A Goretti<sup>8</sup>, M Gromov<sup>10</sup>,  
 C Hagner<sup>5</sup>, E Hungerford<sup>15</sup>, Aldo Ianni<sup>6</sup>, Andrea Ianni<sup>8</sup>,  
 K Jedrzejczak<sup>17</sup>, M Kaiser<sup>5</sup>, V Kobychiev<sup>18</sup>, D Korablev<sup>1</sup>, G Korga<sup>6</sup>,  
 D Kryn<sup>14</sup>, M Laubenstein<sup>6</sup>, B Lehnert<sup>19</sup>, E Litvinovich<sup>13,20</sup>,  
 F Lombardi<sup>6</sup>, P Lombardi<sup>3</sup>, L Ludhova<sup>3</sup>, G Lukyanchenko<sup>13,20</sup>,  
 I Machulin<sup>13,20</sup>, S Manecki<sup>7</sup>, W Maneschg<sup>22</sup>, S Marcocci<sup>11</sup>, E Meroni<sup>3</sup>,  
 M Meyer<sup>5</sup>, L Miramonti<sup>3</sup>, M Misiaszek<sup>17,6</sup>, M Montuschi<sup>23</sup>,  
 P Mosteiro<sup>8</sup>, V Muratova<sup>12</sup>, B Neumair<sup>2</sup>, L Oberauer<sup>2</sup>,  
 M Obolensky<sup>14</sup>, F Ortica<sup>24</sup>, M Pallavicini<sup>9</sup>, L Papp<sup>2</sup>, L Perasso<sup>9</sup>,  
 A Pocar<sup>26</sup>, G Ranucci<sup>3</sup>, A Razeto<sup>6</sup>, A Re<sup>3</sup>, A Romani<sup>24</sup>, R Roncin<sup>6,14</sup>,  
 N Rossi<sup>6</sup>, S Schönert<sup>2</sup>, D Semenov<sup>12</sup>, H Simgen<sup>22</sup>,  
 M Skorokhvatov<sup>13,20</sup>, O Smirnov<sup>1</sup>, A Sotnikov<sup>1</sup>, S Sukhotin<sup>13</sup>,  
 Y Suvorov<sup>27,13</sup>, R Tartaglia<sup>6</sup>, G Testera<sup>9</sup>, J Thurn<sup>19</sup>, M Toropova<sup>13</sup>,  
 E Unzhakov<sup>12</sup>, R B Vogelaar<sup>7</sup>, F von Feilitzsch<sup>2</sup>, H Wang<sup>27</sup>,  
 S Weinz<sup>28</sup>, J Winter<sup>28</sup>, M Wojcik<sup>17</sup>, M Wurm<sup>28</sup>, Z Yokley<sup>7</sup>,  
 O Zaimidoroga<sup>1</sup>, S Zavatarelli<sup>9</sup>, K Zuber<sup>19</sup> and G Zuzel<sup>17</sup> (Borexino  
 Collaboration)

<sup>1</sup> Joint Institute for Nuclear Research, 141980 Dubna, Russia

<sup>2</sup> Physik-Department and Excellence Cluster Universe, Technische Universität München, 85748 Garching, Germany

<sup>3</sup> Dipartimento di Fisica, Università degli Studi e INFN, 20133 Milano, Italy

<sup>4</sup> Chemical Engineering Department, Princeton University, Princeton, NJ 08544, USA

<sup>5</sup> Institut für Experimentalphysik, Universität, 22761 Hamburg, Germany

<sup>6</sup> INFN Laboratori Nazionali del Gran Sasso, 67010 Assergi (AQ), Italy

<sup>7</sup> Physics Department, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061, USA

<sup>8</sup> Physics Department, Princeton University, Princeton, NJ 08544, USA

<sup>9</sup> Dipartimento di Fisica, Università degli Studi e INFN, Genova 16146, Italy

<sup>10</sup> Lomonosov Moscow State University Skobeltsyn Institute of Nuclear Physics, 119234 Moscow, Russia

<sup>11</sup> Gran Sasso Science Institute (INFN), 67100 L'Aquila, Italy

<sup>12</sup> St. Petersburg Nuclear Physics Institute NRC Kurchatov Institute, 188350 Gatchina, Russia

<sup>13</sup> NRC Kurchatov Institute, 123182 Moscow, Russia

<sup>29</sup> Presenter. To whom any correspondence should be addressed.



<sup>14</sup> AstroParticule et Cosmologie, Université Paris Diderot, CNRS/IN2P3, CEA/IRFU, Observatoire de Paris, Sorbonne Paris Cité, 75205 Paris Cedex 13, France

<sup>15</sup> Department of Physics, University of Houston, Houston, TX 77204, USA

<sup>16</sup> Institute for Theoretical and Experimental Physics, 117218 Moscow, Russia

<sup>17</sup> M. Smoluchowski Institute of Physics, Jagiellonian University, 30059 Krakow, Poland

<sup>18</sup> Kiev Institute for Nuclear Research, 06380 Kiev, Ukraine

<sup>19</sup> Department of Physics, Technische Universität Dresden, 01062 Dresden, Germany

<sup>20</sup> National Research Nuclear University MEPhI (Moscow Engineering Physics Institute), Kashirskoe highway 31, Moscow 115409, Russia

<sup>21</sup> Kepler Center for Astro and Particle Physics, Universität Tübingen, 72076 Tübingen, Germany

<sup>22</sup> Max-Planck-Institut für Kernphysik, 69117 Heidelberg, Germany

<sup>23</sup> Dipartimento di Fisica e Scienze della Terra Università degli Studi di Ferrara e INFN, Via Saragat 1-44122, Ferrara, Italy

<sup>24</sup> Dipartimento di Chimica, Università e INFN, 06123 Perugia, Italy

<sup>25</sup> Physics Department, Queen's University, Kingston ON K7L 3N6, Canada

<sup>26</sup> Amherst Center for Fundamental Interactions and Physics Department, University of Massachusetts, Amherst, MA 01003, USA

<sup>27</sup> Physics and Astronomy Department, University of California Los Angeles (UCLA), Los Angeles, California 90095, USA

<sup>28</sup> Institute of Physics and Excellence Cluster PRISMA, Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany

E-mail: \*the2cherry2orchard@gmail.com

**Abstract.** The new limit on the electron lifetime is obtained from data of the Borexino experiment. The expected signal from the  $e \rightarrow \gamma\nu$  decay mode is a 256 keV photon detected in liquid scintillator. Because of the extremely low radioactive background level in the Borexino detector it was possible to improve the previous measurement by two orders of magnitude.

## 1. Introduction

The electric charge conservation law is a fundamental physical principle. There are no hints for violation of this law neither in theory within the Standard model nor in any experiment. Since the electric charge non-conservation (CNC) is admitted in exotic theories such as extra-dimensional theories [1], investigation of such processes is an evident way to search for physics beyond the Standard model.

The most frequently searched for CNC processes are decays of the electron into neutral particles. Two decay modes are usually accounted for experimentally:

$$e \rightarrow \gamma\nu, \quad (1)$$

where a monoenergetic 256 keV photon is searched for, and

$$e \rightarrow \nu\nu\nu, \quad (2)$$

where only effects due to the electron disappearance would be observed. However, the impossibility of occurrence of such processes is presented in [2], where it is shown that such decays would be followed by a huge amount of low-energy bremsstrahlung photons. For the process (1) it would mean the absence of 256 keV photon while the electron disappearance is more model-independent and the corresponding atomic effects in the case (2) would remain the same. Thus one can see that observing the 256 keV photon from the electron decay would mean not only CNC but also going beyond the Standard model.

## 2. Overview of experiments

Study of the electron stability has long experimental history. The list of the experiments in which the electron decay was being searched for is presented in table 1. There are also plans for

**Table 1.** Experimental tests for the electron stability.

year	material	limit for $e \rightarrow \gamma\nu$	limit for $e \rightarrow \nu\nu\nu$	CL	reference
1959	NaI	$10^{19}$	$10^{17}$	68%	[3]
1965	NaI	$4 \times 10^{22}$	$2 \times 10^{21}$	68%	[4]
1975	Ge	—	$5.3 \times 10^{21}$	68%	[5]
1979	NaI	$3.5 \times 10^{23}$	—	68%	[6]
1983	Ge	$3 \times 10^{23}$	$2 \times 10^{22}$	68%	[7]
1986	Ge	$1.5 \times 10^{25}$	—	68%	[8]
1993	Ge	$1.63 \times 10^{25}$	—	68%	[9]
1995	Ge	$2.1 \times 10^{25}$	$2.6 \times 10^{23}$	90%	[10]
1996	Xe	$2 \times 10^{25}$	$1.5 \times 10^{23}$	68%	[11]
1999	NaI	—	$(1.5 - 2.4) \times 10^{23}$	90%	[12]
1999	NaI	—	$2.4 \times 10^{24}$	90%	[13]
2000	Xe	$2 \times 10^{26}$	—	90%	[14]
2002	PXE	$4.6 \times 10^{26}$	—	90%	[15]
2007	Ge	$1.93 \times 10^{26}$	—	90%	[16]
2012	NaI	—	$1.2 \times 10^{24}$	90%	[17]

providing analogous studies at present and future experiments [18, 19, 20].

### 2.1. NaI detectors

Experiments based on NaI detectors were the first to provide the limits on the electron stability [3, 4]. The expected signal for the mode (2) is a photon with maximal energy of 33.2 keV emitted while filling the vacancy caused by the electron disappearance from the iodine K-shell. The decay to a photon and a neutrino is investigated by searching for the 256 keV photon. Various coincidence techniques are also applied in such detectors. First was the search of simultaneous 256 keV and 33.2 keV photons occurrence [4]. Another approach based on the electron capture by a nucleus without the consequent atomic number change was considered recently in [17]. Simultaneous observation of the 33.2 keV photons and the nucleus deexcitation (417.9 keV) would mean the electron disappearance.


### 2.2. Ge detectors

The electron stability is widely studied using germanium detectors. The main advantage of such detectors is good energy resolution (about 1 keV). In addition, the background level in the region of interest is lower than that in NaI detectors. The expected signal is a photon of energy 11.1 keV for the mode (2) and a 256 keV photon for the mode (1), respectively.

### 2.3. Liquid scintillators

The strongest limits on the electron lifetime with respect to the decay mode (1) during the last fifteen years have been obtained with liquid scintillation detectors. Their main advantages are large mass and a possibility of purification from radioactive contaminations. The first one was

Fit result for the electron decay rate = 1.23 cpd/100 tons



**Figure 1.** Borexino spectrum composition.

DAMA/LXe experiment [11, 14]. This detector contains 6.5 kg ( $\sim 2$  litres) of liquid xenon. This apparatus has rather low energy threshold and is sensitive to both electron decay modes.

The second one is the prototype of the Borexino detector, CTF-II [21]. Its main goal was to test the purification techniques developed for Borexino. During the tests various scintillators were used. CTF-II was filled with 4 tons of PXE (phenylxylylethane) which has less ionization quenching in comparison with PC (pseudocumene) used in Borexino. Large mass and extremely low background level made it possible to obtain a stronger limit of  $4.6 \times 10^{26}$  years (90% confidence level) on the electron lifetime in shorter exposure time. This result remained the best until the same study was performed in Borexino.

### 3. The Borexino detector

Borexino is a large volume scintillation detector located deep underground in the Laboratori Nazionali del Gran Sasso [22]. Its active media contains 278 tons of organic liquid scintillator, namely, pseudocumene (1,2,4-trimethylbenzene) with admixture of PPO (2,5-diphenyloxazole) at a concentration of 1.5 g/l. Borexino has extremely low background level in the region of interest, namely  $0.15 \text{ day}^{-1} \text{ ton}^{-1} \text{ keV}^{-1}$ . The energy threshold is above 50 keV so Borexino is not sensitive to the disappearance mode. By comparing sensitivity of CTF and Borexino the expected electron lifetime limit is estimated, which exceeds the previous one at two orders of magnitude.

## 4. The electron decay search

### 4.1. Analysis approach

The data set used in the analysis were acquired from January 2012 to May 2013 (Borexino Phase 2). This data set was obtained after the purification campaign [23] which reduced in particular the contamination of  $^{85}\text{Kr}$  and  $^{210}\text{Bi}$  which give a significant contribution in the low-energy region.

The same 408 days data set is successfully used in the measurement of solar pp-neutrino flux [24]. In this analysis the same energy range (150–600 keV) and parameters used in the fitting procedure are considered. The only difference is the addition of the 256 keV photon line in the fitting function. The sample of spectral fit is shown on Fig. 4.1. One can see that the sought-for peak (marked by arrow) is shifted to the lower energies due to ionization quenching.

### 4.2. Constraint on the pp-neutrino event rate

The 256 keV photon occurrence is strongly correlated with pp-neutrino event rate. Therefore treating the pp-neutrino rate as a free parameter in the fit leads to non-physical values at the limit. Indeed, the 256 keV photon event rate corresponding to the 90% confidence level ( $\simeq 12$  cpd/100 tons) corresponds to zero pp-neutrino rate which is not consistent with observations by radiochemical experiments [25]. As far as the latter ones are not sensitive to the electron decay it is reasonable to use their results to constrain the pp-neutrino event rate. This constraint gives the limit on the event rate of 1.23 cpd/100 tons. The lifetime limit is expressed as  $\tau \geq \epsilon N_e T / S_{\text{lim}}$ , where  $N_e$  is the total number of electrons in the detector,  $\epsilon$  is the fraction of electrons survived after the fiducial volume cut,  $T$  is the exposure time, and  $S_{\text{lim}}$  is the event rate limit. It gives the electron lifetime of  $\tau \geq 7.2 \times 10^{28}$  years.

### 4.3. Systematic errors study and final results

The main sources of the systematic errors in this study are the following. The most important is the precision of the scintillator light yield measurement ( $\sim 1\%$ ). It strongly influences the peak position and therefore affects the sensitivity. Another source of systematic errors is the fiducial mass measurement precision, which gives negligible effect. Choice of the energy estimator can also affect the result. In the present study two variables are used as energy estimators, namely, number of PMTs hit in the time intervals of 230 ns and 400 ns. After having accounted for all these effects the lifetime limit has become weaker and the final result for the electron lifetime limit is  $\tau_{e \rightarrow \gamma \nu} \geq 6.6 \times 10^{28}$  years at the 90% confidence level. This study is described in more details in [26].

## Acknowledgments

Russian colleagues from MEPhI acknowledge partial support from MEPhI Academic Excellence Project (contract No. 02.a03.21.0005, 27.08.2013).

## References

- [1] Dubovsky S L, Rubakov V A and Tinyakov P G 2000 *JHEP* **0008** 041
- [2] Okun L B 1989 *Sov. Phys. Usp.* **32** 543
- [3] Feinberg G and Goldhaber M 1959 *Proc. Natl. Acad. Sci. US* **45** 1301
- [4] Moe M K and Reines F 1965 *Phys. Rev.* **140** 992
- [5] Steinberg R I *et al.* 1975 *Phys. Rev. D* **12** 2582
- [6] Koval'chuk E L, Pomanskii A A and Smol'nikov A A 1979 *JETP Lett.* **29** 145
- [7] Belotti E *et al.* 1983 *Phys. Lett. B* **124** 435
- [8] Avignone III F T, Brodzinski R L, Hensley W K, Miley H S and Reeves J H 1986 *Phys. Rev. D* **34** 97
- [9] Balysh A *et al.* 1993 *Phys. Lett. B* **298** 278
- [10] Aharonov Y *et al.* 1995 *Phys. Lett. B* **353** 168
- [11] Belli P *et al.* 1996 *Astropart. Phys.* **5** 217-9

- [12] Belli P *et al.* 1999 *Phys. Rev. C* **60** 065501
- [13] Belli P *et al.* 1999 *Phys. Lett. B* **460** 236
- [14] Belli P *et al.* 2000 *Phys. Rev. D* **61** 117301
- [15] Back H O *et al.* 2002 *Phys. Lett. B* **525** 29
- [16] Klapdor-Kleingrothaus H V, Krivosheina I V and Titkova I V 2007 *Phys. Lett. B* **644** 109
- [17] Bernabei R *et al.* 2012 *Eur. Phys. J. C* **72** 1920
- [18] Aalseth C E *et al.* 2004 *Phys. Atom. Nucl.* **67** 2002
- [19] Ardito R *et al.* 2005 *arXiv:0501010*
- [20] Wurm M *et al.* 2012 *Astropart. Phys.* **35** 685-732
- [21] Alimonti G *et al.* 1998 *Nucl. Instrum. Meth. A* **406** 411
- [22] Alimonti G *et al.* 2009 *Nucl. Instrum. Meth. A* **600** 568-93
- [23] Bellini G *et al.* 2014 *Phys. Rev. D* **89** 112007
- [24] Bellini G *et al.* 2014 *Nature* **512** 383
- [25] Abdurashitov J N *et al.* 2009 *Phys. Rev. C* **80** 015807
- [26] Agostini M *et al.* 2015 *arXiv:1509.01223*