

Sleep deprivation alters the effective connectivity of central executive and salience network with the right intraparietal sulcus


E. Bettazzi^{1,2}, A. Silchenko¹, D. Elmenhorst^{1,3,4}, S. Eickhoff^{1,2}, M. Tahmasian^{1,2}, F. Hoffstaedter^{1,2}

¹Research Center Juelich GmbH, Juelich, Germany, ²Heinrich Heine University Düsseldorf, Düsseldorf, Germany, ³University Hospital Cologne, University of Cologne, Cologne, Germany, ⁴Rheinische Friedrich-Wilhelms-University Bonn, Bonn, Germany

BACKGROUND

Cognitive impairments observed after sleep deprivation (SD) mostly affect attention maintenance and emotional salience¹. Impairments, such as attention lapses, can be associated to the disrupted activity of the central executive network (CEN), while the impaired saliency discrimination is associated to the altered activity of the salience network (SN)¹. The intraparietal sulcus is a parietal region densely connected to the frontoparietal networks (as CEN), which has been referred to as a "connectivity hub" with roles in orientation, attention and salience-detection². Results from an ALE meta-analysis on 31 studies found the right intraparietal sulcus (rIPS) to be consistently hypoactive in total sleep deprivation³, but it is not clear how this region interacts with the above networks.


How do these brain areas influence each other?



To answer that, the effective connectivity can be investigated. An effective connectivity analysis provides information on how regions influence each other.

Positive/negative connectivity value (Hz) → **Increased/decreased influence of region A on region B**

METHODS



RESULTS


Pairwise correlation between difference matrices related to SD conditions

Central Executive Network (CEN)				Salience Network (SN)			
Acute Partial SD	1			Acute Partial SD	1		
Chronic Partial SD	0	1		Chronic Partial SD	0.17	1	
Total SD	0.03	-0.02	1	Total SD	-0.06	0.07	1

Whole-network connectivity changes across SD conditions

Network	Metric	Acute Partial SD	Chronic Partial SD	Total SD
CEN + rIPS	Impact	2.52	0.91	1.35
	Balance	+0.11	+0.04	-0.22
SN + rIPS	Impact	0.84	0.24	0.89
	Balance	+0.23	+0.09	+0.26

rIPS connectivity changes across SD conditions



CONCLUSIONS

- Sleep deprivation impacts the resting-state effective connectivity differently depending on the amount sleep debt.
- For both salience and central executive networks:
- rIPS connectivity altered in all three sleep deprivation conditions.
- rIPS influence on the network decreases at increasing sleep pressure.

- The hypoactivity of the rIPS is reflected in its decreased influence on the resting-state networks, salience and central executive networks. Such reduction in connectivity seems to be dose-dependent on the amount of sleep debt.
- We suggest that sleep deprivation disrupts the role of rIPS as a connectivity hub for resting-state networks, possibly resulting in cognitive dysfunctions as attentional lapses and emotional hyperarousal.

REFERENCES

[1] Krause et al., *Nat Rev Neurosci*, 2017. [2] Brown et al., *IJCHP*, 2023. [3] Javaheripour et al., *Sleep Med Rev*, 2019. [4] Nilsson et al., *Sci Rep*, 2017. [5] Chu et al., *Journal of Neuroscience*, 2023. [6] Schaefer et al., *Cerebral Cortex*, 2018. [7] Razi et al., *Netw Neurosci*, 2017.

CONTACT

e.bettazzi@fz-juelich.de
m.tahmasian@fz-juelich.de
f.hoffstaedter@fz-juelich.de

AFFILIATIONS

INM7, Research Center Juelich GmbH, Juelich, Germany
INM2, Research Center Juelich GmbH, Juelich, Germany