



Global aviation contrail climate effects from 2019 to 2021

Roger Teoh¹, Zebediah Engberg², Ulrich Schumann³, Christiane Voigt^{3,4}, Marc Shapiro², Susanne Rohs⁵ and Marc E.J. Stettler¹

¹Department of Civil and Environmental Engineering, Imperial College London, London, SW7 2AZ, United Kingdom

5 ²Breakthrough Energy, 4110 Carillon Point, Kirkland, WA 98033, United States

³Institute of Atmospheric Physics, Deutsches Zentrum für Luft- und Raumfahrt, 82234 Oberpfaffenhofen, Germany

⁴Institute of Atmospheric Physics, University Mainz, 55099 Mainz, Germany

⁵IEK-8 Troposphäre, Institut für Energie und Klimaforschung, Forschungszentrum Jülich GmbH, Jülich, Germany

Correspondence to: Marc E.J. Stettler (m.stettler@imperial.ac.uk)

10 **Abstract.** The global annual mean radiative forcing (RF) attributable to contrail cirrus is comparable to the RF from aviation's cumulative CO₂ emissions. Here, we simulate the global contrail climate forcing for 2019–2021 using reanalysis weather data and improved engine emission estimates along actual flight trajectories derived from Automatic Dependent Surveillance–Broadcast telemetry. Our 2019 global annual mean contrail net RF (62.1 mW m⁻²) is 44% lower than current best estimates for 2018 (111 [33, 189] mW m⁻²). Regionally, the contrail net RF is largest over Europe (876 mW m⁻²) and the US (414 mW
15 m⁻²), while the RF over East Asia (64 mW m⁻²) and China (62 mW m⁻²) are close to the global mean value because fewer flights in these regions form contrails as a result of lower cruise altitudes and limited ice supersaturated regions in the subtropics due to the Hadley Circulation. Globally, COVID-19 reduced the flight distance flown and contrail net RF in 2020 (-43% and -56% respectively vs. 2019) and 2021 (-31% and -49% respectively) with significant regional variation. Around 14% of all flights form a contrail with a net warming effect, yet only 2% of all flights account for 80% of the annual contrail energy
20 forcing. The spatiotemporal patterns of the most strongly warming and cooling contrail segments can be attributed to flight scheduling factors, aircraft–engine particle number emissions, tropopause height, background cloud and radiation fields, and albedo. Our contrail RF estimates are most sensitive to corrections applied to the global humidity fields, followed by assumptions on the aircraft-engine particle number emissions, and is least sensitive to radiative heating effects on the contrail plume and contrail-contrail overlapping. Accounting for the sensitivity analysis, we estimate a 2019 global contrail net RF of
25 62.1 [34.8, 74.8] mW m⁻².

1 Introduction

Aviation contributes to significant social and economic benefits, but also emits CO₂ and non-CO₂ pollutants that cause global warming and degrade air quality. In particular, aviation's cumulative CO₂ emissions account for one-third of its overall climate forcing, while the remaining two-thirds are estimated to arise from non-CO₂ components such as contrail cirrus, nitrogen
30 oxides (NO_x), particulate matter, and stratospheric water vapour emissions (Lee et al., 2021). When taken together, aviation



was responsible for ~3.5% of the global anthropogenic radiative forcing (RF) in 2018 with contrail cirrus estimated to contribute more than half of the aviation-induced RF (Lee et al., 2021; Kärcher, 2018).

Contrails form when flights traverse air masses with ambient temperatures below the Schmidt-Appleman criterion (SAC) threshold temperature (T_{SAC} , typically < 230 K). Aircraft particle emissions, which consist of non-volatile particulate matter (nvPM), metallic compounds, semi-volatile organic and sulfuric particles (Petzold et al., 2005), and ambient natural aerosols can activate into droplets and freeze to form contrail ice crystals (Schumann, 1996; Kärcher, 2018; Kleine et al., 2018). The nvPM acts as the primary source of condensation nuclei in the “soot-rich” regime, defined when the soot number emissions index (EI_n) exceeds 10^{13} kg^{-1} , while ambient aerosols, organic and sulfuric particles can nucleate under “soot-poor” conditions ($EI_n < 10^{13} \text{ kg}^{-1}$). Most kerosene-burning aircraft engines typically have nvPM EI_n of 10^{14} – 10^{16} kg^{-1} (EASA, 2021; Petzold et al., 1999; Moore et al., 2017; Durdina et al., 2017), and for these aircraft types, in-situ measurements and modelling studies show that the nvPM EI_n influences various contrail properties and associated climate forcing (Voigt et al., 2021; Bräuer et al., 2021b; Teoh et al., 2022a; Jeßberger et al., 2013; Kärcher, 2016). However, there is a small but increasing share of aircraft types powered by staged combustors with nvPM EI_n as low as $\sim 10^{11} \text{ kg}^{-1}$ (EASA, 2021; Boies et al., 2015), for which the initial contrail properties need further investigation (Voigt et al., 2022).

Over time, contrails formed in ice supersaturated regions (ISSR) can persist, spread, mix with natural cirrus, and transition into contrail cirrus clusters with observed lifetimes of up to 19 h (Haywood et al., 2009). The contrail spreading rate and coverage area is predominantly determined by horizontal wind components, wind shear and ice crystal sedimentation, while contrail lifetime is dependent on the ambient relative humidity with respect to ice (RHi), atmospheric turbulence, and the rate of ice crystal losses (Schumann and Heymsfield, 2017; Lewellen et al., 2014; Lewellen, 2014; Li et al., 2023). Contrail cirrus also interacts with solar and terrestrial radiation in two distinct ways. Firstly, it reflects incoming shortwave (SW) radiation and contribute to a cooling effect, while trapping and re-emitting outgoing longwave (LW) radiation causing a warming effect at all times (Meerkötter et al., 1999). Secondly, the absorption of SW and LW radiation heat up the contrail, which can drive plume-internal turbulence and local updraft, thereby changing the plume RHi and sublimation rate of contrail ice crystals (Jensen et al., 1998; Schumann et al., 2010; Schumann and Heymsfield, 2017; Lewellen, 2014; Unterstrasser and Gierens, 2010).

Previous studies have utilised air traffic data for 2002 (Eyers et al., 2005) and 2006 (Wilkerson et al., 2010) to estimate the global annual mean contrail cirrus net RF. Using the European Center/Hamburg General Circulation Model Version 4 (ECHAM4), Burkhardt & Kärcher (2011) estimated the 2002 global annual mean contrail net RF to be 37.5 mW m^{-2} . Bock & Burkhardt (2016a) updated the representation of contrail microphysical and optical properties within the ECHAM5 climate model (Lohmann et al., 2008; Bock and Burkhardt, 2016b) and estimated the global contrail net RF for 2002 (35 mW m^{-2}) and 2006 (56 mW m^{-2}). A follow-up study lowered the 2006 estimate by 22% (from 56 to 44 mW m^{-2}) because the initial contrail ice crystal numbers from Bock & Burkhardt (2016a) did not account for: (i) the lower nvPM activation rate when ambient temperatures are close to T_{SAC} ; and (ii) ice crystal losses in the wake vortex phase (Bier and Burkhardt, 2022). Chen & Gettelman (2013) applied the Community Atmosphere Model (CAM5) to obtain a 2006 global contrail net RF of $13 \pm 10 \text{ mW}$



65 m^{-2} , but this RF value was later revised to 57 mW m^{-2} after the simulation was re-run with initial contrail properties that are consistent with in-situ measurements (Lee et al., 2021). Schumann et al. (2015) coupled the contrail cirrus prediction model (CoCiP) with CAM3 to account for humidity exchange between contrails and the background air, estimating the 2006 global contrail net RF to be 63 mW m^{-2} (or 74 mW m^{-2} without humidity exchange).

Lee et al. (2021) compiled results from these studies and used the growth in annual flight distance flown to extrapolate the
70 2006 global annual mean contrail net RF to 2018 ($111 [33, 189] \text{ mW m}^{-2}$, 95% confidence interval). However, the extrapolation could lead to inaccuracies because: (i) the formation and climate forcing of contrails has a spatiotemporal dependence (Lamquin et al., 2012; Schumann et al., 2012; Bier and Burkhardt, 2022); (ii) air traffic growth was not uniform across the globe (ICAO, 2014, 2016); and (iii) there can be significant interannual variability in the contrail climate forcing (Wilhelm et al., 2021; Teoh et al., 2022a). In addition, existing global contrail studies generally assume a constant particle number
75 emissions which do not account for differences in nvPM EI_n between aircraft-engine types, and therefore, were unable to evaluate the variabilities in the contrail climate forcing that arise from individual flights and identify the set of flights with strongly warming/cooling contrails. Regional studies have also found that 80% of the contrail climate forcing were caused by 2–12% of all flights (Teoh et al., 2020, 2022a), but the applicability of these findings on the global level remain unknown.

In this study, we use a new global aviation emissions inventory based on Automatic Dependent Surveillance–Broadcast (ADS-
80 B) telemetry (GAIA) (Teoh et al., 2023) to: (i) quantify the global contrail properties and climate forcing for 2019–2021; (ii) identify the set of conditions that causes flights to form strongly warming/cooling contrails; (iii) evaluate the sensitivity of the simulated contrail climate forcing to aircraft emissions, meteorology, and contrail model parameters; and (iv) compare our global contrail RF estimates with existing studies.

2 Materials and methods

85 This section outlines the datasets and models used to achieve the stated research objectives. Section 2.1 describes a new global aviation emissions inventory, which includes actual trajectories, aircraft performance parameters and nvPM emissions from individual flights (Teoh et al., 2023). Section 2.2 provides an overview of the European Centre for Medium-Range Weather Forecast (ECMWF) ERA5 high-resolution realisation (HRES) reanalysis dataset (ECMWF, 2021; Hersbach et al., 2020) and extends a methodology to address known limitations in the ERA5 HRES humidity fields. Section 2.3 describes the CoCiP
90 contrail model (Schumann, 2012; Schumann et al., 2012), Section 2.4 summarises the climate forcing metrics used in this study, while Section 2.5 sets out the difference in model setup that is used to conduct a sensitivity analysis. Further information not included in the main text can be found in the Supporting Information (SI).

2.1 Global aviation emissions inventory

Global airspace surveillance systems have been transitioning towards the ADS-B standard, which enables real-time tracking
95 of flights at high spatiotemporal resolutions and over remote regions that previously lack radar coverage (ICAO, 2021;



EUROCONTROL, 2021). Recently, Teoh et al. (2023) used global ADS-B telemetry data to derive historical flight trajectories and develop a new aviation emissions inventory for 2019–2021. The dataset, known as the global aviation emissions inventory based on ADS-B (GAIA), captures 103.7 million unique flight and contains the: (i) flight metadata, including the unique flight identifier, origin and destination airports, and aircraft-engine type; and (ii) flight-waypoint data provided at time intervals of 40–60 s, including the 3D position, time, fuel consumption, aircraft mass, overall propulsion efficiency, and the nvPM EI_n. All flights are assumed to be powered by conventional Jet A-1 fuel.

Individual aircraft types can be powered by different engine options (Quadros et al., 2022; Teoh et al., 2023), where their nvPM EI_n can vary by up to five orders of magnitude and influence various contrail properties (Schumann, 1996; EASA, 2021; Teoh et al., 2022a). However, due to the lack of data, previous contrail studies (Teoh et al., 2020; Schumann et al., 2021; Teoh et al., 2022a) assigned a default aircraft-engine combination that is provided by the Base of Aircraft Data (BADA) aircraft performance model. GAIA partially addresses this limitation by using the registered aircraft tail number to extract the specific aircraft variant and engine model from a global fleet database (Cirium, 2022) whenever possible, covering 59% of all flights or 79% of flights with jet aircraft (Teoh et al., 2023).

For each flight, GAIA uses: (i) BADA 4.2 and 3.15 to estimate the fuel consumption and overall propulsion efficiency (EUROCONTROL, 2016, 2019); (ii) regional monthly passenger load factors to estimate the fuel requirements and initial aircraft mass; (iii) the ICAO Aircraft Engine Emissions Databank (EDB) to construct the nvPM emissions profile for each engine type (EASA, 2021); and three approaches to estimate the nvPM EI_n including (iv) the T_4/T_2 methodology (Teoh et al., 2022a, b, 2023), which uses the ratio of turbine-inlet temperature (T_4) to compressor-inlet temperature (T_2) to interpolate the nvPM emissions profile provided by the ICAO EDB (82% of the total flight distance flown); (v) the fractal aggregates model (Teoh et al., 2019, 2020) for older engine types without engine-specific nvPM measurements (11% of flight distance flown); and (vi) a constant value of 10^{15} kg^{-1} for remaining flights without engine-specific data (Teoh et al., 2020; Schumann et al., 2015). Further information on GAIA is detailed in Teoh et al. (2023).

2.2 Meteorology

Global meteorological and radiation data is provided by the ERA5 HRES reanalysis (Hersbach et al., 2020), which can be publicly downloaded from the ECMWF Copernicus Climate Data Store at a longitude-latitude grid resolution of $0.25^\circ \times 0.25^\circ$ over 37 pressure levels and at a 1 h time resolution (ECMWF, 2021). At altitudes of 25,000–42,000 feet, we note that the ERA5 HRES reanalysis provides meteorological data at 6 pressure levels which corresponds to a mean vertical resolution of ~3000 feet.

The simulated contrail properties and lifetime are highly sensitive to RH_i (Schumann, 2012; Schumann et al., 2021; Teoh et al., 2022a). However, humidity fields provided by ERA5-products contain several limitations. First, the ERA5-derived ISSR coverage area could be overestimated relative to radiosonde measurements (Agarwal et al., 2022), or underestimated when compared with in-situ measurements (Reutter et al., 2020). Second, the RH_i magnitude within ISSRs are generally weakly supersaturated (RH_i ≈ 100%), rarely exceeding RH_i > 120%, and are inconsistent with in-situ measurements (Reutter et al.,



2020; Gierens et al., 2020; Teoh et al., 2022a). Teoh et al. (2022a) recently developed a humidity correction methodology so that the corrected RH_i from the ERA5 HRES has a probability density function that is consistent with in-situ measurements from the European research infrastructure In-service Aircraft for a Global Observing System (IAGOS) (Petzold et al., 2015; Boulanger et al., 2022),

$$\text{RH}_{i,\text{corrected}} = \begin{cases} \frac{\text{RH}_i}{a_{\text{opt}}} & \text{for } \left(\frac{\text{RH}_i}{a_{\text{opt}}}\right) \leq 1 \\ \min\left(\left(\frac{\text{RH}_i}{a_{\text{opt}}}\right)^{b_{\text{opt}}}, \text{RH}_{i,\text{max}}\right) & \text{for } \left(\frac{\text{RH}_i}{a_{\text{opt}}}\right) > 1 \end{cases}, \quad (1)$$

where $a_{\text{opt}} = 0.9779$, $b_{\text{opt}} = 1.635$ and $\text{RH}_{i,\text{max}} = 1.65$ were calibrated for the North Atlantic region (40–75° N and 10–50° W). However, these coefficients cannot be applied globally because the RH_i errors have a latitude dependence where the ERA5-derived ISSR coverage area could be overpredicted at the tropics and subtropics (0–40° N) and underpredicted at latitudes above 40° N (Table S1).

To simulate contrails globally, we extend Eq. (1) by using the global IAGOS dataset for 2019 (Petzold et al., 2020; Boulanger et al., 2022), consisting of 2,161 flights and 682,308 data points. The IAGOS dataset is split into latitude bands of 10° intervals to avoid oversampling at specific latitudes. For each latitude band, a_{opt} is optimised so that the ISSR occurrence from the ERA5 and IAGOS have a symmetrical false positive and negative rate, while b_{opt} is optimised by minimising the Cramer-von Mises test statistic (Parr and Schucany, 1980) so that the RH_i distribution is consistent with in-situ measurements (Table S2). The optimised a_{opt} and b_{opt} for each latitude band are then fitted with a sigmoid to capture the rapid change in tropopause height between 20–50° N/S (Santer et al., 2003),

$$a_{\text{opt}} = \frac{0.06262}{1 + \exp(0.04589 \times (|\text{lat}| - 39.25))} + 0.9522, \quad (2)$$

$$b_{\text{opt}} = \frac{1.471}{1 + \exp(0.04431 \times (|\text{lat}| - 18.76))} + 1.433. \quad (3)$$

We also revise $\text{RH}_{i,\text{max}}$ to ensure that $\text{RH}_{i,\text{corrected}}$ is thermodynamically realistic, i.e., below water saturation and below the threshold that leads to homogeneous ice nucleation (Pruppacher et al., 2007; Kärcher and Lohmann, 2002; Tompkins et al., 2007),

$$\text{RH}_{i,\text{max}} = \begin{cases} \frac{p_{\text{liq}}(T_w)}{p_{\text{ice}}(T_w)}, & \text{when } T_w > 235 \text{ K} \\ 1.67 + (1.45 - 1.67) \times \frac{(T_w - 190)}{(235 - 190)}, & \text{when } T_w \leq 235 \text{ K} \end{cases}, \quad (4)$$

where T_w is the ambient temperature, and $p_{\text{liq}}(T_w)$ and $p_{\text{ice}}(T_w)$ are the saturation pressure of water vapour over liquid water and ice respectively that is estimated using Eq. (S3) and (S4). Application of the global humidity correction to the ERA5 HRES leads to: (i) a smaller ISSR coverage area around the tropics; (ii) larger ISSR coverage above 40° N and below 40° S; and (iii)



a higher occurrence of localised regions with RHi above 140% (Fig. S5). Further details on the extended humidity correction
 155 are listed in the SI §S1.3.

2.3 Contrail simulation

CoCiP simulates the properties and lifecycle of individual contrail segments along a flight trajectory (Schumann, 2012; Schumann et al., 2012). Two consecutive flight waypoints that satisfy the SAC form a contrail segment (Schumann, 1996), and persistent contrail segments are defined when their post-wake vortex ice water content (IWC) is greater than 10^{-12} kg kg⁻¹. The initial ice crystal number per contrail length ($n_{ice, initial}$) is calculated by,
 160

$$n_{ice, initial} = \max(nvPM EI_n, 10^{13}) \times \dot{m}_{f, dist} \times p_{activation} \times f_{surv}, \text{ where} \quad (5)$$

$$p_{activation} = -0.661 \exp(dT_{SAC}) + 1, \quad (6)$$

$$dT_{SAC} = T_w - T_{SAC} (< 0, \text{ in K}), \text{ and} \quad (7)$$

$$f_{surv} = \frac{IWC_1}{IWC_0}. \quad (8)$$

165 The lower bound of nvPM EI_n is constrained to 10^{13} kg⁻¹ to account for the potential activation of semi-volatile particles and ambient aerosols (Kärcher, 2018). $\dot{m}_{f, dist}$ is the fuel consumption per distance flown, $p_{activation}$ is the proportion of nvPM that activates to form contrail ice crystals (Bräuer et al., 2021a; Teoh et al., 2022a), and f_{surv} is the difference in IWC before and after the wake vortex phase as denoted by the subscripts “0” and “1” respectively (Schumann, 2012). Contrail evolution is simulated using a Runge-Kutta scheme with model time steps of 300 s, a time resolution that is higher than previous CoCiP
 170 studies (1800–3600 s) which were constrained by computational demands (Schumann, 2012; Schumann et al., 2015; Teoh et al., 2020, 2022a). Persistent contrail segments reach their end-of-life when the: (i) contrail ice number concentration falls below background levels of 10^3 m⁻³; (ii) contrail optical depth ($\tau_{contrail}$) is lower than 10^{-6} ; (iii) contrail age exceeds the maximum contrail lifetime that is set to 12 h; or (iv) mid-point of the contrail plume falls outside the defined altitude domain of between 6 and 13 km. The 13 km upper bound in condition (iv) was previously applied in Teoh et al. (2020) and could lead
 175 to a small underestimation of persistent contrail formation by military aircraft and private jets in the tropics, where 0.2% of the global annual flight distance flown in GAIA occurred above 13 km and between -30°S and 30°N. CoCiP is set up in its original form without accounting for humidity exchange between contrails and the atmosphere and without radiative effects from contrail-contrail overlapping (Schumann, 2012; Teoh et al., 2020, 2022a). Previous studies estimated that the annual mean contrail net RF could reduce by: (i) 15–20% when CoCiP was coupled with a general circulation model to account for
 180 the contrail–atmosphere humidity exchange; and (ii) 3% globally, or up to 57% in regions with dense air traffic when the effects from contrail-contrail overlapping are included (Schumann et al., 2015, 2021; Sanz-Morère et al., 2021). CoCiP has also been updated to incorporate the radiative heating effects on the contrail plume, where ice crystal losses are enhanced by the cumulative radiative energy absorbed by the contrail, which increases plume temperature and suppresses ice supersaturation,



and differential heating rate, which drives convective turbulence and vertical mixing (Jensen et al., 1998; Schumann and Graf, 2013; Schumann and Heymsfield, 2017; Schumann et al., 2010). Both quantities are estimated in accordance with Schumann et al. (2010), who developed a parametric model by least squares fit to outputs from the libRadtran radiative transfer model (Mayer and Kylling, 2005). We note that CoCiP algorithm has recently been open sourced and can be accessed via the pycontrails repository on GitHub (Shapiro et al., 2023), with the global contrail simulations in this study conducted using pycontrails v0.30.0. The regional contrail properties and climate forcing are estimated using rectangular spatial bounding boxes (Table S5 and Fig. S7) that are consistent with previous studies (Wilkerson et al., 2010; Hoare, 2014; Teoh et al., 2023). The five different output formats provided by CoCiP are described in the SI §S2.

2.4 Climate forcing metrics

Four different metrics are used to report the contrail climate forcing, the: (i) local contrail RF (RF'), i.e., the change in radiative flux over the area covered by the contrail (Schumann et al., 2012); (ii) energy forcing (EF_{contrail}) which provides the cumulative contrail climate forcing over its lifetime (Schumann et al., 2011),

$$EF_{\text{contrail}} [J] = \int_0^T RF'_{\text{net}}(t) \times L(t) \times W(t) dt, \quad (9)$$

where T , L and W are the contrail segment lifetime, length, and width respectively; (iii) annual mean contrail RF which quantifies the change in radiative flux over the globe/region at a given time, and is approximated from the annual EF_{contrail} assuming a linear relationship between contrail cover and RF,

$$\text{Annual mean contrail net RF } [W \text{ m}^{-2}] = \frac{\sum EF_{\text{contrail}} [J]}{S_{\text{region}} [m^2] \times (365 \times 24 \times 60^2 [s])}, \quad (10)$$

where S_{region} is the surface area of the region of interest; and (iv) the global annual mean contrail effective radiative forcing (ERF) which is estimated by assuming an ERF/RF ratio of 0.42 (Lee et al., 2021). As CoCiP simulates the full lifecycle of contrails (from line-shaped clouds to contrail cirrus), our climate forcing estimates are comparable to estimates of RF due to “contrail cirrus” in previous studies.

We also approximate the contrail cirrus global warming potential over a 20-year (GWP_{20}) and 100-year (GWP_{100}) time horizon using the energy forcing metric,

$$\text{Contrail cirrus } GWP_{\text{TH}} = \frac{EF_{\text{contrail}} \times \left(\frac{\text{ERF}}{\text{RF}}\right)}{EF_{\text{CO}_2, \text{TH}}}. \quad (11)$$

The ERF/RF ratio is assumed to be 0.42, and the CO_2 energy forcing ($EF_{\text{CO}_2, \text{TH}}$) is calculated according to Teoh et al. (2020),

$$EF_{\text{CO}_2, \text{TH}} [J] = \int_0^{\text{TH}} RF_{\text{CO}_2} dt \times S_{\text{Earth}} \approx AGWP_{\text{CO}_2, \text{TH}} \times (365 \times 24 \times 60^2) \times m_{\text{CO}_2} \times S_{\text{Earth}}, \quad (12)$$



210 where S_{Earth} is the Earth surface area ($5.101 \times 10^{14} \text{ m}^2$), m_{CO_2} is the total CO_2 emissions, and the CO_2 absolute global warming potential over 20-years ($\text{AGWP}_{\text{CO}_2,20}$) and 100-years ($\text{AGWP}_{\text{CO}_2,100}$) are assumed to be 2.39×10^{-14} and $88.0 \times 10^{-15} \text{ yr W m}^{-2} \text{ kg}^{-1}$ respectively (Gaillot et al., 2023).

2.5 Sensitivity analysis

215 Earlier studies suggest that the simulated contrail properties are highly sensitive to the humidity fields, aircraft performance (fuel consumption and overall propulsion efficiency), nvPM particle number emissions, and contrail model parameters (Schumann et al., 2021; Teoh et al., 2022a; Bier and Burkhardt, 2022). However, these sensitivity studies are limited because the simulations: (i) only cover a specific region; (ii) were re-run using different constants of soot EI_n (Bier and Burkhardt, 2022); or (iii) assume the default aircraft-engine assignment that is provided by BADA (Schumann et al., 2021; Teoh et al., 2022a, 2020).

We perform a sensitivity analysis by re-running the global contrail simulation for 2019 with seven distinct set-ups: (i) a simulation without humidity corrections applied to the ERA5 HRES (Section 2.2); (ii) a simulation using a constant humidity correction that was adopted in earlier studies (Schumann, 2012; Schumann et al., 2015; Teoh et al., 2020; Schumann et al., 2021), where the ERA5-derived RH_i fields were uniformly increased by dividing it with a factor of 0.95; (iii) a simulation that 225 uses the default aircraft-engine combination from BADA3 (EUROCONTROL, 2019), instead of the specific aircraft variant and engine model provided by a fleet database (Cirium, 2022); (iv) two simulations where a constant nvPM EI_n of 10^{15} and 10^{14} kg^{-1} is assumed for all waypoints; (v) a simulation without the effects of radiative heating interactions with the contrail plume; and (vi) a simulation that approximates the change in contrail climate forcing due to contrail-contrail overlapping (methodology detailed in the SI §S4.3).

230 3 Results and discussion

Section 3.1 presents the simulated global and regional contrail properties, including the annual statistics for 2019 (Section 3.1.1) and 2020–2021 (Section 3.1.2), seasonal effects (Section 3.1.3), and the spatiotemporal patterns and set of conditions that causes strongly warming/cooling contrail segments (Section 3.1.4). Section 3.2 evaluates the sensitivity of the 2019 contrail climate forcing to different input and model parameters, while Section 3.3 compares our results with existing studies.

235 3.1 Global contrail simulation

3.1.1 2019 global and regional contrail climate forcing

In 2019, 40.2 million flights collectively flew $60.9 \times 10^9 \text{ km}$, of which 24% of flights and 5% of the annual distance flown formed persistent contrails (Table 1). The fleet-aggregated mean nvPM EI_n and nvPM per flight distance flown are 1.02×10^{15}



kg⁻¹ and $4.7 \times 10^{12} \text{ m}^{-1}$ respectively, and around 53% of the nvPM activated to form contrail ice crystals and persisted after the
wake vortex phase ($2.5 \times 10^{12} \text{ m}^{-1}$) (Table 1). We estimate that these persistent contrail segments have a mean: (i) lifetime of
2.4 h; (ii) dimension of 9900 m (width) $\times$ 800 m (depth); (iii) ice particle volume mean radius (r_{ice}) of $\sim 10 \mu\text{m}$; and (iv) τ_{contrail}
of 0.14 respectively. Globally, contrail cirrus covers 0.06% of the sky area and the annual mean LW RF (126 mW m^{-2}) is
around a factor of two larger than the SW RF (-63.7 mW m^{-2}), yielding a net RF of 62.1 mW m^{-2} (Table 1 and Fig. 1).

Regional variabilities in the annual mean contrail cirrus net RF can be explained by differences in the: (i) annual flight distance
flown; (ii) percentage of flight distance forming persistent contrails (p_{contrail}); and (iii) $\text{EF}_{\text{contrail}}$ per unit length of contrail (Fig.
2). The US and Europe have the largest contrail climate forcing, accounting for 21% and 18% of the global annual $\text{EF}_{\text{contrail}}$
respectively, because they have the highest air traffic activity (27% and 15% of the global annual flight distance flown
respectively) (Table 2). We note that the annual mean contrail net RF over Europe (876 mW m^{-2}) is around two times larger
than the US (414 mW m^{-2}) because: (i) Europe is situated at a higher latitude which likely caused p_{contrail} (7.3%) to be larger
than the US (5.0%) (Table 2); and (ii) the contrail forcing in Europe is concentrated over a smaller domain area (c.f. Eq. (10)
and Fig. S7).

The North Atlantic has a significantly higher share of annual $\text{EF}_{\text{contrail}}$ (11%) relative to its flight distance flown (4.9%) because
flights are predominantly flown at cruising altitudes (Fig. S9), and the influence of warm conveyor belts in transporting humid
air to cruise altitudes which can lead to larger ISSR coverage area in this region (Voigt et al., 2017). Both factors likely caused
 p_{contrail} ($\sim 10\%$) in the North Atlantic to be two times higher than the global average ($\sim 5\%$). In contrast, the share of annual
 $\text{EF}_{\text{contrail}}$ over China and India (6%) is significantly lower than their flight distance flown (19%), and their contrail net RF (62--
 64 mW m^{-2}) are close to the global net RF (62.1 mW m^{-2}) (Table 2). This phenomenon is likely caused by the Hadley circulation
(where warm and moist air around the surface of the equator rises to the upper troposphere, moves poleward, becomes drier
and cooler, and sinks at the subtropics) and lower cruising altitudes (Fig. S9), both of which likely reduced the flight distance
flown in ISSRs (Lau and Kim, 2015; Reutter et al., 2020) and causes p_{contrail} in the subtropics ($\sim 2\%$) to be lower than the global
average ($\sim 5\%$) (Fig. 3).




Figure 1: The 2019 global: (a) annual mean contrail cirrus net RF; and (b) annual probability of warming contrails, where the hourly contrail net RF is greater than zero when persistent contrails are present in the grid cell. The global annual mean contrail cirrus SW and LW RF, and the annual mean ratio of contrail LW-to-SW RF are shown in Fig. S8 in the SI §S3. Basemap plotted using Cartopy 0.21.1 © Natural Earth; license: public domain.

Table 1: Global annual air traffic activity, emissions, and contrail properties from 2019 to 2021.

Annual statistics	Units	2019	2020	2021	2019 vs. 2020	2019 vs. 2021
-------------------	-------	------	------	------	---------------	---------------




Number of flights	-	40,220,293	27,910,301	35,576,165	-31%	-12%
Annual flight distance flown	10^9 km	60.94	34.50	41.90	-43%	-31%
Annual fuel burn	10^9 kg	280	145	165	-48%	-41%
Fuel burn per flight distance	kg km ⁻¹	4.596	4.200	3.926	-8.6%	-15%
Annual CO ₂ emissions	10^9 kg	885	458	520	-48%	-41%
Mean overall propulsion efficiency, η	-	0.297	0.285	0.289	-4.0%	-2.7%
Mean nvPM EI _n	10^{15} kg ⁻¹	1.021	1.016	1.021	-0.5%	0.0%
Mean nvPM per flight distance	10^{12} m ⁻¹	4.693	4.265	4.009	-9.1%	-15%
Flights forming contrails	%	42.5	34.6	34.2	-19%	-20%
Flights forming persistent contrails ^a	%	23.8	17.7	17.8	-25%	-25%
Annual contrail length	10^9 km	21.35	11.29	14.00	-47%	-34%
Flight distance forming contrails	%	35.0	32.7	33.4	-6.6%	-4.6%
Annual persistent contrail length ^a	10^9 km	3.018	1.403	1.732	-54%	-43%
Flight distance forming persistent contrails ^a	%	4.95	4.07	4.13	-18%	-17%
Initial mean ice particle number per contrail length, $n_{ice, initial}$	10^{12} m ⁻¹	2.50	2.31	2.17	-8%	-13%
Lifetime mean ice particle number per contrail length, n_{ice}	10^{12} m ⁻¹	1.88	1.75	1.64	-6.9%	-13%
Mean contrail lifetime	h	2.43	2.34	2.25	-3.6%	-7.3%
Mean ice particle volume mean radius, r_{ice}	μ m	9.96	10.4	10.7	4.4%	7.4%
Mean contrail segment optical depth, $\tau_{contrail}$	-	0.139	0.127	0.126	-8.4%	-9.2%
Mean contrail width	m	9903	9573	9081	-3.3%	-8.3%
Mean contrail depth	m	803	782	776	-2.6%	-3.4%
Contrail cirrus coverage ^b	%	0.064	0.030	0.038	-53%	-41%
Contrail cirrus coverage, clear sky ^c	%	0.66	0.28	0.33	-57%	-50%
Cloud-contrail overlap	%	90.3	89.3	88.5	-1.1%	-2.0%
Number of flights: warming contrails	-	6,741,548	3,491,439	4,430,717	-48%	-34%
Number of flights: cooling contrails	-	2,821,562	1,458,930	1,904,533	-48%	-33%
Ratio: warming-to-cooling contrails	-	2.39	2.39	2.33	0.2%	-2.6%
Mean contrail cirrus SW RF'	W m ⁻²	-4.15	-3.70	-3.90	-11%	-6.0%
Mean contrail cirrus LW RF'	W m ⁻²	5.36	4.89	5.05	-8.8%	-5.8%
Mean contrail cirrus net RF'	W m ⁻²	1.22	1.18	1.15	-3.3%	-5.7%
Annual mean contrail cirrus SW RF	mW m ⁻²	-63.7	-26.4	-33.0	-59%	-48%
Annual mean contrail cirrus LW RF	mW m ⁻²	126	53.8	64.8	-57%	-49%
Annual mean contrail cirrus net RF	mW m ⁻²	62.1	27.3	31.7	-56%	-49%
Annual mean contrail cirrus net ERF	mW m ⁻²	26.1	11.5	13.3	-56%	-49%
Annual EF _{contrail}	10^{18} J	999	440	510	-56%	-49%
EF _{contrail} per flight distance	10^8 J m ⁻¹	0.164	0.128	0.122	-22%	-26%
EF _{contrail} per contrail length	10^8 J m ⁻¹	3.31	3.14	2.94	-5%	-11%
Flights responsible for 80% EF _{contrail}	%	2.68	1.78	1.73	-34%	-35%
Contrail cirrus GWP ₂₀ ^d	-	1.17	0.99	1.02	-15%	-13%
Contrail cirrus GWP ₁₀₀ ^d	-	0.32	0.27	0.28	-15%	-13%

^a: Persistent contrails are defined when the contrail ice water content after the wake vortex phase is greater than 10^{-12} kg kg⁻¹. We note that contrails do not sublimate immediately when the RH_i is less than 100%, as the ice crystal sublimation rate depends on the background RH_i.

^b: Global contrail cirrus cover as a percentage of sky area, where contrail cirrus is assumed to be present in a grid cell if: (i) their optical depth is greater than 0.1, which is selected to be consistent with the satellite detectability threshold; and (ii) the optical depth of natural cirrus in the grid cell is less than 0.1.

^c: Global contrail cirrus coverage area under clear sky conditions without the presence of natural cirrus.

^d: The contrail cirrus GWP over a 100-year and 20-year time horizon is approximated using the energy forcing metric as described in Eq. (11) and Eq. (12).



275 **Figure 2: The 2019 global: (a) annual flight distance flown; (b) percentage of flight distance forming persistent contrails; and (c) EF_{contrail} per unit length of persistent contrail formed, where the total EF_{contrail} throughout the contrail lifetime is added to the pixel where contrails were initially formed. Basemap plotted using Cartopy 0.21.1 © Natural Earth; license: public domain.**



Table 2: Regional air traffic activity, emissions, and contrail properties for 2019. The statistics for 2020 and 2021 can be found in Tables S6 and S7 (SI §S3).

Regional statistics: 2019	Global	USA	Europe	East Asia	SEA	Latin America	Africa & Middle East	China	India	North Atlantic	North Pacific	Arctic Region
Annual distance flown ($\times 10^9$ km)	60.94	16.30	8.858	8.303	3.989	2.250	4.631	8.946	2.551	2.975	2.393	0.3826
- Percentage relative to global values ^a	-	26.8%	14.5%	13.6%	6.5%	3.7%	7.6%	14.7%	4.2%	4.9%	3.9%	0.6%
Annual dist. flown above FL250 ($\times 10^9$ km)	49.03	11.97	6.779	6.477	3.207	1.803	3.952	7.091	2.134	2.876	2.033	0.3738
- Percentage relative to global values ^a	-	24.4%	13.8%	13.2%	6.5%	3.7%	8.1%	14.5%	4.4%	5.9%	4.1%	0.8%
Air traffic density ($\text{km}^{-1} \text{h}^{-1}$)	0.014	0.116	0.152	0.059	0.029	0.006	0.009	0.047	0.032	0.030	0.012	0.002
Mean nvPM EI _a ($\times 10^{15} \text{ kg}^{-1}$)	1.021	1.361	1.155	1.082	0.983	1.035	0.883	1.083	1.036	0.621	0.649	0.406
Mean nvPM per dist. ($\times 10^{12} \text{ m}^{-1}$)	4.69	4.48	4.79	5.44	4.83	4.37	4.44	5.41	5.36	3.93	4.53	3.18
Persistent contrail length ($\times 10^9$ km)	3.018	0.807	0.646	0.137	0.133	0.073	0.128	0.175	0.045	0.308	0.136	0.025
- Percentage relative to global values ^a	-	26.7%	21.4%	4.5%	4.4%	2.4%	4.2%	5.8%	1.5%	10.2%	4.5%	0.8%
Dist. forming persistent contrails	4.95%	4.95%	7.29%	1.65%	3.34%	3.25%	2.75%	1.96%	1.75%	10.4%	5.68%	6.64%
Area-mean contrail optical depth, τ	0.018	0.065	0.099	0.027	0.020	0.014	0.013	0.026	0.019	0.051	0.023	0.026
Mean contrail age in domain (h)	2.43	1.84	1.87	2.31	2.72	3.44	2.81	2.23	2.25	2.26	2.60	3.51
Contrail cirrus coverage (%)	0.06	0.31	0.95	0.05	0.03	0.01	0.02	0.99	0.02	0.36	0.07	0.05
Contrail cirrus coverage, clear sky (%)	0.66	5.4	8.7	1.0	0.88	0.082	0.14	0.90	0.38	4.0	0.54	0.12
Annual mean SW RF (mW m^{-2})	-63.7	-485	-1160	-88.9	-83.8	-14.7	-20.0	-87.8	-35.6	-300	-55.0	-10.2
Annual mean LW RF (mW m^{-2})	126	900	2038	153	174	33.3	38.7	150	81.2	601	103	29.2
Annual mean Net RF (mW m^{-2})	62.1	414	876	63.9	90.4	18.5	18.6	62.3	45.4	300	47.7	19.0
Ratio: LW/SW RF	1.98	1.86	1.76	1.72	2.08	2.27	1.94	1.71	2.28	2.00	1.87	2.86
EF _{contrail} ($\times 10^{18} \text{ J}$)	999	209	184	32.6	44.2	23.3	35.4	42.5	13.2	109	35.7	12.9
- Percentage relative to global values ^a	-	20.9%	18.4%	3.3%	4.4%	2.3%	3.5%	4.3%	1.3%	10.9%	3.6%	1.3%
EF _{contrail, initial} ($\times 10^{18} \text{ J}$) ^b	999	218	201	32.9	44.8	23.6	37.8	41.4	13.2	116	35.0	10.2
- Percentage relative to global values ^a	-	21.8%	20.1%	3.3%	4.5%	2.4%	3.8%	4.1%	1.3%	11.6%	3.5%	1.0%
Ratio: EF _{contrail} /EF _{contrail, initial} ^c	1.00	0.96	0.92	0.99	0.99	0.99	0.94	1.03	1.00	0.94	1.02	1.26
EF _{contrail} per flight distance ($\times 10^8 \text{ J m}^{-1}$)	0.164	0.134	0.227	0.040	0.112	0.105	0.082	0.046	0.052	0.390	0.146	0.267
EF _{contrail} per contrail length ($\times 10^8 \text{ J m}^{-1}$)	3.31	2.70	3.11	2.40	3.36	3.23	2.96	2.37	2.95	3.76	2.57	4.02

^a: There are some overlapping between regional bounding boxes (Fig. S7), and therefore, the summation of regional statistics does not add up to 100%.

^b: The total EF_{contrail} throughout the contrail lifetime is added back to the location where contrails were initially formed.

^c: A higher ratio indicates that a larger share of contrail climate forcing is from contrails initially formed outside of the region but subsequently advected into the domain.




Figure 3: The percentage of annual flight distance flown that formed persistent contrails (p_{contrail}) by latitude in 2019 (blue line), 2020 (orange line), and 2021 (green line). The large interannual variability in p_{contrail} at high latitudes (above 60°N and below 60°S) can be attributed to a small sample size, where only 0.2–0.3% of the global annual flight distance were flown in these latitudes.



The EF_{contrail} per contrail length also tends to be large over the Atlantic and Indian Ocean, Sahara Desert, Central Europe, and Greenland (Fig. 2c), and these effects can be attributed to: (i) high mean albedos (Fig. S13), which reduces the contrail SW RF; (ii) high surface temperatures and outgoing longwave radiation (OLR) (Fig. S12), which drives the contrail LW RF; and/or (iii) flight scheduling, where long-haul flights tend to fly at night and at higher altitudes (Section 3.1.4). On average, contrails persisting over Greenland and the Sahara Desert have a net warming effect in $\sim 90\%$ of the hourly time periods, while contrails over the Mediterranean Sea and Southern Argentina/Chile are cooling for $\sim 43\%$ of the time (Fig. 1b).

3.1.2 Impacts from COVID-19

COVID-19 caused significant reductions in the global annual mean global annual mean contrail cirrus net RF in 2020 (27.3 mW m^{-2}) and 2021 (31.7 mW m^{-2}) when compared to 2019 (62.1 mW m^{-2}). The percentage reduction in global contrail net RF (-56% in 2020 and -49% in 2021 relative to 2019) is greater than the change in global annual flight distance flown (-43% and -31% respectively) due to an increased share of: (i) general aviation activity below 30,000 feet (Teoh et al., 2023), which likely lowered p_{contrail} from 5.0% in 2019 to 4.1% in 2020 and 2021; and (ii) short-haul flights ($< 3 \text{ h}$) from 83% in 2019 to 88% in 2020 and 2021 (Teoh et al., 2023), where increased usage of narrowbody aircraft lowers the fleet-aggregated mean fuel consumption and nvPM per flight distance flown (-9% in 2020 and -15% in 2021 relative to 2019) and EF_{contrail} per contrail length (-5% and -11% respectively).




Figure 4: Change in the absolute annual flight distance flown when comparing between (a) 2020 vs. 2019 and (b) 2021 vs. 2019, and the annual mean contrail net RF between (c) 2020 vs. 2019 and (d) 2021 vs. 2019. Basemap plotted using Cartopy 0.21.1 © Natural Earth; license: public domain.



A regional comparison between 2019 and 2020 shows that the North Atlantic experienced the largest percentage reduction in annual flight distance flown (-61%) and contrail net RF (-65%) because it has a large share of long-haul international flights which were most impacted by COVID-19. In East Asia and China, the reduction in contrail net RF (50–54%) is more than two times greater than their reduction in flight distance flown (-24%). The higher relative reduction in contrail net RF is most likely due to: (i) a higher share of domestic air traffic in parts of China (Fig. 4a) that led to an 8% increase in flight distance flown at 25,000–30,000 feet where persistent contrails are less likely to form; and (ii) the reduction in international overflights caused a 39% reduction in flight distance flown above 30,000 feet (Fig. S10c). In contrast, the 11% reduction in contrail net RF over the Arctic is significantly smaller than the 58% reduction in flight distance flown because a higher share of contrails in this region were formed elsewhere and subsequently advected into the domain (Table 2 vs. Table S6).

The 2021 annual mean contrail net RF in ten of the eleven regions are 4–52% larger than their 2020 levels (Table S6 vs. Table S7). There is a 44% year-on-year reduction in the contrail net RF over the Arctic which is most likely caused by the interannual variability in meteorology (Teoh et al., 2022a), where the 2020 p_{contrail} (12%) and $\text{EF}_{\text{contrail}}$ per flight distance ($0.6 \times 10^8 \text{ J m}^{-1}$) were outliers that were around two times larger than those recorded in 2019 (6.6% and $0.3 \times 10^8 \text{ J m}^{-1}$ respectively) and 2021 (7.3% and $0.3 \times 10^8 \text{ J m}^{-1}$) (Fig. 3, Tables 2, S6 and S7). Notably, the 2021 annual mean contrail net RF over the Gulf of Mexico and Caribbean Sea (70–100°W, 5–35°N) is 3.8% larger than their 2019 levels (Fig. 4d), and this is most likely caused by the increase in air traffic over Southern US–Mexico (Fig. 4b) where contrails were formed and advected southwards over its lifetime.

3.1.3 Seasonal statistics

The seasonal statistics reported here are biased towards the Northern Hemisphere where 92% of the global annual flight distance was flown (Teoh et al., 2023). In 2019, global air traffic activity peaked in the summer (JJA) and was 14% below peak levels during the winter months (DJF) (Fig. 5a). The seasonal changes in contrail properties shows the opposite effect where p_{contrail} in wintertime is around 1.5 times larger than the summer (5.9% vs. 3.8%, Fig. 5b), and persistent contrails formed during this time have a larger mean lifetime (2.8 vs. 2.4 h in the summer, Fig. 5e), global coverage area (0.07 vs. 0.05% in 2019, Fig. 5f), net RF (86 vs. 40 mW m^{-2} , Fig. 5g) and $\text{EF}_{\text{contrail}}$ per contrail length (10 vs. $7.4 \times 10^7 \text{ J m}^{-1}$, Fig. 5h). The larger contrail occurrence, lifetime, and climate forcing in wintertime can be attributed to: (i) larger seasonal ISSR coverage area in the northern mid-latitudes (30–60°N) (Teoh et al., 2022a), a region accounting for 64% of global annual flight distance flown; (ii) smaller initial RH_i (107 vs. 110% in summertime, Fig. S15d) and dT_{SAC} (-7 vs. -4 K, Fig. S15e) both of which lowers specific humidity and the amount of condensable water available, c.f. Eq. (S2), leading to smaller r_{ice} (11 vs. 13 μm , Fig. 5d) and longer lifetimes (2.8 vs. 2.4 h); (iii) higher percentage of cloud-contrail overlapping (91 vs. 88%, Fig. S15f); and (iv) shorter daylight hours. Factors (iii) and (iv) are expected to lower the contrail SW RF (Teoh et al., 2022a). In contrast, the shorter contrail lifetime during the summer is likely due to persistent contrails forming at warmer temperatures with a smaller dT_{SAC} , thereby reducing $p_{\text{activation}}$ and $n_{\text{ice,initial}}$, c.f. Eq. (5) and (6) and Fig. 5c, which result in larger r_{ice} (Fig. 5d) and ice crystal sedimentation rate. The larger mean overlying natural cirrus optical depth above contrails (τ_{cirrus}) in summertime (0.39 vs.

0.25 in winter, Fig. S15h) also contributes to a smaller contrail climate forcing relative to the winter months (Schumann et al., 2012; Teoh et al., 2022a).




Figure 5: Monthly statistics on the global: (a) flight distance flown; (b) percentage of flight distance forming persistent contrails (p_{contrail}); (c) fraction of nvPM that activates into contrail ice crystals and survive the wake vortex phase; (d) mean contrail ice particle volume mean radius (r_{ice}); (e) mean contrail lifetime; (f) global contrail cirrus coverage as a percentage of sky area; (g) mean net RF; and (h) EF_{contrail} per unit length of persistent contrail formed from January 2019 to December 2021. Additional metrics that are not presented here are available in Fig. S15 in the SI §S3).

Due to the higher relative contrail climate forcing in wintertime, around one-third of the days in 2019 and 2021 (110–121 days) accounted for half of the global annual EF_{contrail} , while two-thirds of the days (236–243 days) caused 80% of the annual EF_{contrail} (Fig. 6a). In 2020, the global contrail climate forcing was further concentrated on a smaller number of days, where 23% (83 days) and 53% (195 days) of the days accounted for 50% and 80% of the annual EF_{contrail} respectively (Fig. 6a),



because the percentage of annual flight distance flown in the first quarter of 2020 (37%), where contrails are expected to be strongly warming, was higher than those recorded in 2019 (23%) and 2021 (19%), and because of the significant reduction in global air traffic activity during the spring and summer (Fig. 5a) where the EF_{contrail} per contrail length is at a minimum (Fig. 5h).




Figure 6: Cumulative density function of the global annual EF_{contrail} from 2019 to 2021 versus the: (a) number of days in a year; and (b) the percentage of all flights that accounted for the proportion of EF_{contrail} . We note that the cumulative density function in (b) exceeds and returns to 1 because of the presence of flights with cooling contrails.

3.1.4 Strongly warming or cooling contrails

In 2019, 24% of all flights formed persistent contrails, of which 70% of these persistent contrail-forming flights have a net warming effect ($EF_{\text{contrail}} > 0$) and their mean magnitude of EF_{contrail} (1.45×10^{14} J) is a factor of 5 larger than the remaining 30% of contrail-forming flights with a net cooling effect (-0.28×10^{14} J). Around 2.7% of all flights (or 11% of contrail-forming flights) accounted for 80% of the global annual EF_{contrail} in 2019 (Fig. 6b). The contrail climate forcing was concentrated on a smaller subset of flights in 2020 and 2021, where 1.7% of all flights or 10% of contrail-forming flights accounted for 80% of the annual EF_{contrail} (Fig 6b), and this is likely due to the larger share in general aviation activity and short-haul flights with flight times below 3 h (Teoh et al., 2023), both of which generally fly at lower altitudes where persistent contrails are less likely to form.

Individual flight segments with the most strongly warming contrails (EF_{contrail} per contrail length $> 15.4 \times 10^8$ J m^{-1} , 95th percentile) are commonly found over the US and North Atlantic (Fig. 7a), and these contrail segments are more likely to be formed by eastbound transatlantic flights and transcontinental flights across the US (Table S9) because they tend to depart during the evenings (Teoh et al., 2022a). In contrast, the most strongly cooling contrails (EF_{contrail} per contrail length $< -2.39 \times 10^8$ J m^{-1} , 5th percentile) are more prevalent over Southeast Asia, North Asia, Europe, and the eastern North Atlantic region (Fig. 7b), and these contrail segments are more commonly formed by short-/medium-haul flights around Southeast and East



Asia, long-haul flights between Oceania, Asia and Europe, and both east- and westbound transatlantic flights (Table S10). The ratio of solar direct radiation (SDR) to OLR also contributes to the spatial distribution of strongly warming and cooling contrail segments: strongly cooling contrails are more prevalent in Southeast Asia because the region has a high SDR-to-OLR ratio which drives the contrail SW RF and limits its LW RF; while the lower SDR-to-OLR ratio contributes to a higher share of strongly warming contrails at latitudes above 30°N (Fig. S14). Both strongly warming and cooling contrails are generally formed above 38,000 feet over the tropics and across a wider altitude range (30,000–40,000 feet) at higher latitudes (Fig. 7c and 7d), and this can likely be attributed to the latitude and seasonal variations in the tropopause height (Santer et al., 2003; Hoinka et al., 1993).




Figure 7: The location (longitude, latitude and altitude) of individual contrail segments that are strongly warming (EF_{contrail} per contrail length $> 15.4 \times 10^8 \text{ J m}^{-1}$, 95th percentile) and strongly cooling (EF_{contrail} per contrail length $< -2.39 \times 10^8 \text{ J m}^{-1}$, 5th percentile), shown in subplots (a) to (d); and probability density functions showing their respective day of year; (f) time of day; (g) nvPM emissions; (h) the dT_{SAC} when these contrails were initially formed; and their mean (i) effective albedo; and (j) outgoing longwave radiation (OLR); and (k) τ_{cirrus} throughout their lifetime. Basemap plotted using Cartopy 0.21.1 © Natural Earth; license: public domain.



The most strongly warming contrail segments tend to occur during the winter and spring and between 16:00 and 03:00 local time, while strongly cooling contrails are more prevalent in spring and between 03:00 and 12:00 (Fig. 7e and 7f). Both strongly warming and cooling contrails are formed by aircraft-engine types with mean nvPM emissions (6.9 and $5.3 \times 10^{12} \text{ m}^{-1}$ respectively) that are larger than the 2019 fleet-aggregated values ($4.7 \times 10^{12} \text{ m}^{-1}$, Table 1), and this is consistent with an earlier study that found a positive correlation between nvPM number emissions and the absolute magnitude and variability of $\text{EF}_{\text{contrail}}$ (Teoh et al., 2022a). Strongly cooling contrails are more likely to have shorter lifetimes (mean of 5.6 h) relative to warming contrails (6.7 h) because of their smaller nvPM emissions and dT_{SAC} (Fig. 7g and 7h), and their formation time (03:00–12:00 local time) and lifetime also suggest that these contrail segments spread during daylight hours and sublimate before dusk, thus maximising their SW RF.

The surface conditions and background cloud fields also influence the contrail climate forcing (Schumann et al., 2012; Teoh et al., 2022a). Strongly warming contrail segments generally have a: (i) larger effective albedo relative to cooling contrails (0.39 vs. 0.31); (ii) larger OLR (211 vs. 196 W m^{-2}); and (iii) smaller mean τ_{cirrus} (0.25 vs. 0.57) (Fig. 7i to 7k), and these are indicative of optically thick low-level water clouds, snowy and sandy terrains, and/or warmer surface temperatures. In contrast, contrails are more likely to be strongly cooling when formed over the dark ocean surface and/or below optically-thick high-level cirrus clouds that tends to reduce the contrail LW RF' more strongly than the SW RF' (Teoh et al., 2022a).

3.2 Sensitivity analysis

We re-run the 2019 global contrail simulation to assess the sensitivity of the simulated contrail climate forcing to humidity corrections applied to the ERA5 HRES (Section 3.2.1), assumptions in aircraft-engine assignments and emissions (Section 3.2.2), and contrail model parameters (Section 3.2.3). Fig. 8 and Table S11 summarises the change in global aviation emissions and contrail properties from the different simulation runs.

3.2.1 Humidity corrections

Globally, the baseline simulation with the extended global humidity correction, c.f. Eq. (1) to (4), produces an annual mean contrail net RF (62.1 mW m^{-2}) that is 78% larger than the simulation without humidity correction applied to the ERA5 HRES (34.8 mW m^{-2}). The change in contrail net RF between the two simulations is largest at latitudes above 35°N (+96%, 76.4 mW m^{-2} vs. 38.9 mW m^{-2} without humidity correction), followed by the tropics (+59%, 52.3 vs. 32.9 mW m^{-2}), and is smallest in the subtropics at around $30^\circ\text{N/S} \pm 5^\circ$ (+2.2%, 84.2 vs. 82.4 mW m^{-2}) (Fig. 8a).

Alternatively, the use of a constant humidity correction that was adopted in earlier studies (Schumann, 2012; Schumann et al., 2015; Teoh et al., 2020; Schumann et al., 2021), c.f. Eq. (S5) where $\text{RH}_{\text{ic}} = 0.95$, causes the global annual mean contrail net RF to be 4% larger than the baseline simulation (64.5 vs. 62.1 mW m^{-2}). However, the constant humidity correction approach does not capture the latitude-dependent errors in the ERA5-derived ISSR (Table S1), and therefore, could overestimate the annual mean contrail net RF in the tropics and subtropics and underestimate it at high latitudes (Fig. 8b).




Figure 8: Absolute change in the 2019 global annual mean contrail cirrus net RF (in mW m^{-2}) when comparing the baseline scenario with the simulation: (a) without humidity corrections applied to the ERA5 HRES; (b) with a constant humidity correction of $\text{RH}_i/\text{RH}_{ic}$, where $\text{RH}_{ic} = 0.95$; (c) with the default aircraft-engine assignment from BADA3; (d) with a constant nvPM EI_n of 10^{15} kg^{-1} for all waypoints; (e) without radiative heating effects; and (f) that approximates the radiative effects of contrail-contrail overlapping. Basemap plotted using Cartopy 0.21.1 © Natural Earth; license: public domain.

3.2.2 Aircraft-engine assignment and emissions

The simulation with default aircraft-engine assignments from BADA3 causes the global annual mean contrail net RF to be 18% larger than the baseline simulation (73.1 vs. 62.1 mW m^{-2}). This is because BADA3 assumes that some widely used aircraft types (i.e., Airbus A320, A320neo, and Boeing 787 families) are powered by engines with nvPM EI_n that are up to four orders of magnitude larger than their alternative engine options (Teoh et al., 2023), which leads to a larger 2019 global mean nvPM EI_n (1.39 vs. $1.02 \times 10^{15} \text{ kg}^{-1}$ in the baseline simulation, +36%) and contrail net RF (Fig. 8c).



The simulation with a constant nvPM EI_n of 10^{15} kg^{-1} for all waypoints leads to a global contrail net RF of 74.8 mW m^{-2} (+20% relative to the baseline). Regionally, this approach could underestimate the contrail net RF over Europe and the US (Fig. 8d) likely because a higher proportion of flights are short-haul and utilises the Airbus A320 family, where the nvPM EI_n from one of the engine options ($3\text{--}7 \times 10^{15} \text{ kg}^{-1}$) is significantly larger than the assumed 10^{15} kg^{-1} (Teoh et al., 2023; EASA, 2021). A change in the assumed nvPM EI_n from 10^{15} to 10^{14} kg^{-1} leads to a smaller mean τ_{contrail} (-50%), lifetime (-35%), and coverage area (-70%), which in turn, produces a global contrail net RF of 13.7 mW m^{-2} (-82%) (Table S11). The simulation with an EI_n of 10^{14} kg^{-1} demonstrates the potential of fleet-wide reductions in engine particle emissions by an order of magnitude, but it is not included in our range of contrail RF estimates for 2019.

3.2.3 Contrail model parameters

The global contrail net RF from the baseline simulation (with radiative heating effects) is 8% smaller than the simulation without radiative heating effects (62.1 vs. 66.8 mW m^{-2}). This is because solar and terrestrial radiation heats up the contrail plume and reduces the mean contrail lifetime by 23% (2.4 vs. 3.0 h without radiative heating), which in turn, lowers the contrail net RF over: (i) regions with a higher fraction of aged contrails (Fig. S19); and (ii) Europe as less contrails are advected into the region via the North Atlantic jet stream (Fig. 8e). However, radiative heating also increases the vertical mixing rate and τ_{contrail} and causes the contrail net RF to be larger along established flight corridors (Fig. 8e). Seasonally, differences in the mean contrail net RF are largest in spring and summer (47.7 vs. 53.1 mW m^{-2} without radiative heating, -11%) and smallest in wintertime (84.0 vs. 87.3 mW m^{-2} , -3.9%) (Fig. S17) because the impacts of radiative heating on the contrail lifetime depends on the magnitude of SDR and OLR.

The effects of contrail-contrail overlapping reduces the global annual mean contrail net RF by 5% (from 62.1 mW m^{-2} in the baseline to 59.1 mW m^{-2}), which is consistent with an earlier study that estimated a 3% reduction in the global contrail net RF due to contrail-contrail overlapping (Sanz-Morère et al., 2021). Regionally, the change in contrail climate forcing depends on the magnitude of the annual mean contrail net RF, OLR, and the ratio of SW-to-LW RF (Fig. S20). The largest reduction in contrail climate forcing occurs in regions with dense air traffic such as the US (-9.7%) and Europe (-9.4%); while the contrail net RF is increased in areas with a higher prevalence of strongly cooling contrails, such as the eastern North Atlantic and North Asia (Fig. 7b), because the change in SW RF is larger than the LW RF (Fig. 8f and Table S12).

3.3 Comparison with other studies

When taken together, the sensitivity analysis gives a 2019 global contrail cirrus net RF range of 62.1 [34.8 , 74.8] mW m^{-2} . Lee et al. (2021) used the 2006 global annual mean contrail cirrus net RF estimates from three different contrail models (SI §S5) and extrapolated the contrail cirrus net RF to 2018 levels (111 [33 , 189] mW m^{-2}) assuming that the growth in contrail climate forcing is proportional to the growth in global annual flight distance flown. The nominal 2019 global contrail cirrus net RF from this study (62.1 mW m^{-2}) is 44% lower than the 2018 central estimate from Lee et al. (2021) (111 mW m^{-2}), and we expect that this discrepancy is due to a higher air traffic growth rate in the subtropics (+12% per annum in China and India vs.



+6% globally) between 2006 and 2018 (World Bank, 2023), where persistent contrails are less likely to form (Fig. 3). Differences in the contrail modelling approaches (i.e., general circulation models vs. Lagrangian models), model inputs, and parameter settings are also likely to contribute to discrepancies in the contrail net RF estimates between studies. The EF_{contrail} and annual CO_2 emissions derived from this study are also used to estimate the 2019–2021 annual mean contrail cirrus GWP_{20} and GWP_{100} , c.f. Eq. (11) and (12) and Table 1. Our estimates (1.06 for GWP_{20} and 0.29 for GWP_{100}) are 54% smaller than Lee et al. (2021) (2.32 for GWP_{20} and 0.63 for GWP_{100}) owing to the lower contrail cirrus RF for the reasons discussed above. We note that the 2019–2021 annual CO_2 emissions (c.f. Table 1) used to calculate the contrail GWPs in this study are 0.8–0.9% lower than those provided by GAIA (Teoh et al., 2023) because the flight waypoints that were flown above 13 km were not included in the contrail simulation (Section 2.3), and our estimate of the 2019 annual CO_2 emissions (885 Tg) is ~14% lower than that of Lee et al. (2021) (1034 Tg in 2018) due to differences between top-down and bottom-up estimates.




Figure 9: Comparison of the simulated contrail properties from CoCiP with in-situ, remote sensing and satellite observations from the contrail library database (COLI) (Schumann et al., 2017) versus the contrail segment age. The contrail properties compared includes the contrail: (a) ice particle number concentration in the plume ($n_{\text{ice,conc}}$); (b) r_{ice} ; (c) depth; (d) τ_{contrail} ; (e) width; and (f) total extinction, i.e., the integral of the optical extinction over the contrail cross-sectional area, which influences the contrail RF'. The red data points are from in-situ measurements, blue data points are from remote sensing, and the black lines in (d) represent the 0th, 10th, 50th, 90th and 100th percentiles from the Meteosat Second Generation (MSG) satellites and the automatic contrail



tracking algorithm (ACTA) (Vázquez-Navarro et al., 2015). The grey data points are the simulated CoCiP contrail properties from a subset of results within 6-9 UTC in 1-January-2021 and 15-July-2021.

485 Recently, Bier & Burkhardt (2022) improved parameterisations of the initial contrail ice nucleation and particle losses in the ECHAM general circulation model and lowered their 2006 global contrail net RF from 56 mW m^{-2} (Bock and Burkhardt, 2016a) to 43.7 mW m^{-2} . Our 2019 global annual mean contrail net RF (62.1 mW m^{-2}) is 42% larger than their 2006 estimates (43.7 mW m^{-2}). The derived average annual growth rate of the global contrail net RF for 2006–2019 (+2.7% per annum) is lower than the growth in flight distance flown during the same period (+3.6% per annum), and could be explained by: (i) the
 490 higher share of air traffic growth in the subtropics where p_{contrail} is smaller than the global average (Fig. 3); and (ii) improvements in aircraft engine technology which reduced the fuel consumption per distance travelled by ~6% (from 4.87 kg km^{-1} in 2006 to 4.60 kg km^{-1} in 2019) and, in turn, is expected to lower the nvPM (c.f. Eq. (5)) and $\text{EF}_{\text{contrail}}$ per flight distance flown (Teoh et al., 2022a).

Regionally, Teoh et al. (2022a) used CoCiP without radiative heating effects and estimated a 2019 annual mean contrail net
 495 RF over the North Atlantic (235 mW m^{-2}) that is 22% smaller than this study's estimates (300 mW m^{-2}). The contrail net RF from Teoh et al. (2022a) are likely underestimated because: (i) their air traffic dataset only recorded flights that enter the Shanwick Flight Information Region ($10\text{--}40^\circ\text{W}$), thereby capturing ~80% of the North Atlantic oceanic traffic (Molloy et al., 2022); and (ii) contrails that were formed outside the North Atlantic and subsequently advected into the domain were not accounted for. Meijer et al. (2022) used geostationary satellite imagery and a machine learning algorithm to estimate the 2018–
 500 19 annual mean contrail cirrus cover over the United States (0.15%), which is around 50% smaller than our 2019 estimates (0.31%, Table 2). These differences may be due to the reduced probability for satellites detecting: (i) freshly formed contrails with sub-pixel width; (ii) aged contrails that have lost their line-shaped structure; (iii) $\tau_{\text{contrail}} < 0.1$; and (iv) contrails that overlap with natural cirrus (Mannstein et al., 2010; Vázquez-Navarro et al., 2015). Fig. 9 also shows that our simulated contrail properties from CoCiP are generally in good agreement with in-situ, remote sensing and satellite observations from the contrail
 505 library database (COLI) (Schumann et al., 2017).

4 Conclusions

We simulated the global contrail climate forcing for 2019–2021 using historical flight trajectories derived from ADS-B telemetry, improved nvPM EI_n that accounts for specific aircraft-engine types, the ERA5 HRES reanalysis with global corrections applied to the humidity fields, and CoCiP. Table 3 summarises the global annual mean contrail net RF from the
 510 different simulation runs performed in this study.

Our 2019 global annual mean contrail cirrus net RF estimate (62.1 mW m^{-2}) is: (i) 44% lower than the central estimate from Lee et al. (2021), where the 2006 global contrail net RF from three studies were extrapolated to 2018 levels ($111 [33, 189] \text{ mW m}^{-2}$) assuming the global growth in flight distance flown and contrail net RF are proportional; and (ii) 43% higher than the 2006 estimate from Bier & Burkhardt (2022) (43.7 mW m^{-2}), where the derived 2006–2019 average annual growth rate of



the global contrail net RF (+2.7% per annum) is lower than the growth in flight distance flown (+3.6% per annum). These discrepancies are most likely caused by the higher relative air traffic growth rate in the subtropics where persistent contrails are less likely to form because aircraft fly at lower altitudes and the Hadley Circulation limits the ISSR coverage area. Regionally, we estimate that Europe (876 mW m^{-2}), the US (414 mW m^{-2}) and North Atlantic (300 mW m^{-2}) have the largest contrail net RF in 2019, while the forcing in East Asia (63.9 mW m^{-2}) and China (62.3 mW m^{-2}) are close to the global mean value. Policy response to COVID-19 impacted global aviation operations and fleet composition, lowering the 2020 and 2021 annual mean contrail net RF to 27.3 and 31.7 mW m^{-2} respectively. Globally, the 2019–2021 annual mean contrail cirrus GWP₁₀₀ and GWP₂₀ are estimated to be 0.29 and 1.06 respectively (Table 1).

Around 20% of all flights formed persistent contrails in 2019–2021 (i.e. contrails that survive the wake vortex phase), of which: (i) 70% of the contrail-forming flights have a warming effect ($\text{EF}_{\text{contrail}} > 0$); and (ii) 10% of the contrail-forming flights (or 2% of all flights) were responsible for 80% the global annual $\text{EF}_{\text{contrail}}$ (Fig. 6). The most strongly warming and cooling contrail segments are generally formed: (i) at higher latitudes and in specific regions (i.e., North Atlantic and Southeast Asia), depending on the spatiotemporal variations in meteorology, radiation, and tropopause height; (ii) in specific origin–destination routes that is related to flight scheduling factors; and (iii) by aircraft-engine types with high nvPM number emissions that are larger than the mean fleet-aggregated values (Fig. 7, Tables S9 and S10). These results indicate the potential to significantly reduce the contrail climate forcing by targeting the mitigating actions to a small subset of flights and regions associated with strongly warming contrails.

Table 3: Summary of the global contrail simulations performed in this study.

	Engine assignment	Humidity correction	Radiative heating effects	Contrail overlapping effects	Global annual mean contrail net RF (mW m^{-2})
Baseline simulations (2019 – 2021)					
2019	Cirium	Eq. (1) – (4)	✓	×	62.1
2020	Cirium	Eq. (1) – (4)	✓	×	27.3
2021	Cirium	Eq. (1) – (4)	✓	×	31.7
Sensitivity: Humidity correction (2019)					
No humidity correction	Cirium	N/A	✓	×	34.8
Constant humidity correction	Cirium	Eq. (S5), $\text{RH}_{\text{ic}} = 0.95$	✓	×	64.5
Sensitivity: Aircraft performance and emissions (2019)					
Default aircraft-engine assignment	BADA default	Eq. (1) – (4)	✓	×	73.1
Constant nvPM EI_n (10^{15} kg^{-1})	N/A	Eq. (1) – (4)	✓	×	74.8
Constant nvPM EI_n (10^{14} kg^{-1})	N/A	Eq. (1) – (4)	✓	×	13.7
Sensitivity: Contrail model parameters (2019)					
No radiative heating effects	Cirium	Eq. (1) – (4)	×	×	66.8
Contrail-contrail overlapping	Cirium	Eq. (1) – (4)	×	✓	59.1

The 2019 global annual mean contrail net RF is most sensitive to the humidity corrections applied to the ERA5 HRES (34.8 mW m^{-2} without humidity correction, -44% relative to the baseline simulation), followed by assumptions on the aircraft-engine assignment and nvPM emissions ($73.1\text{--}74.8 \text{ mW m}^{-2}$, +18–20%), and is least sensitive to the radiative heating (66.8 mW m^{-2} ,



+7.6%) and contrail-contrail overlapping effects (59.1 mW m^{-2} , -4.8%) (Table 3). When taken together, our sensitivity analysis yields a 2019 global contrail net RF range of $62.1 [34.8, 74.8] \text{ mW m}^{-2}$.

Future work should be prioritised towards: (i) performing inter-model comparison studies to understand differences in the estimated contrail climate forcing from CoCiP versus those provided by other contrail models (Chen and Gettelman, 2013; Bier and Burkhardt, 2022; Fritz et al., 2020); (ii) evaluating the overall uncertainty in the simulated contrail climate forcing by propagating uncertainties in various input parameters, including meteorology, aircraft–engine performance and particle number emissions, ice particle habits and radiation transfer scheme; (iii) systematically comparing simulated contrail properties with in-situ measurements and observations from ground-based cameras, lidar and satellites; (iv) evaluating the ERF/RF ratio at regional scale and the effects of contrail cirrus climate forcing on surface temperature change; and (v) advancing mitigation of contrail climate forcing via forecasting and flight trajectory optimisation, alternative fuels, and cleaner-burning engines.

Author contributions

RT, US, CV and MEJS conceptualised the study, developed the methodology and undertook the investigation. RT, ZE, US, SR and MS were responsible for software development and data curation. RT, MEJS and US created or sourced the figures. RT wrote the original manuscript. RT, ZE, US, CV, MS and MEJS reviewed and edited the manuscript. MEJS, MS and CV acquired funding. All authors have read, edited, and reviewed the manuscript, and agreed upon the published version of the paper.

Acknowledgements

We thank Google for providing cloud credits on the Google Cloud Platform, which were utilised to simulate contrails globally to produce the results included in this paper. RT and MEJS are supported by Innovate UK (10049507). CV has been supported by DFG under contracts no. VO 1504/7-1 and VO 1504/9-1 and by the EU Horizon 2020 under the Advancing the Science for Aviation and Climate (ACACIA) project, grant no. 875036. We thank Andreas Petzold for reviewing an earlier version of the paper.

Data availability

The Global Aviation emissions Inventory based on ADS-B (GAIA) (Teoh et al., 2023) is provided as gridded outputs on Zenodo (<https://doi.org/10.5281/zenodo.7969631>). Flight trajectory and aircraft fuel consumption data are commercially sensitive, and the flight-waypoint and flight-summary outputs can be made available for scientific research upon reasonable request. The pycontrails repository that contains the CoCiP algorithm has recently been published and publicly available at



565 <https://doi.org/10.5281/zenodo.7776686>. IAGOS data were created with support from the European Commission, national agencies in Germany (BMBF), France (MESR), and the UK (NERC), and the IAGOS member institutions (<http://www.iagos.org/partners>). The participating airlines (Deutsche Lufthansa, Air France, Australian, China Airlines, Iberia, Cathay Pacific, Air Namibia, Sabena) have supported IAGOS by carrying the measurement equipment free of charge since 1994. The data are available at <https://doi.org/10.25326/20> and <https://doi.org/10.25326/06>. This document used elements of
570 Base of Aircraft Data (BADA) Family 4 Release 4.2 which has been made available by EUROCONTROL to Imperial College London. EUROCONTROL has all relevant rights to BADA. ©2019 The European Organisation for the Safety of Air Navigation (EUROCONTROL). EUROCONTROL shall not be liable for any direct, indirect, incidental, or consequential damages arising out of or in connection with this document, including the use of BADA. This document contains Copernicus Climate Change Service information 2023. Neither the European Commission nor ECMWF is responsible for any use of the
575 Copernicus information.

Competing interests

There are no conflicts of interest and all funding sources have been acknowledged. All figures are our own. None of the authors has any competing interests.

References

- 580 Agarwal, A., Meijer, V. R., Eastham, S. D., Speth, R. L., and Barrett, S. R. H.: Reanalysis-driven simulations may overestimate persistent contrail formation by 100-250%, *Environ. Res. Lett.*, 17, 1–14, <https://doi.org/10.1088/1748-9326/AC38D9>, 2022.
- Bier, A. and Burkhardt, U.: Impact of Parametrizing Microphysical Processes in the Jet and Vortex Phase on Contrail Cirrus Properties and Radiative Forcing, *J. Geophys. Res. Atmos.*, 127, e2022JD036677, <https://doi.org/10.1029/2022JD036677>, 2022.
- Bock, L. and Burkhardt, U.: Reassessing properties and radiative forcing of contrail cirrus using a climate model, *J. Geophys. Res. Atmos.*,
585 121, 9717–9736, <https://doi.org/10.1002/2016JD025112>, 2016a.
- Bock, L. and Burkhardt, U.: The temporal evolution of a long-lived contrail cirrus cluster: Simulations with a global climate model, *J. Geophys. Res. Atmos.*, 121, 3548–3565, 2016b.
- Boies, A. M., Stettler, M. E. J., Swanson, J. J., Johnson, T. J., Olfert, J. S., Johnson, M., Eggersdorfer, M. L., Rindlisbacher, T., Wang, J., and Thomson, K.: Particle emission characteristics of a gas turbine with a double annular combustor, *Aerosol Sci. Technol.*, 49, 842–855,
590 <https://doi.org/10.1080/02786826.2015.1078452>, 2015.
- Boulanger, D., Bundke, U., Gallagher, M., Gerbig, C., Hermann, M., Nédélec, P., Rohs, S., Sauvage, B., Ziereis, H., Thouret, V., and Petzold, A.: IAGOS Time series [Data set], <https://doi.org/10.25326/06>, 2022.
- Bräuer, T., Voigt, C., Sauer, D., Kaufmann, S., Hahn, V., Scheibe, M., Schlager, H., Diskin, G. S., Nowak, J. B., DiGangi, J. P., Huber, F., Moore, R. H., and Anderson, B. E.: Airborne Measurements of Contrail Ice Properties—Dependence on Temperature and Humidity,



- 595 Geophys. Res. Lett., 48, e2020GL092166, <https://doi.org/10.1029/2020GL092166>, 2021a.
- Bräuer, T., Voigt, C., Sauer, D., Kaufmann, S., Hahn, V., Scheibe, M., Schlager, H., Huber, F., Le Clercq, P., Moore, R., and Anderson, B.: Reduced ice number concentrations in contrails from low aromatic biofuel blends, *Atmos. Chem. Phys.*, 21, 16817–16826, <https://doi.org/https://doi.org/10.5194/acp-2021-582>, 2021b.
- Burkhardt, U. and Kärcher, B.: Global radiative forcing from contrail cirrus, *Nat. Clim. Chang.*, 1, 54–58, <https://doi.org/10.1038/nclimate1068>, 2011.
- 600 Chen, C. C. and Gettelman, A.: Simulated radiative forcing from contrails and contrail cirrus, *Atmos. Chem. Phys.*, 13, 12525–12536, <https://doi.org/10.5194/acp-13-12525-2013>, 2013.
- Cirium: Aircraft fleets and values: the world's largest independent aircraft database, 2022.
- Durdina, L., Brem, B. T., Setyan, A., Siegerist, F., Rindlisbacher, T., and Wang, J.: Assessment of Particle Pollution from Jetliners: from Smoke Visibility to Nanoparticle Counting, *Environ. Sci. Technol.*, 51, 3534–3541, <https://doi.org/10.1021/acs.est.6b05801>, 2017.
- 605 EASA: ICAO Aircraft Engine Emissions Databank (07/2021): <https://www.easa.europa.eu/domains/environment/icao-aircraft-engine-emissions-databank>, 2021. Last access: 17 August 2021.
- ECMWF: The Copernicus Programme: Climate Data Store, <https://cds.climate.copernicus.eu/#!/home>, 2021. Last access: 15 February 2022.
- EUROCONTROL: User Manual for the Base of Aircraft Data (BADA) Family 4, 2016.
- 610 EUROCONTROL: User Manual for the Base of Aircraft Data (BADA) Revision 3.15, EUROCONTROL Experimental Centre (EEC), 2019.
- EUROCONTROL: Integrating space-based ADS-B into EUROCONTROL's network operations system brings major predictability gains and will unlock future capacity: <https://www.eurocontrol.int/press-release/integrating-space-based-ads-b-eurocontrols-network-operations-system>, 2021. Last access: 26 August 2022.
- Eyers, C. J., Addleton, D., Atkinson, K., Broomhead, M. J., Christou, R., Elliff, T., Falk, R., Gee, I., Lee, D. S., Marizy, C., Michot, S., Middel, J., Newton, P., Norman, P., Plohr, M., Raper, D., and Stanciou, N.: AERO2K global aviation emissions inventories for 2002 and 2025, *QinetiQ Eur. Comm. under Contract No. G4RD-CT-2000-00382*, Farnborough, Hampshire, GU14 0LX, pp., 2005., 2005.
- 615 Fritz, T. M., Eastham, S. D., Speth, R. L., and Barrett, S. R. H.: The role of plume-scale processes in long-term impacts of aircraft emissions, *Atmos. Chem. Phys.*, 20, 5697–5727, <https://doi.org/10.5194/ACP-20-5697-2020>, 2020.
- Gaillot, T., Beauchet, S., Lorne, D., and Krim, L.: The impact of fossil jet fuel emissions at altitude on climate change: A life cycle assessment study of a long-haul flight at different time horizons, *Atmos. Environ.*, 311, 119983, <https://doi.org/10.1016/J.ATMOSENV.2023.119983>, 2023.
- 620 Gierens, K., Matthes, S., and Rohs, S.: How Well Can Persistent Contrails Be Predicted?, 7, 169, <https://doi.org/10.3390/AEROSPACE7120169>, 2020.
- Haywood, J. M., Allan, R. P., Bornemann, J., Forster, P. M., Francis, P. N., Milton, S., Rädcl, G., Rap, A., Shine, K. P., and Thorpe, R.: A case study of the radiative forcing of persistent contrails evolving into contrail-induced cirrus, *J. Geophys. Res. Atmos.*, 114, <https://doi.org/10.1029/2009JD012650>, 2009.



- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Hoare, G.: Country bounding boxes, <https://gist.github.com/graydon/11198540>, 2014. Last access: 22 December 2022.
- Hoinka, K. P., Reinhardt, M. E., and Metz, W.: North Atlantic air traffic within the lower stratosphere: Cruising times and corresponding emissions, *J. Geophys. Res. Atmos.*, 98, 23113–23131, <https://doi.org/10.1029/93JD02262>, 1993.
- 635 ICAO: The World of Air Transport in 2013: <https://www.icao.int/annual-report-2013/pages/the-world-of-air-transport-in-2013.aspx>, 2014. Last access: 22 February 2022.
- ICAO: The World of Air Transport in 2015, <https://www.icao.int/annual-report-2015/pages/the-world-of-air-transport-in-2015.aspx>, 2016. Last access: 22 February 2023.
- ICAO: Overview of Automatic Dependent Surveillance-Broadcast (ADS-B) Out, <https://www.icao.int/NACC/Documents/Meetings/2021/ADSB/P01-OverviewAD>, 2021. Last access: 22 February 2023.
- 640 Jensen, E. J., Ackerman, A. S., Stevens, D. E., Toon, O. B., and Minnis, P.: Spreading and growth of contrails in a sheared environment, *J. Geophys. Res. Atmos.*, 103, 31557–31567, <https://doi.org/10.1029/98JD02594>, 1998.
- Jeßberger, P., Voigt, C., Schumann, U., Sölch, I., Schlager, H., Kaufmann, S., Petzold, A., Schäuble, D., and Gayet, J.-F.: Aircraft type influence on contrail properties, *Atmos. Chem. Phys.*, 13, 11965–11984, <https://doi.org/10.5194/acp-13-11965-2013>, 2013.
- 645 Kärcher, B.: The importance of contrail ice formation for mitigating the climate impact of aviation, *J. Geophys. Res. Atmos.*, 121, 3497–3505, <https://doi.org/10.1002/2015JD024696>, 2016.
- Kärcher, B.: Formation and radiative forcing of contrail cirrus, *Nat. Commun.*, 9, 1824, <https://doi.org/10.1038/s41467-018-04068-0>, 2018.
- Kärcher, B. and Lohmann, U.: A parameterization of cirrus cloud formation: Homogeneous freezing of supercooled aerosols, *J. Geophys. Res. Atmos.*, 107, AAC 4-1, <https://doi.org/10.1029/2001JD000470>, 2002.
- 650 Kleine, J., Voigt, C., Sauer, D., Schlager, H., Scheibe, M., Jurkat-Witschas, T., Kaufmann, S., Kärcher, B., and Anderson, B. E.: In Situ Observations of Ice Particle Losses in a Young Persistent Contrail, *Geophys. Res. Lett.*, 45, 13,553–13,561, <https://doi.org/10.1029/2018GL079390>, 2018.
- Lamquin, N., Stubenrauch, C. J., Gierens, K., Burkhardt, U., and Smit, H.: A global climatology of upper-tropospheric ice supersaturation occurrence inferred from the Atmospheric Infrared Sounder calibrated by MOZAIC, *Atmos. Chem. Phys.*, 12, 381–405, <https://doi.org/10.5194/ACP-12-381-2012>, 2012.
- 655 Lau, W. K. M. and Kim, K. M.: Robust Hadley circulation changes and increasing global dryness due to CO₂ warming from CMIP5 model projections, *Proc. Natl. Acad. Sci. U. S. A.*, 112, 3630–3635, https://doi.org/10.1073/PNAS.1418682112/SUPPL_FILE/PNAS.201418682SI.PDF, 2015.



- Lee, D. S., Fahey, D. W., Skowron, A., Allen, M. R., Burkhardt, U., Chen, Q., Doherty, S. J., Freeman, S., Forster, P. M., Fuglestad, J.,
660 Gettelman, A., De León, R. R., Lim, L. L., Lund, M. T., Millar, R. J., Owen, B., Penner, J. E., Pitari, G., Prather, M. J., Sausen, R., and
Wilcox, L. J.: The contribution of global aviation to anthropogenic climate forcing for 2000 to 2018, *Atmos. Environ.*, 244, 117834,
<https://doi.org/10.1016/J.ATMOENV.2020.117834>, 2021.
- Lewellen, D. C.: Persistent contrails and contrail cirrus. Part II: Full lifetime behavior, *J. Atmos. Sci.*, 71, 4420–4438,
<https://doi.org/10.1175/JAS-D-13-0317.1>, 2014.
- 665 Lewellen, D. C., Meza, O., Huebsch, W. W., Lewellen, D. C., Meza, O., and Huebsch, W. W.: Persistent Contrails and Contrail Cirrus. Part
I: Large-Eddy Simulations from Inception to Demise, *J. Atmos. Sci.*, 71, 4399–4419, <https://doi.org/10.1175/JAS-D-13-0316.1>, 2014.
- Li, Y., Mahnke, C., Rohs, S., Bundke, U., Spelten, N., Dekoutsidis, G., Groß, S., Voigt, C., Schumann, U., Petzold, A., and Krämer, M.:
Upper-tropospheric slightly ice-subsaturated regions: frequency of occurrence and statistical evidence for the appearance of contrail cirrus,
Atmos. Chem. Phys., 23, 2251–2271, <https://doi.org/10.5194/ACP-23-2251-2023>, 2023.
- 670 Lohmann, U., Spichtinger, P., Jess, S., Peter, T., and Smit, H.: Cirrus cloud formation and ice supersaturated regions in a global climate
model, *Environ. Res. Lett.*, 3, 045022, <https://doi.org/10.1088/1748-9326/3/4/045022>, 2008.
- Mannstein, H., Brömser, A., and Bugliaro, L.: Ground-based observations for the validation of contrails and cirrus detection in satellite
imagery, *Atmos. Meas. Tech.*, 3, 655–669, <https://doi.org/10.5194/amt-3-655-2010>, 2010.
- Mayer, B. and Kylling, A.: Technical note: The libRadtran software package for radiative transfer calculations - description and examples
675 of use, *Atmos. Chem. Phys.*, 5, 1855–1877, <https://doi.org/10.5194/acp-5-1855-2005>, 2005.
- Meerkötter, R., Schumann, U., Doelling, D. R., Minnis, P., Nakajima, T., and Tsushima, Y.: Radiative forcing by contrails, *Ann. Geophys.*,
17, 1080–1094, <https://doi.org/10.1007/s00585-999-1080-7>, 1999.
- Meijer, V. R., Kulik, L., Eastham, S. D., Allroggen, F., Speth, R. L., Karaman, S., and Barrett, S. R. H.: Contrail coverage over the United
States before and during the COVID-19 pandemic, *Environ. Res. Lett.*, 17, 034039, <https://doi.org/10.1088/1748-9326/AC26F0>, 2022.
- 680 Molloy, J., Teoh, R., Harty, S., Koudis, G., Schumann, U., Poll, I., and Stettler, M. E. J.: Design Principles for a Contrail-Minimizing Trial
in the North Atlantic, 9, 375, <https://doi.org/10.3390/AEROSPACE9070375>, 2022.
- Moore, R. H., Thornhill, K. L., Weinzierl, B., Sauer, D., D’Ascoli, E., Kim, J., Lichtenstern, M., Scheibe, M., Beaton, B., and Beyersdorf,
A. J.: Biofuel blending reduces particle emissions from aircraft engines at cruise conditions, *Nature*, 543, 411–415,
<https://doi.org/10.1038/nature21420>, 2017.
- 685 Parr, W. C. and Schucany, W. R.: Minimum distance and robust estimation, *J. Am. Stat. Assoc.*, 75, 616–624,
<https://doi.org/10.1080/01621459.1980.10477522>, 1980.
- Petzold, A., Döpelheuer, A., Brock, C. A., and Schröder, F.: In situ observations and model calculations of black carbon emission by aircraft
at cruise altitude, *J. Geophys. Res. Atmos.*, 104, 22171–22181, <https://doi.org/10.1029/1999JD900460>, 1999.
- Petzold, A., Gysel, M., Vancassel, X., Hittenberger, R., Puxbaum, H., Vrochitcky, S., Weingartner, E., Baltensperger, U., and Mirabel, P.:
690 On the effects of organic matter and sulphur-containing compounds on the CCN activation of combustion particles, *Atmos. Chem. Phys.*, 5,
3187–3203, <https://doi.org/10.5194/ACP-5-3187-2005>, 2005.



- Petzold, A., Thouret, V., Gerbig, C., Zahn, A., Brenninkmeijer, C. A. M., Gallagher, M., Hermann, M., Pontaud, M., Ziereis, H., Boulanger, D., Marshall, J., Nédélec, P., Smit, H. G. J., Friess, U., Flaud, J. M., Wahner, A., Cammas, J. P., Volz-Thomas, A., Thomas, K., Rohs, S., Bundke, U., Neis, P., Berkes, F., Houben, N., Berg, M., Tappertzhofen, M., Blomel, T., Pätz, W., Filges, A., Boschetti, F., Verma, S., Baum, S., Athier, G., Cousin, J. M., Sauvage, B., Blot, R., Clark, H., Gaudel, A., Gressent, A., Auby, A., Fontaine, A., Gautron, B., Bennouna, Y., Petetin, H., Karcher, F., Abonne, C., Dandin, P., Beswick, K., Wang, K. Y., Rauthe-Schöch, A., Baker, A. K., Riede, H., Gromov, S., Zimmermann, P., Thorenz, U., Scharffe, D., Koepfel, C., Slemr, F., Schuck, T. J., Umezawa, T., Ditas, J., Cheng, Y., Schneider, J., Williams, J., Neumaier, M., Christner, E., Fischbeck, G., Safadi, L., Petrelli, A., Gehrlein, T., Heger, S., Dyroff, C., Weber, S., Assmann, D., Rubach, F., Weigelt, A., Stratmann, G., Stock, P., Penth, L., Walter, D., Heue, K. P., Allouche, Y., Marizy, C., Hermira, J., Bringtown, S., Saueressig, G., Seidel, N., Huf, M., Waibel, A., Franke, H., Klaus, C., Stosius, R., Baumgardner, D., Braathen, G., Paulin, M., and Garnett, N.: Global-scale atmosphere monitoring by in-service aircraft – current achievements and future prospects of the European Research Infrastructure IAGOS, *Tellus, Ser. B Chem. Phys. Meteorol.*, 67, 1–24, <https://doi.org/10.3402/TELLUSB.V67.28452>, 2015.
- Petzold, A., Neis, P., Rütimann, M., Rohs, S., Berkes, F., G. J. Smit, H., Krämer, M., Spelten, N., Spichtinger, P., Nédélec, P., and Wahner, A.: Ice-supersaturated air masses in the northern mid-latitudes from regular in situ observations by passenger aircraft: Vertical distribution, seasonality and tropospheric fingerprint, *Atmos. Chem. Phys.*, 20, 8157–8179, <https://doi.org/10.5194/ACP-20-8157-2020>, 2020.
- Pruppacher, H. R., Klett, J. D., and Wang, P. K.: *Microphysics of Clouds and Precipitation*, 28, 381–382, <https://doi.org/10.1080/02786829808965531>, 2007.
- Quadros, F. D. A., Snellen, M., Sun, J., and Dedoussi, I. C.: Global Civil Aviation Emissions Estimates for 2017–2020 Using ADS-B Data, *J. Aircr.*, 1–11, <https://doi.org/10.2514/1.C036763>, 2022.
- Reutter, P., Neis, P., Rohs, S., and Sauvage, B.: Ice supersaturated regions: Properties and validation of ERA-Interim reanalysis with IAGOS in situ water vapour measurements, *Atmos. Chem. Phys.*, 20, 787–804, <https://doi.org/10.5194/ACP-20-787-2020>, 2020.
- Santer, B. D., Sausen, R., Wigley, T. M. L., Boyle, J. S., AchutaRao, K., Doutriaux, C., Hansen, J. E., Meehl, G. A., Roeckner, E., Ruedy, R., Schmidt, G., and Taylor, K. E.: Behavior of tropopause height and atmospheric temperature in models, reanalyses, and observations: Decadal changes, *J. Geophys. Res. Atmos.*, 108, ACL 1-1, <https://doi.org/10.1029/2002JD002258>, 2003.
- Sanz-Morère, I., Eastham, S. D., Allroggen, F., Speth, R. L., and Barrett, S. R. H.: Impacts of multi-layer overlap on contrail radiative forcing, *Atmos. Chem. Phys.*, 21, 1649–1681, <https://doi.org/10.5194/ACP-21-1649-2021>, 2021.
- Schumann, U.: On conditions for contrail formation from aircraft exhausts, *Meteorol. Zeitschrift*, 5, 4–23, <https://doi.org/10.1127/metz/5/1996/4>, 1996.
- Schumann, U.: A contrail cirrus prediction model, *Geosci. Model Dev.*, 5, 543–580, <https://doi.org/10.5194/gmd-5-543-2012>, 2012.
- Schumann, U. and Graf, K.: Aviation-induced cirrus and radiation changes at diurnal timescales, *J. Geophys. Res. Atmos.*, 118, 2404–2421, <https://doi.org/10.1002/jgrd.50184>, 2013.
- Schumann, U. and Heymsfield, A. J.: On the lifecycle of individual contrails and contrail cirrus, *Meteorol. Monogr.*, 58, 3.1-3.24, <https://doi.org/10.1175/amsmonographs-d-16-0005.1>, 2017.
- Schumann, U., Mayer, B., Hamann, U., Graf, K.: Radiative heating in contrail cirrus: <https://elib.dlr.de/67811/>, 2010. Last access: 5



725 December 2022.

Schumann, U., Graf, K., and Mannstein, H.: Potential to reduce the climate impact of aviation by flight level changes, in: 3rd AIAA Atmospheric Space Environments Conference, <https://doi.org/10.2514/6.2011-3376>, 2011.

Schumann, U., Mayer, B., Graf, K., and Mannstein, H.: A parametric radiative forcing model for contrail cirrus, *J. Appl. Meteorol. Climatol.*, 51, 1391–1406, <https://doi.org/10.1175/JAMC-D-11-0242.1>, 2012.

730 Schumann, U., Penner, J. E., Chen, Y., Zhou, C., and Graf, K.: Dehydration effects from contrails in a coupled contrail–climate model, *Atmos. Chem. Phys.*, 15, 11179–11199, <https://doi.org/10.5194/acp-15-11179-2015>, 2015.

Schumann, U., Baumann, R., Baumgardner, D., Bedka, S., Duda, D., Freudenthaler, V., Gayet, J.-F., Heymsfield, A. J., Minnis, P., and Quante, M.: Properties of individual contrails: A compilation of observations and some comparisons, *Atmos. Chem. Phys.*, 17, 1–62, <https://doi.org/10.5194/acp-17-403-2017>, 2017.

735 Schumann, U., Poll, I., Teoh, R., Koelle, R., Spinielli, E., Molloy, J., Koudis, G. S., Baumann, R., Bugliaro, L., Stettler, M., and Voigt, C.: Air traffic and contrail changes over Europe during COVID-19: A model study, *Atmos. Chem. Phys.*, 21, 7429–7450, <https://doi.org/10.5194/ACP-21-7429-2021>, 2021.

Shapiro, M., Engberg, Z., Teoh, R., and Dean, T.: pycontrails: Python library for modeling aviation climate impacts, <https://doi.org/10.5281/zenodo.7776686>, 2023.

740 Teoh, R., Stettler, M. E. J., Majumdar, A., Schumann, U., Graves, B., and Boies, A.: A methodology to relate black carbon particle number and mass emissions, *J. Aerosol Sci.*, 132, 44–59, <https://doi.org/10.1016/J.JAEROSCI.2019.03.006>, 2019.

Teoh, R., Schumann, U., Majumdar, A., and Stettler, M. E. J.: Mitigating the Climate Forcing of Aircraft Contrails by Small-Scale Diversions and Technology Adoption, *Environ. Sci. Technol.*, 54, 2941–2950, <https://doi.org/10.1021/acs.est.9b05608>, 2020.

745 Teoh, R., Schumann, U., Gryspeerdt, E., Shapiro, M., Molloy, J., Koudis, G., Voigt, C., and Stettler, M.: Aviation Contrail Climate Effects in the North Atlantic from 2016–2021., 22, 10919–10935, <https://doi.org/https://doi.org/10.5194/acp-2022-169>, 2022a.

Teoh, R., Schumann, U., Voigt, C., Schripp, T., Shapiro, M., Engberg, Z., Molloy, J., Koudis, G., and Stettler, M. E. J.: Targeted Use of Sustainable Aviation Fuel to Maximise Climate Benefits, *Environ. Sci. Technol.*, 56, 17246–17255, <https://doi.org/https://doi.org/10.1021/acs.est.2c05781>, 2022b.

750 Teoh, R., Engberg, Z., Shapiro, M., Dray, L., and Stettler, M. E. J.: A high-resolution Global Aviation emissions Inventory based on ADS-B (GAIA) for 2019 - 2021 [preprint], <https://doi.org/doi.org/10.5194/egusphere-2023-724>, 2023.

Tompkins, A. M., Gierens, K., and Rädcl, G.: Ice supersaturation in the ECMWF integrated forecast system, *Q. J. R. Meteorol. Soc.*, 133, 53–63, <https://doi.org/10.1002/qj.14>, 2007.

Unterstrasser, S. and Gierens, K.: Numerical simulations of contrail-to-cirrus transition-Part 2: Impact of initial ice crystal number, radiation, stratification, secondary nucleation and layer depth, *Atmos. Chem. Phys.*, 10, 2037–2051, <https://doi.org/10.5194/ACP-10-2037-2010>, 2010.

755 Vázquez-Navarro, M., Mannstein, H., and Kox, S.: Contrail life cycle and properties from 1 year of MSG/SEVIRI rapid-scan images, *Atmos. Chem. Phys.*, 15, 8739–8749, <https://doi.org/10.5194/acp-15-8739-2015>, 2015.



- Voigt, C., Schumann, U., Minikin, A., Abdelmonem, A., Afchine, A., Borrmann, S., Boettcher, M., Buchholz, B., Bugliaro, L., Costa, A., Curtius, J., Dollner, M., Dörnbrack, A., Dreiling, V., Ebert, V., Ehrlich, A., Fix, A., Forster, L., Frank, F., Fütterer, D., Giez, A., Graf, K., Groß, J. U., Groß, S., Heimerl, K., Heinold, B., Hüneke, T., Järvinen, E., Jurkat, T., Kaufmann, S., Kenntner, M., Klingebiel, M., Klimach, T., Kohl, R., Krämer, M., Krisna, T. C., Luebke, A., Mayer, B., Mertes, S., Molleker, S., Petzold, A., Pfeilsticker, K., Port, M., Rapp, M., Reutter, P., Rolf, C., Rose, D., Sauer, D., Schäfler, A., Schlage, R., Schnaiter, M., Schneider, J., Spelten, N., Spichtinger, P., Stock, P., Walser, A., Weigel, R., Weinzierl, B., Wendisch, M., Werner, F., Wernli, H., Wirth, M., Zahn, A., Ziereis, H., and Zöger, M.: ML-CIRRUS: The Airborne Experiment on Natural Cirrus and Contrail Cirrus with the High-Altitude Long-Range Research Aircraft HALO, *Bull. Am. Meteorol. Soc.*, 98, 271–288, <https://doi.org/10.1175/BAMS-D-15-00213.1>, 2017.
- 760 Voigt, C., Kleine, J., Sauer, D., Moore, R. H., Bräuer, T., Le Clercq, P., Kaufmann, S., Scheibe, M., Jurkat-Witschas, T., Aigner, M., Bauder, U., Boose, Y., Borrmann, S., Crosbie, E., Diskin, G. S., DiGangi, J., Hahn, V., Heckl, C., Huber, F., Nowak, J. B., Rapp, M., Rauch, B., Robinson, C., Schripp, T., Shook, M., Winstead, E., Ziemba, L., Schlager, H., and Anderson, B. E.: Cleaner burning aviation fuels can reduce contrail cloudiness, *Commun. Earth Environ.* 2021 21, 2, 1–10, <https://doi.org/10.1038/s43247-021-00174-y>, 2021.
- Voigt, C., Sauer, D., Markl, R., Hahn, V., Kaufmann, S., Dischl, R., Sakellariou, F., Harlaß, T., Scheibe, M., Marsing, A., Tomsche, L., Rottger, C., Dornbrack, A., Roiger, A., Le Clercq, P., Rauch, B., Seeliger, K., LeChenadec, G., Feuillerat, E., Moreau, J., O’Nions, D., Delhay, D., Greslin, E., Darbois, L., Mambre, P., and Renard, C.: Inflight measurements of particle emissions and contrails from an A319neo with lean burn engine technology, in: TAC-5 Conference, 2022.
- 770 Wilhelm, L., Gierens, K., and Rohs, S.: Weather Variability Induced Uncertainty of Contrail Radiative Forcing, *Aerosp.* 2021, Vol. 8, Page 332, 8, 332, <https://doi.org/10.3390/AEROSPACE8110332>, 2021.
- Wilkerson, J. T., Jacobson, M. Z., Malwitz, A., Balasubramanian, S., Wayson, R., Fleming, G., Naiman, A. D., and Lele, S. K.: Analysis of emission data from global commercial aviation: 2004 and 2006, *Atmos. Chem. Phys.*, 10, 6391–6408, <https://doi.org/10.5194/acp-10-6391-2010>, 2010.
- World Bank: Air transport, passengers carried: <https://data.worldbank.org/indicator/IS.AIR.PSGR>, 2023. Last access: 14 May 2023.