

Reviews of Geophysics®



REVIEW ARTICLE

10.1029/2023RG000828

Key Points:

- Atmospheric forcings, land use and management, and soil processes and mechanisms explain how and why soil moisture memory emerges in ecosystems
- Nonlocality of moisture memory, its spread across different regions, and its interaction with large-scale climate phenomena are underexplored
- Further advances in land surface models and closer integration of model simulations and observations are needed to better characterize moisture memory

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

M. Rahmati,
mehdirmti@gmail.com;
m.rahmati@fz-juelich.de

Citation:

Rahmati, M., Amelung, W., Brogi, C., Dari, J., Flammini, A., Bogen, H., et al. (2024). Soil moisture memory: State-of-the-art and the way forward. *Reviews of Geophysics*, 62, e2023RG000828. <https://doi.org/10.1029/2023RG000828>

Received 2 OCT 2023

Accepted 2 MAY 2024















Author Contributions:

Conceptualization: Mehdi Rahmati, Wulf Amelung, Harry Vereecken
Formal analysis: Mehdi Rahmati
Investigation: Mehdi Rahmati, Cosimo Brogi, Jacopo Dari, Alessia Flammini, Heye Bogen, Luca Brocca, Hao Chen, Jannis Groh, Shirin Moradi, Arash Rahi, Farnaz Sharghi S., Harry Vereecken
Methodology: Mehdi Rahmati, Harry Vereecken
Resources: Mehdi Rahmati
Software: Mehdi Rahmati
Supervision: Mehdi Rahmati

© 2024. The Authors.

This is an open access article under the terms of the [Creative Commons Attribution License](#), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Soil Moisture Memory: State-Of-The-Art and the Way Forward

Mehdi Rahmati^{1,2} , Wulf Amelung^{1,3}, Cosimo Brogi¹ , Jacopo Dari^{4,5} , Alessia Flammini⁴ , Heye Bogen¹ , Luca Brocca⁵ , Hao Chen^{1,6}, Jannis Groh^{1,3,7} , Randal D. Koster⁸ , Kaighin A. McColl^{9,10} , Carsten Montzka¹ , Shirin Moradi¹ , Arash Rahi^{1,4} , Farnaz Sharghi S.³ , and Harry Vereecken¹ 

¹Agrosphere Institute IBG-3, Forschungszentrum Jülich GmbH, Jülich, Germany, ²Department of Soil Science and Engineering, University of Maragheh, Maragheh, Iran, ³Institute of Crop Science and Resource Conservation (INRES)-Soil Science and Soil Ecology, University of Bonn, Bonn, Germany, ⁴Department of Civil and Environmental Engineering, University of Perugia, Perugia, Italy, ⁵Research Institute for Geo-Hydrological Protection, National Research Council, Perugia, Italy, ⁶School of Geographic and Environmental Sciences, Tianjin Normal University, Tianjin, China, ⁷Isotope Biogeochemistry and Gasfluxes, Landscape Functioning, ZALF, Müncheberg, Germany, ⁸Global Modeling and Assimilation Office, NASA Goddard Space Flight Center, Greenbelt, MD, USA, ⁹Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA, USA, ¹⁰School of Engineering and Applied Sciences, Harvard University, Cambridge, MA, USA

Abstract Soil moisture is an essential climate variable of the Earth system. Understanding its spatiotemporal dynamics is essential for predicting weather patterns and climate variability, monitoring and mitigating the effects and occurrence of droughts and floods, improving irrigation in agricultural areas, and sustainably managing water resources. Here we review in depth how soils can remember information on soil moisture anomalies over time, as embedded in the concept of soil moisture memory (SMM). We explain the mechanisms underlying SMM and explore its external and internal drivers; we also discuss the impacts of SMM on different land surface processes, focusing on soil-plant-atmosphere coupling. We explore the spatiotemporal variability, seasonality, locality, and depth-dependence of SMM and provide insights into both improving its characterization in land surface models and using satellite observations to quantify it. Finally, we offer guidance for further research on SMM.

Plain Language Summary Our review paper takes an in-depth look at soil moisture memory, which is how soil records its moisture history over time and space. Analogous to human psychology, which seeks to understand how a person's/society's memory influences his/her present and future behavior, understanding soil moisture memory encourages consideration of how such memory determines present state and might determine future behavior of soils exposed to environmental disturbances. Soil moisture memory can be affected by a variety of factors, both external (e.g., weather extremes) and internal (soil's unique properties). It affects everything from the air to the way our landscapes respond to disasters like droughts, wildfires, and floods. We also studied how this phenomenon affects the balance of water and energy in our environment, the health of our plants, and even how it communicates with the atmosphere. We show how it can change depending on where you are on the planet, the time of year, and how deep you dig into the soil. We offer scientists insights into how weather and land surface models can become more accurate by accounting for soil moisture memory. Its understanding not only helps us predict and manage our environment, but also provides opportunities for exciting scientific discoveries.

1. Introduction

Soil as upper “skin” of the Earth's surface harbors plant roots and provides habitat to (soil) biota. The rooting depth can be variable, from a few centimeters up to 2 m, as in reanalysis data such as GLDAS-NOAH (Beau-
doing, 2016; Rodell, Houser, Jambor, et al., 2004; Rodell, Houser, Peters-Lidard, et al., 2004), ERA5-Land (Hersbach et al., 2020), and GLEAM (Martens et al., 2017; Miralles et al., 2011). Hence, soils are considered here as a biologically active region on soils, exceeding common classification depths of 1–2 m, but encompassing the whole area of root growth, contact to groundwater, and annual freeze-thaw cycles in permafrost areas (Chesworth, 2007). Soils thus supply the water that is transpired by plants or evaporated directly from the soil surface, with or without connection to the groundwater. It should be noted here, however, that this term does not

Validation: Mehdi Rahmati,
Wulf Amelung, Randal D. Koster, Kaighin
A. McColl, Carsten Montzka,
Harry Vereecken

Visualization: Mehdi Rahmati
Writing – original draft: Mehdi Rahmati,
Wulf Amelung, Cosimo Brogi,
Jacopo Dari, Alessia Flammini,
Heye Bogena, Luca Brocca, Hao Chen,
Jannis Groh, Randal D. Koster, Kaighin
A. McColl, Carsten Montzka,
Shirin Moradi, Arash Rahi, Farnaz Sharghi
S., Harry Vereecken

Writing – review & editing:
Mehdi Rahmati, Wulf Amelung,
Cosimo Brogi, Jacopo Dari,
Alessia Flammini, Heye Bogena,
Luca Brocca, Hao Chen, Jannis Groh,
Randal D. Koster, Kaighin A. McColl,
Carsten Montzka, Shirin Moradi,
Arash Rahi, Farnaz Sharghi S.,
Harry Vereecken

include the full vadose or critical zone, which extends down to the geological weathering front and might not be reached by plants and related symbiotic organisms. In general, at the ecosystem scale, 60% to 80% (with a global mean value of $61 \pm 15\%$) of the global terrestrial evapotranspiration (~ 567 mm per year (Elnashar et al., 2021)) occurs in the form of transpiration and the remaining occurs in the form of evaporation (by ignoring the interception loss) (Schlesinger & Jasechko, 2014). Soils can regulate the storage of water and its support for plants and groundwater recharge (Vereecken et al., 2016). Hence, soil provides important ecosystem services to society.

Soil moisture, as a key observable variable of soil system, serves as a vital link between the atmosphere, plants, and the subsurface, and thus plays a critical role in several land-surface and ecological processes. Soil moisture is defined as the amount of the water in the active (rooted) layer of the soil, typically in the top 1–2 m soil layer (Robock, 2003), which has a large interaction with the atmosphere (Orth & Seneviratne, 2012) and is “connected” in the sense of Good et al. (2015) (though not, of course, including surface intercepted water or plant vascular water); it is usually measured as volumetric or gravimetric water content and related to the soil water potential through the water retention characteristic. Soil moisture directly affects agricultural productivity, as well as the overall terrestrial water cycle, related climate patterns, and ecosystem dynamics (Robock, 2003). Soil moisture is one of the most important drivers of land surface greenness by providing water for irrigation and natural vegetation (Robock, 2003), and thus impacting food and environmental security. It affects cloud formation and thus the occurrence of precipitation and consequently the occurrence and duration of droughts through the supply or lack of water that evaporates from the soil and transpired by vegetation (Robock, 2003; Seneviratne et al., 2010; Tuller et al., 2023). It also controls soil surface temperature, thereby influencing the occurrence of wildfires and heat waves by affecting the distribution of available energy at the surface into sensible and latent heat (Robock, 2003; Seneviratne et al., 2010; Tuller et al., 2023). Soil moisture also influences the occurrence of floods by determining how much precipitation or snowmelt flows directly into rivers and streams (Robock, 2003).

Fundamental to understanding soil moisture's connection to and control over these aspects of the climate system is its “memory”—the fact that a wet or dry soil moisture anomaly can persist over a long time, sometimes weeks to months. Relative to atmospheric quantities (wind speed, temperatures, etc.), soil moisture varies slowly, effectively imposing a time filter on the higher-frequency processes that modify it. This imposes a slowly varying forcing on the processes that soil moisture in turn affects. A proper understanding of any natural process that involves soil moisture thus relies on a proper characterization of soil moisture memory (SMM). SMM is in fact critically important in the prediction arena (for e.g., Dirmeyer et al., 2015; Mariotti et al., 2020; Robertson & Vitart, 2001)—knowledge of a current soil moisture anomaly implies some knowledge of the anomaly weeks hence and thus some knowledge of the processes that this future anomaly will affect. Considering that the SMM is so central to our understanding of the role of soil moisture in the climate system, we believe that an overview of the current state of research on this topic is urgently needed.

Soil as a complex system has many state variables which are interconnected and influence each other while they are evolving through time. Soil moisture is one of the observable state variables of such a complex system. These variables as well as external forcings may leave an imprint on soil moisture dynamics that can be described by SMM. Typically, variables of complex systems are subjected to reversible oscillations as well as irreversible decays that paradoxically coexist and lead to formation of memory in these systems (Kenkre, 2021). The co-existence of oscillations and decays also applies to soil moisture. In this context, Delworth and Manabe (1988) considered soil moisture evolution as a first-order Markovian process and used it to analyze soil moisture decay. Later, Koster and Suarez (2001) showed that the inverse of the decay rate introduced by Delworth and Manabe (1988) determines the SMM timescale. There is already sufficient evidence that the SMM timescale is important from various points of view and has implications for various land surface processes, from surface energy balance and drought occurrence and severity to biogeochemical cycles. Therefore, in this paper, we provide a comprehensive review of previous research on SMM, examining its drivers and impacts on land surface processes and discussing the current state of research in this area. The article is organized as follows with Section 2 first defines the concept of SMM and quantification of its timescale and discusses the different terminologies used for SMM. Section 3 comments on the length of the SMM timescale as reported in the literature and discusses its temporal variability. Section 4 discusses the spatial variability of the SMM timescale. In Section 5, we first provide information on the coupling of soil moisture with land surface processes and the hotspots of soil moisture-atmosphere coupling, and then address the factors controlling SMM and the impact of SMM on various land surface processes. Section 6 discusses how SMM is integrated and represented by models. Section 7 provides a discussion on how SMM can be observed from space. In Section 8, we discuss how the concept of

SMM can be used for soil moisture prediction and the downscaling of large-scale soil moisture products. Finally, Section 9 discusses current issues in the field and prospects for future research, and Section 10 provides a summary and outlook for the paper.

2. SMM: Soil Moisture Memory

2.1. Concept

The term SMM can be traced back to the seminal work of Koster and Suarez (2001), who built on the work of Hasselmann (1976) and Delworth and Manabe (1988). Koster and Suarez (2001) defined SMM as the time required for the soil column to “forget” a perturbation, which might have arisen from an extreme precipitation event or from an anomalously dry period. Hasselmann (1976) proposed a concept that emphasizes the ability of a particular component within the climate system, characterized by high-frequency fluctuations, to influence another component, resulting in low-frequency fluctuations. Building on this, Frankignoul and Hasselmann (1977) provided a practical demonstration of this theory by showing how short-term atmospheric forcings can trigger long-term anomalies in sea surface temperatures, which in turn can be attributed to the response of the oceanic surface layer. Similarly, Shukla and Mintz (1982) also effectively discussed SMM: “In the extratropics, with its large seasonal changes, the soil plays a role analogous to that of the ocean. The ocean stores some of the radiational energy it receives in summer and uses it to heat the atmosphere over the ocean in winter. The soil stores some of the precipitation it receives in winter and uses it to humidify the atmosphere in summer.” In this analogy, the soil functions similarly to the ocean by taking the random precipitation and producing a time series of anomalies in soil moisture (Delworth & Manabe, 1988). We should note, however, that soil moisture variability generally occurs on shorter timescales than sea surface temperature variability, and this variability is characterized by the interactions between soil moisture and atmosphere as influenced by the energy and water balance of the land surface (Timbal et al., 2002).

More recently, Song et al. (2019) approached the definition of SMM from a novel perspective, viewing it as the period wherein detectable moisture anomalies hold the potential to influence the atmosphere. Gao et al. (2018) explained this concept by pointing to the link between positive and negative soil moisture anomalies and corresponding rainfall surplus or deficits, thus triggering a domino effect on subsequent periods of increased or decreased evapotranspiration, then on the water and energy balances of the land surface and from there again the atmospheric state. Encompassing a broader perspective, Ruscica et al. (2014) assumed that anomalous soil moisture impacts the atmospheric state through complicated land surface feedback mechanisms that span across diurnal to seasonal timescales. The multifaceted nature of SMM finds expression in the explanation offered by He et al. (2023), who propose two distinct but not independent descriptions: one represents the SMM as the temporal duration required for a perturbation to manifest and decay in the time domain (irreversible changes), while the second definition relates to the time taken for soil moisture to regain equilibrium following a perturbation (reversible changes). In any case, the perturbations considered so far encompass a diverse array of wet anomalies like precipitation or dry anomalies like drought. Sörensson and Berbery (2015) presented SMM as a gauge of the temporal span during which a moisture anomaly retains detectability and sustains the potential to exert influence upon the atmosphere.

Drawing from cognitive analogies, Asharaf and Ahrens (2013) expressed memory as a complicated process of encoding and recalling information, whereby the power of memory stems from intrinsic changes within the system. These system changes are not necessarily included in the definitions noted above. However, such a notion of soil memory has a major impact on the predictability of weather and climate events (Santanello Jr et al., 2018), thus enriching our understanding of the temporal variability that governs our climate system on Earth.

2.2. Quantification

A typical framework used in the literature to analyze SMM is the 1D soil moisture balance equation for a homogeneous soil (Delworth & Manabe, 1988; McColl, Wang, et al., 2017):

$$C_s \frac{dS(t)}{dt} = P(t) - L(S(t)) = P(t) - [D(S(t)) + ET(S(t)) + Q(S(t))] \quad (1)$$

where $S(t)$ is soil saturation degree (dimensionless) at time t (T), $P(t)$ is the rainfall rate (LT^{-1}) and $L(S(t))$ is the soil water loss rate (LT^{-1}). The components of loss term include $Q(S(t))$ —surface runoff rate (LT^{-1}), $D(S(t))$ —the drainage rate (LT^{-1}), and $ET(S(t))$ —evapotranspiration (LT^{-1}); all as a function of $S(t)$. The quantity C_s is soil water storage capacity (L), which is defined as $C_s = n\Delta z$, where n is soil porosity (L^3L^{-3}) and Δz is soil rooting depth or active layer (L). The $S(t)$ term is also defined as $\theta(t)/\theta_{\text{sat}}$ where $\theta(t)$ is volumetric soil moisture content (L^3L^{-3}) at the time t and θ_{sat} is the saturated moisture content of soil (L^3L^{-3}).

Delworth and Manabe (1988), building on the pioneering work of Hasselmann (1976) who applied first-order Markov processes to explore the dependencies between white noise (short-term variation) and red noise spectra of sea surface temperatures, explored the temporal spectrum of soil moisture anomalies. They showed that soil moisture dynamics as described by Equation 1 can be formulated as a first-order Markov process:

$$\frac{dW(t)}{dt} = -\lambda W(t) + \omega(t) \quad (2a)$$

$$\omega(t) = \text{rainfall} + \text{snowmelt} - \text{runoff} \quad (2b)$$

where $W(t)$ represents soil moisture (L) in the soil root zone as a function of time t (T). As defined above, $W(t) = C_s S(t)$. The term $\omega(t)$ represents the white noise (LT^{-1}) at time t , and λ (T^{-1}) is a constant defined as $\lambda = E_0/W_{\text{FC}}$, where E_0 is potential evapotranspiration (LT^{-1}) and W_{FC} is soil moisture at field capacity (L). The quantity $1/\lambda$ denotes the decay timescale (T) of the autocorrelation function, later defined as the timescale of SMM by Koster and Suarez (2001). The approach assumes that (a) anomalies of effective precipitation (precipitation minus runoff) can be represented as a white noise process and (b) anomalies of evapotranspiration can be approximated as a linear function of soil moisture.

Inspired by the above formalisms, several approaches have been proposed to quantify the timescale of SMM based on the analysis of time series data of soil moisture; these approaches include computing the e-folding autocorrelation, integral timescale, soil moisture variance spectrum, and decorrelation time as well as employing a hybrid stochastic-deterministic model, as detailed further below. However, to date, the research conducted by McColl, Alemohammad, et al. (2017) is, to the best of our knowledge, almost the only investigation that evaluates comprehensively the advantages and disadvantages of these metrics when it comes to quantifying the memory timescale of soil moisture. They mentioned three aspects in which memory metrics may differ: timescale definition, anomaly reference state, and consideration of positive or negative anomalies. They state that commonly used autocorrelation-based metrics, such as e-folding and integral timescales, are fine to the extent that the time series is approximated as red noise. While this is often a reasonable approximation at monthly or longer time scales, it is often invalid at shorter time scales. In addition, they argue that autocorrelation-based measurement techniques ignore the sign of the soil moisture anomaly and thus neglect valuable information. It is argued that the manifestation of positive peaks in soil moisture is caused by rapid, irregular precipitation events, whereas negative anomalies of soil moisture content are caused by more gradual, quasi-deterministic mechanisms exemplified by the complicated interplay of evapotranspiration processes. McColl, Alemohammad, et al. (2017) suggest that it would be beneficial to quantify the dissipation timescales of these fast and slow processes separately. They also considered metrics that have been proposed to overcome the above limitations, including mean persistence time, which measures the average amount of time that the soil moisture time series spends above or below a fixed threshold, such as soil moisture at the wilting point. They caution, however, that while this approach considers positive and negative anomalies separately, it still depends on the choice of threshold. Although the scope of the study by McColl, Alemohammad, et al. (2017) (focusing on six metrics) is appreciated, a more comprehensive analysis that incorporates all identified metrics seems necessary to select the most appropriate metric for the memory time scale. We thus call for such a comprehensive analysis in future research that systematically compares the performance and applicability of all the different metrics identified in this review.

Before diving into the details of the SMM timescale metrics, we would like to point out that while some references use τ as the notation for the SMM timescale, we suggest here the use of t_{SMM} (with a dimension of [T]) instead given that τ also refers to time lag in these formulations. We will also continue to use the notation SMM when referring to the concept of soil moisture memory (rather than its timescale) as an emergent property of the complex soil system (“emergent” in the sense that it arises from various interacting components in the earth/soil system without belonging to the individual components themselves).

2.2.1. Autocorrelation Timescales

t_{SMM} is usually defined as the time lag at which autocorrelation in soil moisture data is reduced to its e-folding (Delworth & Manabe, 1988; Vinnikov & Yesserkepova, 1991; Wu & Dickinson, 2004) or it is reduced to zero (Ghannam et al., 2016). Delworth and Manabe (1988) (with further reformulations by Vinnikov and Yesserkepova (1991), Robock et al. (1995), Vinnikov et al. (1996), and Dirmeyer et al. (2016)) defined the autocorrelation function, $r(\tau)$, of a time series of soil moisture measurements as follows, based on a first-order statistical model of the Markov process:

$$r(\tau) = 1 - \alpha \quad \tau = 0 \quad (3)$$

$$r(\tau) = \exp(-\lambda\tau) - \alpha \quad \tau \neq 0 \quad (4)$$

where τ is the lag (T), λ with a dimension of $1/T$ is the constant from Equation 2a, and α is part of the variance that is attributable to random processes without autocorrelation being ascribed to the random error of the measurements (Vinnikov & Yesserkepova, 1991). The autocorrelation expressed by above equations provide the sum of the red (with a variance of unity for pure red-noise spectrum) and white noises (Robock et al., 1995; Vinnikov & Yesserkepova, 1991). Later, Robock et al. (1995) using observational soil moisture data approximated the α to be 0.1.

According to above formulations, the t_{SMM} can be defined in three ways: (a) the first-time lag (τ) at which $r(\tau)$ drops to $1/e \approx 0.37$ (e-folding) of its initial value ($=1$), (b) the first-time lag (τ) at which $r(\tau)$ crosses zero (Ghannam et al., 2016), (c) or the first time lag at which it drops below the autocorrelation corresponding to the 95% or 99% confidence level (Dirmeyer et al., 2009; MahfuzurRahman & Lu, 2015; Ruscica et al., 2014), given the sample size. The latter corresponds to the lag value at which the autocorrelation reaches the lowest significant ($p = 0.05$ or 0.01) values. Alternatively, one computes the area under the $r(\tau)$ -curve (Ghannam et al., 2016; Katul et al., 2007; McColl, Alemohammad, et al., 2017) obtained from Equation 4 to produce an integral timescale:

$$t_{\text{SMM}} = \int_0^{+\infty} r(\tau) d\tau \quad (5)$$

The above formulation assumes that $r(\tau)$ decays to zero as τ tends to infinity. In order to determine the autocorrelation of the data, the seasonal cycle must be removed from the data for all the methods mentioned above before calculations are performed (Vinnikov & Yesserkepova, 1991). However, Ghannam et al. (2016) argue and show that the de-trending treatments cause losses in the variances of the soil moisture time series, which can consequently lead to losses in the memory of the time series, especially when estimated through the integral timescale. This is because de-trending changes the spectrum of soil moisture at low frequencies. Interestingly, applying different detrending methods (e.g., de-trending by local monthly or seasonal averages) can result in different ranges of autocorrelation and clearly different memory timescales (Ghannam et al., 2016). Therefore, t_{SMM} estimates should be interpreted with caution when the soil moisture time series is subjected to different statistical treatments.

Several researchers (Koster & Suarez, 2001; Orth & Seneviratne, 2012, 2013; Seneviratne, Koster, et al., 2006; Seneviratne & Koster, 2012; Wei et al., 2006) have also used interannual autocorrelation over a particular lag to quantify t_{SMM} . To do this, one needs to find the correlation between soil moisture data of day n from all years and the data from day $n + \tau$ from all years. The largest τ value that results in a significant autocorrelation at a 95% confidence level is treated as a measure of t_{SMM} (Rahman et al., 2015) (Figure 1).

Vinnikov et al. (1999) followed by Entin et al. (2000) showed that there might be two different timescales for a particular climate system (Hasselmann, 1976). This is particularly the case when rainfall is not climatologically random or when excessive runoff occurs (Delworth & Manabe, 1988). In this regard, Entin et al. (2000) separated the temporal variance of soil moisture into two components: (a) one at a small temporal scale, determined by land surface type (soil characteristics, topography, vegetation, and root structure), and (b) one at a large temporal scale, reflecting atmospheric forcing. For both components, time remains the measurement unit. They characterized the small-scale component of soil moisture variance in time as white noise and the large-scale component as red noise. The basic idea behind this concept is that the nature of the soil surface affects the direct infiltration of water




Figure 1. Calculation of soil moisture memory timescale (t_{SMM}) from time series data of soil moisture (represented by filled black circles) based on the interannual e-folding method. The pale dots in the above figure mean that the data of a particular year can be excluded from the analysis during different iterations to examine the effects of that specific year on long-term t_{SMM} . The notations of r_1 , r_2 , r_τ , and $r_{\tau+1}$ donate autocorrelation at time lags of 1, 2, τ , and $\tau + 1$, respectively, and $r_{0.95}$ donates for significant autocorrelation at a 95% confidence level.

into and through the soil and the amount of water that the soil can store, while the atmospheric component is responsible for the amount of water available to the soil through rain or snowmelt and for the rate at which water is released through evapotranspiration. Accordingly, the total estimated variance of soil moisture, denoted as $\text{var}(\theta)$, is:

$$\text{var}(\theta) = \text{var}_{\text{sur}}(\theta) + \text{var}_{\text{atm}}(\theta) \quad (6)$$

where $\text{var}_{\text{sur}}(\theta)$ and $\text{var}_{\text{atm}}(\theta)$ denote soil moisture variance induced by land surface-related variability and atmosphere-related variability, respectively. Accordingly, Entin et al. (2000) expressed the estimates of the temporal autocorrelation of soil moisture as below:

$$r(\tau) = \text{var}_{\text{sur}}(\theta) \exp\left(-\frac{\tau}{t_{\text{SMM}}^{\text{sur}}}\right) + \text{var}_{\text{atm}}(\theta) \exp\left(-\frac{\tau}{t_{\text{SMM}}^{\text{atm}}}\right) \quad (7)$$

where $r(\tau)$ is the temporal covariance function, τ is the time lag, and $t_{\text{SMM}}^{\text{sur}}$ and $t_{\text{SMM}}^{\text{atm}}$ are the scales of temporal autocorrelation, t_{SMM} , derived by land surface-related variability and atmosphere-related variability, respectively. The smaller timescale, $t_{\text{SMM}}^{\text{sur}}$, is assumed to be of the order of a few days (Entin et al., 2000) and therefore can be ignored when using soil moisture data with temporal resolution of larger than a day (e.g., weekly, or monthly data). However, the larger timescale, $t_{\text{SMM}}^{\text{atm}}$, is assumed to be of the order of months (Entin et al., 2000).

To determine the atmospheric forcing's timescale, autocorrelations are calculated for different time lags (a few days up to a few months, when the autocorrelation approaches zero). Then, the natural logarithm of the autocorrelation estimates is plotted against the applied lag values, and a line of best fit is found. The negative inverse of its slope will provide the atmospheric forcing's temporal timescale, and the y-intercept will provide the variance

induced by red noise (Entin et al., 2000). For the timescale associated with land surface-related variability, the autocorrelations among different locations should be averaged together for each lag value before the same plotting process is applied (Entin et al., 2000).

2.2.2. Soil Moisture Variance Spectrum

The t_{SMM} can also be determined from the normalized temporal spectrum of soil moisture, $E_{\text{ns}}(f)$, where f is the number of cycles per unit time (frequency) (Ghannam et al., 2016; Katul et al., 2007; Nakai et al., 2014). In fact, the $E_{\text{ns}}(f)$ is the Fourier transform of $r(\tau)$, also known as the Wiener-Khinchin theorem, which states that the autocorrelation function of a long-range stationary random process has a spectral decomposition given by the power spectrum of that process (Chatfield, 2003). The $E_{\text{ns}}(f)$ is formulated as follows (Ghannam et al., 2016):

$$E_{\text{ns}}(f) = 2 \int_{-\infty}^{+\infty} r(\tau) e^{-i2\pi f\tau} d\tau \quad (8)$$

Ghannam et al. (2016) used an ad hoc extrapolation of the spectral behavior of $\theta(t)$ when f tends to zero to estimate t_{SMM} as follows:

$$E_{\text{ns}}(0) = 4 \int_0^{+\infty} r(\tau) d\tau = 4t_{\text{SMM}} \rightarrow t_{\text{SMM}} = \frac{E_{\text{ns}}(0)}{4} = \int_0^{+\infty} r(\tau) d\tau \quad (9)$$

The above formulation is identical to the integral timescale.

2.2.3. Decorrelation Time

Von Storch and Zwiers (2002) used “decorrelation time” as a measure of t_{SMM} . According to them, decorrelation time refers to a physical time scale representing the interval between successive uncorrelated observations. It is derived from the lag-1 autocorrelation coefficient (ρ) as follows (Gao et al., 2018; Von Storch and Zwiers, 2002):

$$T_d = \frac{1 + \rho}{1 - \rho} \quad (10)$$

where T_d , the decorrelation time, serves as a measure of t_{SMM} .

2.2.4. Hybrid Stochastic-Deterministic Model

McColl et al. (2019) argued that the theoretical basis for the e-folding autocorrelation timescale (i.e., using a red noise process to approximate soil water balance) is fundamentally suitable for coarse scales (both temporal and spatial) and is thus not applicable at finer spatial and temporal resolutions, as might be encountered with modern satellite observations and models. Therefore, they reconceptualized the SMM and introduced a new hybrid stochastic-deterministic model including a deterministic component for dry conditions and a stochastic component for wet conditions. Finally, they used the occurrence of precipitation to separate the deterministic and stochastic components (Figure 2). The new hybrid model has been formulated as follows (McColl et al., 2019):

$$\frac{d\theta(t)}{dt} = -\frac{\theta(t) - \theta_w}{t_{\text{SMM}}^L} \text{ if precipitation} = 0 \text{ in the interval } [t - \Delta t, t] \quad (11a)$$

$$\frac{d\theta(t)}{dt} = -\frac{\theta(t) - \bar{\theta}}{t_{\text{SMM}}^S} + \varepsilon(t) \text{ if precipitation} > 0 \text{ in the interval } [t - \Delta t, t] \quad (11b)$$

where θ_w is the minimum soil moisture value for the given location, $\bar{\theta}$ is the time average of soil moisture, $\varepsilon(t)$ is an independent and equally distributed random variable with an expected mean value of zero, t is time, and Δt is the time interval of data observations. The quantity t_{SMM}^L is referred to as long-term memory, which is controlled by stage-II evapotranspiration (where the evapotranspiration rate decreases due to the decrease of soil moisture) resolved by the observations, while t_{SMM}^S is referred to as short-term memory, which is determined by a




Figure 2. Soil moisture, $\theta(t)$, drydowns at different timescales. When soil moisture data are collected at sufficiently high frequencies, drydowns can be fully resolved, approximating drying phases with a fast drainage timescale (the short-term memory t_{SMM}^S) and a slower ET timescale (the long-term memory t_{SMM}^L). If the sampling frequency is not high enough, the drydowns are only partially resolved (only the later phases of the drydown). If the sampling frequency is very low (e.g., for older models on a timescale of weeks to months), almost all the drydowns will not be resolved - figure and caption are adopted from McColl et al. (2019).

combination of unresolved processes (especially, but not exclusively, by drainage). Figure 2, adapted from McColl et al. (2019), clearly shows the short- and long-term t_{SMM} for fully and partially resolved and unresolved processes. It should be noted that when the hybrid model is applied to monthly data (“ $\Delta t = 30$ days”), the model reduces to the original red noise model as introduced by the previous metrics. This is because precipitation is non-zero for all time blocks, so that in the reduced form of the hybrid model, t_{SMM}^L is zero and t_{SMM}^S is equivalent to t_{SMM} obtained by the previous metrics.

Calculating t_{SMM}^S and t_{SMM}^L from the hybrid model requires a binary precipitation variable that is significantly flawed when extracted from remote sensing data (McColl et al., 2019). Therefore, McColl et al. (2019) provided two other alternative formulations for t_{SMM}^S and t_{SMM}^L calculations to avoid introducing a separate precipitation time series into the analysis. For brevity, we refrain from providing more information on these alternatives, instead referring the reader to their study.

2.3. Similar Terminologies

Two other terms in the literature refer to the concept of SMM but from different perspectives, namely (a) Anomaly Persistence of Soil Moisture (APSM) and (b) Soil Moisture Drydowns (SMD). The APSM predates the SMM in the literature as it is primarily used in drought characterization research (Oladipo & Hare, 1986). As Oladipo and Hare (1986) reported, Namias (1960) was probably among the first researchers to provide evidence of drought persistence (anomalous moisture conditions) when he showed the persistence of drought from one summer to the next in the continental United States of America. This finding was later evidenced by Walker and Rowntree (1977) in Africa; they noted that once the land was wet or dry, it remained in that condition for at least several weeks. This was also later confirmed by Kraus (1977) and Katz (1978). The more modern concept of the APSM regards it as a measure of the distribution of periods when soil moisture is above or below a certain threshold (e.g., water stress to plants) (Ghannam et al., 2016). In general terms, the notion of persistence in a stochastic field $\phi(x,t)$, oscillating around its ensemble mean $\langle \phi(x,t) \rangle$ under a given set of dynamics, is defined at a fixed point as the probability that the quantity $\text{sgn}[\phi(x,t) - \langle \phi(x,t) \rangle]$ does not change until time t (Ghannam et al., 2016; Perlekar et al., 2011). In the context of soil moisture dynamics, the ensemble mean can be replaced by a certain threshold, as mentioned above (Ghannam et al., 2016).

Although researchers have used the terms SMM and APSM interchangeably, they are not identical. Ghannam et al. (2016) examined the differences between SMM and APSM timescales (t_{SMM} and t_{APSM} , respectively) for root zone soil moisture. They made a clear distinction between t_{SMM} and t_{APSM} , characterizing t_{SMM} as an essentially quasi-deterministic timescale that is largely determined by evapotranspiration and drainage (water

losses from the soil column), and t_{APSM} as an inherently probabilistic scale that is primarily determined by precipitation and represents a distribution of periods when soil moisture is above or below a certain threshold. Ghannam et al. (2016) interpreted SMM and APSM as encoding different information about soil moisture dynamics in the root zone, making them relevant to different problems. For example, SMM is more relevant to land-atmosphere interaction schemes used in climate models because these schemes rely on SMM to improve their predictive ability for seasonal forecasts (Seneviratne, Koster, et al., 2006). However, as a measure of the strength of land-atmosphere coupling, APSM (an indicator of wet or dry conditions) may be more relevant than SMM (correlation timescale) because the wetness or dryness of the soil column largely controls surface energy fluxes (Ghannam et al., 2016). Several metrics have been introduced to quantify t_{APSM} , as listed in Supporting Information S1.

The term SMD refers to the quasi-exponential decrease in soil moisture immediately following the occurrence of precipitation (McColl, Wang, et al., 2017). During this period, Equation 1 can be rewritten as follows, neglecting drainage and runoff fluxes (McColl, Wang, et al., 2017):

$$\frac{d\theta}{dt} = -\frac{ET(\theta, t)}{\Delta z} = -\beta(\theta) \frac{E_0}{\Delta z} \quad (12)$$

where $\beta(\theta)$ is a dimensionless function equal to 1 for intermediate moist soils ($\theta_c < \theta < \theta_{\text{FC}}$) and defined as below for dry soils ($\theta_{\text{WP}} < \theta < \theta_c$):

$$\beta(\theta) = \frac{\theta(t) - \theta_{\text{WP}}}{\theta_c - \theta_{\text{WP}}} \quad (13)$$

where θ_{FC} and θ_{WP} are the soil moisture at field capacity and wilting point, respectively, and θ_c is the critical soil moisture beyond which soil moisture is not a limiting factor for evapotranspiration. McColl, Wang, et al. (2017) rearranged Equation 13 for dry soils to obtain the SMD timescale as follows:

$$-\frac{\theta(t) - \theta_{\text{WP}}}{\text{SMD}} = -\beta(\theta) \frac{E_0}{\Delta z} \rightarrow \text{SMD} = \frac{\Delta z(\theta(t) - \theta_{\text{WP}})}{E_0} \quad (14)$$

where SMD timescale is a measure of t_{SMM} . Comparing the formula for t_{SMM} given by Delworth and Manabe (1988) as $t_{\text{SMM}} = W_{\text{FC}}/E_0$, where $W_{\text{FC}} = \Delta z\theta_{\text{FC}}$, with the formula given by McColl, Wang, et al. (2017) in Equation 14, we can see that they are almost identical, differing only by the soil moisture level considered.

To quantify the SMD timescale, Shellito et al. (2016) and McColl, Wang, et al. (2017) first identified the individual drydowns in the soil moisture time series and then modeled them by fitting the following exponential model for each drydown:

$$\theta(t) = \Delta\theta \exp\left(-\frac{t}{\text{SMD}}\right) + \hat{\theta}_{\text{WP}} \quad (15)$$

where $\theta(t)$ is the soil moisture content (L^3L^{-3}) observed t days after the onset of desiccation, $\Delta\theta$ is the positive increase in soil moisture (L^3L^{-3}) preceding desiccation, $\hat{\theta}_{\text{WP}}$ is the effective wilting point (the estimated lower limit of soil moisture (L^3L^{-3}), which is likely to be less than the actual wilting point). Finally, the median of the estimated SMD for all drydowns is considered as the final estimate of SMD for the respective pixel/point.

Note that all current considerations assume that soil moisture dynamics are fully reversible. Hence, t_{SMM} is conceptually linked to concepts of resilience, which consider the return of a system to its original properties after an external perturbation.

3. The SMM Timescale and Its Temporal Variability

In general, the t_{SMM} is reported to be a couple of days to several months (from 1 month up to 12 months) (Amenu et al., 2005; Delworth & Manabe, 1988; Liu & Avissar, 1999; MacDonald & Huffman, 2004; McColl, Ale-mohammad, et al., 2017; McColl, Wang, et al., 2017; Orth & Seneviratne, 2012; Rowntree & Bolton, 1983;

Seneviratne et al., 2010; Simmonds & Hope, 1998; Walker & Rowntree, 1977; Yasunari, 2007; Yeh et al., 1984) or even more than 1 year (Amenu et al., 2005; Song et al., 2019; Stahle & Cleaveland, 1988), which is confirmed by both observational data (Entin et al., 2000; Ganeshi et al., 2023; Ghannam et al., 2016; Orth & Seneviratne, 2012; Orth et al., 2013; Seneviratne & Koster, 2012; Shinoda & Nandintsetseg, 2011; Vinnikov & Yeserkepova, 1991; Vinnikov et al., 1996) and model simulated data (Gao et al., 2018; Koster et al., 2000; Koster & Suarez, 2001; Koster et al., 2010; Schlosser & Milly, 2002; Seneviratne, Koster, et al., 2006; Seneviratne & Koster, 2012; Wu & Dickinson, 2004). This is also confirmed with both theoretical (calculation of W_f/E_0 ratio) and empirical (fitting Equations 2a and 2b to measured data) estimation methods (Vinnikov & Yeserkepova, 1991).

t_{SMM} varies in time. Delworth and Manabe (1988) highlighted that the seasonal cycle of potential evaporation at mid- and high latitudes results in shorter t_{SMM} in summer and longer t_{SMM} in winter. Entin et al. (2000) and Douville et al. (2007) confirmed the existence of such seasonal variations in t_{SMM} . Shinoda and Nandintsetseg (2011) found for the Mongolian steppe that t_{SMM} can last 5.5–8.2 months in autumn and winter, while spring and summer showed t_{SMM} of 1.5–3.0 months. In the forest-steppe zone, t_{SMM} was even longer in autumn and winter (6.0–7.0 months), but again longer than in spring and summer (3.0–1.8 months) (Nandintsetseg & Shinoda, 2014). Liu et al. (2014) confirmed that t_{SMM} lasted longer during spring (around 3.0–4.0 months) than during summer (around 2.0–3.0 months) and autumn (2.0 months) and this was especially the case in mid-latitudes. According to Dirmeyer et al. (2009), t_{SMM} is largest in wetter and/or colder seasons as well as in areas covered by snow or in dry regions. However, when comparing the t_{SMM} values from different studies, the frequency of soil moisture sampling must be considered. This is because—as we discuss in more detail later in Section 5—the longer intervals can naturally lead to longer t_{SMM} estimates.

However, the earlier work of Wu and Dickinson (2004) does not confirm the strong control of seasonality on t_{SMM} and argues that the mechanisms controlling its timescales are likely more complex. The authors considered four belts including equatorial, subtropical, midlatitude, and high latitude in the Northern Hemisphere and determined the belt-averaged autocorrelation coefficient profiles with depth (3.5 m deep) and across seasons; they found that t_{SMM} was not necessarily longer in winter than in summer as reported by, for example, Delworth and Manabe (1988). Contrary to previous reports, Orth and Seneviratne (2012) even found t_{SMM} in Europe to be weakest in spring and then increasing until fall. Based on these studies, both the timescale and seasonality of t_{SMM} seem to be site-specific and dependent on local hydrological settings. In this regard, Hagemann and Stacke (2015) reported that the simulated t_{SMM} in global climate models is generally elevated during the dry season when a soil moisture buffer exists below the root zone, but that t_{SMM} tends to be shortened where bare soil evaporation has increased; this is more common in semi-arid regions and wet seasons. In some areas, the increased evaporation can be offset by reduced transpiration which in turn also offsets the shortening of the t_{SMM} (Hagemann & Stacke, 2015). A conceptualization of the underlying mechanisms for these variable responses, however, is still lacking. Nevertheless, there appears to be an interaction between the t_{SMM} and the climatic regimes as well as the vegetation cover and local hydrological settings. Overall, it is difficult to isolate the individual effects of the several factors on t_{SMM} , given their potentially complex interaction. A good example of such a challenging interaction would be the effect of transpiration and evaporation on t_{SMM} , as the effects of an increase in one can be offset by a decrease in the other (Hagemann & Stacke, 2015).

4. Spatial Variability of SMM

t_{SMM} not only varies in time but also in space. On the global scale, Yeh et al. (1984) employed a model with idealized geography and found that the persistence of soil moisture anomalies depended significantly on latitude. Delworth and Manabe (1988) also confirmed a latitudinal dependence of soil moisture anomaly persistence, with the persistence increasing from tropical areas to high latitudes. The authors assume that this reflects an overall dependency of t_{SMM} on geographically varying climate parameters, yet, without going more into detail. They showed that the geographic dependence of the temporal variability of memory timescale is rooted in the spatial dependence of potential evaporation and soil field capacity. Physically, the lower the latitude, the greater the available radiation for evaporation and thus the greater the potential evaporation rate. As a result, soil moisture anomalies dissipate faster, and the memory timescale is shorter (Delworth & Manabe, 1988). However, we would like to point out that the incoming radiation (shortwave and longwave) at the surface is not only influenced by latitude, but also by cloud cover. Therefore, regions under subtropical high pressure may receive more shortwave radiation than tropical regions affected by, for example, the Intertropical Convergence Zone (ITCZ) with

persistent cloud cover. Accordingly, the reasoning of Delworth and Manabe (1988) for shorter t_{SMM} at lower latitudes may face some limitations. Liu and Avissar (1999) analyzed the spatial distribution of the memory timescale in the land–atmosphere system using simulated data. The authors found that soil moisture has strong persistence with 1-month autocorrelation coefficients of over 30% everywhere on Earth (an average of about 60% at the global scale). The authors confirmed that t_{SMM} increases at high latitudes and is intimately related to the extent of aridity in the regions. They found greater persistence (indicated by greater autocorrelations) and associated prolonged t_{SMM} in arid regions, where soil moisture variations are less severe and infrequent than in humid regions. They supported this result with observations from China.

McColl, Alemohammad, et al. (2017) concluded that consistently shorter t_{SMM} in the tropics is due to intense rainfall as well as rapid evapotranspiration and drainage fluxes. The authors explained that these short residence times in soil water reflect the rapid overturning of the terrestrial hydrologic cycle at the land surface, with, for example, most inflows from precipitation leaving the topsoil within 3 days. Conversely, the t_{SMM} was highest in mid-latitudes, particularly in northern Africa, parts of the Middle East, central Asia, and northern China as well as the western United States, because in these regions, the terrestrial hydrologic cycle is overturned only slowly at the land surface. The analysis was confirmed by Liu et al. (2014) who showed that land surface memory for soil moisture anomalies is longer in midlatitudes (ca. 2–3 months) and shorter in the Tropics (1.0–2.0 months). Similarly, Ruscica et al. (2014) report minimum t_{SMM} (0–5 days) over northern Uruguay, southern Brazil, and some points in Argentina and Paraguay where precipitation is persistent and high, while maximum t_{SMM} (30 days) occurred in northwestern areas of South America that experience low precipitation persistence.

Several studies analyzed the spatial variability in t_{SMM} for specific climate regions or continents. Asharaf and Ahrens (2013) examined the Indian summer monsoon season and showed that simulated memory lengths were longer in the western region than in the eastern region (14 and 9 days, respectively, at 34 cm soil layer depth), thus following the higher rainfall in the west than in the east. Also, the t_{SMM} increased with soil depth. MacLeod et al. (2016) reported that in general, memory increases with soil depth (and, thus, increasing mean residence time of soil water), though with significant spatial differences and depending on the start date of the modeling.

According to Orth and Seneviratne (2013) SMM serves as a kind of upper bound for the memory found in other hydrological processes like streamflow and evapotranspiration. The stronger the coupling between SMM and streamflow or evapotranspiration, the stronger their respective memory. The authors also found significant SMM in all examined catchments in Europe. The largest SMM, quantified by applying the interannual autocorrelation over a 30-day lag to daily data, was found in central Europe (Germany, eastern France), while it was low in mountainous regions (Alps, Massif Central, Scandinavian mountains).

Instead of a simple rationale for the latitudinal dependence of spatial variability in t_{SMM} , Orth et al. (2013) linked it to several factors by showing that t_{SMM} decreases with elevation and with increasing topography and aridity, with elevation being the most important, followed by topography and the aridity index.

He et al. (2023) found that the short-term memory SMM_7^S , as defined by McColl et al. (2019), lasted longer in arid regions (i.e., the Midwest of the United States and central Australia). In contrast, the long-term memory SMM_L^L is longer over wet areas. This is linked to the spatial distribution of soil hydraulic properties, allowing water from precipitation to drain rapidly into deeper soil in wet soils with higher hydraulic conductivities.

5. SMM and Soil-Plant-Atmosphere Interactions

In this section, we briefly present how soil moisture dynamics and therewith SMM impact processes in the soil-plant-atmosphere (SPA) system, resulting in feedback loops in which various processes influence SMM, and SMM, in turn, influences these processes. Figure 3 illustrates the processes involved in this feedback loop.

In general, the interactions between soil moisture and land surface processes can be considered from various angles, including water and energy balances, vegetation dynamics, climate feedback, and SPA interactions (Seneviratne et al., 2010). From the water balance equation, Equation 1, it is clear that available soil moisture is linked to the different components of the water balance equation which also affect atmosphere and land surface processes (Daly & Porporato, 2005; Ghannam et al., 2016; Katul et al., 2012; Seneviratne et al., 2010). Similarly, considering the soil energy balance equation, Equation 16 (Seneviratne et al., 2010); soil moisture affects the partitioning of net surface radiation into sensible heat, latent heat, and soil heat flux. Generally, outside of energy-




Figure 3. Representation of the effect of soil moisture memory (SMM) on processes involved in the coupling (black arrows) of land, plant, and atmosphere processes in the soil-plant-atmosphere system. The size of the red dots indicates those processes that are influenced by SMM and that are supported by previous research (indicated by a purple halo; the larger the halo, the more phenomena studied according to the number of references cited in Tables 1 and 2) or postulated by us and/or other researchers but not yet underpinned by findings documented in the literature (no halo). As an example, SMM can have an impact on precipitation through its effect on evapotranspiration and surface energy partitioning which is documented in literature. This may lead to changes that can then impact air circulation and cloud formation which then will finally impact precipitation (Yao et al., 2023). This feedback loop occurs when the soil that is excessively wet from a precipitation event continues to experience above-average evaporation in subsequent weeks, triggering additional precipitation (Koster et al., 2003). Conversely, a precipitation deficit can also trigger a feedback loop in which evaporation rates reduced by the lack of rain can further reduce subsequent precipitation (Koster et al., 2003). The lagged effects of soil moisture on evaporation have also been documented more recently (Rahmati et al., 2023a; Yao et al., 2023) which nicely fits into the memory concept of soil moisture feedback on evapotranspiration.

limited evaporation regimes, moist soils have a higher evaporation rate, resulting in higher latent heat flux and lower surface temperatures and therefore leading to a cooler surface (Humphrey et al., 2021). Conversely, dry soils result in higher sensible heat flux, higher surface temperatures, and a warmer land surface (Humphrey et al., 2021).

$$\frac{dH}{dt} = R_n(t) - \lambda ET(t) - SH(t) - G(t) \quad (16)$$

where dH/dt is the energy change within the surface soil layer considered, t is time, $R_n(t)$ is the net radiation, λET is the latent heat flux, SH is the sensible heat flux, and G is the soil heat flux.

The feedback loop between soil moisture and soil water and energy balances (as shown in Figure 3) can well explain the emergence of SMM and its effects on various processes in the SPA system. However, an important consideration here is the strength of the coupling between soil, plant, and atmospheric processes. There are regions where the coupling is strong and others where it is weaker, which should be considered when dealing with SMM investigations. In this regard, the term “hot spots” designates specific terrestrial, where a strong coupling between soil moisture and the atmosphere exists (Koster et al., 2004). To identify such hot spots, we must consider the strength of the coupling between soil moisture and a given atmospheric variable (e.g., air

temperature, relative humidity, or vapor pressure deficit) in relation to all other boundary conditions that affect this variable (Koster et al., 2004). Many studies related to soil moisture-atmosphere coupling tend to focus on these areas (Barcellos et al., 2018; Bu et al., 2023; Giles et al., 2023; Sangelantoni et al., 2023; Yin et al., 2023). However, it is worth noting that the location of the hotspots also depends on the metric used to define them. For example, the hotspots of predictive skill do not coincide with the classical hotspots of land-atmosphere coupling (Koster et al., 2010, 2011, 2016).

Koster et al. (2004) considered the strength of coupling between soil moisture and precipitation—the coupling strength comes from a study with several models based on model ensemble statistics, as it is not a directly measurable variable and therefore could not be validated—and identified hot spots of soil moisture and atmosphere in the central Great Plains of North America, the Sahel, equatorial Africa, and India. Less intensive couplings between soil moisture and precipitation were found in South America, Central Asia, and China. The authors argued that the hot spots are in transition zones between dry and humid regions, which comprise regions where boundary layer moisture can trigger moist convection. In these regions, evaporation is high but still sensitive to soil moisture and, therefore, can transfer the effects of soil moisture to the atmosphere (precipitation). Wet regions in contrast feature evapotranspiration rates (and thus precipitation rates) that vary little with soil moisture, and in dry regions, the evapotranspiration rates, while sensitive to soil moisture, are too low to have a significant impact. The occurrence of hot spots in transition zones was later confirmed by Seneviratne et al. (2010), who showed that such a strong coupling between soil moisture and atmosphere prevails only in transition zones having both a strong dependence of evapotranspiration on soil moisture and large mean evapotranspiration.

Exploration of soil moisture and atmospheric hot spots has also focused on the coupling between soil moisture and air temperature (e.g., Dirmeyer, 2011b; Koster et al., 2006; Miralles et al., 2012). Such investigations have confirmed that the hot spots occur in transition climatic regions; they also tend to show that the coupling is a bit stronger than that between soil moisture and precipitation. However, several new hot spots were discovered (Mueller & Seneviratne, 2012) where a strong coupling of soil moisture and temperature was later confirmed by remote sensing data, albeit with some underestimations (Hirschi et al., 2014).

In the following subsections, we focus on the driving factors of t_{SMM} and then on the implications of SMM obtained from the literature.

5.1. Controlling Factors of SMM and t_{SMM}

In general, the SMM, and more specifically its timescale (t_{SMM}), is controlled by seasonal variations in the atmosphere and their coupling with soil moisture, as well as by the dependence of evaporation and runoff on soil moisture (Douville et al., 2007). However, there may be other controlling factors, such as variability in soil properties.

The following autocorrelation expression, originally introduced by Koster & Suarez (2001) and then improved by Seneviratne and Koster (2012), allows an examination of the factors influencing the autocorrelation of soil moisture and thus the t_{SMM} :

$$\rho(W_{n+1}, W_n) = \frac{\sigma_{W_n}(1 - \alpha_n) + \sigma_{\phi_n}\rho(W_n, \phi_n)}{\sqrt{\sigma_{W_n}^2(1 - \alpha_n)^2 + 2\sigma_{W_n}(1 - \alpha_n)\sigma_{\phi_n}\rho(W_n, \phi_n) + \sigma_{\phi_n}^2}} \quad (17)$$

where ρ , σ , and σ^2 represent autocorrelation, standard deviation, and variance, respectively, and w_n and w_{n+1} implies degrees of soil saturation at the period n and $n + 1$. ϕ_n is an atmospheric forcing term combining the net effects on the water balance (based on climatological E/R_{net} and Q/P ratios, where E is the total evaporation (i.e., transpiration, bare soil evaporation, interception loss), R_{net} is net radiation, Q is runoff, and P is precipitation) of the accumulated fluxes of precipitation and net radiation over the period n . The coefficient α_n combines the sensitivity of the total evaporation to soil moisture (specifically, c_n , where $E/R_{\text{net}} = c_n W + d_n$) and runoff sensitivity to soil moisture (specifically, a_n , where $Q/P = a_n W + b_n$) as follows:

$$\alpha_n = \frac{c_n \bar{R}_n}{C_s} + \frac{a_n \bar{P}_n}{C_s} \quad (18)$$

where C_s is the water storage capacity of the column, and \bar{R}_n and \bar{P}_n are the long-term mean values of accumulated net radiation and precipitation over period n , respectively. We should note that c_n , the sensitivity of total evaporation to soil moisture, will reflect whether an area is in a water-limited or energy-limited evaporative regime (being close to zero in the latter case). The coefficient a_n will vary with the character of the local topography, with steeper slopes, for example, more amenable to producing runoff for a given level of soil moisture. Soil texture and vegetation characteristics will also affect the local values of c_n and a_n .

According to the above expression, the t_{SMM} (Seneviratne & Koster, 2012) is controlled by five factors: (a) the variability of initial soil moisture, as reflected in σ_{W_n} , (b) the variability of the forcing (as reflected in σ_{ϕ_n}), (c) the correlation between the initial soil moisture and the forcing, as reflected in $\rho(W_n, \phi_n)$, (d) the sensitivity of total evaporation to soil moisture, as reflected in $\frac{c_n \bar{R}_n}{C_s}$, and (e) the sensitivity of runoff to soil moisture, as reflected in $\frac{a_n \bar{P}_n}{C_s}$.

Seneviratne and Koster (2012) interpreted the contribution of those five controls under two conditions: with and without feedback between soil moisture and the forcing variables. In the absence of any impact of soil moisture on either evapotranspiration, runoff, or atmospheric forcing, Equation 17 simplifies to a simple function of the relative variability of the initial soil moisture and the atmospheric forcing:

$$\rho = \frac{\kappa_1}{\sqrt{\kappa_1^2 + 1}} \quad (19)$$

where

$$\kappa_1 = \frac{\sigma_{W_n}}{\sigma_{\phi_n}} \quad (20)$$

Based on how σ_{W_n} and σ_{ϕ_n} compare to each other, three situations can be distinguished (Seneviratne & Koster, 2012): (a) $\sigma_{W_n} \ll \sigma_{\phi_n}$ and $\kappa_1 \ll 1$, which indicates low memory; (b) $\sigma_{W_n} \gg \sigma_{\phi_n}$ and $\kappa_1 \gg 1$, which indicates high memory; and (c) $\sigma_{W_n} \approx \sigma_{\phi_n}$ and $\kappa_1 \sim 1$ which indicates moderate memory. There is, so far, no direct coupling between soil moisture and its forcing formulated, but these simplifications already allow us to classify memory based on comparisons of variability. That is, the larger the (scaled) atmospheric variability relative to the initial soil moisture variability, the smaller the t_{SMM} will be.

When soil moisture does affect either the total evaporation or runoff, one can see that $\frac{c_n \bar{R}_n}{C_s}$ and $\frac{a_n \bar{P}_n}{C_s}$ decrease the t_{SMM} because, for a given level of forcing, these terms would act to decrease the distinction between different soil moisture levels (Seneviratne & Koster, 2012). A positive correlation between initial soil moisture and atmospheric forcing terms, $\rho(W_n, \phi_n)$, would act to increase the t_{SMM} (Seneviratne & Koster, 2012). Conversely, a negative $\rho(W_n, \phi_n)$ would decrease it (Seneviratne & Koster, 2012).

Although not directly mentioned by either Koster and Suarez (2001) or Seneviratne and Koster (2012), the above expressions indirectly relate the contribution of soil properties to t_{SMM} through the soil water storage capacity, that is, the C_s parameter. When C_s is large, it compensates for the negative contribution of both total evaporations, $c_n \bar{R}_n$, and runoff, $a_n \bar{P}_n$, to t_{SMM} . Conversely, a small C_s value will amplify these negative effects. Therefore, any change in C_s due to external or internal forces will affect the anomalies of soil moisture and thus the t_{SMM} . A change in C_s can be triggered, for instance, by changes in soil structure and soil particle arrangement, changes in soil organic matter content, and all related effects induced by changes in land use, climatic conditions (e.g., droughts), vegetation, soil microbial and faunal activity, or soil compaction.

When the overall literature is screened for factors that control t_{SMM} , we find eight factors: (a) atmospheric forcings, (b) anthropogenic activities, (c) soil hydrological forcings, (d) soil properties, (e) groundwater dynamics, (f) vegetation properties, (g) sampling frequency, and (h) data sources. These factors, listed in Table 1 (which serves to summarize the t_{SMM} control factors identified in the literature and their effects), are all represented, either directly or indirectly, in the autocorrelation representation, Equation 17. For example, vegetation affects evapotranspiration and runoff generation and can thus also contribute to changes in soil water storage, and

Table 1

List of Factors (Forcings, Properties, Observational Characteristics) That Impact Soil Moisture Memory (SMM) and Its Timescale (t_{SMM}) and Related Effects

Factors	Effect
Atmospheric forcings	<ol style="list-style-type: none"> 1. Potential Evapotranspiration: It contributes to the attenuation of soil moisture anomalies and plays an important role in shaping SMM (Delworth & Manabe, 1988; Rahman et al., 2015). The amount of radiant energy absorbed by the soil surface affects the length of t_{SMM} by affecting evapotranspiration (Yeh et al., 1984). 2. Precipitation: As one of the water sources in the system, it leads to positive soil moisture anomalies and its absence leads to negative soil moisture anomalies and by that shapes its memory (Delworth & Manabe, 1988; McColl, Alemohammad, et al., 2017; Rahman et al., 2015; Small & Papuga, 2002; Song et al., 2019; Yeh et al., 1984). Note that rainfall has an asymmetric effect on SMM (discussed further in item 4 below). 3. Snowmelt and soil freezing: Snowmelt acts as another source of water and from there impacts SMM (Delworth & Manabe, 1988; Shinoda, 2001). Winter soil freezing and low snow depth can preserve soil moisture anomalies from fall to next spring and extend t_{SMM} (Shinoda, 2001; Shinoda & Nandintsetseg, 2011). Areas with longer snowpack duration have longer t_{SMM} compared to regions with shorter snowpack duration (Delworth & Manabe, 1988). 4. Extreme events: Extreme events such as heavy rainfall, droughts, or temperature fluctuations have profound effects on the condition of the soil (Bao et al., 2023), as well as on soil water storage (Mahanama & Koster, 2003; Orth et al., 2013) and by that they can affect SMM. Both extremely dry and wet soils lead to long t_{SMM} (McColl, Wang, et al., 2017; Orth & Seneviratne, 2012) due to increases in soil moisture variability and correlation with precipitation (Orth & Seneviratne, 2012). However, drier conditions tend to have longer t_{SMM} compared to wet conditions (Rahman et al., 2015). The elongated t_{SMM} under dry conditions can be related to changes in physical soil properties that may make the soil more water-repellent, thereby prolonging a drought anomaly (Orth & Seneviratne, 2012). On the other hand, a greater increase in t_{SMM} under extremely dry conditions compared to extremely wet conditions is reasonable because dry periods can potentially be more extreme than wet periods (Orth & Seneviratne, 2012). Heavy rainfall, for its part, effectively acts as a reset button for memory.
Anthropogenic activities	<ol style="list-style-type: none"> 1. Deforestation: Forests play a critical role in regulating soil moisture and surface temperature by intercepting precipitation as well as the cooling effects due to its higher evapotranspiration (Hesslerová et al., 2019). Deforestation removes vegetation cover, disrupts soil moisture regulation (Guo et al., 2002), reduces infiltration, accelerates runoff (Peili & Wenhua, 2001), and potentially shortens t_{SMM} by reducing the soil's ability to retain moisture over time. 2. Land use change: This can lead to both lengthening and shortening of t_{SMM} depending on which land use change is imposed. However, a detailed investigation into this is missing. 3. Irrigation: Conceptually, irrigation can contribute to wet soil moisture anomalies that likely prolong t_{SMM} (Yeh et al., 1984). However, improper irrigation can lead to waterlogging and poor drainage (Gebrehiwot, 2018; Khalil et al., 2021) which can limit soil's ability to store water for future use by weakening the soil condition, thus potentially shortening the t_{SMM}. This requires further investigation in future. 4. Other activities: Human activities like urbanization, soil sealing, overgrazing, and accelerated soil erosion presumably impact soil dynamics (Feng et al., 2023) and therefore t_{SMM}, but research on this is lacking. For example, it is easy to assume that plowing can greatly alter the soil's ability to retain water and affect infiltration as it breaks up the surface crusting, or that changes in composition through the addition of organic matter and fertilizers can also alter the interaction between soil and water and thus affect SMM. To the best of our knowledge, however, such impacts still need to be researched.
Soil hydrological forcings	<ol style="list-style-type: none"> 1. Actual evapotranspiration: This is the main coupler between the atmosphere and soil (especially in transition zones) and is a key factor in controlling the storage of soil moisture and thus the extent of SMM (Bonan & Stillwell-Soller, 1998; Liu & Avissar, 1999; Wu & Dickinson, 2004). Higher actual evapotranspiration potentially leads to shorter t_{SMM} (Liu & Avissar, 1999). 2. Runoff and drainage: It attenuates soil moisture anomalies (mostly in wet regions) and shortens the duration of positive anomalies, thus decreasing t_{SMM} (Delworth & Manabe, 1988; Yeh et al., 1984), more possibly the short-term t_{SMM}. 3. Variations in the antecedent soil moisture: It, as an indicator of abnormal conditions, contributes to t_{SMM} (Song et al., 2019). Dry anomalies decay more slowly than moist anomalies under similar atmospheric conditions and thus potentially result in a longer t_{SMM} (Song et al., 2019).
Soil properties	<ol style="list-style-type: none"> 1. Soil water storage: Soil water storage is an important controlling factor of SMM as it affects the impacts of evapotranspiration and runoff (Orth & Seneviratne, 2012; Seneviratne, Koster, et al., 2006). 2. Soil field capacity ($n(\Delta z)$), porosity (n), and depth (Δz): The lower the field capacity, the shorter the t_{SMM} (Delworth & Manabe, 1988; Orth et al., 2013; Yeh et al., 1984). As field capacity is used directly in the autocorrelation expression of soil moisture (Koster & Suarez, 2001; Seneviratne & Koster, 2012), it can be a good candidate for studying the effects of other soil properties on SMM. The t_{SMM} increases with greater soil depth (Amenu et al., 2005; Asharaf & Ahrens, 2013; Douville et al., 2007; He et al., 2023; MacDonald & Huffman, 2004; Martínez-Fernández et al., 2021; Ruscica et al., 2014; Song et al., 2019; Wu et al., 2002), as deeper layers exhibit higher organic and clay contents (Martínez-Fernández et al., 2021), larger magnitudes of soil moisture spectra (Asharaf & Ahrens, 2013), and slower drying times after precipitation events. 3. Soil particle-size distribution: Although the effect of soil separates (specifically sand content) on SMM (directly and indirectly) is evaluated through several recent investigations (Akbar et al., 2018; Groh et al., 2020; Shellito et al., 2018) and no clear conclusion has been made yet, it seems that coarse-textured soils (sandy soils) exhibit shorter t_{SMM} due to easier water release via evapotranspiration and drainage (Martínez-Fernández et al., 2021; McColl, Wang, et al., 2017). However, some research contradicts this (McColl, Alemohammad, et al., 2017).

Table 1
Continued

Factors	Effect
	<ol style="list-style-type: none"> 4. Soil structure and pore system: Although there is no direct link between SMM and the soil structure and pore system, it has been postulated that larger pores with lower suction can lead to faster attenuation of water from soil system (McColl, Wang, et al., 2017) and therefore can potentially engender in shorter t_{SMM}. Since soil structure also directly affects the soil pore system, we postulate that it is also a key controller of SMM. 5. Organic matter content: Higher organic matter content is associated with increased water retention capacity and thus longer t_{SMM} (Martínez-Fernández et al., 2021). 6. Soil bulk density: Although bulk density indirectly reflects soil porosity, which affects water holding capacity and thus SMM (Koster & Suarez, 2001; Seneviratne & Koster, 2012), no significant effect of soil bulk density on SMM has been reported (Martínez-Fernández et al., 2021).
Groundwater dynamics	<p>Although more research is needed here, shallow groundwater tables can significantly affect soil moisture behavior by altering the dependence of soil moisture on precipitation and decoupling it from the atmosphere, which in turn affects SMM (Martínez-de la Torre & Miguez-Macho, 2019). It is also the case that groundwater contributes to evapotranspiration (Hou et al., 2023) and from there can contribute to SMM. However, the range in which groundwater contributes to evapotranspiration through capillary rise strongly depends on the soil hydraulic properties (Groh et al., 2016; Soyulu et al., 2011). On the other hand, it is argued that SMM has the potential to contribute to climate prediction on multi-year time scales by using information stored in slowly changing components of the soil system such as groundwater (Bellucci et al., 2015; Bierkens & van den Hurk, 2007; Fan & Miguez-Macho, 2010; Langford et al., 2014). Although not explicitly mentioned, this implies that groundwater, as part of the soil water storage, has a clear role in shaping SMM. However, the full extent of groundwater's influence on SMM and from there on climate predictability has yet to be fully assessed due to challenges related to long-term measurements, limited spatial representation, and current limitations of LSMs (Song et al., 2019).</p>
Vegetation properties	<ol style="list-style-type: none"> 1. Land cover: Forested areas have higher transpiration rates and often buffer soil moisture variations and exhibit weaker memory compared to nearby grasslands (Orth & Seneviratne, 2012), indicating that land cover affects SMM dynamics (Laio et al., 2001; Porporato et al., 2001; Ruscica et al., 2014; Teuling et al., 2006). Some others (McColl, Wang, et al., 2017; Small & Papuga, 2002) challenge that the existence of a clear relationship between land cover type and SMM. 2. Vegetation density: If the external forcings are strong, denser vegetation (forest) tends to have longer t_{SMM} and slower recovery from anomalies while a weakening of external forcing can lead to a longer t_{SMM} in grassland and deserts (Wei et al., 2006). 3. Soil-atmosphere coupling: Vegetation affects SMM by influencing precipitation and the coupling between the soil and atmosphere. Vegetation-rich areas (forests) can enhance rainfall due to increased evapotranspiration (Spracklen et al., 2012). Vegetation dynamics also influence the condensation of water vapor and atmospheric pressure in the lower atmosphere (Makarieva & Gorshkov, 2007; Makarieva et al., 2013). 4. Root structure: Root structure can affect the relationship between soil moisture and evapotranspiration under anomalous conditions and thus can affect t_{SMM} (Entin et al., 2000). Vegetation types with shallower root systems can be more sensitive to atmospheric forcings (Rahmati et al., 2023a), possibly resulting in shorter t_{SMM}.
Topography and subsurface geology	<p>Research shows a clear link between t_{SMM} and soil properties such as karst distribution (Dirmeyer & Norton, 2018), as karst formation can lead to a shortened t_{SMM} due to its contribution to preferential water flow in the soil. This can clearly have implications for modeling distributed land surfaces, as preferential water fluxes through geologic formations are absent from typical modeling frameworks (Dirmeyer & Norton, 2018).</p> <p>Regarding the effect of topography on the SMM, it is worth noting that the contribution of red and white noise to the variance of soil moisture and thus to the t_{SMM}—as linked in Equations 3 and 4—can vary with topographic conditions and land surface characteristics (Vinnikov et al., 1996). On flat, homogeneous plots, the white noise component contributes less to the variance of soil moisture and represents only random measurement errors. In natural landscapes with different vegetation and soil types and complex topography, however, the contribution of white noise to the variance of soil moisture is much greater (Vinnikov et al., 1996). One can also view the topographic dependence of SMM through the reported latitudinal dependency (Delworth & Manabe, 1988; Liu & Avissar, 1999; Yeh et al., 1984)—as discussed more in detail in Section 4. However, a clearer relationship between topographic features (e.g., slope and aspect) and SMM has yet to be explored.</p>
Sampling frequency	<p>A higher sampling frequency of soil moisture data enables the detection of rapid changes in soil moisture and ensures that short-term fluctuations are not overlooked in the calculation of t_{SMM}. Conversely, lower soil moisture sampling frequency reduces the likelihood that rapid drying of soil moisture is captured, which can lead to an overestimation of the memory timescales (Martínez-Fernández et al., 2021; McColl, Alemohammad, et al., 2017; McColl, Wang, et al., 2017). The effect of sampling frequency on t_{SMM} can be understood with the help of signal processing, where the irregular sampling acts like a low-pass filter on the data and smooths out rapid fluctuations. Investigating the relationship between sampling frequency and the dissipation rate of the soil moisture anomaly could be a valuable starting point for further methodological investigations to understand and quantify SMM dynamics. This would increase the clarity of our understanding and highlight an important methodological research need in this area.</p>
Data sources	<ol style="list-style-type: none"> 1. In-situ data: In-situ data provide valuable insight into SMM (Entin et al., 2000; Koster & Suarez, 2001; Martínez-Fernández et al., 2021; Seneviratne, Koster, et al., 2006; Seneviratne & Koster, 2012; Shellito et al., 2016; Vinnikov & Yesserkepova, 1991), but the lack of global coverage, sampled soil volume, areal representativeness issues, and uncertainty in

Table 1
Continued

Factors	Effect
	global soil databases must be carefully considered (McColl et al., 2019). The consequences of in-situ for validation of model grid cell t_{SMM} is examined in Dirmeyer et al. (2016) as t_{SMM} is sensitive to aggregation (averaging over a spatial scale).
2.	Model simulations and uncertainty: Model simulations offer alternative approaches but are subject to uncertainty due to the impacts of model-specific parameterizations – different models will provide different estimates of t_{SMM} (Delworth & Manabe, 1988; Liang & Yuan, 2021; Rind, 1982; Rowntree & Bolton, 1983; Yeh et al., 1984). Dirmeyer et al. (2016) argue that although models capture well the large-scale variability of soil moisture across climate regimes, they fail to reproduce the observed patterns at smaller scales (hundreds of kilometers or less). One could also pay attention to the differences between the results of “offline” land models driven by atmospheric data and coupled land-atmosphere models (which include feedback), as only the latter allow the feedback itself to imprint itself on the memory.
3.	Space-based observations: Spaceborne soil moisture data are also used for quantitative analysis of t_{SMM} (McColl, Alemohammad, et al., 2017). However, satellite-derived soil moisture data may exhibit faster drying processes, potentially leading to shorter t_{SMM} compared to in-situ measurements (Champagne et al., 2016; Chan et al., 2016; Rondinelli et al., 2015; Shellito et al., 2016). Differences in spatial resolution and penetration depth between satellite and in-situ observations can contribute to these discrepancies (Dai et al., 2019; Jackson et al., 2016; Martínez-Fernández et al., 2021; Owe and Van de Griend, 1998).

the sampling frequency can affect the length of the quantification period. Jacobs et al. (2020) showed that stochastic rainfall plays a crucial role in memory and persistence of regional soil moisture. The frequency of rainfall was identified as the primary factor determining persistence across the region, while variations in land cover and soil properties had a secondary impact.

5.2. Implications of SMM

In this section, we explore the effects of SMM on different land surface processes. The reviewed literature shows that SMM has implications for weather variations and forecasts, land surface energy balances, monitoring and forecasting of droughts, floods, and heat waves, water use efficiency, biogeochemical cycles, groundwater predictions, and climate phenomena. Table 2 summarizes the processes, events and phenomena controlled by SMM and the potential impacts identified in the literature.

6. SMM Representation by Models

An accurate representation of SMM by LSMs requires a reliable parameterization of evapotranspiration and its dependence on soil moisture (Daly & Porporato, 2005; Seneviratne et al., 2010). Evapotranspiration is coupled to energy, water, and carbon balance processes (Daly & Porporato, 2005), and plays a crucial role in determining the intensity of the greening-induced boundary forcing (Zeng et al., 2016). In the so-called hotspot regions, soil moisture variability is the most important controlling factor of evapotranspiration variability (Koster et al., 2004; Seneviratne et al., 2010).

One can argue that accurate parameterization is also important for other processes. Such is the case of drainage and runoff, which are as important as evapotranspiration for determining SMM given the comparable roles that these play in the determination of the evolution of soil moisture in the land system (Entekhabi & Rodriguez-Iturbe, 1994; Koster & Milly, 1997; McColl et al. (2019). Additional aspects of land surface behavior, such as microbial moisture response curves used in the carbon cycle, may also prove important in the simulation of SMM. Here, for the sake of brevity, we focus on the evapotranspiration component. In the hybrid stochastic-deterministic model of McColl et al. (2019), long-term memory of soil moisture is controlled by stage-II evapotranspiration and short-term memory by drainage and runoff. However, the control of drainage and runoff on SMM is of short duration, typically a few hours, whereas the control of evapotranspiration is much longer, typically from a few days to several months. On the other hand, drainage- and runoff-induced SMM are neglected due to the lower resolution of the data (in many cases daily and higher), especially at large scale investigations. For this reason, we prefer to focus on evapotranspiration parameterization only rather than drainage and runoff.

Over time, the representation of the interrelationship between evapotranspiration and soil moisture in the field of climate modeling has evolved considerably through improved understanding of relevant complex processes and

Table 2
List of Processes, Events, and Phenomena Controlled by Soil Moisture Memory (SMM) and the Corresponding Impact

Processes, events, phenomena	Effect
Atmospheric and Climate Variability and Predictability	In cases of strong land-atmosphere coupling, weather conditions can be influenced by SMM, resulting in significant implications for seasonal and long-term forecasts (Douveille & Chauvin, 2000; Douville, 2004; Koster et al., 2010; Mahanama & Koster, 2003; Martínez-de la Torre & Miguez-Macho, 2019; Namias, 1959, 1963; Nicolai-Shaw et al., 2016; Notaro, 2008; Orłowsky & Seneviratne, 2010; Ruscica et al., 2014; Wang et al., 2014). Such a role can be twofold: (a) direct effects on energy and water budgets, influencing a range of extremes, and (b) the memory aspect that translates to persistence in atmospheric and land hydrology variables. Soil moisture serves as a repository of anomalies within the water budget of the land surface, and from there, through SMM, it exerts a lasting impact on the atmosphere above, primarily through the exchange of heat and moisture via land surface fluxes (Shinoda & Yamaguchi, 2003). SMM apparently affects climate and atmospheric variability (Delworth & Manabe, 1988). In fact, SMM has a possible impact on surface air temperature, surface pressure, and precipitation (Alfieri et al., 2008; Koster et al., 2003; Liu et al., 2014), especially in tropics and extratropics (Shukla & Mintz, 1982). Such an impact is also confirmed over Africa ⁵⁷ , the Sahel (Douveille et al., 2007), and Europe (Rowntree & Bolton, 1983). The effects of SMM on local rainfall are also well-documented—the higher the persistence of wet anomalies, the higher the local rainfall amount in the following period (Pal & Eltahir, 2001; Rind, 1982; Rowntree & Bolton, 1983; Shukla & Mintz, 1982). Such an impact can also occur non-locally in adjacent areas through teleconnections (Pal & Eltahir, 2002, 2003).
Land surface energy balance	<ol style="list-style-type: none"> 1. Surface heat balance: Variations in soil moisture impact the partitioning of outgoing heat fluxes into latent and sensible heat fluxes (Delworth & Manabe, 1988; Ganeshi et al., 2023; Yeh et al., 1984). Increased soil moisture enhances latent heat flux and reduces sensible heat flux, regulating energy exchange at the land surface and affecting surface air temperature variability (Amenu et al., 2005; Yeh et al., 1984). The heat capacity of the soil varies with soil moisture, so that soil moisture memory affects the memory for heat storage, which in turn has its own effects on the distribution of net surface radiation between sensible, latent and soil heat fluxes (Xue et al., 2021). 2. Surface temperature: Moist soil dissipates excess radiation through latent heat fluxes, keeping the soil cool. Dry or vegetation-less soil absorbs excess energy, gradually warming and dissipating it through sensible heat fluxes, impacting the thermal state of the surrounding atmosphere (Rind, 1982). 3. Atmospheric circulation: Soil moisture anomalies affect the thermal state of the atmosphere and overall atmospheric circulation (Koster et al., 2016; Miralles et al., 2019; van den Hurk et al., 2009; Wey et al., 2015; Yeh et al., 1984).
Drought events	<p>Drought is a multi-faceted phenomenon and accordingly has many definitions, depending in part on the impact considered. Agricultural drought (or soil moisture drought) is commonly tied to soil moisture deficits and is thus tied strongly to SMM. Due to the current length of the paper, we do not plan to elaborate on different drought definitions here; the discussion in this table focuses on soil moisture drought.</p> <ol style="list-style-type: none"> 1. Drought predictions: Soils characterized by extensive dry anomaly persistence (as an indicator of SMM) are frequently affected by prolonged and persistent droughts (Abolafia-Rosenzweig et al., 2023; Entekhabi et al., 1992; Soulsby et al., 2021); although extensive wet anomaly persistence can also mitigate the effects of droughts (Stahle & Cleaveland, 1988; Tjeldeman & Menzel, 2021). In this context, SMM, in conjunction with land-atmosphere interactions, can possibly improve the ability to predict drought (more specifically soil moisture drought) on seasonal to decadal timescales by converting a weak precipitation signal into a more predictable soil moisture signal (Esit et al., 2021). 2. Resilience against droughts: Elevated SMM makes soils resistant to drought events or can prolong soil moisture drought, influencing the severity and impact of droughts (Nicholson, 2000; Rahmati et al., 2023c). Local meteorological conditions and the presence of sufficient storage capacity in the root zone can prevent soil moisture drought even during severe drought years (Tjeldeman & Menzel, 2021). 3. Predicting flash droughts: Manipulating initial soil moisture anomalies in forecasting models enables accurate simulation of flash drought (Liang & Yuan, 2021), which are characterized by rapid intensification and severe impacts (Otkin et al., 2018; Yuan et al., 2018). 4. Influence on climate extremes: SMM impacts climate extremes by modulating droughts and influencing hot and cold extremes (Liu et al., 2014). Dry anomalies in soil moisture contribute to the maintenance of drought conditions over time (Hong & Kalnay, 2000), leading to prolonged and intensified drought events.
Flood events	<ol style="list-style-type: none"> 1. Runoff predictability and flood forecasting: Variability and uncertainty in SMM significantly affect runoff predictability and flood forecasting as they play a role in precipitation and runoff generation as well as evapotranspiration (MacLeod et al., 2016; Orth & Seneviratne, 2013). It has been shown that delayed extreme soil wetness in spring can delay the annual peak runoff, which has great implications for flood monitoring and management (Xu et al., 2021). 2. Flood duration and intensity: Persistence in wet soil moisture anomalies (which can be read as lengthened t_{SMM}) in flood-prone regions can contribute to prolonged flooding of greater intensity (Bonan & Stillwell-Soller, 1998; Liu et al., 2014; Pal & Eltahir, 2002).
Heatwave events	<ol style="list-style-type: none"> 1. Heatwave occurrence: SMM has implications for the occurrence of heatwaves (Difffenbaugh et al., 2007; Fischer, Seneviratne, Lüthi, & Schär, 2007; Fischer, Seneviratne, Vidale, et al., 2007; Haarsma et al., 2009; Hirschi et al., 2011; Jaeger & Seneviratne, 2011; Seneviratne, Lüthi, et al., 2006; Vautard et al., 2007). For example, spring soil moisture anomalies can persist into the summer season, altering heat fluxes and significantly affecting the occurrence of hot days and heatwaves (Wu & Zhang, 2015).

Table 2
Continued

Processes, events, phenomena	Effect
	<ol style="list-style-type: none"> Heatwave predictability: Soil moisture conditions in spring can serve as useful predictors for summer heat extremes (Miralles et al., 2014; Quesada et al., 2012; Wu & Zhang, 2015) as it can alter latent and sensible heat fluxes (Wu & Zhang, 2015). Benson and Dirmeyer (2023) also linked heatwave prediction skills at longer lead times to SMM in NOAA forecast models. Heatwave duration and intensity: The persistence of heatwaves can be influenced by SMM (Lorenz et al., 2010). Simulations with interactive soil moisture (with memory) exhibit higher heatwave persistence compared to simulations with fixed or preset soil moisture (without memory) (Lorenz et al., 2010). Anomalies of soil moisture can also act as an amplifying/dampening factor for heatwaves (Lorenz et al., 2010).
Wildfire events	The long-term memory stored in deep soil moisture and groundwater, spanning multiple seasons to multiple years, plays a role in predicting hydroclimate features like wildfire at seasonal to decadal timescales (Esit et al., 2021). Wild fire events affect soil properties, for example, alter the soil water storage capacity (Agbeshie et al., 2022) as well as vegetation properties (Lloret & Zedler, 2009; Verma et al., 2017), which may also impacts SMM.
Water use efficiency and plant physiological responses	Dry anomalies of soil moisture and their persistence have a 1- to 12-month (depending on vegetation type and region) lagged effect on Water use efficiency in terrestrial ecosystems showing both negative and positive impacts depending on vegetation type (Ji et al., 2021). Vidale et al. (2021), assessing the impact of the relationship between soil moisture stress and vegetation function using the JULES LSM model, showed that the past soil moisture deficit observed in April 2003 leads to a suppression of vegetation growth primary production (GPP) during the following months of June, July, and August. A wealth of literature linking soil moisture to vegetation health and productivity is indeed available; a full survey is beyond the scope of this review.
Biogeochemical processes	<ol style="list-style-type: none"> Carbon source and sink: Soil moisture anomalies are the main cause for most of the interannual variation in global carbon uptake mainly through their impact on photosynthesis (Green et al., 2019; Humphrey et al., 2021). This is mainly due to the amplification of temperature and vapor pressure deficit anomalies (in semi-arid and tropical regions) and the amplification of the direct effects of soil water stress (in temperate and tropical biomes) through the soil moisture–atmosphere coupling (Green et al., 2019; Humphrey et al., 2021). In fact, dry anomalies of soil moisture can lead to vegetation stomatal closure and reduce photosynthesis and consequently can lead to decreased land uptake of carbon dioxide (CO₂) (Green et al., 2019). Carbon decomposition and microbial responses: SMM can influence microbial responses in the carbon cycle. Soils with wetter climate histories exhibit higher respiration rates (probably higher decomposition rate of organic carbon) compared to soils from drier areas, indicating the importance of considering SMM in understanding microbial responses and carbon dynamics (Evans et al., 2022; Hawkes et al., 2017). Nitrous oxide emissions: Anomalous soil moisture conditions affect the production and consumption of nitrous oxide (N₂O), a potent greenhouse gas. Soil moisture variations influence the balance between N₂O and N₂ emissions and impact the availability of oxygen in the soil. Excessive soil moisture can lead to oxygen deficiency, promoting anaerobic conditions that encourage denitrification and higher N₂O emissions (Rubol, 2010).
Groundwater	Like feedback loop between SMM and other forcings (e.g., precipitation, evapotranspiration, and runoff), a feedback loop may also exist between SMM and groundwater, and thus SMM can be expected to impact groundwater. However, the reasons limiting research on the full extent of groundwater influence on SMM (Song et al., 2019) may also be the reason for the lack of research on SMM impacts on groundwater. There is indeed a body of literature that establishes a connection between soil moisture and groundwater; a detailed overview of the relationship between soil moisture and groundwater is beyond the scope of this paper.
Global climatic phenomena	<ol style="list-style-type: none"> Climate-ENSO connection: Evidence shows that soil moisture crucially impacts the El Niño-Southern Oscillation (ENSO)-based statistical seasonal forecasting (Amenu et al., 2005; Timbal et al., 2002). For example, it is shown that the SMM can persist the in-phase relationship between Southern Oscillation Index (SOI) and precipitation and can be critical for the lagged relationship between SOI and surface temperature (Timbal et al., 2002). West African monsoon: SMM contributes to the spatial extent and temporal evolution of soil moisture anomalies in the West African monsoon region, influencing the annual cycle and inter-seasonal persistence of water and heat fluxes between the surface and atmosphere (Fontaine et al., 2007). However, other studies argue that such a soil moisture/monsoon regulation mechanism does not exist, as soil moisture does not act as a memory for rainfall anomalies (Douville et al., 2007; Shinoda & Yamaguchi, 2003; Xue et al., 2012). Similarly, the analyses of van den Hurk and van Meijgaard (2010) also shows that atmospheric mechanisms strongly limit the influence of soil moisture on the regional water cycle. Monsoon rainfall predictability: SMM influences monsoon rainfall predictability through a positive feedback loop between soil moisture and rainfall (Douville et al., 2007; Yasunari, 2007). However, it seems that SMM diminishes rapidly during dry seasons and does not provide a significant contribution to monsoon rainfall predictability in summer (Douville et al., 2007). Meiyu event in East Asian summer monsoon: It has been shown (Dong et al., 2023) that SMM allowed the negative soil moisture anomalies in May 2020 over the Indo-China Peninsula (which is subjected to increased surface temperature and sensible heat flux) to persist into the Meiyu period (characterized by heavy rainfall) during the East Asian summer monsoon in 2020, leading to the manifestation of the Super Meiyu Event.

Table 3

Modeling Aspects of Soil Moisture (SM)—Evapotranspiration (ET) Relationship in First to Third Generations of Land Surface Models (LSMs)

Models	Modeling aspects and possible drawbacks
First-generation LSMs: bucket-type parameterization (Sellers et al., 1997; Seneviratne et al., 2010)	<ul style="list-style-type: none"> • Simple parametrization of ET and SM. • Typically employing two thresholds (namely critical SM and the wilting point), where ET is unrestricted until the SM falls below critical SM, beyond which ET will linearly decrease by a further decrease in SM and reach zero when SM falls below the wilting point. • Not accurately capturing trends in SMM because: <ul style="list-style-type: none"> ◦ They tend to overestimate ET relative to other land surface systems. This is primarily because they overlook additional factors besides soil moisture that limit plant transpiration. ◦ They typically consider only a single soil store and fail to account for interception storage and spatial variations in soil and vegetation parameters, and they provide an oversimplified representation of runoff formation, temperature conduction, and soil freezing.
Second-generation LSMs: biophysical models (Sellers et al., 1997; Seneviratne et al., 2010)	<ul style="list-style-type: none"> • Incorporate more detailed representations of land surface processes. • Employ soil moisture models that consider the actual water content of the soil, rather than relying only on fixed thresholds. • Simulate a gradual decrease in ET as SM decreases. • Include a clearly defined upper layer of the canopy, soil with multiple layers, and the incorporation of key physical phenomena occurring within the plant canopy and soil. • Higher ability to regulate ET through stomatal resistance, considering the physiological factors involved. • Evaporation can originate from four distinct sources: potential evaporation from the interception layer, evaporation from exposed soil, transpiration from vegetation, and snow sublimation. • Vegetation cover can draw water from the deep root zone for transpiration, contributing to long-term climate memory. <ul style="list-style-type: none"> ◦ Better representation of SMM compared to bucket models, because they distinguish between soil and root zone evapotranspiration, which are separate moisture reservoirs with different memory characteristics and corresponding effects on surface fluxes. ◦ They include geographic detail regarding variations in soil and vegetation parameters, particularly factors such as water-holding capacity and rooting depth, which contribute to improved model representation despite some uncertainty regarding their specification. ◦ They include the interception reservoir that allows for fast evaporation which is significant in different regions around the world.
Third generation LSMs: physiological models (Fisher & Koven, 2020; Seneviratne et al., 2010)	<ul style="list-style-type: none"> • Further refined representation of the interactions between ET and SM. • More advanced land surface schemes that included multiple soil layers to capture vertical variability in SM. • Including explicit parameterizations to account for the effects of soil texture, vegetation type, and root distribution on ET. • Incorporate various aspects of plant photosynthesis, such as carbon assimilation and nutrient uptake, enzyme kinetics, electron transport, and the absorption of light by chloroplasts in plant leaves. • Including the feedback mechanisms between SM and the atmosphere allows for a more dynamic representation of the ET process. • Considering the potential effects of CO₂ concentrations on plant water use efficiency and, consequently, changes in the relationship between SM and ET under elevated CO₂. • Using the biophysical responses of plants to increase CO₂ levels to mitigate the effects of climate change, including drought and wildfires, although these biophysical responses can be affected by nutrient limitations that inhibit plant growth, which means that this interaction is not adequately accounted for, and the memory effect may not be fully represented.

the advent of unprecedented computational capabilities (Seneviratne et al., 2010). In fact, the different generations of climate models have developed increasingly sophisticated approaches to capture this relationship. Table 3 summarizes such representations (along with their possible advancements and drawbacks) in the first through third generation of LSMs—the evolution and progress of models characterized by their complexity, capabilities, and the scientific understanding they incorporate (Sellers et al., 1997). Here, only the current state-of-the-art climate models, and how SMM is represented by LSMs will be addressed in detail. The newest generations of LSMs (fourth generation onwards, including dynamic vegetation and vegetation demographics) see improvements in the representation of key hydrological processes (Zeng et al., 2016) such as the movement of water through the soil profile, surface runoff, groundwater recharge, and the treatment of subgrid-scale soil moisture variability. In parallel, the inclusion of complex feedback between the land surface and the atmosphere allows for a more realistic representation of the hydrologic cycle (Zeng et al., 2016). For example, LSMs can now mimic the so-called greening of the Earth (Mahowald et al., 2015) in which leaf area index (LAI) and stomatal conductance

increase, thus affecting evapotranspiration rates. Despite such progress, it is unclear whether the overestimation of key features of evaporative drought (also known as Edrought, which includes regularly occurring dry seasons and abnormal dry periods) undermines the ability of models to simulate realistic drought responses to climate change, which has broader implications, for example, in the study of heatwaves (Ukkola et al., 2016). There are also concerns over the sensitivity of LSMs to changes in atmospheric and hydrologic factors (including soil moisture availability) when characterizing global variability in soil carbon uptake (Humphrey et al., 2021). Additional uncertainties in mean surface temperature and variability, probably related to the coupling between evapotranspiration and soil moisture in different models, have been reported (Berg & Sheffield, 2018, 2019). Further advancement in Earth system forecasting models is required. Several research pathways have been suggested such as the combination of models and data for Earth system forecasting to better capture the interconnected systems of our planet (Gettelman et al., 2022).

Rind (1982) was among the first to investigate the importance of soil moisture anomalies in model predictions, who investigated the influence of SMM on summertime model predictability over North America. He showed that a reduction in early summer soil moisture resulted in a significantly higher surface air temperature and lower precipitation and cloud cover during summertime. The same methodology, albeit with different applications, has been used in several studies to date (Georgescu et al., 2003; Liang & Yuan, 2021; Zhao et al., 2019) and many have investigated SMM by integrating observations with LSMs and atmospheric general circulation models (GCMs), such as Dirmeyer, (1999, 2000); Dirmeyer and Brubaker (2007); Dirmeyer et al. (2009); Douville et al. (2001); Douville (2002) and Koster et al. (2009) among the others (see the comprehensive review by Dirmeyer (2011a)).

These studies have generally focused on regional to global scales (Seneviratne et al., 2013; Tjeldeman & Menzel, 2021; Wu & Dickinson, 2004). For example, Rowntree and Bolton (1983) assessed the importance of initial soil moisture anomalies to short-term changes in climate and hydrology. Also, Yeh et al. (1984) examined the latitudinal dependence of climatic and hydrologic response to soil moisture anomalies caused by large-scale irrigation. Delworth and Manabe (1988) examined the effects of soil moisture variability on the atmosphere by performing a long-term GCM integration, manipulating the boundary conditions and the hydrologic interaction between the atmosphere and the land surface. Mahanama and Koster (2003) contrasted the memory behavior of two LSMs and found that the differences between the models were related to differences in water holding capacity and ET and runoff parameterizations. Other similar studies showed the dependency between the initial wet or dry conditions and the subsequent model predictions (Sörensson & Berbery, 2015), which points to the need for detailed land-surface representations when modeling certain particular regions.

Despite the potential of these methods, generalized conclusions may be model-dependent due to the varying complexity of different models (Asharaf & Ahrens, 2013; Seneviratne, Koster, et al., 2006; Song et al., 2019). This was first investigated by Seneviratne, Koster, et al. (2006) who found, among relatively similar global SMM patterns, local differences between model results due to different water-holding capacity or biases in radiation forcing. Other studies have since compared SMM across models because SMM can be used to characterize the temporal variability of soil moisture and serve as a proxy for assessing land-atmosphere flux exchange in LSMs (He et al., 2023). For instance, SMM during dry periods can be greater when a multi-layer soil moisture scheme is used in place of a single layer (Hagemann & Stacke, 2015). Similarly, t_{SMM} can increase with increasing soil depth (Asharaf & Ahrens, 2013). Further, LSMs generally simplify or ignore lateral flow or groundwater table fluctuations, resulting in non-realistic spatial distributions of groundwater that affect SMM predictions (Martínez-de la Torre & Miguez-Macho, 2019).

The uncertainty of model outputs and parameterization schemes has also been investigated. For example, in their global sensitivity analysis, MacLeod et al. (2016) argued that the dependence of SMM uncertainties on the uncertainty of model parameters (e.g., soil hydraulic properties) is still unclear. They showed that a more deterministic parameter of the model could result in a narrower range of simulated SMM. With respect to model complexity and resulting uncertainty in SMM estimates, there are sometimes different viewpoints among the studies reviewed here. On the one hand, some authors, for example, MacLeod et al. (2016), argue that forecasting the reliability of SMM using a process-based model could be enhanced by explicitly incorporating parameter uncertainty into the land-surface hydrology equations. Others have suggested that LSMs and GCMs are sometimes too complex and thus unsuited for certain mechanistic studies for which simpler models prove to be adequately efficient (Wei et al., 2006). Overall, there are several reports (He et al., 2023; McColl et al., 2019; Seneviratne, Koster, et al., 2006) that show large differences in SMM between individual models that largely

reflect differences in model parameterizations (e.g., soil hydraulic properties) and, to a lesser degree, soil layer depth and simulation framework (i.e., online vs. offline). There is also some agreement, for example, refer to He et al. (2023); McColl et al. (2019) that LSMs overestimate t_{SMM} . Similarly, Wei et al. (2010) showed that the GLACE GCMs may overestimate the strength of the land-atmosphere coupling and the SMM due to model biases with respect to low-frequency precipitation variability compared to observations. The overestimation of t_{SMM} is also indirectly supported for some (but not all) reanalysis and coupled land-atmosphere models by Dirmeyer et al. (2018).

Overall, in discussing the limitations of LSMs, particularly in the simulation of SMM, we should point out that certain LSM structures or parameters are likely to be consistently challenging. For example, the representation and parameterization of soil-related processes within the framework of classical ordinary and partial differential equations might not be able to account for past states and trajectories of soil moisture (adopted from Rahmati, Or, Amelung, Bauke, et al. (2023)) and therefore poorly represent the impact of SMM on the evolution of soil moisture. In addition, the simplified representation of soil physics and hydrology, such as the simplified parameterization of soil texture, hydraulic conductivity, and soil moisture retention curve as well as the vertical discretization and its effects on for example, infiltration, may not accurately capture the complex interactions between soil, vegetation and atmosphere and therefore may lead to a distortion of soil moisture and its memory. Furthermore, LSMs often struggle to accurately simulate soil moisture feedback mechanisms because soil moisture interacts with various land surface components, such as vegetation dynamics, surface runoff, and groundwater recharge, and any failure to capture this feedback can lead to discrepancies between simulated and observed soil moisture dynamics. As shown in the previous sections, systematic biases in LSM simulations can be identified, indicating clear trends or patterns in model deficits in different regions or under different climatic conditions, which may result from differences in model structures, parameterizations and forcing data. In addition, LSM performance may vary depending on climatic conditions, for example, between dry and humid regions or between temperate and tropical climates. In arid regions, LSMs may struggle to capture the dynamics of sparse vegetation and limited soil moisture availability, while in humid regions they may be confronted with excessive precipitation and runoff processes.

7. SMM From Space

One way to assess the ability of models to represent SMM at the regional to global scale, particularly when in-situ data are sparse, is to benchmark models against satellite-based surface soil moisture products such as those from the Soil Moisture and Ocean Salinity (SMOS) or Soil Moisture Active Passive (SMAP) (Montzka et al., 2017) missions or direct retrieval of soil moisture from multispectral active and passive satellites (Babaeian et al., 2016, 2019; Hassanpour et al., 2020; Mohanty et al., 2017; Rahmati et al., 2015).

However, many satellite products lack the necessary temporal resolution, and this can affect the t_{SMM} results, especially when relevant processes occur within the satellite revisiting period (He et al., 2023). For multi-decadal analyses, which are possible with the multi-mission European Space Agency (ESA) Climate Change Initiative (CCI) Soil Moisture product dating back to 1978, early observations are not available in daily intervals. Nevertheless, their potential at relevant scales is undisputed. In more recent versions of CCI, starting with V7, an “interruption-adjusted” product was developed to minimize the effects of discontinuities and moment shifts when different satellites go online and offline (Preimesberger et al., 2020)—these artifacts could interfere with delayed autocorrelation and SMM calculations. In addition, satellite-derived surface soil moisture products harbor random and periodic errors that impact the estimates of land-atmosphere coupling and therefore shorten the estimated SMM (Seo & Dirmeyer, 2022). In fact, the t_{SMM} is intrinsically linked to the slope of the power density spectrum of soil moisture in $\log(\text{frequency})$: $\log(\text{power})$ phase space, which flattens at high wavenumbers due to random observational noise (based on application of the framework of the first-order Markov process model to assess soil moisture observational data), which is particularly problematic in remote sensing (Kumar et al., 2018). Even if the notion of the validity of a first-order Markov model for soil moisture at daily resolution is rejected (as discussed earlier), the effects of noise on soil moisture readings must be accounted for by applying other routines (e.g., Abdolghafoorian & Dirmeyer, 2022). Another limitation is that satellite observations based on microwave emissions or backscatter can effectively measure soil moisture and its variability only up to a depth of 2–5 cm from the surface, even though they can effectively capture dynamics relevant to deeper layers, up to 10–15 cm (Feldman et al., 2023). This impedes their use in examining t_{SMM} as a function of depth or, for that matter, for a bulk depth representing transpiration processes (MacLeod et al., 2016; Wu & Dickinson, 2004; Yang &

Zhang, 2016). Therefore, it becomes crucial to understand how the temporal and spatial dynamics of the upper layer being observed from space relate to those of the lower layers. Here, the integration of remote sensing and modeling by data assimilation can provide support. For example, the SMAP Level-4 (Reichle et al., 2017) soil moisture product is based on the assimilation of SMAP observations into the Catchment land surface model and includes surface soil moisture (0–5 cm vertical average) as well as root-zone soil moisture (0–100 cm vertical average). However, the integration of multiple satellite data sources with different resolutions or with different acquisition techniques (e.g., active, and passive microwaves) has an impact on the accuracy of the t_{SMM} estimates. It is strongly recommended to integrate data from a single technique (e.g., active microwaves), a single band (e.g., C-band), and with the same algorithm. This minimizes the potential impact of such inconsistencies on the t_{SMM} estimate. While recommended, however, such an approach is not always practical, as data sources are limited. Data sets such as the soil moisture product of the ESA Climate Change Initiative were created to obtain a long-term SM time series from 1978 onwards by statistically combining the data from several satellite missions. Here, the characteristics of the different observation strategies (microwave frequency, active/passive, retrieval algorithm) were harmonized with reduced uncertainty to estimate the t_{SMM} . Earlier data with a lower temporal resolution may, however, be sparser. Alternative methods to estimate root zone soil moisture are P-band radar measurements able to deeper penetrate the soil (15–20 cm) (Tabatabaenejad et al., 2020), or statistical scaling of surface soil moisture time series to the root zone by an exponential filter (Wagner et al., 1999). Other attempts (e.g., Hassanpour et al., 2020) are also underway to determine soil moisture in the root zone from remote sensing data that can be used to determine SMM for deeper depths.

SMM can also be highly variable in space due to land cover or soil texture heterogeneity. To investigate this further, higher spatial resolution soil moisture needs to be considered. Here, the SMAP/Sentinel-1 combined Radiometer/Radar data at 3 km (Das et al., 2019) or the Copernicus Global Land Service Sentinel-1 1 km data (Bauer-Marschallinger et al., 2018) can be utilized.

The first global study attempting to characterize SMM from NASA's SMAP mission was carried out by McColl, Alemohammad, et al. (2017). Several studies have performed additional analyses to characterize t_{SMM} from satellite soil moisture products and their relationship with precipitation (Akbar et al., 2018; Short Gianotti et al., 2019). Kim and Lakshmi (2019) compared multiple satellite soil moisture products and reanalysis in this regard, also investigating the impact of the observed layer depth and temporal frequency. Indeed, memory derived from remote sensing data may be limited to the top layer of the soil profile. This might be different from for example, soil moisture characterizing the whole root zone and its memory as simulated by models. In their study, McColl et al. (2019) proposed and validated a method relying on SMAP observations to estimate t_{SMM} under different soil and climate conditions. The authors found that the use of the Catchment-LSM model to simulate near-surface soil moisture overestimated t_{SMM} related to water limitations, while it underestimated t_{SMM} related to energy-limiting conditions. In a similar study, He et al. (2023) evaluated the hydrometeorological behavior of four widely used global LSMs by comparing them to 5-year t_{SMM} from SMAP observations. They confirmed the findings by McColl, Alemohammad, et al. (2017). Koster et al. (2018) evaluated surface SMM in the Catchment LSM using SMAP data and found it to be deficient; they then used the SMAP data to improve the LSM's parameterizations, thereby improving the simulated memory. In summary, when comparing t_{SMM} from modeling and satellite observations it is possible to improve the structure and the parameterization of LSMs. Nevertheless, future practices using satellite soil moisture data sets with higher temporal frequency, spatial resolution, and longer temporal coverage are expected and urgently needed. In fact, in the near future, new satellite missions dedicated to soil moisture measurements will be launched, such as the National Aeronautics and Space Administration-Indian Space Research Organization (NASA-ISRO) Synthetic Aperture Radar (NISAR) mission, the Radar Observation System for Europe at L-band (ROSE-L) Synthetic Aperture Radar (SAR) mission, and the EUMETSAT Polar System-Second Generation (EPS-SG) missions with the new Scatter-meter (SCA) sensors onboard. The spatiotemporal sampling and accuracy of soil moisture will also be significantly improved through the integration of the guaranteed launch of the Sentinel-1C and Sentinel-1D missions into the Sentinel-1 observation strategy, providing more accurate satellite-based estimates of SMM.

8. Utilizing SMM to Predict and Scale Soil Moisture

The impact of SMM extends beyond its influence on hydrologic processes and can also affect the quality of soil moisture prediction and downscaling of large-scale remote sensing products. Researchers have explored several approaches to improve spatial downscaling of soil moisture data. Mao et al. (2022) used SMM and mass

conservation to improve the spatial downscaling performance of soil moisture provided in SMAP products and for developing high-resolution soil moisture information. To this end, the random forest algorithm was applied by adding three- and 7-day lagged soil moisture as a predictor to represent SMM, along with other regular predictors in routine downscaling studies. However, we believe that the SMM time scale and all lagged soil moisture contents within this time scale could have been used as additional predictors in the model instead of defining the time lags more arbitrarily. In this regard, in the studies of Pal et al. (2016) and Pal and Maity (2019) all lagged soil moisture contents at the target depth that fall within a given time scale of p (referred to as the memory component order), along with current and lagged soil moisture contents of the overlying layer that fall within a given time scale of q (referred to as the forcing component order), were used to predict the soil moisture content of the target depth at a given time. Although this remains to be investigated in future research, the use of SMM and lagged soil moisture content to downscale the large-scale remote sensing products may vary in effectiveness due to the diversity of environmental conditions in different ecosystems. Overall, factors such as soil composition, vegetation cover, topography, and climate variability may play a significant role in the generalizability of these methods in different ecosystems. The application of these methods in ecosystems with homogeneous soil composition and vegetation cover, and consequently with uniform hydrological processes, may be more effective. However, their predictive accuracy may be impaired in ecosystems characterized by high heterogeneity of soil types, vegetation distribution or topographic features. Furthermore, the applicability of these methods may be limited in ecosystems that are subject to rapid land-use change or disturbance, as these changes can significantly affect soil moisture dynamics and challenge the assumptions underlying the downscaling models.

The initialization of soil moisture states in climate models is crucial for accurate hydrological predictions. Walker and Houser (2001) proposed a data assimilation approach using remotely sensed soil moisture to initialize soil moisture states in the NASA NSIPP climate model. By considering the long-term persistence of soil moisture, this method significantly improves model performance in hydrological predictions.

Incorporating soil moisture history and teleconnection indices, Nicolai-Shaw et al. (2016) investigated temporal variations in soil moisture using regression analysis. They found that the predictability of soil moisture decreases with increasing lead time. The influence of previous states of soil moisture on the predictability of its states at any given time depends on the region and season, with higher predictability in dry regions due to minimal atmospheric noise. However, in dry regions, the soil moisture anomaly is only dissipated by evapotranspiration, so noise rarely occurs.

9. The Way Forward

9.1. SMM Emergence

Building on the literature reviewed, this section discusses how SMM develops in soil due to climatic influences and other mediating factors. Figure 4 illustrates the emergence and evolution of SMM in soil, its driving forces, carriers, and effects, adopted from Rahmati, Or, Amelung, Bauke, et al. (2023) and modified upon conducted review in previous sections.

Past research on SMM has been strongly embedded in the field of climate research looking at the fingerprints of SMM on climatic processes but with less attention in providing underlying mechanistic explanations for the occurrence of SMM. Future research should focus on examining the fundamentals that control the emergence, the spatial and temporal extent, and the strength of SMM. To advance this, we propose to classify the controlling factors of SMM into three groups (see Figure 5): (a) atmospheric forcings, (b) land use and management, and (c) soil processes and mechanisms and their properties. Grouping drivers of SMM into these three main groups, we try to elaborate on “how” and “why” SMM emerges in terrestrial ecosystems.

The atmospheric forcings (Group 1) determine the inputs and outputs of information (in the form of anomalies) fed into soil systems, and from there influence the strength of the SMM and the length of its timescale (t_{SMM}). However, it should be noted that Equation 1 and the current equations used to derive SMM ignore important fluxes such as capillary rise, lateral fluxes, irrigation, and miscellaneous non-rainfall water (e.g., dew). Capillary rise is important for conditions where for example, the groundwater level is close to the active soil root zone. The findings by Martínez-de la Torre and Miguez-Macho (2019) have so far been the only research that linked groundwater table variations to the timescale of the memory, thus calling for the continued inclusion of groundwater dynamics in modeling approaches for better predictions of soil moisture




Figure 4. Soil moisture memory (SMM), its drivers, and implications (being adapted from Rahmati et al. ¹¹ and modifications according to conducted review).

dynamics, hydrological processes, and of the interactions between land surface and atmosphere. Although not directly related to SMM, the importance of considering groundwater when addressing soil moisture dynamics is also highlighted by Soyulu and Bras (2022). With respect to lateral fluxes, Rodriguez-Iturbe et al. (2001) argue that although the effects on soil moisture dynamics are local in flat areas, in regions with significant topographic features or in river basins with a complicated drainage network and associated gradient system, lateral fluxes prove to be a crucial determinant of the spatiotemporal distribution of soil moisture dynamics. It is unclear whether the inputs of non-rainfall water, more specifically dew, can contribute enough anomaly to affect SMM. Depending on location, the non-rainfall water inputs can range from 1 to >100% of the monthly precipitation (Xiao et al., 2009) and typically ranges between 4% and 19% of the annual precipitation (Aguirre-Gutiérrez et al., 2019; Groh et al., 2018; Hanisch et al., 2015); however, much of the dewfall presumably takes the form of interception loss and never infiltrates the soil. Another important issue to consider when analyzing SMM is the uncertainty of precipitation measurements with standard rain gauges, which in some cases lead to a very significant underestimation of precipitation (Gebler et al., 2015; Schnepfer et al., 2023). Further research is needed to address all these potential drivers of SMM.




Figure 5. Drivers of soil moisture memory (SMM). The $f(\text{soil})$ implies the role of soil properties and mechanisms that through a feedback loop mediate soil water storage and redistribution and thereby impact SMM. The loop arrow for “Input” refers to the input of information (in the form of moisture anomalies) into the soil system and the arrow for “Output” refers to the dissipation of the anomaly condition.

Finally, SMM is the result of a complex interplay of physical, biological, and hydrological processes and soil properties (Group 3) (Rahmati, Or, Amelung, Bauke, et al., 2023). In fact, SMM is rooted in the integrative nature of soil moisture as a water reservoir (Orth & Seneviratne, 2013) which can be influenced by multiple processes (Figure 3), including soil infiltration, soil water redistribution and storage, root water uptake, capillary rise, and drainage. This review shows that the literature, in general, considers soil depth and soil porosity (as it appears in the autocorrelation expression) to be the main soil properties controlling SMM. While we recognize the valuable contributions of previous efforts such as the SoilWat initiatives (e. g., Aliku & Oshunsanya, 2018; Andrews & Bradford, 2016; Oyeogbe & Oluwasemire, 2013), we maintain that additional consideration should be given to pore size distribution, soil mineral composition (e.g., type and amount of clay), soil organic carbon, and other such properties, as these can control water retention, hydraulic conductivity, and diffusivity and accordingly can influence SMM. In addition, the importance of “hydraulic redistribution” by roots (Dawson, 1993), which is of prominent importance during dry periods by bringing water from deep reservoirs to the near surface soil (Caldwell et al., 1998; Jackson et al., 2000), needs to be emphasized in future research. Hagemann and Stacke (2015) have already shown that hydraulic redistribution by a wide range of plant species is significant in many different biomes around the globe and has implications for SMM.

9.2. Modeling Considerations

The reviewed literature shows that while noteworthy progress has been made in evaluating SMM as captured by LSMs, challenges remain. The lack of long-term measurements and limited simulation power of LSMs for long-term soil moisture variability currently hinder comprehensive analysis. Of course, we recognize that there are ever-growing networks for soil moisture monitoring as additional in situ monitoring stations and new networks come online every year. The oldest has been in continuous operation for more than 30 years. Networks such as the Global Soil Moisture Data Bank (Robock et al., 2000), the International Soil Moisture Network (Dorigo et al., 2011), COSMOS-Europe (Bogena et al., 2022), and the North American Soil Moisture Database (Quiring et al., 2016) are a tremendous resource and have been used for SMM estimation (e.g., Dirmeyer et al., 2016; Seneviratne, Lüthi, et al., 2006) and that will likely become even more useful over time. We also note that there are some valuable efforts, such as Baker et al. (2022), aimed at improving the fidelity of soil moisture monitoring to improve ground truthing of quantities like SMM. Also, isotope tracing studies are rare in truly quantifying water partitioning and the stored precipitation fraction across scales and for model validation. In addition, generalizing conclusions across different models is difficult due to differences in model complexity and parameter uncertainties. Future research efforts should focus on overcoming these challenges to improve the reliability and understanding of SMM in climate models. By means of a synergistic fusion of computational model simulations, empirical observations, and meticulous joint analyses with state-of-the-art satellite-based products, researchers can improve our basic understanding of SMM and its profound impacts on the complicated interplay between Earth's water and energy cycles. Continued efforts to refine models and improve data availability will contribute to more accurate predictions and a better understanding of the influence of SMM on climate dynamics. Several researchers (e.g., MacLeod et al., 2016) have pointed out that the uncertainty in current memory estimates is not clear and that it is not obvious to what extent they depend on model parameterization uncertainties. Sensitivity analyses indicate that memory estimates and their uncertainty depend to a significant extent on key hydraulic parameters used to parameterize various processes in land surface models, suggesting that the models do not represent the memory as exists. On the other hand, soil hydraulic parameters in large-scale land surface, hydrology, and crop models are usually approximated by pedotransfer functions (PTFs), and recent evaluations show that the choice of PTFs is important for simulating soil water balance fluxes (Weihermüller et al., 2021) and probably for SMM estimates.

Subsurface water vapor fluxes are rarely ever considered when simulating terrestrial energy, water, and carbon budgets using LSMs (Garcia Gonzalez et al., 2012). By including isothermal and thermal water vapor transfer, Garcia Gonzalez et al. (2012) evaluated the Joint UK Environment Simulator LSM (known as JULES for short) to simulate the key soil variables and showed that such inclusion contributes significantly to water and heat transfer in the upper soil layers in semiarid and temperate arid climates. Similar attempts have also been made to parameterize soil evaporation by coupled transport of moisture and heat for arid and semiarid regions by considering mechanisms associated with the transportation, condensation, and evaporation of water vapor in the soil matrix (Meng et al., 2023); note that Milly (1984) concluded some time back that neglecting thermal effects

on vapor diffusion in the soil has minimal impact on computed evaporation rates. However, no study to date (to the authors' knowledge) has investigated the impact of soil water vapor specifically on SMM. Further Community research in this area may be necessary in the future.

Again, Equation 1 is typically used to analyze SMM. Recent developments in data-driven analysis using for example, machine learning or deep learning methods provide new opportunities to study and analyze hydrological processes (De Lavenne et al., 2022; Lees et al., 2021; Ma et al., 2021; Sungmin & Orth, 2021). These data-driven analyses typically do not account for the specifics of hydrological dynamics. In a recent paper, De la Fuente et al. (2023) developed an improved machine learning approach based on Long Short-Term Memory (LSTM) that is adapted to the specific system dynamics of hydrological processes and considers the importance of trends and patterns in data. They exploited the similarity between Equation 1 and the underlying equations used in LSTM to develop this framework. They obtained a similar performance as compared to standard LSTM approaches but provided better interpretability of hydrological processes observed in 588 catchments across the US. This proposed framework and the ongoing developments in data-driven approaches can serve as a basis for further exploration of SMM as well as its interactions with other terrestrial processes. However, it should be mentioned that while data-driven approaches offer promising avenues for SMM research, there are still challenges in their usage such as incorporating physical understanding, ensuring data quality, improving generalization, improving interpretability, and reducing overfitting. There are a variety of ways to overcome these challenges. For example, hybrid models that combine data-driven techniques with mechanistic models can improve interpretability and robustness. One can ensure data quality and quantity by using data preprocessing techniques to deal with missing or erroneous data, and by using techniques such as data augmentation or ensemble learning to improve model performance and robustness. There are also already well-documented techniques to increase the interpretability of the data-driven models (e.g., feature importance analysis, sensitivity analysis or visualization methods) and to prevent overfitting (e.g., dropout, L1/L2 regularization or early stopping). One other possible pathway to analyze SMM that has not yet been explored is to use mathematical formalisms applied to signal processing and dynamical systems with memory, as proposed by Rahmati, Or, Amelung, Bauke, et al. (2023) in the case of soil memory as a whole. These mathematical formalisms may include, among others, fractional differential equations (Khalighi et al., 2022) that can store information about past states and trajectories of a dynamical system. Indeed, it is worth noting that SMM is an emergent property of the coupled land-atmosphere systems that is the result of many other factors, and therefore its proper study requires new frameworks that go beyond the conventional evaporative fraction and soil moisture relationship (e.g., Haghighi et al., 2018). An initiative by Rahmati, Or, Amelung, and Vereecken, (2023) that uses fractional differential equations to redefine a hydrologic model by including a memory term showed that SMM can mitigate and amplify the effects of drought.

9.3. SMM Under Extreme Events and Future Climate Projection

Studying SMM under the bottleneck of extreme conditions is a promising way to gain deep insight into the complicated behavior and responsiveness of soil dynamics during extreme events. Orth and Seneviratne (2012) shed light on the critical importance of excluding extreme periods from analytical consideration while illuminating the potential role of soil physical properties in regulating SMM under extreme drought. Recent research (e.g., Rahmati et al., 2020) shows that increasing drought has implications for the long-term lagged relationship (representative of the memory effect) between soil moisture and evapotranspiration as a key variable linking soil moisture to the atmosphere. Therefore, exploring the physical processes underlying SMM in these extremes, whether drought, flood, or wildfire, will strengthen our predictive power and enable us to skillfully manage the uncertainties in the predictability of extreme events, as well as to better forecast their role in future regional climate. It can be argued that the focus on extremes is driven by societal demands rather than scientific necessity. Since extremes are the tails of a distribution, there is no difference to SMM processes in extremes. However, unlike a mere compilation of existing land surface features, extreme events often trigger nonlinear responses in land surface processes (including SMM) that can deviate significantly from typical patterns. For example, the loss of land-atmosphere coupling, as studied by Wu and Dirmeyer (2020), can occur as one of several mechanisms during extreme events and lead to shifts in regional climate dynamics. The methods used in the literature to analyze SMM after extreme events are summarized in Table 4.

The projected impact of future climate on SMM in the coming decades is also an important aspect of SMM research that is missing from the literature. This is even more important when we consider climate change and land use intensification. Therefore, we suggest that this could be explored in detail in future research.

Table 4

Approaches Used in Literature to Analyze Soil Moisture Memory (SMM) in Relation to Extreme Events

Methodology	Description
Periods with On-off extreme events	The impact of extreme events on SMM can be analyzed by excluding the periods where these extreme conditions occur (Orth & Seneviratne, 2012). SMM can then be compared between the original and the truncated data. This methodology is particularly useful for analyzing extreme events at seasonal or shorter scales by applying the internal autocorrelation metric.
Regions with and without extreme events	In this method, the SMM of regions with and without extreme events were compared (Ashraf & Ahrens, 2013). The authors divided the study area into two subregions with and without extreme events (e.g., low rainfall and heavy and frequent rainfall).
Conducting joint control-sensitivity experiment	The relationship between SMM and extreme events (such as wildfires and drought) can also be analyzed by conducting control experiments along with sensitivity experiments in a model environment (Lorenz et al., 2010). A control experiment is defined by coupled soil moisture-atmosphere and a sensitivity experiment is a coupled simulation with prescribed soil moisture in which soil moisture is fixed at some preset values (e.g., soil moisture being fixed at some preset values such as field capacity or wilting point).
Manipulated initial soil moisture anomalies	Manipulating initial soil moisture anomalies is also a common method used to establish relationships between SMM and extreme events (Abolafia-Rosenzweig et al., 2023; Liang & Yuan, 2021; Nicholson, 2000; Stahle & Cleaveland, 1988; Tisdeman & Menzel, 2021).

9.4. Investigations Into the Spatial Component of SMM

As reviewed in Section 3, the temporal variation of memory timescale exhibits complex dynamics influenced by seasonality, availability of radiant energy, hydrological factors, and geographic dependencies. Divergent findings pervade scientific debates, with certain investigations supporting the idea of a prolonged memory timescale in winter and a shortened one in summer (Delworth & Manabe, 1988; Dirmeyer et al., 2009; Douville et al., 2007; Entin et al., 2000; Liu et al., 2014; Shinoda & Nandintsetseg, 2011). However, a counter-narrative emerges from other scientific investigations (Hagemann & Stacked, 2015; Orth & Seneviratne, 2012; Wu & Dickinson, 2004), casting doubt on this idea. Consequently, there is an undeniable need for further research to gain a deeper understanding of the intricate regulatory mechanisms that govern differences in memory timescales across regions and different climatic contexts. Note that spatial variations in SMM are influenced by a combination of factors (e.g., latitude, elevation, drought, soil depth, topography, and hydraulic properties (He et al., 2023; Orth et al., 2013)) that also affect its timescale. SMM estimation is sensitive to uncertainties in hydraulic parameters (e.g., MacLeod et al., 2016), and several of these hydraulic parameters show very high spatial heterogeneity.

In examining the spatial variability of SMM, examples can be found where unexpected patterns or contradictions to prevailing theories have been observed that may challenge our understanding. For example, the assumption of a correlation between t_{SMM} and latitude, as proposed by Delworth and Manabe (1988), where a shorter t_{SMM} is attributed to a lower latitude due to greater potential evaporation rates and faster dissipation of moisture anomalies, overlooks the effects of cloud cover on incoming radiation. This is because subtropical regions can receive more shortwave radiation than tropical regions (belonging to lower latitudes) with higher cloud cover within the Intertropical Convergence Zone (ITCZ). The recognition and inclusion of these examples of unexpected patterns or inconsistencies emphasizes the importance of continued research to refine existing theories and models to capture the full range of spatial variability in the SMM.

In the context of the spatiotemporal variations that characterize SMM, an examination of the existing literature reveals a perplexing observation: compared to the temporal aspect of SMM, the spatial aspect—the ability of SMM in one location to affect climate variables in another—has remained conspicuously unexplored. To date, no clear spatial component (non-local effects) has been established for SMM, although Seneviratne et al. (2010) nicely brought this to the attention of the community by mentioning the possibility of large-scale and non-local impacts of the soil moisture (e.g., the impacts of soil moisture on large-scale circulation patterns). This is only indirectly investigated by Koster et al. (2014) and Koster et al. (2016), who explore the mechanisms that allow the state of soil moisture in one region to influence atmospheric conditions in another, and more recently, Giles et al. (2023) reported a non-local coupling mechanism between soil moisture and the atmosphere in South

America. Thess initiative needs to be followed with similar studies as the question of whether the memory of a particular point in space can affect surrounding areas has not been clearly answered. Another good example of non-local impacts of SMM is provided by Dong et al. (2023), who showed that the negative soil moisture anomalies in May 2020 over the Indo-China Peninsula in Southeast Asia contributed to the Meiyu period in East Asia during the East Asian summer monsoon in 2020 (see Table 2 for details). The question of how changing conditions in neighboring areas can lead to the modification of memory at any point in space has also not been resolved, although some teleconnections have been made between the occurrence of SMM and ENSO events (Amenu et al., 2005; Timbal et al., 2002). By performing further research into this spatial component of SMM, scientists can gain a better understanding of how SMM propagates across different regions. Further investigations on teleconnections between the occurrence of SMM and events such as ENSO can shed light on how large-scale climate phenomena interact with local SMM. Research can also focus on scaling up SMM from point observations to larger areas. By integrating (effectively, upscaling) data from multiple points, researchers can analyze the collective impact of SMM on a broader scale. However, it is worth noting that the spatial component of SMM is not always distinct or easy to identify. Factors such as regional and temporal variations, methodological challenges, feedback mechanisms, and data limitations can make it difficult to understand it. For example, depending on the region, there may be consistent or inconsistent relationships between SMM and climate variables. On the other hand, the strength and duration of non-local SMM impacts may also vary over time. For example, SMM responses to climate drivers may fluctuate on an interannual or decadal timescale, introducing uncertainty into the understanding of its spatial component. Methodological challenges in quantifying and attributing the effects of the SMM in space and the presence of feedback mechanisms between the SMM and local climate conditions can lead to further uncertainties. Finally, limited observational data in certain regions or over certain time periods can make it even more difficult to identify non-local impacts of SMM.

9.5. SMM Links to Community Oriented UPHs

Recently, 23 major unsolved problems in hydrology (UPHs) have been identified through a community initiative hoping to help and guide research efforts in the coming years (Blöschl et al., 2019). Therefore, in this section we attempt to wrap up the links between SMM and those UPHs calling for future research topics. In fact, SMM, as the capacity of the soil retaining the memory of past moisture conditions, is relevant to several of the UPHs. Among them, the following questions are particularly important:

- “*Variability of extremes—question 11. “Why, how, and when do rain-on-snow events produce exceptional runoff?”*”: Rain-on-snow events can lead to rapid snowmelt and increased runoff, particularly in regions with significant SMM—regions with persistent wet anomalies—as water infiltration into soils with high moisture content may be limited and therefore excess water contributes to surface runoff. Such an understanding is likely to be of greater importance for the prediction and management of flood risk in cold regions. These phenomena need to be further investigated in future research.
- “*Modeling methods—question 20. How can we disentangle and reduce model structural/parameter/input uncertainty in hydrological prediction?”*”: As discussed in this review, the SMM is a vital component of the hydrological cycle. Therefore, its correct representation in hydrological models can potentially improve their ability to simulate the interactions between soil, vegetation, and atmosphere and to capture the effects of land surface changes on water resources. Such integration may involve the use of observational data, remote sensing techniques and process-based modeling approaches to capture the complex interactions between soil properties, vegetation dynamics and atmospheric factors. A proper integration of SMM into hydrological models is also in line with the perspectives for the future of land surface models with respect to the representative complex terrestrial systems as presented by Fisher and Koven (2020). In particular, we believe that improved representation of SMM in hydrological models and land surface models is likely to help address two of the three “grand challenges” identified by Fisher and Koven (2020) (i.e., managing process complexity and representing land surface heterogeneity). However, this still needs to be explored.

10. Summary and Outlook

In this paper, we reviewed the state of the art in analyzing and characterizing SMM in the Earth system. We analyzed the role of SMM on key terrestrial system processes and identified the factors that affect SMM. Atmospheric forcings, water storage and movement, soil hydraulic properties, and vegetation as well as anthropogenic activities influence the character of SMM. Extreme events such as heavy precipitation, drought, and

wildfire can alter the soil over time, thus additionally affecting the link between past and current soil moisture conditions. Also, the depth and properties of the active soil layer and plant root development contribute to the manifestation of SMM.

We examined the factors that control the timescale of SMM. The SMM timescale is influenced by several factors, including seasonal variations in the atmospheric conditions, soil hydrology, occurrence and their severity of extreme event, anthropogenic activities, soil properties and their variability in space and time, groundwater levels, vegetation, sampling frequency, and data sources. We suggest grouping these controlling factors into three groups to help organize SMM research: (a) atmospheric forcings, (b) land use and management, and (c) soil processes and soil properties. Some of the key processes that control soil moisture dynamics and thus SMM at the field to catchment scale such as capillary rise, groundwater dynamics and lateral fluxes should receive more attention.

Our literature analysis shows that SMM has significant implications for weather variability, surface energy balance, drought and flood monitoring, water use efficiency, biogeochemical cycling, groundwater prediction, and climate impacts. Excluding extreme periods from SMM quantification reduces the time scale of SMM, especially under drought conditions. Further research should investigate the mechanisms, regional impacts, and relationship between soil properties and SMM under extreme conditions to support decision-making during extreme weather events.

Several approaches have been identified in the literature to quantify memory timescale and its strength. These metrics include autocorrelation timescale, variance spectrum, and the fraction of precipitation stored, among others. Using these metrics, published literature reports that the magnitude of the SMM ranges from weeks to over a year. Examination of the reported spatiotemporal variability of SMM indicates that the memory timescale of soil moisture varies throughout the year and is influenced by seasonal changes, availability of radiant energy, and hydrologic factors. Some studies suggest longer memory timescales in winter and shorter timescales in summer, whereas others find more complex behavior. Geographic dependencies and soil depth also contribute to temporal variations in memory timescales. Further scientific research is required to gain a much-needed deeper understanding of these complicated dynamics in different climatic environments. SMM also exhibits considerable spatial variability, with memory timescales increasing from tropical regions to high latitudes and influenced by spatially varying potential evapotranspiration rates. In arid regions, the memory timescale is longer due to smaller variations in soil moisture. Spatial variation in memory timescale is also related to factors such as precipitation duration, runoff, and evapotranspiration. However, estimates of the memory timescale are limited by uncertainties in hydraulic parameters, indicating the need for further research.

We also investigated how SMM is represented by LSMs. In this respect it is important to recognize that a correct description of the coupling of soil moisture, atmosphere, and land surface processes is critical for quantifying SMM, especially in regions where soil moisture strongly influences evapotranspiration. Climate models have evolved to better represent this relationship, with advances in parameterizing evapotranspiration and in the treatment of vegetation and soil dynamics. However, challenges remain, including the overestimation of soil moisture drought, highlighting the need for further progress and a closer integration of models and observations. Improved characterization of SMM may also be reached by assimilating observational data into an LSM system. In this regard, satellite observations can effectively estimate surface soil moisture, but their depth effect is limited. Obtaining soil moisture at deeper depths is important as several studies have shown that SMM is depth-dependent and typically increases with soil depth. We also pointed out the possibilities of using data-driven approaches and mathematical methods such as fractional mathematics as a basis for further research on SMM, as well as on its interactions with other terrestrial processes.

Finally, we have identified four avenues to further explore and quantify the role of SMM based on a better understanding of the underlying mechanisms and processes that influence it. These are: understanding the underlying mechanisms and processes that determine the character of SMM, improving the treatment of SMM in land models, exploring the physical processes underlying SMM during extreme events, and exploring the spatial component (non-local effect) of SMM.

Data Availability Statement

Data availability does not apply to this article, as no new data was created or analyzed in this study.

Acknowledgments

MR and CM gratefully acknowledge funding by the German Federal Ministry for Digital and Transport (BMDV) for the HABKIS2 project (19F2264B). HV, WA, CM, and HB acknowledge support by the Deutsche Forschungsgemeinschaft—SFB 1502/1–2022—Projektnummer 450058266. HV, WA, CM, and HB acknowledge support from the Deutsche Forschungsgemeinschaft under Germany's Excellence Strategy, EXC-2070—390732324. JG was supported by the Deutsche Forschungsgemeinschaft (project no. 460817082). We would like to express our sincere gratitude to Prof. Dr. Jan Vanderborcht (Forschungszentrum Jülich GmbH, Germany), Prof. Dr. Dani Or (ETH Zurich, Switzerland and University of Nevada, USA) and Prof. Dr. Dara Entekhabi (Massachusetts Institute of Technology) for their invaluable support and insightful discussions throughout the course of this research. In addition, we are very grateful to three reviewers, two anonymous and Prof. Dr. Paul Dirmeyer, whose constructive feedback significantly improved the quality of this work, as well as to the editor (Prof. Dr. Alan Robock) for their guidance and expertise in shaping the final manuscript. Open Access funding enabled and organized by Projekt DEAL.

References

- Abdolghafoorian, A., & Dirmeyer, P. A. (2022). Accounting for the effect of noise in satellite soil moisture data on estimates of land-atmosphere coupling using information theoretical metrics. *Journal of Hydrometeorology*, 23(10), 1587–1605. <https://doi.org/10.1175/Jhm-D-21-0232.1>
- Abolafia-Rosenzweig, R., He, C., Chen, F., Ikeda, K., Schneider, T., & Rasmussen, R. (2023). High resolution forecasting of summer drought in the western United States. *Water Resources Research*, 59(3), e2022WR033734. <https://doi.org/10.1029/2022wr033734>
- Agbeshie, A. A., Abugre, S., Atta-Darkwa, T., & Awuah, R. (2022). A review of the effects of forest fire on soil properties. *Journal of Forestry Research*, 33(5), 1419–1441. <https://doi.org/10.1007/s11676-022-01475-4>
- Aguirre-Gutiérrez, C. A., Holwerda, F., Goldsmith, G. R., Delgado, J., Yezpe, E., Carbajal, N., et al. (2019). The importance of dew in the water balance of a continental semiarid grassland. *Journal of Arid Environments*, 168, 26–35. <https://doi.org/10.1016/j.jaridenv.2019.05.003>
- Akbar, R., Short Gianotti, D., McColl, K. A., Haghighi, E., Salvucci, G. D., & Entekhabi, D. (2018). Hydrological storage length scales represented by remote sensing estimates of soil moisture and precipitation. *Water Resources Research*, 54(3), 1476–1492. <https://doi.org/10.1002/2017wr021508>
- Alfieri, L., Claps, P., D'Odorico, P., Laio, F., & Over, T. M. (2008). An analysis of the soil moisture feedback on convective and stratiform precipitation. *Journal of Hydrometeorology*, 9(2), 280–291. <https://doi.org/10.1175/2007jhm863.1>
- Aliku, O., & Oshunsanya, S. O. (2018). Assessment of the SOILWAT model for predicting soil hydro-physical characteristics in three agro-ecological zones in Nigeria. *International Soil and Water Conservation Research*, 6(2), 131–142. <https://doi.org/10.1016/j.iswcr.2018.01.003>
- Amenu, G. G., Kumar, P., & Liang, X. Z. (2005). Interannual variability of deep-layer hydrologic memory and mechanisms of its influence on surface energy fluxes. *Journal of Climate*, 18(23), 5024–5045. <https://doi.org/10.1175/Jcli3590.1>
- Andrews, C. M., & Bradford, J. B. (2016). SOILWAT: A mechanistic ecohydrological model for ecosystem classification and prediction. In *World conference on natural resource modeling*.
- Asharaf, S., & Ahrens, B. (2013). Soil-moisture memory in the regional climate model COSMO-CLM during the Indian summer monsoon season. *Journal of Geophysical Research-Atmospheres*, 118(12), 6144–6151. <https://doi.org/10.1002/jgrd.50429>
- Babaeian, E., Homaee, M., Montzka, C., Vereecken, H., Norouzi, A., & van Genuchten, M. T. (2016). Soil moisture prediction of bare soil profiles using diffuse spectral reflectance information and vadose zone flow modeling. *Remote Sensing of Environment*, 187, 218–229. <https://doi.org/10.1016/j.rse.2016.10.029>
- Babaeian, E., Sadeghi, M., Jones, S. B., Montzka, C., Vereecken, H., & Tuller, M. (2019). Ground, proximal, and satellite remote sensing of soil moisture. *Reviews of Geophysics*, 57(2), 530–616. <https://doi.org/10.1029/2018rg000618>
- Baker, C. B., Cosh, M., Bolton, J., Brunsberg, M., Caldwell, T., Connolly, S., et al. (2022). Working toward a National coordinated soil moisture monitoring network: Vision, progress, and future directions. *Bulletin of the American Meteorological Society*, 103(12), E2719–E2732. <https://doi.org/10.1175/Bams-D-21-0178.1>
- Bao, L., Yu, L., Li, Y., Yan, F., Lyne, V., & Ren, C. (2023). Climate change impacts on agroecosystems in China: Processes, mechanisms and prospects. *Chinese Geographical Science*, 33(4), 583–600. <https://doi.org/10.1007/s11769-023-1362-0>
- Barcellos, D., O'Connell, C., Silver, W., Meile, C., & Thompson, A. (2018). Hot spots and hot moments of soil moisture explain fluctuations in iron and carbon cycling in a humid tropical forest soil. *Soil Systems*, 2(4), 59. <https://doi.org/10.3390/soilsystems2040059>
- Bauer-Marschallinger, B., Freeman, V., Cao, S., Paulik, C., Schaufler, S., Stachl, T., et al. (2018). Toward global soil moisture monitoring with Sentinel-1: Harnessing assets and overcoming obstacles. *IEEE Transactions on Geoscience and Remote Sensing*, 57(1), 520–539. <https://doi.org/10.1109/TGRS.2018.2858004>
- Beaudoin, H. a. M. R. N. G. H. (2016). GLDAS NOAA land surface model L4 3 hourly 0.25 x 0.25 degree V2.1. In M. Greenbelt (Ed.). Goddard Earth Sciences Data and Information Services Center (GES DISC). <https://doi.org/10.5067/E7TYRXPJKWOQ>
- Bellucci, A., Haarsma, R., Bellouin, N., Booth, B., Cagnazzo, C., van den Hurk, B., et al. (2015). Advancements in decadal climate predictability: The role of nonoceanic drivers. *Reviews of Geophysics*, 53(2), 165–202. <https://doi.org/10.1002/2014rg000473>
- Benson, D. O., & Dirmeyer, P. A. (2023). The soil moisture–surface flux relationship as a factor for extreme heat predictability in subseasonal to seasonal forecasts. *Journal of Climate*, 36(18), 6375–6392. <https://doi.org/10.1175/jcli-d-22-0447.1>
- Berg, A., & Sheffield, J. (2018). Soil moisture–evapotranspiration coupling in CMIP5 models: Relationship with simulated climate and projections. *Journal of Climate*, 31(12), 4865–4878. <https://doi.org/10.1175/Jcli-D-17-0757.1>
- Berg, A., & Sheffield, J. (2019). Historic and projected changes in coupling between soil moisture and evapotranspiration (ET) in CMIP5 models confounded by the role of different ET components. *Journal of Geophysical Research-Atmospheres*, 124(11), 5791–5806. <https://doi.org/10.1029/2018jd029807>
- Bierkens, M. F. P., & van den Hurk, B. J. J. M. (2007). Groundwater convergence as a possible mechanism for multi-year persistence in rainfall. *Geophysical Research Letters*, 34(2). <https://doi.org/10.1029/2006gl028396>
- Blöschl, G., Bierkens, M. F., Chambel, A., Cudennec, C., Destouni, G., Fiori, A., et al. (2019). Twenty-three unsolved problems in hydrology (UPH)—a community perspective. *Hydrological Sciences Journal*, 64(10), 1141–1158. <https://doi.org/10.1080/02626667.2019.1620507>
- Bogena, H., Schrön, M., Jakobi, J., Ney, P., Zacharias, S., Andreasen, M., et al. (2022). COSMOS-Europe: A European network of cosmic-ray neutron soil moisture sensors. *Earth System Science Data*, 14(3), 1125–1151. <https://doi.org/10.5194/essd-14-1125-2022>
- Bonan, G. B., & Stillwell-Soller, L. M. (1998). Soil water and the persistence of floods and droughts in the Mississippi River Basin. *Water Resources Research*, 34(10), 2693–2701. <https://doi.org/10.1029/98wr02073>
- Bu, L. L., Zuo, Z., Zhang, K., & Yuan, J. (2023). Impact of evaporation in Yangtze River valley on heat stress in North China. *Journal of Climate*, 36(12), 4005–4017. <https://doi.org/10.1175/Jcli-D-22-0573.1>
- Caldwell, M. M., Dawson, T. E., & Richards, J. H. (1998). Hydraulic lift: Consequences of water efflux from the roots of plants. *Oecologia*, 113(2), 151–161. <https://doi.org/10.1007/s004420050363>
- Champagne, C., Rowlandson, T., Berg, A., Burns, T., L'Heureux, J., Tetlock, E., et al. (2016). Satellite surface soil moisture from SMOS and Aquarius: Assessment for applications in agricultural landscapes. *International Journal of Applied Earth Observation and Geoinformation*, 45, 143–154. <https://doi.org/10.1016/j.jag.2015.09.004>
- Chan, S. K., Bindlish, R., O'Neill, P. E., Njoku, E., Jackson, T., Colliander, A., et al. (2016). Assessment of the SMAP passive soil moisture product. *IEEE Transactions on Geoscience and Remote Sensing*, 54(8), 4994–5007. <https://doi.org/10.1109/Tgrs.2016.2561938>
- Chatfield, C. (2003). *The analysis of time series: An introduction*. Chapman and hall/CRC. <https://doi.org/10.4324/9780203491683>
- Chesworth, W. (2007). *Encyclopedia of soil science*. Springer Science & Business Media. <https://doi.org/10.1007/978-1-4020-3995-9>
- Dai, Y. J., Shanguan, W., Wei, N., Xin, Q., Yuan, H., Zhang, S., et al. (2019). A review of the global soil property maps for Earth system models. *Soils*, 5(2), 137–158. <https://doi.org/10.5194/soil-5-137-2019>
- Daly, E., & Porporato, A. (2005). A review of soil moisture dynamics: From rainfall infiltration to ecosystem response. *Environmental Engineering Science*, 22(1), 9–24. <https://doi.org/10.1089/ees.2005.22.9>

- Das, N. N., Entekhabi, D., Dunbar, R. S., Chaubell, M. J., Colliander, A., Yueh, S., et al. (2019). The SMAP and Copernicus Sentinel 1A/B microwave active-passive high resolution surface soil moisture product. *Remote Sensing of Environment*, 233, 111380. <https://doi.org/10.1016/j.rse.2019.111380>
- Dawson, T. E. (1993). Hydraulic lift and water use by plants: Implications for water balance, performance and plant-plant interactions. *Oecologia*, 95(4), 565–574. <https://doi.org/10.1007/BF00317442>
- De la Fuente, L. A., et al. (2023). Towards interpretable LSTM-based modelling of hydrological systems. *EGU sphere*, 2023, 1–29. <https://doi.org/10.5194/hess-28-945-2024>
- De Lavenne, A., Andréassian, V., Crochemore, L., Lindström, G., & Arheimer, B. (2022). Quantifying multi-year hydrological memory with catchment forgetting curves. *Hydrology and Earth System Sciences*, 26(10), 2715–2732. <https://doi.org/10.5194/hess-26-2715-2022>
- Delworth, T. L., & Manabe, S. (1988). The influence of potential evaporation on the variabilities of simulated soil wetness and climate. *Journal of Climate*, 1(5), 523–547. [https://doi.org/10.1175/1520-0442\(1988\)001%3C0523:TIOPEO%3E2.0.CO;2](https://doi.org/10.1175/1520-0442(1988)001%3C0523:TIOPEO%3E2.0.CO;2)
- Diffenbaugh, N. S., Pal, J. S., Giorgi, F., & Gao, X. (2007). Heat stress intensification in the Mediterranean climate change hotspot. *Geophysical Research Letters*, 34(11). <https://doi.org/10.1029/2007gl030000>
- Dirmeyer, P. A. (1999). Assessing GCM sensitivity to soil wetness using GSWP data. *Journal of the Meteorological Society of Japan*, 77(1b), 367–385. https://doi.org/10.2151/jmsj.1965.77.1B_367
- Dirmeyer, P. A. (2000). Using a global soil wetness dataset to improve seasonal climate simulation. *Journal of Climate*, 13(16), 2900–2922. [https://doi.org/10.1175/1520-0442\(2000\)013<2900:Uagswd>2.0.Co;2](https://doi.org/10.1175/1520-0442(2000)013<2900:Uagswd>2.0.Co;2)
- Dirmeyer, P. A. (2011a). A history and review of the global soil wetness project (GSWP). *Journal of Hydrometeorology*, 12(5), 729–749. <https://doi.org/10.1175/Jhm-D-10-05010.1>
- Dirmeyer, P. A. (2011b). The terrestrial segment of soil moisture-climate coupling. *Geophysical Research Letters*, 38(16). <https://doi.org/10.1029/2011gl048268>
- Dirmeyer, P. A., & Brubaker, K. L. (2007). Characterization of the global hydrologic cycle from a back-trajectory analysis of atmospheric water vapor. *Journal of Hydrometeorology*, 8(1), 20–37. <https://doi.org/10.1175/Jhm557.1>
- Dirmeyer, P. A., Chen, L., Wu, J., Shin, C. S., Huang, B., Cash, B. A., et al. (2018). Verification of land-atmosphere coupling in forecast models, reanalyses and land surface models using flux site observations. *Journal of Hydrometeorology*, 19(No 2), 375–392. <https://doi.org/10.1175/JHM-D-17-0152.1>
- Dirmeyer, P. A., et al. (2015). Land-atmosphere interactions and the water cycle. In *Seamless prediction of the Earth system. From minutes to months* (pp. 145–154). WMO.
- Dirmeyer, P. A., & Norton, H. E. (2018). Indications of surface and sub-surface hydrologic properties from SMAP soil moisture retrievals. *Hydrology*, 5(3), 36. <https://doi.org/10.3390/hydrology5030036>
- Dirmeyer, P. A., Schlosser, C. A., & Brubaker, K. L. (2009). Precipitation, recycling, and land memory: An integrated analysis. *Journal of Hydrometeorology*, 10(1), 278–288. <https://doi.org/10.1175/2008jhm1016.1>
- Dirmeyer, P. A., Wu, J., Norton, H. E., Dorigo, W. A., Quiring, S. M., Ford, T. W., et al. (2016). Confronting weather and climate models with observational data from soil moisture networks over the United States. *Journal of Hydrometeorology*, 17(4), 1049–1067. <https://doi.org/10.1175/JHM-D-15-0196.1>
- Dong, Y. S., Chen, H., & Dong, X. (2023). Impact of antecedent soil moisture anomalies over the Indo-China Peninsula on the super Meiyu event in 2020. *Journal of Meteorological Research*, 37(2), 234–247. <https://doi.org/10.1007/s13351-023-2144-4>
- Dorigo, W. A., Wagner, W., Hohensinn, R., Hahn, S., Paulik, C., Xaver, A., et al. (2011). The International soil moisture network: A data hosting facility for global in situ soil moisture measurements. *Hydrology and Earth System Sciences*, 15(5), 1675–1698. <https://doi.org/10.5194/hess-15-1675-2011>
- Douville, H. (2002). Influence of soil moisture on the Asian and African monsoons. Part II: Interannual variability. *Journal of Climate*, 15(7), 701–720. [https://doi.org/10.1175/1520-0442\(2002\)015<0701:Iosmot>2.0.Co;2](https://doi.org/10.1175/1520-0442(2002)015<0701:Iosmot>2.0.Co;2)
- Douville, H. (2004). Relevance of soil moisture for seasonal atmospheric predictions: Is it an initial value problem? *Climate Dynamics*, 22(4), 429–446. <https://doi.org/10.1007/s00382-003-0386-5>
- Douville, H., & Chauvin, F. (2000). Relevance of soil moisture for seasonal climate predictions: A preliminary study. *Climate Dynamics*, 16(10–11), 719–736. <https://doi.org/10.1007/s003820000080>
- Douville, H., Chauvin, F., & Broqua, H. (2001). Influence of soil moisture on the Asian and African monsoons. Part I: Mean monsoon and daily precipitation. *Journal of Climate*, 14(11), 2381–2403. [https://doi.org/10.1175/1520-0442\(2001\)014<2381:Iosmot>2.0.Co;2](https://doi.org/10.1175/1520-0442(2001)014<2381:Iosmot>2.0.Co;2)
- Douville, H., Conil, S., Tyteca, S., & Voldoire, A. (2007). Soil moisture memory and West African monsoon predictability: Artefact or reality? *Climate Dynamics*, 28(7–8), 723–742. <https://doi.org/10.1007/s00382-006-0207-8>
- Elnashar, A., Wang, L., Wu, B., Zhu, W., & Zeng, H. (2021). Synthesis of global actual evapotranspiration from 1982 to 2019. *Earth System Science Data*, 13(2), 447–480. <https://doi.org/10.5194/essd-13-447-2021>
- Entekhabi, D., & Rodriguez-Iturbe, I. (1994). Analytical framework for the characterization of the space-time variability of soil moisture. *Advances in Water Resources*, 17(1–2), 35–45. [https://doi.org/10.1016/0309-1708\(94\)90022-1](https://doi.org/10.1016/0309-1708(94)90022-1)
- Entekhabi, D., Rodriguez-Iturbe, I., & Bras, R. L. (1992). Variability in large-scale water balance with land surface-atmosphere interaction. *Journal of Climate*, 5(8), 798–813. [https://doi.org/10.1175/1520-0442\(1992\)005<0798:vilswb>2.0.co;2](https://doi.org/10.1175/1520-0442(1992)005<0798:vilswb>2.0.co;2)
- Entin, J. K., Robock, A., Vinnikov, K. Y., Hollinger, S. E., Liu, S., & Namkhai, A. (2000). Temporal and spatial scales of observed soil moisture variations in the extratropics. *Journal of Geophysical Research*, 105(D9), 11865–11877. <https://doi.org/10.1029/2000jd900051>
- Esit, M., Kumar, S., Pandey, A., Lawrence, D. M., Rangwala, I., & Yeager, S. (2021). Seasonal to multi-year soil moisture drought forecasting. *Npj Climate and Atmospheric Science*, 4(1), 16. <https://doi.org/10.1038/s41612-021-00172-z>
- Evans, S., D. Allison, S., & V. Hawkes, C. (2022). Microbes, memory and moisture: Predicting microbial moisture responses and their impact on carbon cycling. *Functional Ecology*, 36(6), 1430–1441. <https://doi.org/10.1111/1365-2435.14034>
- Fan, Y., & Miguez-Macho, G. (2010). Potential groundwater contribution to Amazon evapotranspiration. *Hydrology and Earth System Sciences*, 14(10), 2039–2056. <https://doi.org/10.5194/hess-14-2039-2010>
- Feldman, A. F., Short Gianotti, D. J., Dong, J., Akbar, R., Crow, W. T., McColl, K. A., et al. (2023). Remotely sensed soil moisture can capture dynamics relevant to plant water uptake. *Water Resources Research*, 59(2), e2022WR033814. <https://doi.org/10.1029/2022WR033814>
- Feng, T. F., Shen, Y., Wang, F., Chen, Q., & Ji, K. (2023). Spatiotemporal variability and driving factors of the shallow soil moisture in North China during the past 31 years. *Journal of Hydrology*, 619, 129331. <https://doi.org/10.1016/j.jhydrol.2023.129331>
- Fischer, E. M., Seneviratne, S. I., Lüthi, D., & Schär, C. (2007). Contribution of land-atmosphere coupling to recent European summer heat waves. *Geophysical Research Letters*, 34(6). <https://doi.org/10.1029/2006gl029068>
- Fischer, E. M., Seneviratne, S. I., Vidale, P. L., Lüthi, D., & Schär, C. (2007). Soil moisture - Atmosphere interactions during the 2003 European summer heat wave. *Journal of Climate*, 20(20), 5081–5099. <https://doi.org/10.1175/Jcli4288.1>

- Fisher, R. A., & Koven, C. D. (2020). Perspectives on the future of land surface models and the challenges of representing complex terrestrial systems. *Journal of Advances in Modeling Earth Systems*, 12(4), e2018MS001453. <https://doi.org/10.1029/2018MS001453>
- Fontaine, B., Louvet, S., & Roucou, P. (2007). Fluctuations in annual cycles and inter-seasonal memory in West Africa: Rainfall, soil moisture and heat fluxes. *Theoretical and Applied Climatology*, 88(1–2), 57–70. <https://doi.org/10.1007/s00704-006-0246-4>
- Frankignoul, C., & Hasselmann, K. (1977). Stochastic climate models, Part II Application to sea-surface temperature anomalies and thermocline variability. *Tellus*, 29(4), 289–305. <https://doi.org/10.1111/j.2153-3490.1977.tb00740.x>
- Ganeshi, N. G., Mujumdar, M., Takaya, Y., Goswami, M. M., Singh, B. B., Krishnan, R., & Terao, T. (2023). Soil moisture revamps the temperature extremes in a warming climate over India. *npj Climate and Atmospheric Science*, 6(1), 12. <https://doi.org/10.1038/s41612-023-00334-1>
- Gao, C. J., Chen, H., Sun, S., Ongoma, V., Hua, W., Ma, H., et al. (2018). A potential predictor of multi-season droughts in Southwest China: Soil moisture and its memory. *Natural Hazards*, 91(2), 553–566. <https://doi.org/10.1007/s11069-017-3140-8>
- Garcia Gonzalez, R., Verhoef, A., Luigi Vidale, P., & Braud, I. (2012). Incorporation of water vapor transfer in the JULES land surface model: Implications for key soil variables and land surface fluxes. *Water Resources Research*, 48(5). <https://doi.org/10.1029/2011WR011811>
- Gebler, S., Hendricks Franssen, H. J., Pütz, T., Post, H., Schmidt, M., & Vereecken, H. (2015). Actual evapotranspiration and precipitation measured by lysimeters: A comparison with eddy covariance and tipping bucket. *Hydrology and Earth System Sciences*, 19(5), 2145–2161. <https://doi.org/10.5194/hess-19-2145-2015>
- Gebrehiwot, K. A. (2018). A review on waterlogging, salinization and drainage in Ethiopian irrigated agriculture. *Sustainable Water Resources Management*, 4(1), 55–62. <https://doi.org/10.1007/s40899-017-0121-8>
- Georgescu, M., Weaver, C. P., Avissar, R., Walko, R. L., & Miguez-Macho, G. (2003). Sensitivity of model-simulated summertime precipitation over the Mississippi River Basin to the spatial distribution of initial soil moisture. *Journal of Geophysical Research*, 108(D22). <https://doi.org/10.1029/2002jd003107>
- Gettelman, A., Geer, A. J., Forbes, R. M., Carmichael, G. R., Feingold, G., Posselt, D. J., et al. (2022). The future of Earth system prediction: Advances in model-data fusion. *Science Advances*, 8(14), eabn3488. <https://doi.org/10.1126/sciadv.abn3488>
- Ghannam, K., Nakai, T., Paschalis, A., Oishi, C. A., Kotani, A., Igarashi, Y., et al. (2016). Persistence and memory timescales in root-zone soil moisture dynamics. *Water Resources Research*, 52(2), 1427–1445. <https://doi.org/10.1002/2015wr017983>
- Giles, J. A., Menéndez, C. G., & Ruscica, R. C. (2023). Nonlocal impacts of soil moisture variability in South America: Linking two land-atmosphere coupling hot spots. *Journal of Climate*, 36(1), 227–242. <https://doi.org/10.1175/JCLI-D-21-0510.1>
- Good, S. P., Noone, D., & Bowen, G. (2015). WATER RESOURCES. Hydrologic connectivity constrains partitioning of global terrestrial water fluxes. *Science*, 349(6244), 175–177. <https://doi.org/10.1126/science.aaa5931>
- Green, J. K., Seneviratne, S. I., Berg, A. M., Findell, K. L., Hagemann, S., Lawrence, D. M., & Gentile, P. (2019). Large influence of soil moisture on long-term terrestrial carbon uptake. *Nature*, 565(7740), 476–479. <https://doi.org/10.1038/s41586-018-0848-x>
- Groh, J., Slawitsch, V., Herndl, M., Graf, A., Vereecken, H., & Pütz, T. (2018). Determining dew and hoar frost formation for a low mountain range and alpine grassland site by weighable lysimeter. *Journal of Hydrology*, 563, 372–381. <https://doi.org/10.1016/j.jhydrol.2018.06.009>
- Groh, J., Vanderborght, J., Pütz, T., & Vereecken, H. (2016). How to control the lysimeter bottom boundary to investigate the effect of climate change on soil processes? *Vadose Zone Journal*, 15(7), 1–15. <https://doi.org/10.2136/vzj2015.08.0113>
- Groh, J., Vanderborght, J., Pütz, T., Vogel, H. J., Gründling, R., Rupp, H., et al. (2020). Responses of soil water storage and crop water use efficiency to changing climatic conditions: A lysimeter-based space-for-time approach. *Hydrology and Earth System Sciences*, 24(3), 1211–1225. <https://doi.org/10.5194/hess-24-1211-2020>
- Guo, D., Mou, P., Jones, R. H., & Mitchell, R. J. (2002). Temporal changes in spatial patterns of soil moisture following disturbance: An experimental approach. *Journal of Ecology*, 90(2), 338–347. <https://doi.org/10.1046/j.1365-2745.2001.00667.x>
- Haarsma, R. J., Selten, F., Hurk, B. v., Hazeleger, W., & Wang, X. (2009). Drier Mediterranean soils due to greenhouse warming bring easterly winds over summertime central Europe. *Geophysical Research Letters*, 36(4). <https://doi.org/10.1029/2008gl036617>
- Hagemann, S., & Stacke, T. (2015). Impact of the soil hydrology scheme on simulated soil moisture memory. *Climate Dynamics*, 44(7–8), 1731–1750. <https://doi.org/10.1007/s00382-014-2221-6>
- Haghighi, E., Short Gianotti, D. J., Akbar, R., Salvucci, G. D., & Entekhabi, D. (2018). Soil and atmospheric controls on the land surface energy balance: A generalized framework for distinguishing moisture-limited and energy-limited evaporation regimes. *Water Resources Research*, 54(3), 1831–1851. <https://doi.org/10.1002/2017wr021729>
- Hanisch, S., Lohrey, C., & Buerkert, A. (2015). Dewfall and its ecological significance in semi-arid coastal south-western Madagascar. *Journal of Arid Environments*, 121, 24–31. <https://doi.org/10.1016/j.jaridenv.2015.05.007>
- Hassanpour, R., Zarehaghi, D., Neyshabouri, M. R., Feizizadeh, B., & Rahmati, M. (2020). Modification on optical trapezoid model for accurate estimation of soil moisture content in a maize growing field. *Journal of Applied Remote Sensing*, 14(3), 034519. <https://doi.org/10.1117/1.Jrs.14.034519>
- Hasselmann, K. (1976). Stochastic climate models part I. Theory. *Tellus*, 28(6), 473–485. <https://doi.org/10.3402/tellusa.v28i6.11316>
- Hawkes, C. V., Waring, B. G., Rocca, J. D., & Kivlin, S. N. (2017). Historical climate controls soil respiration responses to current soil moisture. *Proceedings of the National Academy of Sciences of the USA*, 114(24), 6322–6327. <https://doi.org/10.1073/pnas.1620811114>
- He, Q., Lu, H., & Yang, K. (2023). Soil moisture memory of land surface models utilized in major reanalyses differ significantly from SMAP observation. *Earth's Future*, 11(4). <https://doi.org/10.1029/2022EF003215>
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., et al. (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730), 1999–2049. <https://doi.org/10.1002/qj.3803>
- Hesslerová, P., et al. (2019). Wetlands and forests regulate climate via evapotranspiration. In S. An & J. Verhoeven (Eds.), *Wetlands: Ecosystem services, restoration and wise use* (pp. 63–93). Springer. https://doi.org/10.1007/978-3-030-14861-4_4
- Hirschi, M., Mueller, B., Dorigo, W., & Seneviratne, S. (2014). Using remotely sensed soil moisture for land-atmosphere coupling diagnostics: The role of surface vs. root-zone soil moisture variability. *Remote Sensing of Environment*, 154, 246–252. <https://doi.org/10.1016/j.rse.2014.08.030>
- Hirschi, M., Seneviratne, S. I., Alexandrov, V., Boberg, F., Boroneant, C., Christensen, O. B., et al. (2011). Observational evidence for soil-moisture impact on hot extremes in southeastern Europe. *Nature Geoscience*, 4(1), 17–21. <https://doi.org/10.1038/Ngeo1032>
- Hong, S. Y., & Kalnay, E. (2000). Role of sea surface temperature and soil-moisture feedback in the 1998 Oklahoma-Texas drought. *Nature*, 408(6814), 842–844. <https://doi.org/10.1038/35048548>
- Hou, X. L., Yang, H., Cao, J., Feng, W., & Zhang, Y. (2023). A review of advances in groundwater evapotranspiration research. *Water*, 15(5), 969. <https://doi.org/10.3390/w15050969>
- Humphrey, V., Berg, A., Ciais, P., Gentile, P., Jung, M., Reichstein, M., et al. (2021). Soil moisture-atmosphere feedback dominates land carbon uptake variability. *Nature*, 592(7852), 65–69. <https://doi.org/10.1038/s41586-021-03325-5>

- Jackson, R. B., Sperry, J. S., & Dawson, T. E. (2000). Root water uptake and transport: Using physiological processes in global predictions. *Trends in Plant Science*, 5(11), 482–488. [https://doi.org/10.1016/S1360-1385\(00\)01766-0](https://doi.org/10.1016/S1360-1385(00)01766-0)
- Jackson, T., et al. (2016). Calibration and validation for the L2/3_SM_P version 3 data products, SMAP project, JPL D-93720. In *Jet propulsion laboratory*.
- Jacobs, E. M., Bertassello, L. E., & Rao, P. S. C. (2020). Drivers of regional soil water storage memory and persistence. *Vadose Zone Journal*, 19(1), e20050. <https://doi.org/10.1002/vzj2.20050>
- Jaeger, E. B., & Seneviratne, S. I. (2011). Impact of soil moisture-atmosphere coupling on European climate extremes and trends in a regional climate model. *Climate Dynamics*, 36(9–10), 1919–1939. <https://doi.org/10.1007/s00382-010-0780-8>
- Ji, Y. D., Li, Y., Yao, N., Biswas, A., Zou, Y., Meng, Q., & Liu, F. (2021). The lagged effect and impact of soil moisture drought on terrestrial ecosystem water use efficiency. *Ecological Indicators*, 133, 108349. <https://doi.org/10.1016/j.ecolind.2021.108349>
- Katul, G. G., Oren, R., Manzoni, S., Higgins, C., & Parlange, M. B. (2012). Evapotranspiration: A process driving mass transport and energy exchange in the soil-plant-atmosphere-climate system. *Reviews of Geophysics*, 50(3). <https://doi.org/10.1029/2011rg000366>
- Katul, G. G., Porporato, A., Daly, E., Oishi, A. C., Kim, H., Stoy, P. C., et al. (2007). On the spectrum of soil moisture from hourly to interannual scales. *Water Resources Research*, 43(5). <https://doi.org/10.1029/2006wr005356>
- Katz, R. W. (1978). Persistence of subtropical African droughts. *Monthly Weather Review*, 106(7), 1017–1021. [https://doi.org/10.1175/1520-0493\(1978\)106%3C1017:POSAD%3E2.0.CO;2](https://doi.org/10.1175/1520-0493(1978)106%3C1017:POSAD%3E2.0.CO;2)
- Kenkre, V. M. (2021). The memory function formalism: What and why, memory functions, projection operators, and the defect technique. *Lecture Notes in Physics*, 982, 1–20. https://doi.org/10.1007/978-3-030-68667-3_1
- Keune, J., Sulis, M., & Kollet, S. J. (2019). Potential added value of incorporating human water use on the simulation of evapotranspiration and precipitation in a continental-scale bedrock-to-atmosphere modeling system: A validation study considering observational uncertainty. *Journal of Advances in Modeling Earth Systems*, 11(7), 1959–1980. <https://doi.org/10.1029/2019ms001657>
- Khalighi, M., Sommeria-Klein, G., Gonze, D., Faust, K., & Lahti, L. (2022). Quantifying the impact of ecological memory on the dynamics of interacting communities. *PLoS Computational Biology*, 18(6), e1009396. <https://doi.org/10.1371/journal.pcbi.1009396>
- Khalil, M. M., Abotalib, A. Z., Farag, M. H., Rabei, M., Abdelhady, A. A., & Pichler, T. (2021). Poor drainage-induced waterlogging in Saharan groundwater-irrigated lands: Integration of geospatial, geophysical, and hydrogeological techniques. *Catena*, 207, 105615. <https://doi.org/10.1016/j.catena.2021.105615>
- Kim, H., & Lakshmi, V. (2019). Global dynamics of stored precipitation water in the topsoil layer from satellite and reanalysis data. *Water Resources Research*, 55(4), 3328–3346. <https://doi.org/10.1029/2018wr023166>
- Koster, R. D., Chang, Y., & Schubert, S. D. (2014). A mechanism for land-atmosphere feedback involving planetary wave structures. *Journal of Climate*, 27(24), 9290–9301. <https://doi.org/10.1175/Jcli-D-14-00315.1>
- Koster, R. D., Chang, Y., Wang, H., & Schubert, S. D. (2016). Impacts of local soil moisture anomalies on the atmospheric circulation and on remote surface meteorological fields during Boreal summer: A comprehensive analysis over North America. *Journal of Climate*, 29(20), 7345–7364. <https://doi.org/10.1175/Jcli-D-16-0192.1>
- Koster, R. D., Dirmeyer, P. A., Guo, Z., Bonan, G., Chan, E., Cox, P., et al. (2004). Regions of strong coupling between soil moisture and precipitation. *Science*, 305(5687), 1138–1140. <https://doi.org/10.1126/science.1100217>
- Koster, R. D., Guo, Z., Yang, R., Dirmeyer, P. A., Mitchell, K., & Puma, M. J. (2009). On the nature of soil moisture in land surface models. *Journal of Climate*, 22(16), 4322–4335. <https://doi.org/10.1175/2009jcli2832.1>
- Koster, R. D., Liu, Q., Mahanama, S. P. P., & Reichle, R. H. (2018). Improved hydrological simulation using SMAP data: Relative impacts of model calibration and data assimilation. *Journal of Hydrometeorology*, 19(4), 727–741. <https://doi.org/10.1175/JHM-D-17-0228.1>
- Koster, R. D., Mahanama, S. P. P., Yamada, T. J., Balsamo, G., Berg, A. A., Boisserie, M., et al. (2010). Contribution of land surface initialization to subseasonal forecast skill: First results from a multi-model experiment. *Geophysical Research Letters*, 37(2). <https://doi.org/10.1029/2009gl041677>
- Koster, R. D., Mahanama, S. P. P., Yamada, T. J., Balsamo, G., Berg, A. A., Boisserie, M., et al. (2011). The second phase of the global land-atmosphere coupling experiment: Soil moisture contributions to subseasonal forecast skill. *Journal of Hydrometeorology*, 12(5), 805–822. <https://doi.org/10.1175/2011jhm1365.1>
- Koster, R. D., & Milly, P. (1997). The interplay between transpiration and runoff formulations in land surface schemes used with atmospheric models. *Journal of Climate*, 10(7), 1578–1591. [https://doi.org/10.1175/1520-0442\(1997\)010%3C1578:TIBTAR%3E2.0.CO;2](https://doi.org/10.1175/1520-0442(1997)010%3C1578:TIBTAR%3E2.0.CO;2)
- Koster, R. D., & Suarez, M. J. (2001). Soil moisture memory in climate models. *Journal of Hydrometeorology*, 2(6), 558–570. [https://doi.org/10.1175/1525-7541\(2001\)002<0558:Smmicm>2.0.Co;2](https://doi.org/10.1175/1525-7541(2001)002<0558:Smmicm>2.0.Co;2)
- Koster, R. D., Suarez, M. J., & Heiser, M. (2000). Variance and predictability of precipitation at seasonal-to-interannual timescales. *Journal of Hydrometeorology*, 1(1), 26–46. [https://doi.org/10.1175/1525-7541\(2000\)001<0026:Vapopa>2.0.Co;2](https://doi.org/10.1175/1525-7541(2000)001<0026:Vapopa>2.0.Co;2)
- Koster, R. D., Suarez, M. J., Higgins, R. W., & Van den Dool, H. M. (2003). Observational evidence that soil moisture variations affect precipitation. *Geophysical Research Letters*, 30(5). <https://doi.org/10.1029/2002gl016571>
- Koster, R. D., Sud, Y. C., Guo, Z., Dirmeyer, P. A., Bonan, G., Oleson, K. W., et al. (2006). Glace: The global land-atmosphere coupling experiment. *Journal of Hydrometeorology*, 7(4), 590–610. <https://doi.org/10.1175/JHM510.1>
- Kraus, E. B. (1977). Subtropical droughts and cross-equatorial energy transports. *Monthly Weather Review*, 105(8), 1009–1018. [https://doi.org/10.1175/1520-0493\(1977\)105<1009:Sdacee>2.0.Co;2](https://doi.org/10.1175/1520-0493(1977)105<1009:Sdacee>2.0.Co;2)
- Kumar, S. V., Dirmeyer, P. A., Peters-Lidard, C. D., Bindlish, R., & Bolten, J. (2018). Information theoretic evaluation of satellite soil moisture retrievals. *Remote Sensing of Environment*, 204, 392–400. <https://doi.org/10.1016/j.rse.2017.10.016>
- Laio, F., Porporato, A., Fernandez-Illescas, C., & Rodriguez-Iturbe, I. (2001). Plants in water-controlled ecosystems: Active role in hydrologic processes and response to water stress: IV. Discussion of real cases. *Advances in Water Resources*, 24(7), 745–762. [https://doi.org/10.1016/S0309-1708\(01\)00007-0](https://doi.org/10.1016/S0309-1708(01)00007-0)
- Langford, S., Stevenson, S., & Noone, D. (2014). Analysis of low-frequency precipitation variability in CMIP5 historical simulations for southwestern North America. *Journal of Climate*, 27(7), 2735–2756. <https://doi.org/10.1175/Jcli-D-13-00317.1>
- Lees, T., Reece, S., Kratzert, F., Klotz, D., Gauch, M., De Bruijn, J., et al. (2021). Hydrological concept formation inside long short-term memory (LSTM) networks. *Hydrology and Earth System Sciences Discussions*, 2021(12), 1–37. <https://doi.org/10.5194/hess-26-3079-2022>
- Liang, M. L., & Yuan, X. (2021). Critical role of soil moisture memory in predicting the 2012 Central United States flash drought. *Frontiers in Earth Science*, 9, 615969. <https://doi.org/10.3389/feart.2021.615969>
- Liu, D., Wang, G., Mei, R., Yu, Z., & Yu, M. (2014). Impact of initial soil moisture anomalies on climate mean and extremes over Asia. *Journal of Geophysical Research-Atmospheres*, 119(2), 529–545. <https://doi.org/10.1002/2013jd020890>
- Liu, Y. Q., & Avissar, R. (1999). A study of persistence in the land-atmosphere system using a general circulation model and observations. *Journal of Climate*, 12(8), 2139–2153. [https://doi.org/10.1175/1520-0442\(1999\)012<2139:ASOPT>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<2139:ASOPT>2.0.CO;2)

- Lloret, F., & Zedler, P. H. (2009). The effect of forest fire on vegetation, In *Fire effects on soils and restoration strategies*. (pp. 273–312). CRC Press. <https://doi.org/10.1201/9781439843338-c9>
- Lorenz, R., Jaeger, E. B., & Seneviratne, S. I. (2010). Persistence of heat waves and its link to soil moisture memory. *Geophysical Research Letters*, 37(9). <https://doi.org/10.1029/2010gl042764>
- Ma, Y. L., Montzka, C., Bayat, B., & Kollet, S. (2021). Using Long Short-Term Memory networks to connect water table depth anomalies to precipitation anomalies over Europe. *Hydrology and Earth System Sciences*, 25(6), 3555–3575. <https://doi.org/10.5194/hess-25-3555-2021>
- MacDonald, L. H., & Huffman, E. L. (2004). Post-fire soil water repellency: Persistence and soil moisture thresholds. *Soil Science Society of America Journal*, 68(5), 1729–1734. <https://doi.org/10.2136/sssaj2004.1729>
- MacLeod, D., Cloke, H., Pappenberger, F., & Weisheimer, A. (2016). Evaluating uncertainty in estimates of soil moisture memory with a reverse ensemble approach. *Hydrology and Earth System Sciences*, 20(7), 2737–2743. <https://doi.org/10.5194/hess-20-2737-2016>
- Mahanama, S. P. P., & Koster, R. D. (2003). Intercomparison of soil moisture memory in two land surface models. *Journal of Hydrometeorology*, 4(6), 1134–1146. [https://doi.org/10.1175/1525-7541\(2003\)004<1134:losmmi>2.0.Co;2](https://doi.org/10.1175/1525-7541(2003)004<1134:losmmi>2.0.Co;2)
- MahfuzurRahman, M., & Lu, M. (2015). Characterizing soil moisture memory by soil moisture autocorrelation. *Journal of Water Resource and Hydraulic Engineering*, 3(1), 85–94. <https://doi.org/10.5963/JWRHE0401007>
- Mahowald, N., Lo, F., Zheng, Y., Harrison, L., Funk, C., & Lombardozzi, D. (2015). Leaf area index in Earth system models: Evaluation and projections. *Earth Syst. Dyn. Discuss*, 6, 761–818. <https://doi.org/10.5194/esdd-6-761-2015>
- Makarieva, A. M., & Gorshkov, V. G. (2007). Biotic pump of atmospheric moisture as driver of the hydrological cycle on land. *Hydrology and Earth System Sciences*, 11(2), 1013–1033. <https://doi.org/10.5194/hess-11-1013-2007>
- Makarieva, A. M., Gorshkov, V. G., Sheil, D., Nobre, A. D., & Li, B. L. (2013). Where do winds come from? A new theory on how water vapor condensation influences atmospheric pressure and dynamics. *Atmospheric Chemistry and Physics*, 13(2), 1039–1056. <https://doi.org/10.5194/acp-13-1039-2013>
- Mao, T. N., Shangguan, W., Li, Q., Li, L., Zhang, Y., Huang, F., et al. (2022). A spatial downscaling method for remote sensing soil moisture based on random forest considering soil moisture memory and mass conservation. *Remote Sensing*, 14(16), 3858. <https://doi.org/10.3390/rs14163858>
- Mariotti, A., Baggett, C., Barnes, E. A., Becker, E., Butler, A., Collins, D. C., et al. (2020). Windows of opportunity for skillful forecasts subseasonal to seasonal and beyond. *Bulletin of the American Meteorological Society*, 101(5), E608–E625. <https://doi.org/10.1175/Bams-D-18-0326.1>
- Martens, B., Miralles, D. G., Lievens, H., van der Schalie, R., de Jeu, R. A. M., Fernández-Prieto, D., et al. (2017). GLEAM v3: Satellite-based land evaporation and root-zone soil moisture. *Geoscientific Model Development*, 10(5), 1903–1925. <https://doi.org/10.5194/gmd-10-1903-2017>
- Martínez-de la Torre, A., & Míguez-Macho, G. (2019). Groundwater influence on soil moisture memory and land-atmosphere fluxes in the Iberian Peninsula. *Hydrology and Earth System Sciences*, 23(12), 4909–4932. <https://doi.org/10.5194/hess-23-4909-2019>
- Martínez-Fernández, J., González-Zamora, A., & Almendra-Martín, L. (2021). Soil moisture memory and soil properties: An analysis with the stored precipitation fraction. *Journal of Hydrology*, 593, 125622. <https://doi.org/10.1016/j.jhydrol.2020.125622>
- McColl, K. A., Alemohammad, S. H., Akbar, R., Konings, A. G., Yueh, S., & Entekhabi, D. (2017). The global distribution and dynamics of surface soil moisture. *Nature Geoscience*, 10(2), 100–104. <https://doi.org/10.1038/Ngeo2868>
- McColl, K. A., He, Q., Lu, H., & Entekhabi, D. (2019). Short-term and long-term surface soil moisture memory time scales are spatially anti-correlated at global scales. *Journal of Hydrometeorology*, 20(6), 1165–1182. <https://doi.org/10.1175/Jhm-D-18-0141.1>
- McColl, K. A., Wang, W., Peng, B., Akbar, R., Short Gianotti, D. J., Lu, H., et al. (2017). Global characterization of surface soil moisture drydowns. *Geophysical Research Letters*, 44(8), 3682–3690. <https://doi.org/10.1002/2017gl072819>
- Meng, C. L., Jin, H., & Jin, B. (2023). Parameterization of soil evaporation and coupled transport of moisture and heat for arid and semiarid regions. *Frontiers in Earth Science*, 11, 1151405. <https://doi.org/10.3389/feart.2023.1151405>
- Milly, P. C. D. (1984). A simulation analysis of thermal effects on evaporation from soil. *Water Resources Research*, 20(8), 1087–1098. <https://doi.org/10.1029/WR020i008p01087>
- Miralles, D. G., Gentile, P., Seneviratne, S. I., & Teuling, A. J. (2019). Land-atmospheric feedbacks during droughts and heatwaves: State of the science and current challenges. *Annals of the New York Academy of Sciences*, 1436(1), 19–35. <https://doi.org/10.1111/nyas.13912>
- Miralles, D. G., Holmes, T. R. H., De Jeu, R. A. M., Gash, J. H., Meesters, A. G. C. A., & Dolman, A. J. (2011). Global land-surface evaporation estimated from satellite-based observations. *Hydrology and Earth System Sciences*, 15(2), 453–469. <https://doi.org/10.5194/hess-15-453-2011>
- Miralles, D. G., Teuling, A. J., van Heerwaarden, C. C., & Vilà-Guerau de Arellano, J. (2014). Mega-heatwave temperatures due to combined soil desiccation and atmospheric heat accumulation. *Nature Geoscience*, 7(5), 345–349. <https://doi.org/10.1038/Ngeo2141>
- Miralles, D. G., van den Berg, M. J., Teuling, A. J., & de Jeu, R. A. M. (2012). Soil moisture-temperature coupling: A multiscale observational analysis. *Geophysical Research Letters*, 39(21). <https://doi.org/10.1029/2012gl053703>
- Mohanty, B. P., Cosh, M. H., Lakshmi, V., & Montzka, C. (2017). Soil moisture remote sensing: State-of-the-Science. *Vadose Zone Journal*, 16(1), 1–9. <https://doi.org/10.2136/vzj2016.10.0105>
- Montzka, C., Bogena, H., Zreda, M., Monerris, A., Morrison, R., Muddu, S., & Vereecken, H. (2017). Validation of spaceborne and modelled surface soil moisture products with cosmic-ray neutron probes. *Remote Sensing*, 9(2), 103. <https://doi.org/10.3390/rs9020103>
- Mueller, B., & Seneviratne, S. I. (2012). Hot days induced by precipitation deficits at the global scale. *Proceedings of the National Academy of Sciences of the United States of America*, 109(31), 12398–12403. <https://doi.org/10.1073/pnas.1204330109>
- Nakai, T., Katul, G. G., Kotani, A., Igarashi, Y., Ohta, T., Suzuki, M., & Kumagai, T. (2014). Radiative and precipitation controls on root zone soil moisture spectra. *Geophysical Research Letters*, 41(21), 7546–7554. <https://doi.org/10.1002/2014gl061745>
- Namias, J. (1959). Persistence of mid-tropospheric circulations between adjacent months and seasons, the atmosphere and the sea in motion. 240, 248.
- Namias, J. (1960). *Factors in the initiation, perpetuation and termination of drought* (Vol. 51, pp. 81–94). International Association of Scientific Hydrology Commission on Surface Waters Publication.
- Namias, J. (1963). Surface-atmosphere interactions as fundamental causes of droughts and other climatic fluctuations. *Arid Zone Research*, 20, 345–359.
- Nandintsetseg, B., & Shinoda, M. (2014). Multi-decadal soil moisture trends in Mongolia and their relationships to precipitation and evapo-transpiration. *Arid Land Research and Management*, 28(3), 247–260. <https://doi.org/10.1080/15324982.2013.861882>
- Nicholson, S. (2000). Land surface processes and Sahel climate. *Reviews of Geophysics*, 38(1), 117–139. <https://doi.org/10.1029/1999rg900014>
- Nicolai-Shaw, N., Gudmundsson, L., Hirschi, M., & Seneviratne, S. I. (2016). Long-term predictability of soil moisture dynamics at the global scale: Persistence versus large-scale drivers. *Geophysical Research Letters*, 43(16), 8554–8562. <https://doi.org/10.1002/2016GL069847>

- Notaro, M. (2008). Statistical identification of global hot spots in soil moisture feedbacks among IPCC AR4 models. *Journal of Geophysical Research*, 113(D9). <https://doi.org/10.1029/2007JD009199>
- Oladipo, E. O., & Hare, F. K. (1986). Persistence of anomalous moisture conditions in the interior Plains of North-America. *Journal of Climatology*, 6(5), 485–494. <https://doi.org/10.1002/joc.3370060504>
- Orlowsky, B., & Seneviratne, S. I. (2010). Statistical analyses of land-atmosphere feedbacks and their possible pitfalls. *Journal of Climate*, 23(14), 3918–3932. <https://doi.org/10.1175/2010JCLI3366.1>
- Orth, R., Koster, R. D., & Seneviratne, S. I. (2013). Inferring soil moisture memory from streamflow observations using a simple water balance model. *Journal of Hydrometeorology*, 14(6), 1773–1790. <https://doi.org/10.1175/Jhm-D-12-099.1>
- Orth, R., & Seneviratne, S. I. (2012). Analysis of soil moisture memory from observations in Europe. *Journal of Geophysical Research*, 117(D15). <https://doi.org/10.1029/2011Jd017366>
- Orth, R., & Seneviratne, S. I. (2013). Propagation of soil moisture memory to streamflow and evapotranspiration in Europe. *Hydrology and Earth System Sciences*, 17(10), 3895–3911. <https://doi.org/10.5194/hess-17-3895-2013>
- Otkin, J. A., Svoboda, M., Hunt, E. D., Ford, T. W., Anderson, M. C., Hain, C., & Basara, J. B. (2018). FLASH DROUGHTS A review and assessment of the challenges imposed by rapid-onset droughts in the United States. *Bulletin of the American Meteorological Society*, 99(5), 911–919. <https://doi.org/10.1175/Bams-D-17-0149.1>
- Owe, M., & Van de Griend, A. A. (1998). Comparison of soil moisture penetration depths for several bare soils at two microwave frequencies and implications for remote sensing. *Water Resources Research*, 34(9), 2319–2327. <https://doi.org/10.1029/98wr01469>
- Oyeogbe, A., & Oluwasemire, K. (2013). Evaluation of SOILWAT model for predicting soil water characteristics in southwestern Nigeria. *International Journal of Soil Science*, 8(2), 58–67. <https://doi.org/10.3923/ijss.2013.58.67>
- Pal, J. S., & Eltahir, E. A. B. (2001). Pathways relating soil moisture conditions to future summer rainfall within a model of the land-atmosphere system. *Journal of Climate*, 14(6), 1227–1242. [https://doi.org/10.1175/1520-0442\(2001\)014<1227:Prsmct>2.0.Co;2](https://doi.org/10.1175/1520-0442(2001)014<1227:Prsmct>2.0.Co;2)
- Pal, J. S., & Eltahir, E. A. B. (2002). Teleconnections of soil moisture and rainfall during the 1993 midwest summer flood. *Geophysical Research Letters*, 29(18), 12-11–12-14. <https://doi.org/10.1029/2002gl014815>
- Pal, J. S., & Eltahir, E. A. B. (2003). A feedback mechanism between soil-moisture distribution and storm tracks. *Quarterly Journal of the Royal Meteorological Society*, 129(592), 2279–2297. <https://doi.org/10.1256/qj.01.201>
- Pal, M., & Maity, R. (2019). Development of a spatially-varying Statistical Soil Moisture Profile model by coupling memory and forcing using hydrologic soil groups. *Journal of Hydrology*, 570, 141–155. <https://doi.org/10.1016/j.jhydrol.2018.12.042>
- Pal, M., Maity, R., & Dey, S. (2016). Statistical modelling of vertical soil moisture profile: Coupling of memory and forcing. *Water Resources Management*, 30(6), 1973–1986. <https://doi.org/10.1007/s11269-016-1263-4>
- Peili, S., & Wenhua, L. (2001). Influence of forest cover change on hydrological process and watershed runoff. *Journal of Natural Resources*, 16(5), 481–487. <https://doi.org/10.11849/zrzyxb.2001.05.014>
- Perlekar, P., Ray, S. S., Mitra, D., & Pandit, R. (2011). Persistence problem in two-dimensional fluid turbulence. *Physical Review Letters*, 106(5), 054501. <https://doi.org/10.1103/PhysRevLett.106.054501>
- Porporato, A., Laio, F., Ridolfi, L., & Rodriguez-Iturbe, I. (2001). Plants in water-controlled ecosystems: Active role in hydrologic processes and response to water stress: III. Vegetation water stress. *Advances in Water Resources*, 24(7), 725–744. [https://doi.org/10.1016/S0309-1708\(01\)00006-9](https://doi.org/10.1016/S0309-1708(01)00006-9)
- Preimesberger, W., Scanlon, T., Su, C. H., Gruber, A., & Dorigo, W. (2020). Homogenization of structural breaks in the global ESA CCI soil moisture multisatellite climate data record. *IEEE Transactions on Geoscience and Remote Sensing*, 59(4), 2845–2862. <https://doi.org/10.1109/tgrs.2020.3012896>
- Quesada, B., Vautard, R., Yiou, P., Hirschi, M., & Seneviratne, S. I. (2012). Asymmetric European summer heat predictability from wet and dry southern winters and springs. *Nature Climate Change*, 2(10), 736–741. <https://doi.org/10.1038/Nclimate1536>
- Quiring, S. M., Ford, T. W., Wang, J. K., Khong, A., Harris, E., Lindgren, T., et al. (2016). THE NORTH AMERICAN SOIL MOISTURE DATABASE development and applications. *Bulletin of the American Meteorological Society*, 97(8), 1441–1459. <https://doi.org/10.1175/Bams-D-13-00263.1>
- Rahman, M. M., Lu, M., & Kyi, K. H. (2015). Variability of soil moisture memory for wet and dry basins. *Journal of Hydrology*, 523, 107–118. <https://doi.org/10.1016/j.jhydrol.2015.01.033>
- Rahmati, M., Graf, A., Bayat, B., Montzka, C., Poppe, C., Vanderborght, J., et al. (2023). Lagged correlation between soil water content and evapotranspiration along recent decades and across different land cover types of Europe. In *EGU general assembly 2023*. Copernicus Meetings. <https://doi.org/10.5194/egusphere-egu23-3771>
- Rahmati, M., Groh, J., Graf, A., Pütz, T., Vanderborght, J., & Vereecken, H. (2020). On the impact of increasing drought on the relationship between soil water content and evapotranspiration of a grassland. *Vadose Zone Journal*, 19(1), e20029. <https://doi.org/10.1002/vzj2.20029>
- Rahmati, M., Or, D., Amelung, W., Bauke, S. L., Bol, R., Hendricks Franssen, H. J., et al. (2023). Soil is a living archive of the Earth system. *Nature Reviews Earth & Environment*, 4(7), 421–423. <https://doi.org/10.1038/s43017-023-00454-5>
- Rahmati, M., Or, D., Amelung, W., & Vereecken, H. (2023). Soil moisture memory mitigates or amplifies drought effects. In *Second TERENO-OZCAR Conference, edited, French OZCAR and German TERENO research networks, Bonn, Germany*.
- Rahmati, M., Oskouei, M., Neyshabouri, M., Walker, J., Fakherifard, A., Ahmadi, A., & Mousavi, S. (2015). Soil moisture derivation using triangle method in the lighvan watershed, north western Iran. *Journal of Soil Science and Plant Nutrition*, 15(1), 167–178. <https://doi.org/10.4067/S0718-95162015005000014>
- Reichle, R. H., De Lannoy, G. J. M., Liu, Q., Ardizzone, J. V., Colliander, A., Conaty, A., et al. (2017). Assessment of the SMAP level-4 surface and root-zone soil moisture product using in situ measurements. *Journal of Hydrometeorology*, 18(10), 2621–2645. <https://doi.org/10.1175/Jhm-D-17-0063.1>
- Rind, D. (1982). The influence of ground moisture conditions in north-America on summer climate as modeled in the Giss Gcm. *Monthly Weather Review*, 110(10), 1487–1494. [https://doi.org/10.1175/1520-0493\(1982\)110<1487:Tiogmc>2.0.Co;2](https://doi.org/10.1175/1520-0493(1982)110<1487:Tiogmc>2.0.Co;2)
- Robertson, A., & Vitart, F. (2001). *Sub-seasonal to seasonal prediction: The gap between weather and climate forecasting*. Elsevier. <https://doi.org/10.1016/C2016-0-01594-2>
- Robock, A. (2003). HYDROLOGY | soil moisture. In J. R. Holton (Ed.), *Encyclopedia of atmospheric Sciences* (pp. 987–993). Academic Press. <https://doi.org/10.1016/b0-12-227090-8/00169-x>
- Robock, A., Vinnikov, K. Y., Schlosser, C. A., Speranskaya, N. A., & Xue, Y. (1995). Use of midlatitude soil-moisture and meteorological observations to validate soil-moisture simulations with biosphere and bucket models. *Journal of Climate*, 8(1), 15–35. [https://doi.org/10.1175/1520-0442\(1995\)008<0015:Uomsma>2.0.Co;2](https://doi.org/10.1175/1520-0442(1995)008<0015:Uomsma>2.0.Co;2)
- Robock, A., Vinnikov, K. Y., Srinivasan, G., Entin, J. K., Hollinger, S. E., Speranskaya, N. A., et al. (2000). The global soil moisture Data Bank. *Bulletin of the American Meteorological Society*, 81(6), 1281–1299. [https://doi.org/10.1175/1520-0477\(2000\)081<1281:Tgsmdb>2.3.Co;2](https://doi.org/10.1175/1520-0477(2000)081<1281:Tgsmdb>2.3.Co;2)

- Rodell, M., Houser, P. R., Peters-Lidard, C., Kato, H., Kumar, S., Gottschalk, J., et al. (2004). NASA/NOAA'S global land data assimilation system (GLDAS): Recent results and future plans. Paper presented at proceedings ECMWF/ELDAS Workshop on Land Surface Assimilation.
- Rodell, M., Houser, P. R., Jambor, U., Gottschalk, J., Mitchell, K., Meng, C. J., et al. (2004). The global land data assimilation system. *Bulletin of the American Meteorological Society*, 85(3), 381–394. <https://doi.org/10.1175/Bams-85-3-381>
- Rodriguez-Iturbe, I., Porporato, A., Laio, F., & Ridolfi, L. (2001). Plants in water-controlled ecosystems: Active role in hydrologic processes and response to water stress - I. Scope and general outline. *Advances in Water Resources*, 24(7), 695–705. [https://doi.org/10.1016/S0309-1708\(01\)00004-5](https://doi.org/10.1016/S0309-1708(01)00004-5)
- Rondinelli, W. J., Hornbuckle, B. K., Patton, J. C., Cosh, M. H., Walker, V. A., Carr, B. D., & Logsdon, S. D. (2015). Different rates of soil drying after rainfall are observed by the SMOS satellite and the south fork in situ soil moisture network. *Journal of Hydrometeorology*, 16(2), 889–903. <https://doi.org/10.1175/Jhm-D-14-0137.1>
- Rowntree, P. R., & Bolton, J. A. (1983). Simulation of the atmospheric response to soil-moisture anomalies over Europe. *Quarterly Journal of the Royal Meteorological Society*, 109(461), 501–526. <https://doi.org/10.1002/qj.49710946105>
- Rubol, S. (2010). *The influence of redox dynamics on nitrogen cycling and nitrous oxide emissions from soils*. University of.
- Ruscica, R. C., Sörensson, A. A., & Menéndez, C. G. (2014). Hydrological links in southeastern South America: Soil moisture memory and coupling within a hot spot. *International Journal of Climatology*, 34(14), 3641–3653. <https://doi.org/10.1002/joc.3930>
- Sangelantoni, L., Sobolowski, S., Lorenz, T., Hodnebrog, Ø., Cardoso, R. M., Soares, P. M. M., et al. (2023). Investigating the representation of heatwaves from an ensemble of km-scale regional climate simulations within CORDEX-FPS convection. *Climate Dynamics*, 1–37. <https://doi.org/10.1007/s00382-023-06769-9>
- Santanello Jr, J. A., Dirmeyer, P. A., Ferguson, C. R., Findell, K. L., Tawfik, A. B., Berg, A., et al. (2018). Land-atmosphere interactions: The LoCo perspective. *Bulletin of the American Meteorological Society*, 99(6), 1253–1272. <https://doi.org/10.1175/BAMS-D-17-0001.1>
- Schlesinger, W. H., & Jasechko, S. (2014). Transpiration in the global water cycle. *Agricultural and Forest Meteorology*, 189, 115–117. <https://doi.org/10.1016/j.agrformet.2014.01.011>
- Schlosser, C. A., & Milly, P. C. D. (2002). A model-based investigation of soil moisture predictability and associated climate predictability. *Journal of Hydrometeorology*, 3(4), 483–501. [https://doi.org/10.1175/1525-7541\(2002\)003<0483:Ambios>2.0.Co;2](https://doi.org/10.1175/1525-7541(2002)003<0483:Ambios>2.0.Co;2)
- Schnepper, T., Groh, J., Gerke, H. H., Reichert, B., & Pütz, T. (2023). Evaluation of precipitation measurement methods using data from a precision lysimeter network. *Hydrology and Earth System Sciences*, 27(17), 3265–3292. <https://doi.org/10.5194/hess-27-3265-2023>
- Sellers, P. J., Dickinson, R. E., Randall, D. A., Betts, A. K., Hall, F. G., Berry, J. A., et al. (1997). Modeling the exchanges of energy, water, and carbon between continents and the atmosphere. *Science*, 275(5299), 502–509. <https://doi.org/10.1126/science.275.5299.502>
- Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., et al. (2010). Investigating soil moisture-climate interactions in a changing climate: A review. *Earth-Science Reviews*, 99(3–4), 125–161. <https://doi.org/10.1016/j.earscirev.2010.02.004>
- Seneviratne, S. I., & Koster, R. D. (2012). A revised framework for analyzing soil moisture memory in climate data: Derivation and interpretation. *Journal of Hydrometeorology*, 13(1), 404–412. <https://doi.org/10.1175/Jhm-D-11-044.1>
- Seneviratne, S. I., Koster, R. D., Guo, Z., Dirmeyer, P. A., Kowalczyk, E., Lawrence, D., et al. (2006). Soil moisture memory in AGCM simulations: Analysis of global land-atmosphere coupling experiment (GLACE) data. *Journal of Hydrometeorology*, 7(5), 1090–1112. <https://doi.org/10.1175/Jhm533.1>
- Seneviratne, S. I., Lüthi, D., Litschi, M., & Schär, C. (2006). Land-atmosphere coupling and climate change in Europe. *Nature*, 443(7108), 205–209. <https://doi.org/10.1038/nature05095>
- Seneviratne, S. I., Wilhelm, M., Stanelle, T., van den Hurk, B., Hagemann, S., Berg, A., et al. (2013). Impact of soil moisture-climate feedbacks on CMIP5 projections: First results from the GLACE-CMIP5 experiment. *Geophysical Research Letters*, 40(19), 5212–5217. <https://doi.org/10.1002/grl.50956>
- Seo, E., & Dirmeyer, P. A. (2022). Improving the ESA CCI daily soil moisture time series with physically based land surface model datasets using a fourier time-filtering method. *Journal of Hydrometeorology*, 23(3), 473–489. <https://doi.org/10.1175/Jhm-D-21-0120.1>
- Shellito, P. J., Small, E. E., Colliander, A., Bindlish, R., Cosh, M. H., Berg, A. A., et al. (2016). SMAP soil moisture drying more rapid than observed in situ following rainfall events. *Geophysical Research Letters*, 43(15), 8068–8075. <https://doi.org/10.1002/2016gl069946>
- Shellito, P. J., Small, E. E., & Livneh, B. (2018). Controls on surface soil drying rates observed by SMAP and simulated by the Noah land surface model. *Hydrology and Earth System Sciences*, 22(3), 1649–1663. <https://doi.org/10.5194/hess-22-1649-2018>
- Shinoda, M. (2001). Climate memory of snow mass as soil moisture over central Eurasia. *Journal of Geophysical Research*, 106(D24), 33393–33403. <https://doi.org/10.1029/2001jd000525>
- Shinoda, M., & Nandintsetseg, B. (2011). Soil moisture and vegetation memories in a cold, arid climate. *Global and Planetary Change*, 79(1–2), 110–117. <https://doi.org/10.1016/j.gloplacha.2011.08.005>
- Shinoda, M., & Yamaguchi, Y. (2003). Influence of soil moisture anomaly on temperature in the Sahel: A comparison between wet and dry decades. *Journal of Hydrometeorology*, 4(2), 437–447. [https://doi.org/10.1175/1525-7541\(2003\)4<437:Iosmao>2.0.Co;2](https://doi.org/10.1175/1525-7541(2003)4<437:Iosmao>2.0.Co;2)
- Short Gianotti, D. J., Salvucci, G. D., Akbar, R., McColl, K. A., Cuenca, R., & Entekhabi, D. (2019). Landscape water storage and subsurface correlation from satellite surface soil moisture and precipitation observations. *Water Resources Research*, 55(11), 9111–9132. <https://doi.org/10.1029/2019WR025332>
- Shukla, J., & Mintz, Y. (1982). Influence of land-surface evapotranspiration on the Earth's climate. *Science*, 215(4539), 1498–1501. <https://doi.org/10.1126/science.215.4539.1498>
- Simmonds, I., & Hope, P. (1998). Seasonal and regional responses to changes in Australian soil moisture conditions. *International Journal of Climatology*, 18(10), 1105–1139. [https://doi.org/10.1002/\(Sici\)1097-0088\(199808\)18:10<1105::Aid-Joc308>3.0.Co;2-G](https://doi.org/10.1002/(Sici)1097-0088(199808)18:10<1105::Aid-Joc308>3.0.Co;2-G)
- Small, E., & Papuga, S. (2002). *The influence of soil moisture on the surface energy balance in semiarid environments*. AGU Spring Meeting Abstracts.
- Song, Y. M., Wang, Z. F., Qi, L. L., & Huang, A. N. (2019). Soil moisture memory and its effect on the surface water and heat fluxes on seasonal and interannual time scales. *Journal of Geophysical Research-Atmospheres*, 124(20), 10730–10741. <https://doi.org/10.1029/2019jd030893>
- Sörensson, A. A., & Berbery, E. H. (2015). A note on soil moisture memory and interactions with surface climate for different vegetation types in the La Plata basin. *Journal of Hydrometeorology*, 16(2), 716–729. <https://doi.org/10.1175/jhm-d-14-0102.1>
- Soulsby, C., Scheliga, B., Neill, A., Comte, J., & Tetzlaff, D. (2021). A longer-term perspective on soil moisture, groundwater and stream flow response to the 2018 drought in an experimental catchment in the Scottish Highlands. *Hydrological Processes*, 35(6), e14206. <https://doi.org/10.1002/hyp.14206>
- Soylu, M. E., & Bras, R. L. (2022). Global shallow groundwater patterns from soil moisture satellite retrievals. *Ieee Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 15, 89–101. <https://doi.org/10.1109/Jstars.2021.3124892>
- Soylu, M. E., Istanbuloglu, E., Lenters, J. D., & Wang, T. (2011). Quantifying the impact of groundwater depth on evapotranspiration in a semi-arid grassland region. *Hydrology and Earth System Sciences*, 15(3), 787–806. <https://doi.org/10.5194/hess-15-787-2011>

- Spracklen, D. V., Arnold, S. R., & Taylor, C. M. (2012). Observations of increased tropical rainfall preceded by air passage over forests. *Nature*, 489(7415), 282–285. <https://doi.org/10.1038/nature11390>
- Stahle, D. W., & Cleaveland, M. K. (1988). Texas drought history reconstructed and analyzed from 1698 to 1980. *Journal of Climate*, 1(1), 59–74. [https://doi.org/10.1175/1520-0442\(1988\)001<0059:Tdhras>2.0.Co;2](https://doi.org/10.1175/1520-0442(1988)001<0059:Tdhras>2.0.Co;2)
- Sungmin, O., & Orth, R. (2021). Global soil moisture data derived through machine learning trained with in-situ measurements. *Scientific Data*, 8(1), 170. <https://doi.org/10.1038/s41597-021-00964-1>
- Tabatabaenejad, A., Chen, R. H., Burgin, M. S., Duan, X., Cuenca, R. H., Cosh, M. H., et al. (2020). Assessment and validation of AirMOSS P-band root-zone soil moisture products. *IEEE Transactions on Geoscience and Remote Sensing*, 58(9), 6181–6196. <https://doi.org/10.1109/Tgrs.2020.2974976>
- Teuling, A. J., Seneviratne, S. I., Williams, C., & Troch, P. A. (2006). Observed timescales of evapotranspiration response to soil moisture. *Geophysical Research Letters*, 33(23). <https://doi.org/10.1029/2006gl028178>
- Tijdeman, E., & Menzel, L. (2021). Controls on the development and persistence of soil moisture drought across Southwestern Germany. *Hydrology and Earth System Sciences*, 25(4), 2009–2025. <https://doi.org/10.5194/hess-25-2009-2021>
- Timbal, B., Power, S., Colman, R., Viviani, J., & Lirola, S. (2002). Does soil moisture influence climate variability and predictability over Australia? *Journal of Climate*, 15(10), 1230–1238. [https://doi.org/10.1175/1520-0442\(2002\)015%3C1230:DSMICV%3E2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015%3C1230:DSMICV%3E2.0.CO;2)
- Tuller, M., et al. (2023). Proximal sensing of soil moisture. In M. J. Goss & M. Oliver (Eds.), *Encyclopedia of soils in the environment* (pp. 591–599). Academic Press. <https://doi.org/10.1016/b978-0-12-822974-3.00157-9>
- Ukkola, A. M., De Kauwe, M. G., Pitman, A. J., Best, M. J., Abramowitz, G., Haverd, V., et al. (2016). Land surface models systematically overestimate the intensity, duration and magnitude of seasonal-scale evaporative droughts. *Environmental Research Letters*, 11(10), 104012. <https://doi.org/10.1088/1748-9326/11/10/104012>
- van den Hurk, B., Haarsma, R., Selten, F., & Seneviratne, S. (2009). Soil drying in Europe and its impact on atmospheric circulations. Paper presented at Proc ECMWF semin parametrizations subgrid phys process.
- van den Hurk, B. J., & van Meijgaard, E. (2010). Diagnosing land-atmosphere interaction from a regional climate model simulation over West Africa. *Journal of Hydrometeorology*, 11(2), 467–481. <https://doi.org/10.1175/2009JHM1173.1>
- Vautard, R., Yiou, P., D'Andrea, F., de Noblet, N., Viovy, N., Cassou, C., et al. (2007). Summertime European heat and drought waves induced by wintertime Mediterranean rainfall deficit. *Geophysical Research Letters*, 34(7). <https://doi.org/10.1029/2006gl028001>
- Vereecken, H., Schnepf, A., Hopmans, J., Javaux, M., Or, D., Roose, T., et al. (2016). Modeling soil processes: Review, key challenges, and new perspectives. *Vadose Zone Journal*, 15(5), 1–57. <https://doi.org/10.2136/vzj2015.09.0131>
- Verma, S., Singh, D., Mani, S., & Jayakumar, S. (2017). Effect of forest fire on tree diversity and regeneration potential in a tropical dry deciduous forest of Mudumalai Tiger Reserve, Western Ghats, India. *Ecological Processes*, 6, 1–8. <https://doi.org/10.1186/s13717-017-0098-0>
- Vidale, P. L., Egea, G., McGuire, P. C., Todt, M., Peters, W., Müller, O., et al. (2021). On the treatment of soil water stress in GCM simulations of vegetation physiology. *Frontiers in Environmental Science*, 9, 689301. <https://doi.org/10.3389/fenvs.2021.689301>
- Vinnikov, K. Y., Robock, A., Qiu, S., & Entin, J. K. (1999). Optimal design of surface networks for observation of soil moisture. *Journal of Geophysical Research*, 104(D16), 19743–19749. <https://doi.org/10.1029/1999jd900060>
- Vinnikov, K. Y., Robock, A., Speranskaya, N. A., & Schlosser, C. A. (1996). Scales of temporal and spatial variability of midlatitude soil moisture. *Journal of Geophysical Research*, 101(D3), 7163–7174. <https://doi.org/10.1029/95jd02753>
- Vinnikov, K. Y., & Yeserkepova, I. B. (1991). Soil-moisture - empirical-data and model results. *Journal of Climate*, 4(1), 66–79. [https://doi.org/10.1175/1520-0442\(1991\)004<0066:Smedam>2.0.Co;2](https://doi.org/10.1175/1520-0442(1991)004<0066:Smedam>2.0.Co;2)
- Von Storch, H., & Zwiers, F. W. (2002). *Statistical analysis in climate research*. Cambridge university press.
- Wagner, W., Lemoine, G., & Rott, H. (1999). A method for estimating soil moisture from ERS scatterometer and soil data. *Remote Sensing of Environment*, 70(2), 191–207. [https://doi.org/10.1016/S0034-4257\(99\)00036-X](https://doi.org/10.1016/S0034-4257(99)00036-X)
- Walker, J., & Rowntree, P. (1977). The effect of soil moisture on circulation and rainfall in a tropical model. *Quarterly Journal of the Royal Meteorological Society*, 103(435), 29–46. <https://doi.org/10.1002/qj.49710343503>
- Walker, J. P., & Houser, P. R. (2001). A methodology for initializing soil moisture in a global climate model: Assimilation of near-surface soil moisture observations. *Journal of Geophysical Research*, 106(D11), 11761–11774. <https://doi.org/10.1029/2001jd900149>
- Wang, F., Notaro, M., Liu, Z., & Chen, G. (2014). Observed local and remote influences of vegetation on the atmosphere across North America using a model-validated statistical technique that first excludes oceanic forcings. *Journal of Climate*, 27(1), 362–382. <https://doi.org/10.1175/jcli-d-13-00080.1>
- Wei, J., Dirmeyer, P. A., & Guo, Z. (2010). How much do different land models matter for climate simulation? Part II: A decomposed view of the land-atmosphere coupling strength. *Journal of Climate*, 23(11), 3135–3145. <https://doi.org/10.1175/2010JCLI3177.1>
- Wei, J. F., Dickinson, R. E., & Zeng, N. (2006). Climate variability in a simple model of warm climate land-atmosphere interaction. *Journal of Geophysical Research*, 111(G3). <https://doi.org/10.1029/2005jg000096>
- Weiherrmüller, L., Lehmann, P., Herbst, M., Rahmati, M., Verhoef, A., Or, D., et al. (2021). Choice of pedotransfer functions matters when simulating soil water balance fluxes. *Journal of Advances in Modeling Earth Systems*, 13(3), e2020MS002404. <https://doi.org/10.1029/2020ms002404>
- Wey, H. W., Lo, M. H., Lee, S. Y., Yu, J. Y., & Hsu, H. H. (2015). Potential impacts of wintertime soil moisture anomalies from agricultural irrigation at low latitudes on regional and global climates. *Geophysical Research Letters*, 42(20), 8605–8614. <https://doi.org/10.1002/2015gl065883>
- Wu, J. X., & Dirmeyer, P. A. (2020). Drought demise attribution over CONUS. *Journal of Geophysical Research-Atmospheres*, 125(4), e2019JD031255. <https://doi.org/10.1029/2019jd031255>
- Wu, L. Y., & Zhang, J. Y. (2015). The relationship between spring soil moisture and summer hot extremes over North China. *Advances in Atmospheric Sciences*, 32(12), 1660–1668. <https://doi.org/10.1007/s00376-015-5003-0>
- Wu, W., & Dickinson, R. E. (2004). Time scales of layered soil moisture memory in the context of land-atmosphere interaction. *Journal of Climate*, 17(14), 2752–2764. [https://doi.org/10.1175/1520-0442\(2004\)017%3C2752:TSOLSM%3E2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017%3C2752:TSOLSM%3E2.0.CO;2)
- Wu, W. R., Geller, M. A., & Dickinson, R. E. (2002). A case study for land model evaluation: Simulation of soil moisture amplitude damping and phase shift. *Journal of Geophysical Research*, 107(D24). <https://doi.org/10.1029/2001jd001405>
- Xiao, H., Meissner, R., Seeger, J., Rupp, H., & Borg, H. (2009). Effect of vegetation type and growth stage on dewfall, determined with high precision weighing lysimeters at a site in northern Germany. *Journal of Hydrology*, 377(1–2), 43–49. <https://doi.org/10.1016/j.jhydrol.2009.08.006>
- Xu, D. H., Ivanov, V. Y., Li, X., & Troy, T. J. (2021). Peak runoff timing is linked to global warming trajectories. *Earth's Future*, 9(8), e2021EF002083. <https://doi.org/10.1029/2021EF002083>

- Xue, Y., Boone, A., & Taylor, C. M. (2012). Review of recent developments and the future prospective in west African atmosphere/land interaction studies. *International Journal of Geophysics*, 2012, 1–12. <https://doi.org/10.1155/2012/748921>
- Xue, Y. K., Yao, T., Boone, A. A., Diallo, I., Liu, Y., Zeng, X., et al. (2021). Impact of initialized land surface temperature and snowpack on subseasonal to seasonal prediction project, phase I (LS4P-I): Organization and experimental design. *Geoscientific Model Development*, 14(7), 4465–4494. <https://doi.org/10.5194/gmd-14-4465-2021>
- Yang, K., & Zhang, J. Y. (2016). Spatiotemporal characteristics of soil temperature memory in China from observation. *Theoretical and Applied Climatology*, 126(3–4), 739–749. <https://doi.org/10.1007/s00704-015-1613-9>
- Yao, Y., Liao, X., Xiao, J., He, Q., & Shi, W. (2023). The sensitivity of maize evapotranspiration to vapor pressure deficit and soil moisture with lagged effects under extreme drought in Southwest China. *Agricultural Water Management*, 277, 108101. <https://doi.org/10.1016/j.agwat.2022.108101>
- Yasunari, T. (2007). Role of land-atmosphere interaction on Asian monsoon climate. *Journal of the Meteorological Society of Japan*, 85b, 55–75. <https://doi.org/10.2151/jmsj.85B.55>
- Yeh, T. C., Wetherald, R. T., & Manabe, S. (1984). The effect of soil-moisture on the short-term climate and hydrology change - A numerical experiment. *Monthly Weather Review*, 112(3), 474–490. [https://doi.org/10.1175/1520-0493\(1984\)112<0474:Teosmo>2.0.Co;2](https://doi.org/10.1175/1520-0493(1984)112<0474:Teosmo>2.0.Co;2)
- Yin, Z., Findell, K. L., Dirmeyer, P., Shevliakova, E., Malyshev, S., Ghannam, K., et al. (2023). Daytime-only mean data enhance understanding of land-atmosphere coupling. *Hydrology and Earth System Sciences*, 27(4), 861–872. <https://doi.org/10.5194/hess-27-861-2023>
- Yuan, X., Wang, L., & Wood, E. F. (2018). Anthropogenic intensification of southern African flash droughts as exemplified by the 2015/16 season. *Bulletin of the American Meteorological Society*, 99(1), S86–S90. <https://doi.org/10.1175/Bams-D-17-0077.1>
- Zeng, Z. Z., Zhu, Z., Lian, X., Li, L. Z. X., Chen, A., He, X., & Piao, S. (2016). Responses of land evapotranspiration to Earth's greening in CMIP5 earth system models. *Environmental Research Letters*, 11(10), 104006. <https://doi.org/10.1088/1748-9326/11/10/104006>
- Zhao, M., Zhang, H., & Dharssi, I. (2019). On the soil moisture memory and influence on coupled seasonal forecasts over Australia. *Climate Dynamics*, 52(11), 7085–7109. <https://doi.org/10.1007/s00382-018-4566-8>

References From the Supporting Information

- Gao, J., Hu, J., Tung, W. W., Cao, Y., Sarshar, N., & Roychowdhury, V. P. (2006). Assessment of long-range correlation in time series: How to avoid pitfalls. *Physical Review E*, 73(1), 016117. <https://doi.org/10.1103/PhysRevE.73.016117>
- Gold, E. (1929). Note on the frequency of occurrence of sequences in a series of events of two types. *Quarterly Journal of the Royal Meteorological Society*, 55(231), 307–309. <https://doi.org/10.1002/qj.49705523112>
- Hurst, H. E. (1951). Long-term storage capacity of reservoirs. *Transactions of the American Society of Civil Engineers*, 116(1), 770–799. <https://doi.org/10.1061/TACEAT.000651>
- Jenkins, G. M. (1968). *Spectral analysis and its applications*. Holden-Day, Inc.
- Pagano, M., & Halvorsen, K. T. (1981). An algorithm for finding the exact significance levels of $r \times c$ contingency tables. *Journal of the American Statistical Association*, 76(376), 931–934. <https://doi.org/10.2307/2287590>
- Riley, M. A., Bonnette, S., Kuznetsov, N., Wallot, S., & Gao, J. (2012). A tutorial introduction to adaptive fractal analysis. *Frontiers in Physiology*, 3, 371. <https://doi.org/10.3389/fphys.2012.00371>
- Shen, S., Ye, S., Cheng, C., Song, C., Gao, J., Yang, J., et al. (2018). Persistence and corresponding time scales of soil moisture dynamics during summer in the Babao river basin, northwest China. *Journal of Geophysical Research: Atmospheres*, 123(17), 8936–8948. <https://doi.org/10.1029/2018JD028414>