

The role of isotope mass and transport for H-mode access in tritium containing plasmas at JET with ITER-like wall

G. Birkenmeier^{1,2,†}, E.R. Solano³, I.S. Carvalho⁴, J.C. Hillesheim⁵, E. Delabie⁶, E. Lerche⁷, D. Taylor⁵, D. Gallart⁸, M.J. Mantsinen^{8,9}, C. Silva⁴, C. Angioni¹, F. Ryter¹, P. Carvalho⁴, M. Fontana⁵, E. Pawelec¹⁰, S.A. Silburn⁵, P. Sirén⁵, S. Aleiferis⁵, J. Bernardo^{5,4}, A. Boboc⁵, D. Douai¹¹, P. Puglia⁵, P. Jacquet⁵, E. Litherland-Smith⁵, I. Jezu⁵, D. Kos⁵, H.J. Sun⁵, A. Shaw⁵, D. King⁵, B. Viola⁵, R. Henriques⁴, K.K. Kirov⁵, M. Baruzzo⁵, J. Garcia¹¹, A. Hakola¹², A. Huber⁵, E. Joffrin¹¹, D. Keeling⁵, A. Kappatou¹, M. Lennholm⁵, P. Lomas⁵, E. de la Luna³, C.F. Maggi⁵, J. Mailloux⁵, M. Maslov⁵, F.G. Rimini⁵, N. Vianello¹³, G. Verdoolaege^{7,15}, H. Weisen¹⁴, M. Wischmeier¹, and JET Contributors[§]

¹Max Planck Institute for Plasma Physics, Boltzmannstr. 2, 85748 Garching, Germany

²Physik-Department E28, Technische Universität München, James-Franck-Str. 1, 85748 Garching, Germany

³Laboratorio Nacional de Fusión, CIEMAT, Madrid, Spain

⁴Instituto de Plasmas e Fusão Nuclear, Instituto Superior Técnico, Universidade de Lisboa, Portugal

⁵CCFE, Culham Science Centre, Abingdon, Oxfordshire, OX14 3DB, United Kingdom of Great Britain and Northern Ireland

⁶Oak Ridge National Laboratory, Oak Ridge, TN 37831-6169, TN, United States of America

⁷Laboratory for Plasma Physics Koninklijke Militaire School, Ecole Royale Militaire Renaissancelaan 30 Avenue de la Renaissance B-1000, Brussels, Belgium

⁸Barcelona Supercomputing Center (BSC), Barcelona, Spain

⁹ICREA, Barcelona, Spain

¹⁰Institute of Physics, Opole University, Oleska 48, 45-052 Opole, Poland

¹¹CEA, IRFM, F-13108, St-Paul-Lez-Durance, France

¹²VTT, P. O. Box 1000, 02044 VTT, Finland

¹³Consorzio RFX, Padova, Italy

¹⁴Ecole Polytechnique Federale de Lausanne (EPFL), Swiss Plasma Center (SPC), CH-1015 Lausanne, Switzerland

¹⁵Department of Applied Physics, Ghent University, 9000, Ghent, Belgium

E-mail: gregor.birkenmeier@ipp.mpg.de

July 2022

Abstract. The required heating power, P_{LH} , to access the high confinement regime (H-mode) in tritium containing plasmas is investigated in JET with ITER-like wall at a toroidal magnetic field of $B_t = 1.8$ T and a plasma current of $I_p = 1.7$ MA. P_{LH} , also referred to as the L-H power threshold, is determined in plasmas of pure tritium as well as mixtures of hydrogen with tritium (H-T) and mixtures of deuterium with tritium (D-T), and is compared to the L-H power threshold in plasmas of pure hydrogen and pure deuterium. It is found that, for otherwise constant parameters, P_{LH} is not the same in plasmas with the same effective isotope mass, A_{eff} , when they differ in their isotope composition. Thus, A_{eff} is not sufficient to describe the isotope effect of P_{LH} in a consistent manner for all considered isotopes and isotope mixtures. The electron temperature profiles measured at the L-H transition in the outer half of the radius are very similar for all isotopes and isotope mixtures, despite the fact that the L-H power threshold varies by a factor of about six. This finding, together with the observation of an offset linear relation between the L-H power threshold, P_{LH} , and an effective heat diffusivity, χ_{eff} , indicates that the composition-dependent heat transport in the low confinement mode (L-mode) determines, how much power is needed to reach the necessary electron temperatures at the edge, and hence P_{LH} .

1. Introduction

Tokamak fusion reactors with a magnetic configuration similar to ITER [1], are foreseen to be operated in the high confinement regime (H-mode) in order to reach the required confinement properties [2]. The H-mode can be accessed in a plasma of low confinement (L-mode) by exceeding a certain threshold of injected heating power, the L-H power threshold, P_{LH} . A regression of an ITPA multi-machine data base^{||} of deuterium plasmas resulted in a scaling law of the L-H power threshold [3]

$$P_{\text{ITPA}} = 0.049 \bar{n}_e^{0.72} B_t^{0.8} S^{0.94}, \quad (1)$$

which identified the line-averaged core electron density, \bar{n}_e , the toroidal magnetic field, B_t , and the surface of the plasma, S , as some of the main parameters determining the L-H power threshold. Please note that this scaling applies only for the high-density branch, i.e. for densities above $n_{e,\text{min}}$, which indicates the density where P_{LH} is minimum in the typically U-shaped curve of $P_{\text{LH}}(\bar{n}_e)$. Several more parameters were identified to additionally impact P_{LH} like plasma shape, wall material, impurity concentrations, and toroidal rotation [4, 5, 6, 7, 8, 9, 10, 11, 12]. However, the relatively simple parameter dependencies of the ITPA scaling have already proven to strongly impact the design of a future tokamak reactor [13], and more complex dependencies could further restrict the reactor operational point. Therefore, for a reliable and robust tokamak reactor design it is required to predict and understand the L-H power threshold as accurately as possible.

The ITPA scaling P_{ITPA} , which is also used to assess P_{LH} in the nuclear and non-nuclear operation phase of ITER [14], only applies to deuterium plasmas. Thus, for

^{||} ASDEX Upgrade and JET, the two of the largest tokamaks contributing to the multi-machine data base, provided only data of plasmas operated in a carbon wall at the time when the scaling was derived, although both devices are equipped with a metallic wall in the meantime.

the reactor relevant fuel mixture of deuterium-tritium or other isotopes or isotope compositions, the isotope dependence of P_{LH} has to be assessed and equation 1 correspondingly extended. The most common and simplest approach to factor in isotope dependences of P_{LH} is to introduce an effective isotope mass, A_{eff} , which can be determined from the densities of the different isotopes of the plasma as

$$A_{\text{eff}} = \frac{n_H + 2n_D + 3n_T}{n_H + n_D + n_T}, \quad (2)$$

with the densities n_H , n_D , and n_T of hydrogen (H), deuterium (D) and tritium (T), respectively. Data of comparative studies in hydrogen and deuterium plasmas [15, 16, 17, 18] suggest an inversely proportional dependence on the effective isotope mass:

$$P_{\text{LH}} \propto A_{\text{eff}}^{-1}. \quad (3)$$

A similar dependence on A_{eff} was found in tritium containing plasmas in JET with plasma facing components (PFCs) made of carbon [19]. Based on these findings, we assume in the following, that the scaling

$$P_{\text{scal}} = 0.098 \bar{n}_e^{0.72} B_t^{0.8} S^{0.94} A_{\text{eff}}^{-1} \quad (4)$$

is a reasonable and data based parametrization of P_{LH} in the high-density branch, which describes the main dependences including isotope effects properly, and serves as a reference for the data discussed in the following. Please note that the prefactor of the scaling P_{scal} differs from equation 1 by a factor of two in order to make both scalings consistent for deuterium corresponding to $A_{\text{eff}} = 2$.

Similar as done in reference [20], we contrast data from JET with ITER-like wall (ILW) at a toroidal magnetic field of $B_t = 1.8$ T and a plasma current of $I_p = 1.7$ MA collected during the tritium campaign 2020/2021 with P_{scal} in order to identify main differences and similarities of the new data in metallic wall conditions with existing scaling laws. The data set at hand includes two data points of the reactor-relevant fuel mixture of D-T, and it contains a T concentration scan of H-T mixed plasmas allowing for a fine variation of A_{eff} . This enables a direct comparison of P_{LH} in plasmas with the same A_{eff} but different isotope compositions. It is shown below, that A_{eff} is not sufficient to describe the isotope dependence of P_{LH} consistently for the considered isotopes and isotope mixtures. The data rather points to the heat transport in L-mode as a key player determining P_{LH} by defining the relevant loss channel, which counteracts the input heating power to reach the edge temperatures required for the L-H transition.

In the remainder, we introduce in section 2 the data set of tritium containing plasmas in JET with ILW, and the L-H power threshold as well as the radiation fraction is presented for different isotope mixtures. In section 3 we contrast the data with equation 4 revealing that A_{eff} cannot describe the isotope effect of the L-H transition consistently. The profiles at the L-H transition and the possible crucial role of transport is discussed in section 4 before a summary and conclusion is given in section 5.

2. The power threshold in tritium containing plasmas

In order to investigate the effect of the effective isotope mass, A_{eff} , and heat transport on the L-H power threshold P_{LH} , we consider a data set acquired in the tritium campaign 2020/2021 at JET with ILW similar to the data presented in reference [20]. The data was taken at a toroidal magnetic field of $B_t = 1.8$ T and a plasma current of $I_p = 1.7$ MA. The geometry of the divertor is called MkII-HD (see reference [4]), the PFC material was tungsten in the divertor and beryllium at the main chamber wall, and the plasma configuration (favourable configuration with the ion $\mathbf{B} \times \nabla B$ -drift direction towards the active X-point) was a lower single null plasma with the outer strikeline on the horizontal target of the divertor referred to as HT configuration as in reference [8]. The new data differs in the following three main aspects from the data discussed in reference [20]: (i) it contains only data with proper phasing of the ion cyclotron heating (ICRH) antenna, (ii) it contains not only two different concentration levels, but in total seven steps of tritium concentration in mixtures of H-T plasmas, and (iii) it contains two data points of a 50 % / 50 % mixture of D-T. Apart from these differences, the data was taken in the same way from discharges with slow heating ramps and averaging the relevant quantities 70 ms prior to the L-H transition. The L-H transition is identified in diagnostic data by a sudden drop of the D_α signal indicating lowered edge transport, and a simultaneous and sudden increase of the edge temperature, edge density and the energy content of the plasma indicating improved confinement. Examples of diagnostic time traces of L-H transitions are given in reference [20].

A few of the pure T plasmas in the data set were heated with neutral beam injection (NBI) using a T heating beam, while most of the plasmas were heated with ICRH. H plasmas or H-T mixed plasmas with a tritium concentration of up to 95 %, were heated with ICRH at a frequency of 51.4 MHz corresponding to the second harmonic ($\omega = 2\omega_{c,H}$) in hydrogen. For tritium concentrations above 95 % in H-T mixtures as well as for D-T plasmas, hydrogen minority heating at the fundamental frequency ($\omega = \omega_{c,H}$) was employed corresponding to a wave frequency of $f = 32.2$ MHz. Since we consider only plasmas heated with low power T-NBI and ICRH, the data set at hand contains only plasmas with low torque input. The concentrations of hydrogenic species were measured with an optical penning gauge in the subdivertor [21, 22]. For more details about the experimental approach the reader is referred to reference [20].

As usual, we use the total loss power

$$P_{\text{loss}} = P_{\text{Ohm}} + P_{\text{aux}} - \frac{dW_p}{dt} \quad (5)$$

with the Ohmic heating power P_{Ohm} , any other auxiliary heating power P_{aux} and the time derivative of the plasma energy content W_p as estimation of P_{LH} , which was likewise used for the ITPA multi-machine data base [3]. All quantities are averaged over 70 ms prior to the L-H transition, which takes place at time t_{LH} . Because of the typically strong rise of the (diamagnetic) energy content W_p at the time point of the L-H transition, we do not take $\frac{dW_p}{dt}$ at t_{LH} , but 100 ms earlier than t_{LH} in order that P_{loss} is not affected by

the L-H transition induced changes. Due to the relatively high radiation levels at the L-H transitions in JET, it turned out that the data is more consistent if the bulk radiated power P_{rad} , i.e. the radiated power inside the magnetic separatrix, is subtracted. The resulting power is the kinetically transported power through the separatrix

$$P_{\text{sep}} = P_{\text{loss}} - P_{\text{rad}}. \quad (6)$$

P_{loss} of the tritium containing plasmas is shown in figure 1, left. Starting at pure hydrogen plasmas (red circles), which exhibit the highest L-H power thresholds, a stepwise increase of the T content in H-T mixed plasmas (open red diamonds, T concentration given next to the respective symbol in percent) reduced the P_{loss} consecutively. The observation of lower P_{loss} with increasing T concentration, and hence increasing A_{eff} , is qualitatively in agreement with equation 3.




Figure 1. P_{loss} (left) and P_{sep} (right) against line-averaged core density \bar{n}_e for different main ion plasma isotopes and isotope mixtures heated with different heating schemes. The data points labelled as "transient" are dithering L-H transitions and were heated with ICRH. The numbers next to the symbols of the H-T mixed plasmas indicate the tritium concentration in percent.

P_{loss} determined in pure T plasmas (magenta symbols) cover a wide density range and show in general the lowest L-H power thresholds for low and high densities. Pure means in this context a T concentration greater than 95 % with respect to other hydrogenic species, which is believed to be sufficient for the plasma dynamics to be dominated by T. The T concentration cannot be much further increased, since a small amount of H must always be present for the applied minority heating scheme with ICRH. T plasmas heated with NBI (black filled symbols), have a lower P_{loss} than ICRH plasmas (magenta filled diamonds). The lowest values of P_{loss} are found for transitions in Ohmically heated phases (x-symbols), which are a peculiarity of T plasmas at this

field and current at JET, since other isotopes could not access the H-mode with Ohmic heating on a regular basis. The possibility to access H-mode with Ohmic heating only [23] indicates that the requirements to reach H-mode in T are facilitated compared to the other isotopes similar to Ohmic H-modes found at other devices in other type of low-threshold conditions (typically low magnetic field) [24, 25, 26]. In contrast to H and D (blue squares) data, the T data does not show a clear minimum of P_{loss} . But the fact that an increase in density let the plasma transition to H-mode resembles the dynamics of transitions in the low-density branch [20], so that the Ohmic transitions in T can be considered as the low-density branch of the T data.

Both pure tritium plasmas and H-T mixed plasmas with T concentrations above 75 % and heated with ICRH often transitioned into a dithering L-H transition phase (plus signs) instead of a sustained H-mode as described in reference [20]. Dithering transitions are likewise observed in other isotopes, however less frequent, and in contrast to T containing plasmas, a sustained H-mode was easier to achieve in plasmas without T by further increasing the heating power. This points to an unfavourable influence of impurity-induced radiation losses on H-mode robustness, which is obviously more severe in T similar as found in He plasmas [27]. This might be related to its enhanced ability to sputter beryllium (Be) more effectively than H or D [28]. The increased Be concentrations could lead to enhanced sputtering of tungsten and, thus, could increase the bulk radiation.

The two data points of D-T plasmas had a T concentration of about 50 % (golden stars) and are within the error bars in the same range as data points of pure D and pure T in the same density range.

If the bulk radiation is subtracted from P_{loss} , the resulting P_{sep} shown in figure 1, right, exhibit the same trends as discussed for P_{loss} in figure 1, left. There is however one main difference: The data points of pure T plasmas heated with ICRH possess the same P_{sep} as NBI-heated pure T plasmas. In other words, the radiation is higher in ICRH plasmas than in NBI-heated plasmas, so that the radiated power compensated the difference of P_{loss} between these two differently heated plasma types. Hence, P_{sep} in tritium plasmas at 1.8 T does not depend on the torque input. This is a difference to the other isotopes, since P_{sep} in D exhibits a very small [27] and in H a very large dependence on the heating method [8].

As shown in figure 2, the radiation fraction at the L-H transition estimated as the ratio between bulk radiation P_{rad} and input heating power P_{loss} is highest for ICRH T plasmas (about 35 %) and low density D plasmas (up to 50 %). The fact, that NBI-heated T plasmas show very low radiation fractions, reveals, that the presence of T alone is not responsible for high radiation levels. It is rather the combination of high tritium concentrations *together with* ICRH that produces the highest radiation levels, probably related to ICRH-induced electric fields in the scrape-off layer [29].

D-T plasmas suffer less from high radiation fractions than pure T plasmas and show radiation fractions in the same range as D plasmas. H-T plasmas with T concentrations of 19 % to 75 % exhibit radiation levels of about 20 %, which is slightly higher than the




Figure 2. Radiation fraction estimated by the ratio between bulk radiation, P_{rad} , and input heating power, P_{loss} , for different isotopes and heating schemes.

pure H plasmas clustering around 15 %.

3. The role of A_{eff} for P_{LH}

The data of P_{loss} and P_{sep} introduced above is now compared to the reference scaling P_{scal} as given by equation 4. Figure 3, left, shows the measured P_{loss} divided by the scaling P_{scal} on the y-axis and the effective isotope mass on the x-axis. Only data from the high density branch is shown, i.e. data for densities higher than the density, $n_{\text{e,min}}$, where the respective minimum of P_{loss} appears, since the scaling only holds for the high density branch [3].

All data points, except for the pure T heated with ICRH, are around 1.0 or mostly lower. This indicates that the measured values of P_{loss} are lower than the reference scaling. Plasmas of pure H, pure D, and D-T mixtures are around 20 % lower than the scaling. This is a well known effect of the ILW [4] and most probably related to the




Figure 3. P_{loss} (left) and P_{sep} (right) normalized to the scaling P_{scal} (equation 4) against effective isotope mass, A_{eff} . Only data from the high density branch is shown. The power threshold can vary significantly for the same A_{eff} indicating that the isotope effect of P_{LH} is not well described by the effective isotope mass.

fact, that the L-H power threshold is lower in metallic wall conditions than in a carbon wall environment as also found in ASDEX Upgrade [30].

The pure T plasmas heated with ICRH exhibit relatively high normalized power levels $P_{\text{loss}}/P_{\text{scal}}$ due to the high radiation fraction. This is obvious from figure 3, right, which shows $P_{\text{sep}}/P_{\text{scal}}$, i.e. the normalized power threshold, but corrected from radiation effects: The data for NBI and ICRH plasmas do not differ anymore within the pure T, and $P_{\text{sep}}/P_{\text{scal}}$ ranges from about 60 % to 90 %.

In general, there is a quite remarkable scatter of $P_{\text{sep}}/P_{\text{scal}}$ within each isotope mass i.e. for a constant value of A_{eff} , although the dependence of the density \bar{n}_e is taken into account by the normalization to the scaling (B_t and S are constant in this data set). This means, that the scaling with its density dependence $\bar{n}_e^{0.72}$ does not reflect the experimental density dependence at JET very well. If the scaling would perfectly describe our data, there would be no scatter along the ordinate, since the normalization to the scaling would compensate the density dependence, which is obviously not the case.

For the pure isotopes H, D and T, the quantity $P_{\text{sep}}/P_{\text{scal}}$ does not show a monotonic behaviour: it is lowest for D, highest for T, and H is in between. This means that this data set disagrees with any approach, which tries to describe the isotope effect of the L-H power threshold in the form of $P_{\text{LH}} \propto A_{\text{eff}}^\alpha$ with any real number α .

$P_{\text{sep}}/P_{\text{scal}}$ of the H-T mixture with 47 % of T concentration corresponding to $A_{\text{eff}} = 2$ is 65 % higher than the data point of the corresponding D plasma at the same density and likewise $A_{\text{eff}} = 2$. Similarly, the two D-T data points are much

lower than the value of $P_{\text{sep}}/P_{\text{scal}}$ of the corresponding H-T mixture with 74 % T concentration, although all plasmas had an effective isotope mass of about $A_{\text{eff}} = 2.5$. This clearly shows, that the effective isotope mass is not a good parameter to describe the isotope effect of P_{LH} in a sufficient and consistent way. Thus, the actual isotope *composition* and not the effective isotope mass matters for P_{LH} . This means, that any theoretical approach or scaling law aiming at a description of the isotope dependence of P_{LH} solely by means of an effective isotope mass A_{eff} , will not be able to describe our data sufficiently well.

The ion heating contribution estimated from PION [31] including pitch angle effects [32] for the ICRH and using PENCIL for NBI heated discharges [33] for this data set reveal the same trends as discussed in reference [20]: For increasing ion mass the ion heating decreases linearly with the T content when starting from pure H towards pure T plasmas during the concentration scan of the H-T mixes plasmas. This seems to be in line with the idea of a critical ion heat flux inversely proportional to A_{eff} as discussed in reference [20]. However, the absolute ion heating as well as the ion heating fraction relative to the total heating for pure D plasmas is significantly lower than the H-T counterpart with 47 % T concentration. Likewise the ion heating fraction of the D-T plasma is lower than in the H-T mixture with 74 % T concentration. In addition, the D-T mixture exhibits the same ion heating fraction as pure D plasmas although they differ in terms of A_{eff} . Thus, the isotope dependence of the ion heating contribution, interpreted as a proxy for the edge ion heat flux, behaves similarly as the power threshold itself and cannot consistently be described solely with A_{eff} for all isotope mixtures.

4. The role of transport for P_{LH}

As shown before, plasmas of the same A_{eff} can exhibit different P_{LH} *when* the isotope composition differs. One might expect, that the kinetic profiles at the time of the L-H transition in such cases, e.g. profiles of a D plasma compared with profiles of an H-T mixed plasma with 50 % T concentration (both have $A_{\text{eff}} = 2$), are very different, since P_{LH} , and hence the input heating power, is very different. However, it turns out, that the electron density profiles and the electron temperature profiles are very similar (even identical within the errors), when the central line-averaged density is matched sufficiently well in these cases.

It is found that the profiles at the L-H transition are similar even for different A_{eff} , as long as the line-averaged density is the same. This is shown in figure 4, which displays Thomson scattering profiles of the electron density (top left) and the electron temperature (top right) averaged over 200 ms in the L-mode phase prior to the L-H transition for a density range of $\bar{n}_e = 3.0 \cdot 10^{19} \text{ m}^{-3}$ to $\bar{n}_e = 3.2 \cdot 10^{19} \text{ m}^{-3}$. These include effective isotope masses A_{eff} from 1 to 3 and values of P_{loss} ranging from 2.03 MW to 5.15 MW. The density profiles are very similar, and they exhibit only a relative deviation of 10 % with respect to the density profile of the H-T mixed plasma with 50 % T content (JPN100102) serving as a reference profile. This small deviation is

similar to the measurement error (see figure 4, bottom left) indicating good agreement between these density profiles. Similarly, the electron temperature profiles, especially in the region from half the minor radius ($R = 3.4$ m) to the edge ($R = 3.72$ m), agree within 25 % relative errors. This is only marginally larger than the measurement error of the diagnostic (see figure 4, bottom right). Thus, both density profiles and electron temperature profiles measured at the L-H transition agree very well for these very different isotope compositions. The energy contents in this set of discharges are likewise very similar (about 0.7 MJ) indicating that the ion temperature profiles cannot be much different from the electron temperature profiles shown here.

Thus, for the same central line-averaged density, the electron temperature and density profiles at the L-H transition are very similar despite the different P_{LH} and A_{eff} . The finding of similar kinetic profiles at the L-H transition for different isotopes and isotope mixtures is not new and was already described in references [36] and [8]. But it is impressive to see this effect also present in T containing plasmas and over a now extended range of P_{LH} and A_{eff} .




Figure 4. Profiles of electron density (top left) and electron temperature (top right) and their relative deviations (bottom) from the profiles of the H-T mixed plasma with 50 % T content (JPN100102) from the Thomson scattering diagnostic prior to the L-H transition for different isotope compositions in a density range of $\bar{n}_e = 3.0 \cdot 10^{19} \text{ m}^{-3}$ to $\bar{n}_e = 3.2 \cdot 10^{19} \text{ m}^{-3}$ (input powers and JET Pulse numbers, JPN, given in the legend).

For other densities than shown in figure 4, the electron density profiles are, naturally, not the same, since the operation of the plasma was on purpose aiming at different line-averaged densities. However, the electron temperatures at the L-H transition are again very similar for all the different densities and isotope masses of the considered data set. This is shown in figure 5. The left panel shows the total averaged temperature \bar{T} estimated as $\bar{T} = W_p / (3\bar{n}_e V)$ with V the volume of the plasma.

This approach assumes that the electron and ion temperature profiles are the same throughout the plasma, which is a good approximation for higher densities, but fails at lowest densities. Furthermore it assumes that the energy content of the plasma, W_p , is not affected by the fast ion pressure and only determined by the thermal bulk of the plasma.

Despite the input power, and hence P_{LH} , varies by a factor of six, the average temperature \bar{T} is quite similar for most of the cases and ranges between 500 eV and 700 eV with a slight upward trend for higher heating power. The semi-transparent symbols correspond to data points from the low-density branch, i.e. plasmas with densities below $n_{\text{e,min}}$ of the respective isotope. They deviate from the main group of the data, which could be related to finite ICRH-induced fast ion contributions to W_p at low densities or a mismatch of electron and ion temperatures, so that \bar{T} is not a meaningful quantity to draw conclusions about thermal temperature profiles. But for data from the high-density branch, \bar{T} is very well aligned for all isotope masses and over a wide range of densities.




Figure 5. Average temperature, \bar{T} (left), and edge electron temperature from the ECE diagnostic, $T_{\text{e,edge}}$ (right), taken 9.95 cm inside the separatrix. The semi-transparent symbols indicate data from the low density branch.

Figure 5, right, shows the electron temperature $T_{\text{e,edge}}$ measured with the electron-cyclotron emission (ECE) diagnostics at the plasma edge at $R = 3.725$ m, i.e. 9.95 cm inside the separatrix corresponding to a normalized poloidal flux coordinate of $\rho_{\text{pol}} \approx 0.90$. The data is quite scattered, but the main part of the data clusters around 200 eV despite the large variation of heating power. In this case, the data of the low-density branch (semi-transparent symbols) is within the errors of the other data points indicating that the temperatures of the thermal plasma are very similar for the whole

density and power range.

The data of \bar{T} and $T_{e,\text{edge}}$ as depicted in figure 5 suggests that the edge temperature at the L-H transition is very similar for a very broad range of input powers and isotope masses as found above for the profiles at the same density (figure 4). However, the data of \bar{T} and $T_{e,\text{edge}}$ extends this result to the whole range of densities, so that we can conclude that the mechanism of the L-H transition requires or causes very similar electron temperatures at the edge. Comparably similar electron temperatures at the L-H transition in JET were already reported in references [4] and [5] for deuterium plasmas. Obviously, this holds likewise for other isotope masses including tritium and is valid for a wide range of input power and densities. This agrees also with results from ASDEX Upgrade, where similar edge temperatures at the L-H transition were found for a wide range of densities [34] and different isotopes [16], and a regression for the edge electron temperature at the L-H transition [35] predicts 200 eV to 300 eV for the present conditions, which is in good agreement with our data.

The fact that the electron temperature profiles are very similar at the L-H transition despite the large variation of heating power implies that the plasma transport must be very different for the different isotopes and densities. Otherwise the global power balance $P_{\text{sep}} \approx -\bar{n}_e \chi_{\text{eff}} S \nabla T$ with an effective heat diffusivity χ_{eff} assuming a single fluid with $T = T_i = T_e$ and a constant gradient across the radius would be violated. We can estimate χ_{eff} directly from our data set due to the relation

$$\frac{P_{\text{sep}}}{W_p} \approx \frac{1}{\tau_E} = \frac{S}{V} \frac{\chi_{\text{eff}} \nabla T}{\bar{T}} \approx \frac{S}{V} \frac{\chi_{\text{eff}}}{a} \quad (7)$$

with minor plasma radius a . The first approximation in equation 7 is due to the fact, that the bulk radiation is normally not subtracted in the definition of the energy confinement time τ_E . The last approximation makes use of the fact that the temperature fall of length is relatively constant for JET L-mode plasmas over large parts of the radius, i.e. outside of $R = 3.2$ m representing the major part of the plasma volume (see figure 4). With these approximations, we estimate the effective heat diffusivity

$$\chi_{\text{eff}} = (aV P_{\text{sep}}) / (W_p S) \quad (8)$$

from macroscopic quantities only. P_{sep} is determined according to equation 6, and all other parameters are taken from the equilibrium reconstruction.

The relation between the L-H transition power estimated as P_{sep} and the effective heat diffusivity χ_{eff} is shown in figure 6 for the whole considered data set, i.e. all densities including the low-density branch and all isotope mixtures. The data aligns well with a linear offset fit curve $P_{\text{sep}} = 2.0\chi_{\text{eff}} - 1.2$, which is only valid for the present plasma shape, toroidal magnetic field of 1.8 T and plasma current of 1.7 MA. The offset linear relation unifies data from the low-density branch (semi-transparent symbols) with data from plasmas with higher densities, and likewise aligns data with a variation of P_{sep} from below 1 MW for Ohmic T plasmas to maximum 6 MW of ICRH H plasmas. This indicates that the L-H power threshold is strongly linked to the effective heat transport in the L-mode phase prior to the L-H transition. Thus, the isotope effect of L-mode




Figure 6. Relation between L-H power threshold estimated as P_{sep} and the effective one fluid heat diffusivity $\chi_{\text{eff}} = (aVP_{\text{sep}}) / (W_p S)$. The offset linear relation suggests that the L-H power threshold is strongly linked to the transport in L-mode prior to the transition. Semi-transparent symbols indicate data from the low-density branch.

transport [36, 37, 38] indirectly determines the L-H power threshold, since the transport determines the heating power, P_{sep} , needed to reach the necessary temperature profiles at the edge, which are required to let the plasma transition into H-mode.

The linear offset relation between P_{sep} and χ_{eff} explains why the L-H power threshold for the pure deuterium plasma was lower than in the corresponding H-T mixture with 47 % T content (both have $A_{\text{eff}} \approx 2.0$): The transport in the pure deuterium plasma ($\chi_{\text{eff}} = 1.38 \text{ m}^2/\text{s}$) is lower than the transport in the H-T mixture ($\chi_{\text{eff}} = 2.16 \text{ m}^2/\text{s}$), and thus P_{sep} is lower. The same relation holds for the comparison between the D-T plasma ($\chi_{\text{eff}} = 1.245 \text{ m}^2/\text{s}$) and the H-T mixture with 74 % T content ($\chi_{\text{eff}} = 1.85 \text{ m}^2/\text{s}$), which both have $A_{\text{eff}} \approx 2.5$. This data clearly demonstrates, that the heat transport quantified by χ_{eff} can be very different for the same effective isotope mass A_{eff} , and consequently the L-H power threshold is very different. This implies

that A_{eff} is not a good parameter to describe the isotope effect of transport consistently, similarly as concluded for the isotope effect of P_{LH} .

5. Summary and conclusion

For JET plasmas in ILW in HT plasma configuration at a magnetic field of $B_t = 1.8$ T and a plasma current of $I_p = 1.7$ MA we estimated the L-H power threshold, P_{LH} , for an unprecedented variety of hydrogenic isotope compositions including tritium containing plasmas like H-T and D-T mixtures. This allowed for scanning the isotope mass and hence P_{LH} in a wide range for otherwise identical parameters. The main results of the investigation of this data set are the following:

- P_{LH} can differ significantly for plasmas with the same A_{eff} and otherwise constant parameters, when the isotope *composition* is different.
- The (edge) electron temperatures are very similar for a very wide range of densities, isotope masses and input powers.
- There is a linear offset relation between P_{sep} and the effective heat diffusivity χ_{eff} in the L-mode prior to the L-H transition.

This observations suggest that the L-mode transport eventually determines P_{LH} , since it is the transport, that defines how much input power is needed to reach the edge temperatures required for the L-H transition. Due to this, the isotope dependencies of L-mode transport directly pass into the isotope dependencies of P_{LH} . Consequently, P_{LH} is different for the same A_{eff} and otherwise constant parameters due to the L-mode transport, which obviously differed in the considered cases with different isotope composition.

It is important to understand the isotope dependence of the L-H power threshold, in order to correctly extrapolate P_{LH} from data of existing devices mostly operated with hydrogen or deuterium plasmas to future tokamak fusion reactors, which most probably will operate in H-mode with a D-T fuel mixture. Existing approaches for extrapolations or predictions sometimes employ the effective isotope mass, A_{eff} , in scaling laws [19] or for fluid simulations in order take into account isotope effects of P_{LH} . As we demonstrated with our data set of P_{LH} , the approach to rely only on A_{eff} for factoring in isotope effects will fail to accurately predict P_{LH} , especially when isotope mixtures like H-T and D-T are compared with their pure or mixed counterparts with the same effective isotope mass. Our data suggests that A_{eff} is not only insufficient to describe the isotope effect of P_{LH} consistently, but it also fails to correctly describe the isotope effect of transport, since different isotope compositions with the same A_{eff} showed different χ_{eff} .

Based on our findings, we conclude that for a quantitative prediction of P_{LH} it is necessary to (i) understand why the L-H transition is associated with critical temperature profiles, which are very similar for a large variation of densities and isotope masses, (ii) investigate whether and how the critical temperature profiles vary

with magnetic field, plasma current, plasma shape and other parameters, and (iii) quantitatively predict the L-mode transport under these conditions.

Edge turbulence codes might be best suited to provide this information, since they deliver the relevant anomalous transport levels at the edge, and intrinsically contain the physics of the turbulence suppression, which correlates with the confinement improvement at the L-H transition [39]. Recent developments of turbulence codes like XGC [40], GRILLIX [41] and GENE-X [42] raise hope, that a comprehensive understanding of the isotope effect of the L-H transition might be available in the near future. They self-consistently provide the density and temperature profiles at the edge including the radial electric field profile due to their global nature, are able to cope with the complex edge geometry including the X-point and open field lines in the SOL, and include neoclassical effects.

For plasmas of mixed isotopes, the interplay of the different ion species must be taken into account by representing them in separate isotope fluids (instead of just considering a single ion fluid with an effective isotope mass A_{eff}), which might increase the demand for computational power of turbulence simulations, but seems to be essential for correct predictions of the edge transport and, thus, the L-H power threshold. A simulation code being able to reliably provide realistic transport levels at the edge for different isotope composition would not only advance the understanding of L-H transition physics, but would likewise be of great value for the understanding of the isotope effect of transport and, hence, the prediction of burning fusion plasmas in tokamaks.

Acknowledgments

This work has been carried out within the framework of the EUROfusion Consortium, funded by the European Union via the Euratom Research and Training Programme (Grant Agreement No 101052200 – EUROfusion). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Commission. Neither the European Union nor the European Commission can be held responsible for them. G. Birkenmeier received funding from the Helmholtz Association under grant no. VH-NG-1350.

- [1] M. Shimada et al., Nucl. Fusion 47 (2007) S1, <https://iopscience.iop.org/article/10.1088/0029-5515/47/6/S01/meta>
- [2] M. Siccinio et al., Fusion Engineering and Design, Volume 176, March 2022, 113047, <https://doi.org/10.1016/j.fusengdes.2022.113047>
- [3] Y. R. Martin et al., Journal of Physics: Conference Series 123 (2008) 012033
- [4] C. F. Maggi et al., Nucl. Fusion 54 (2014) 023007
- [5] Y. Andrew et al., Plasma Phys. Control. Fusion 50 (2008) 124053
- [6] P. Gohil et al., Nucl. Fusion 50 (2010) 064011
- [7] L. Chen et al., Nucl. Fusion 56 (2016) 056013
- [8] J. Hillesheim et al., “Implications of JET-ILW L-H Transition Studies for ITER”, Preprint: 2016 IAEA Fusion Energy Conference, Kyoto, EX/5-2,

- <https://nucleus.iaea.org/sites/fusionportal/Shared%20Documents/FEC%202018/fec2018-preprints/preprint0346.pdf>
- [9] C. Bourdelle et al., Nucl. Fusion 54 (2014) 022001
 - [10] W. L. Zhong et al., Nucl. Fusion 60 (2020) 082002, <https://iopscience.iop.org/article/10.1088/1741-4326/ab868d/meta>
 - [11] Y. Andrew et al., Plasma Phys. Control. Fusion 48 (2006) 479
 - [12] A. R. Field et al., Plasma Phys. Control. Fusion 46 (2004) 981-1007
 - [13] R. Wenninger et al., Nucl. Fusion 57 (2017) 016011
 - [14] ITER Research Plan within Staged Approach, ITR-Report 18-003 (2018), <https://www.iter.org/technical-reports?id=9>
 - [15] P. Gohil, et al., Nucl. Fusion 51 (2011) 103020
 - [16] F. Ryter et al., Plasma Phys. Control. Fusion 58 (2016) 014007
 - [17] U. Plank et al., Nucl. Fusion 60 (2020) 074001
 - [18] L.M. Shao et al., Nucl. Fusion 61 (2021) 016010, <https://iopscience.iop.org/article/10.1088/1741-4326/abfb34/meta>
 - [19] E. Righi et al., Nucl. Fusion 39 (1999) 309
 - [20] G. Birkenmeier et al., Nucl. Fusion 62 (2022) 086005, <https://doi.org/10.1088/1741-4326/ac6dbb>
 - [21] U. Kruezi et al., JINST 15 (2020) C01032
 - [22] S. Vartanian et al., Fusion Engineering and Design Volume 170, September 2021, 112511, <https://www.sciencedirect.com/science/article/abs/pii/S0920379621002878>
 - [23] E. R. Solano et al., Nucl. Fusion 62 (2022) 076026, <https://iopscience.iop.org/article/10.1088/1741-4326/ac4ed8/meta>
 - [24] F. Ryter et al., Plasma Phys. Control. Fusion 36 (1994) A99, <https://iopscience.iop.org/article/10.1088/0741-3335/36/7A/011>
 - [25] T.N. Carlstrom et al., Plasma Phys. Control. Fusion 36 (1994) A147, <https://iopscience.iop.org/article/10.1088/0741-3335/36/7A/019/meta>
 - [26] J.E. Rice et al., Nucl. Fusion 39 (1999) 1175, <https://iopscience.iop.org/article/10.1088/0029-5515/39/9/310/meta>
 - [27] E. R. Solano et al., Nucl. Fusion 61 (2021) 124001, <https://doi.org/10.1088/1741-4326/ac2b76>
 - [28] C. Guillemaut et al., Phys. Scr. 2016 (2016) 014005, <https://iopscience.iop.org/article/10.1088/0031-8949/2015/T167/014005/meta>
 - [29] E. Lerche et al., Nucl. Fusion 56 (2016) 036022, <https://iopscience.iop.org/article/10.1088/0029-5515/56/3/036022/meta>
 - [30] F. Ryter et al., Nucl. Fusion 53 (2013) 113003
 - [31] L.-G. Eriksson and T. Hellsten, Physica Scripta 52 (1995) 70-79, <https://iopscience.iop.org/article/10.1088/0031-8949/52/1/014/meta>
 - [32] D.M.A. Taylor et al., Plasma Phys. Control. Fusion 64 (2022) 055015, <https://iopscience.iop.org/article/10.1088/1361-6587/ac5e6b>
 - [33] C.D. Challis et al., Nucl. Fusion 29 (1989) 563, <https://iopscience.iop.org/article/10.1088/0029-5515/29/4/002/meta>
 - [34] P. Sauter et al., Nucl. Fusion 52 (2012) 012001, <http://dx.doi.org/10.1088/0029-5515/52/1/012001>
 - [35] W. Suttrop et al., Plasma Phys. Control. Fusion 39 (1997) 2051, <https://iopscience.iop.org/article/10.1088/0741-3335/39/12/008>
 - [36] C. F. Maggi et al., Plasma Phys. Control. Fusion 60 (2018) 014045
 - [37] N. Bonanomi et al., Nucl. Fusion 59 (2019) 126025, <https://iopscience.iop.org/article/10.1088/1741-4326/ab3ecc/meta>
 - [38] P.A. Schneider et al., Nucl. Fusion 57 (2017) 066003, <https://iopscience.iop.org/article/10.1088/1741-4326/aa65b3/meta>
 - [39] F. Wagner, Plasma Phys. Control. Fusion 49 (2007) B1, <https://iopscience.iop.org/article/10.1088/0741-3335/49/12B/S01/pdf>
 - [40] C. Chang, S. Ku, G. Tynan et al., Physical Review Letters 118

- (2017) 175001, ISSN 0031-9007, doi:10.1103/PhysRevLett.118.175001,
<http://link.aps.org/doi/10.1103/PhysRevLett.118.175001>
- [41] W. Zholobenko, A. Stegmeir, T. Body et al., Contributions to Plasma Physics 60, ISSN 0863-1042,
doi:10.1002/ctpp.201900131, <https://onlinelibrary.wiley.com/doi/10.1002/ctpp.201900131>
- [42] D. Michels, P. Ulbl, W. Zholobenko et al., Physics of Plasmas 29 (2022) 032307, ISSN 1070-664X,
doi:10.1063/5.0082413, <https://aip.scitation.org/doi/10.1063/5.0082413>