

Research Articles | Behavioral/Cognitive

Connectivity profile and function of uniquely human cortical areas

<https://doi.org/10.1523/JNEUROSCI.2017-24.2025>

Received: 25 October 2024
Revised: 10 January 2025
Accepted: 15 February 2025

Copyright © 2025 Bryant et al.

This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

This Early Release article has been peer reviewed and accepted, but has not been through the composition and copyediting processes. The final version may differ slightly in style or formatting and will contain links to any extended data.

Alerts: Sign up at www.jneurosci.org/alerts to receive customized email alerts when the fully formatted version of this article is published.

Connectivity profile and function of uniquely human cortical areas

Katherine L. Bryant^{1,2}, Julia Camilleri³, Shaun Warrington⁴, Guilherme Blazquez Freches⁵, Stamatios N. Sotiropoulos^{4,6}, Saad Jbabdi¹, Simon Eickhoff^{3,7}, Rogier B. Mars^{1,5}

¹Wellcome Centre for Integrative Neuroimaging, Centre for Functional MRI of the Brain (FMRIB), Nuffield Department of Clinical Neurosciences, John Radcliffe Hospital, University of Oxford, Oxford, United Kingdom

²Institute for Language, Cognition, and the Brain (ILCB), Université Aix-Marseille, Marseille, France

³Institute of Neuroscience and Medicine: Brain and Behavior (INM-7), Research Centre Jülich, Jülich, Germany

⁴Sir Peter Mansfield Imaging Centre, School of Medicine, University of Nottingham, Nottingham, United Kingdom

⁵Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen, The Netherlands

⁶National Institute for Health Research (NIHR) Nottingham Biomedical Research Centre, Queens Medical Centre, Nottingham, United Kingdom

⁷Institute of Systems Neuroscience, Medical Faculty, Heinrich Heine University Düsseldorf, Düsseldorf, Germany

Abbreviated title: Uniquely human brain organization

2 figures, extended data (6 figures, 2 tables)

Number of words: 110 (abstract), 579 (introduction), 1500 (discussion)

Correspondence

Rogier B. Mars

Wellcome Centre for Integrative Neuroimaging

John Radcliffe Hospital

32 Headley Way
33 Oxford OX3 9DU
34 United Kingdom
35 rogier.mars@ndcn.ox.ac.uk
36

37 **Acknowledgements**

38 The work of RBM and KLM was supported by the Biotechnology and Biological Sciences
39 Research Council UK [BB/N019814/1] to RBM. RBM was also supported by the EPA
40 Cephalosporin Fund. The Wellcome Centre for Integrative Neuroimaging is supported by
41 core funding of the Wellcome Trust [203129/Z/16/Z]. SNS was supported by an ERC
42 Consolidator Grant [ERC CoG 101000969]. For the purposes of Open Access, the author
43 has applied a CC BY public copyright license to any Author Accepted Manuscript version
44 arising from this submission.

45

46 **Abstract**

47

48 Determining the brain specializations unique to humans requires directly comparative
49 anatomical information from other primates, especially our closest relatives. Human
50 (*Homo sapiens*) (m/f), chimpanzee (*Pan troglodytes*) (f), and rhesus macaque (*Macaca*
51 *mulatta*) (m/f) white matter atlases were used to create connectivity blueprints, i.e.,
52 descriptions of the cortical grey matter in terms of the connectivity with homologous white
53 matter tracts. This allowed a quantitative comparative of cortical organization across the
54 species. We identified human-unique connectivity profiles concentrated in temporal and
55 parietal cortices, and hominid-unique organization in prefrontal cortex. Functional
56 decoding revealed human-unique hotspots correlated with language processing and social
57 cognition. Overall, our results counter models that assign primacy to prefrontal cortex for
58 human uniqueness.

59

60

61 **Significance statement**

62

63 Understanding what makes the human brain unique requires direct comparisons with other
64 primates, particularly our closest relatives. Using connectivity blueprints, we compared to
65 cortical organization of the human to that of the macaque and, for the first time, the
66 chimpanzee. This approach revealed human-specific connectivity patterns in the temporal
67 and parietal lobes, regions linked to language and social cognition. These findings
68 challenge traditional views that prioritize the prefrontal cortex in defining human cognitive
69 uniqueness, emphasizing instead the importance of temporal and parietal cortical
70 evolution in shaping our species' abilities.

71

72 Introduction

73

74 Our human behavioral repertoire enables us to spread across the globe into a much
75 greater variety of niches than any other primate. Various behavioral innovations have
76 alternatively been suggested to characterize our abilities, including our collaborative social
77 abilities, tool use, ability for mental time travel, and spoken language (Healy, 2021;
78 Suddendorf et al., 2022; Tomasello and Vaish, 2013).

79

80 Understanding the basis of uniquely human behavior requires a comparison of our brain
81 to that of our closest primate relatives. Such comparisons tend to focus on measures of
82 size, highlighting that the human neocortex or cerebellum is expanded (Barton and
83 Venditti, 2014), that certain areas are preferentially expanded (Donahue et al., 2018), or
84 that the absolute number of neurons in the human brain outstrips that of other primates
85 (Herculano-Houzel, 2012). None of these measures, however, provides a link to the
86 behavior that the brain produces and that, ultimately, is the likely target of selection. In
87 contrast, work in neuroimaging has highlighted measures of brain organization at the level
88 of areal connections that do have predictive value for the function of parts of the brain
89 (Mars et al., 2018a; Saygin et al., 2016). Hence, the level of large-scale connections
90 between brain areas is a more suitable level of between-species comparison of brain
91 organization when one wants to understand the unique abilities of the human brain in the
92 context of other primates.

93

94 Connectivity can now be studied at the whole brain level using diffusion MRI and
95 associated tractography algorithms, offering a new type of data for comparative and
96 evolutionary neuroscience (Thiebaut de Schotten and Forkel, 2022). Recent work has
97 created standardized protocols for reconstructing the major fiber pathways of the primate
98 brain, creating white matter atlases of the human, the developing human, and the
99 macaque monkey brain (Mars et al., 2018b; Warrington et al., 2022). These methods
100 characterize the cortical areas of each species' brain in terms of its connectivity with major
101 white matter bundles, known to be homologous among primates. By describing all cortical
102 areas of all brains in terms of connectivity to homologous tracts, we, in effect, place all the
103 brains within a *common connectivity space*. This allows a quantitative comparison of brain
104 organization across species (Mars et al., 2021). While previous studies focused on
105 comparisons of the human brain with that of the most-often studied primate, the macaque,
106 here we additionally exploit our recently developed comprehensive white matter atlases of
107 the chimpanzee (Bryant et al., 2020), which allows us to directly compare humans with our
108 closest relatives, as well as the macaque. To our knowledge, this is the first time
109 connectional organization of the entire cortex is compared between these species,
110 although earlier comparisons of connections with a few specific tracts have been reported
111 (Hecht et al., 2015; Sierpowska et al., 2022).

112

We described each point on the cortical surface of the human and chimpanzee brain as a vector of connectivity probabilities with 18 white matter fiber bundles that are homologous across species. We can then quantify *which* areas of the human brain diverge in terms of connectivity from those of the other species. Next, we assess *how* the connectivity profile of areas of divergence in humans differs from that of the closest match in the other species by identifying which connections are driving the observed differences in brain organization. Finally, we use meta-analytic data on functional brain activation to investigate the functional *roles* of divergent regions in the human brain, linking the anatomical differences between species' brains to behavior.

Materials and Methods

Human data

Thirty human subjects (16 female, ages 22-35) were selected from the in vivo diffusion MRI data provided by the Human Connectome Project (HCP), WU-Minn Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by 16 NIH Institutes and Centers and the McDonnell Center for Systems Neuroscience at Washington University (Van Essen et al., 2013). Minimally preprocessed datasets from the Q2 public data release were used. Data acquisition and preprocessing methods have been previously described (Glasser et al., 2013; Sotiropoulos et al., 2013). Briefly, 1.25 mm isotropic resolution diffusion-weighted data were collected on a 3T Siemens Skyra scanner with a slice-accelerated gradient echo EPI readout. Q-space sampling included 3 shells at $b = 1000, 2000, \text{ and } 3000 \text{ s/mm}^2$. Ninety diffusion encoding gradient directions and 6 $b = 0$ s were obtained twice for each shell, with the phase-encoding direction reversed. An MPRAGE sequence was used to acquire T1-weighted (T1w) images at .7 mm isotropic resolution, then aligned to diffusion space using the HCP minimal preprocessing pipeline (Glasser et al., 2013). Diffusion-weighted images were processed with FSL, using FMRIB's Diffusion Toolbox and bedpostX (Behrens et al., 2007). A high-resolution 164k surface mesh (~164,000 vertices per hemisphere), as well as a lower-resolution mesh (32k) were generated using the PostFreeSurfer pipeline.

Chimpanzee data

Chimpanzee (*Pan troglodytes*; $n=23, 26 \pm 11 \text{ yrs}$, all female) MR scans were obtained from an archive hosted by the National Chimpanzee Brain Resource. Scans acquired prior to the 2015 implementation of U.S. Fish and Wildlife Service and National Institutes of Health regulations governing research with chimpanzees. All the scans reported here were collected as part of a grant to study aging in female primates, were completed by 2012, and have been used in previous studies (Autrey et al., 2014; Bryant et al., 2020, 2019). Chimpanzees were housed at the Emory National Primate Research Center (ENPRC, Atlanta, Georgia, USA) and all procedures were carried out in accordance with protocols approved by the ENPRC and the Emory University Institutional Animal Care and Use Committee (IACUC approval #YER-2001206).

Following standard ENPRC veterinary procedures, chimpanzee subjects were immobilized with ketamine injections (2–6 mg/kg, i.m.), then anesthetized with an intravenous propofol drip (10 mg/kg/h) prior to scanning. Subjects remained sedated for the duration of the scans as well as the time required for transport between the scanner and their home cage. Primates were housed in a single cage for 6–12 h after scanning to recover from the effects of anesthesia before being returned to their home cage and cage mates. Veterinary and research staff evaluated the well-being of chimpanzees twice daily after the scan for possible post-anesthesia distress.

MR scanning protocols and preprocessing for the chimpanzee dataset have been described in detail previously (Autrey et al., 2014). Briefly, anatomical and diffusion MR scans were acquired in vivo in a Siemens 3T Trio scanner (Siemens Medical System, Malvern, PA, USA). Diffusion-weighted MRI data were collected with a single-shot, pulsed-gradient spin-echo echo-planar imaging sequence. Parameters were as follows: 41 slices were scanned at a voxel size of 1.8 mm³, TR/TE: 5900 ms/86 ms, matrix size: 72x128. Two diffusion-weighted images were acquired for each of 60 diffusion directions ($b=1000$ s/mm²), each with 1 of the possible left–right phase-encoding directions and 8 averages, allowing for correction of susceptibility-related distortion (Andersson et al., 2003). For each averaged diffusion-weighted image, six images without diffusion weighting ($b = 0$ s/mm²) were also acquired. High-resolution T1w images were acquired with a MPRAGE sequence. Diffusion-weighted images were processed using FMRIB's Diffusion Toolbox and bedpost (Behrens et al., 2007). Template generation for chimpanzees (previously described in detail (Li et al., 2010)); involved the PreFreeSurfer pipeline was used to align the T1w and T2w volumes of 29 individual chimpanzees to native anterior commissure-posterior commissure space. Cortical surfaces and registrations to a population-specific chimpanzee template were generated using a modified version of the HCP minimal preprocessing pipeline (Glasser et al., 2013). The PostFreeSurfer pipeline was used to produce a high-resolution 164k surface mesh and a lower-resolution mesh (20k).

Macaque data

Eight post mortem macaque brain scans (*Macaca mulatta*, $n=8$; six male, age range 4-14 years) were acquired using a 7T magnet with an Agilent DirectDrive console (Agilent Technologies, Santa Clara, CA, USA). Acquisition and preprocessing have been detailed previously (Folloni et al., 2019). In brief, a 2D diffusion-weighted spin-echo protocol was implemented (DW-SEMS, TE/TR: 25 ms/10 s; matrix size: 128 x 128; resolution: 0.6 x 0.6 mm; number of slices: 128; slice thickness: 0.6 mm). Nine non-diffusion-weighted ($b=0$ s/mm²) and 131 diffusion-weighted ($b=4000$ s/mm²) volumes were acquired with diffusion directions distributed over the whole sphere. The $b=0$ images were averaged and spatial signal inhomogeneities were restored. Ex vivo tissue usually has reduced diffusivity, necessitating larger b -values to achieve equivalent diffusion contrast to in vivo data; this was achieved here by increasing the diffusion sensitization from $b = 1000$ to 4000 s/mm².

Diffusion-weighted images were processed using the same method as chimpanzees, described above. The cortical surface of one macaque with high quality structural MRI was reconstructed using a modified version of the HCP pipeline, nonlinearly registered to the other brains using FSL's FNIRT, warped to the other macaque brains and transformed to F99 standard space (Van ESSEN, 2002).

Between-species comparison based on white matter tracts

Eighteen major white matter bundles were reconstructed for all three species using probabilistic tractography (Behrens et al., 2007). A set of standardized masks previously developed for the human, chimpanzee, and macaque were used to reconstruct tracts based on objective anatomical landmarks that could be identified in all species. The logic behind this approach is that a set of seed, waypoint, stop, and exclusion masks are used to define the body of any white matter tract; the tractography algorithm is then free to reconstruct the rest of the bundle, including its grey matter termination points. In this way, we have something we can objectively define as homologous across the species (the body of the tract based on anatomical criteria) and something that varies across species and is the target of our investigation (the grey matter terminations) (Mars et al., 2018b; Warrington et al., 2020).

All combinations of seed, waypoint, stop, and exclusion masks are described in detail in previous communications (Mars et al., 2018b; Warrington et al., 2020, 2020). The white matter tracts studied in the present study were the anterior commissure (AC), arcuate fascicle (AF), peri-genual, dorsal, and temporal subdivisions of the cingulum bundle (CBP, CBD, and CBT), corticospinal tract (CST), frontal aslant (FA), forceps major (FMA), forceps minor (FMI), fornix (FX), inferior fronto-occipital fascicle (IFO), inferior longitudinal fascicle (ILF), middle longitudinal fascicle (MdLF), first, second, and third branches of the superior longitudinal fascicle (SLF1, SLF2, and SLF3), uncinate fascicle (UNC), and vertical occipital fascicle (VOF).

To assess the connectivity of each vertex of the cortical surface with each white matter fiber bundle we created (surface) x (tract) matrices which we term *connectivity blueprints*. First, tractography is performed from each vertex of the cortical surface towards all voxels of the whole brain white matter, creating a (brain) x (surface) matrix of connectivity. Then, each tract's tractogram, of the format (brain) x (tract), is multiplied by the transposed (brain) x (surface) matrix, resulting in the (surface) x (tract) connectivity blueprint. The columns of this blueprint represent the surface projection of each tract and the rows of the blueprint present the connectivity profile of each vertex of the cortical surface. This method was first applied by Mars and colleagues (Mars et al., 2018b) and is now implemented in FSL's XTRACT tool (Assimopoulos et al., 2024; Warrington et al., 2022).

Blueprints were averaged across subjects in each species to create a species-specific connectivity blueprint. Connectivity profiles can be compared across species by calculating the (vertex) x (vertex) KL divergence between two common connectivity spaces. The best match of a vertex in one species is then found by finding the vertices with the lowest KL value (<2) in the other species. A spatial map of divergence of connectivity of one brain compared to another can be established by assigning to each vertex of the first brain the smallest KL value (minKL) across all vertices in the second brain.

Functional Decoding

To assess the functional roles of the areas of the human cortex that showed the greatest difference with the chimpanzee and the human we used BrainMap, a publicly available meta-analytic database of functional activation studies (www.brainmap.org) (Fox and Lancaster, 2002). BrainMap uses a structured standardized coding scheme to describe published human functional neuroimaging results. In particular, *Behavioral Domains* are categories and subcategories that aim to classify the cognitive functions likely to be isolated by any experimental contrast.

Functional decoding was done as follows. First, the cortex was divided into distinct regions according to the Glasser parcellation (Glasser et al., 2016). Each region was assigned the maximum within-region divergence score, i.e. the divergence value from the vertex that had the highest minKL value in the region. Second, we queried the BrainMap database in 2019 to assign the functional profile of these regions using forward inference (Eickhoff et al., 2011). Using forward inference, a cluster's functional profile is determined by identifying taxonomic labels for which the probability of finding activation in the respective cluster was significantly higher than the a priori chance (across the entire database) of finding activation in that particular cluster. Significance was established using a binomial test ($p < 0.05$, FDR corrected (Genovese et al., 2002)). In other words, we tested whether the conditional probability of activation given a particular label [$P(\text{Activation}|\text{Task})$] was higher than the baseline probability of activating the brain region in question per se [$P(\text{Activation})$].

Results

Between-species comparison of connectivity blueprints

For the human, chimpanzee, and macaque monkey brain, we established the connectivity of each part or vertex of the cortical surface with each of 18 white matter tracts that were

determined in a homologous fashion in all three species. We term this surface by tract matrix the *connectivity blueprint*. The rows of this matrix describe the profile of connectivity of a given vertex of the cortical surface with each of the white matter tracts. The connectivity profile of any human vertex can be compared to that of each chimpanzee and macaque vertex by calculating the Kullback-Leibler (KL) divergence between connectivity profiles (Mars et al., 2018b). The best matching vertex in the non-human species is the one with the minimum KL value. Overall spatial maps of divergence of the human brain to the other species is then visualized by plotting the minimum KL value for each human vertex. When comparing the human to the chimpanzee brain, this shows large zones of divergence in the middle temporal lobe, temporoparietal cortex, and lateral frontal cortex with a particular hotspot in the dorsal frontal cortex (Fig. 1, left).

The divergence of the human brain from the chimpanzee brain can be compared to the divergence of the human brain with the macaque. The distribution of minimum KL values when comparing the human and the chimpanzee differs from that when comparing the human and the macaque (Kolmogorov-Smirnov test $p < 0.001$ for both hemispheres). Plotting the distribution of minimum KL values based on the union of KL values with the chimpanzee and the macaque indeed shows broader differences between the human and the macaque (Fig 1, middle). Indeed, if we color each human vertex's divergence based on the species in which it was greatest we see increases in divergence in the anterior ventral frontal cortex and posterior parietal cortex (Fig 1, right).

Divergence between the human brain and both the chimpanzee and macaque were evident in the dorsal frontal cortex. The vertices of high divergence overlap with anterior area 6, the inferior 6-8 transition area, and the frontal eye fields (Glasser et al., 2016). The connectivity profile of this area is dominated by the frontal-parietal superior longitudinal fascicle, in particular the second branch (SLF2) (Thiebaut de Schotten et al., 2011) (Fig. 1; see Extended Data Fig. 1 for full connectivity profiles). Using the common connectivity space, we can determine which vertices in the chimpanzee and the macaque have a connectivity profile that is the least different from that of the human. Extracting the connectivity of these vertices shows that even these do not show strong SLF2 connectivity (Fig. 1, Extended Data Fig. 2). We thus conclude that strong SLF2 connectivity in this part of dorsal frontal cortex is driving the divergence in brain organization between the human and the other two primates.

Extensive differences between the human and non-human brains were found in ventral frontal cortex and middle temporal gyrus. Both these hotspots of divergence were driven by more extensive connectivity of the arcuate fascicle (AF) in humans (Fig. 1). Such AF connectivity in the human brain has been shown before (Rilling et al., 2008; Sierpowska et al., 2022), but the comparison of the human with the chimpanzee on the one hand and the chimpanzee and macaque on the other shows a dissociation between frontal and temporal cortex. While the best matching vertices for middle temporal cortex showed a lack of innervation of the AF in both chimpanzees and macaques, the best matching

vertices to the anterior ventral frontal cortex show some AF in the chimpanzee, but none in the macaque. This suggests a scenario where the extension of the AF occurred gradually, with frontal expansions occurring in the ape lineage, preceding temporal expansions into the middle temporal cortex in the human lineage.

On the medial wall we noticed a hotspot of divergence in medial parietal area 7. This divergence seems mostly driven by small changes in multiple tracts (Fig. S1), rather than a clear elaboration of a single tract, as is the case for some of the divergent areas discussed above. However, the strongest connection of this area, SLF1, does seem more focal in the human than in the best matching vertices in the other two species.

Functional decoding of divergent regions

Next, we turned to a database of functional neuroimaging studies (brainmap.org)(Fox et al., 2005) to assess the functional role of these regions. We assessed if, for a given behavioral domain, the probability of finding activation of a region was significantly higher than the *a priori* chance, so-called *forward inference*. This approach allows a functional characterization of the areas we identified as structurally divergent from other primate brains (Fig. 2; Extended Data Table 2-1 and 2-2).

It is important to point out that the specificity of the decoding results can only be as good as the taxonomy of the BrainMap database. Thus, our results should not be taken such that any Behavioral Domain associated with an area constitutes the unique role of that area. Rather, the Behavioral Domain indicates the involvement of the area but does not claim the brain region is limited to that Domain. We provide two tables showing the functional decoding of regions based on high divergence between the human and the chimpanzee (Extended Data Table 2-1) and between the human and the macaque monkey (Extended Data Table 2-2). Behavioral domains for significant decoding and likelihood ratios are reported. Regions are labeled according to the atlas of Glasser and co-workers (2016).

For the three dorsal frontal regions mentioned above, the behavioral domains most likely to activate them include spatial cognition, working memory, and reasoning. Some of these regions have previously been identified as part of the so-called multiple demand network (Assem et al., 2020), a network of mostly parietal and frontal regions that consistently activate for a range of high-level cognitive tasks. Although homologs of this network exist in the macaque, recent comparative work shows that the connections between these regions are much more extensive in the human (Karadachka et al., 2023). As such, it has been suggested that human domain-general knowledge has a precursor in parietal-frontal network originally evolved for visuomotor control in early primates (Genovesio et al., 2014). The current results extend this finding to our nearest animal relative, and directly link

anatomical differences to functional domains associated with the multiple demand network.

Consistent with the role of the AF in human language, functional decoding of both the middle temporal and ventral frontal cortex in the left hemisphere yielded the behavioral domain 'language' prominently. However, it was clear that the AF extension, especially in the temporal cortex, was bilateral. Decoding of the right middle temporal cortex yielded the domain 'emotion'. Although the function of right temporal association cortices is yet not well-characterized in the fMRI literature, lesion studies suggest they play a role in nonverbal semantic social cognition (Binney et al., 2012). Importantly, these results speak against a language-only interpretation of AF extensions in the ape and human brain.

A prominent zone of divergence between the human brain and that of both the chimpanzee and macaque was in the posterior superior temporal cortex and inferior parietal lobule, together often referred to as the temporoparietal junction area (TPJ). This effect was particularly prominent in the right hemisphere. The right posterior TPJ especially has often been associated with the human ability to entertain others' belief states, so-called mentalizing or Theory of Mind (Schurz et al., 2017). The hotspot of divergence overlaps with this area, and functional decoding indeed shows 'social cognition' as its most significant behavioral domain. The human posterior TPJ shows strong connectivity to the inferior longitudinal fascicle (ILF), which is not present in the other two species (Extended Data Fig. 6). The ILF is part of the ventral visual pathway but extends into parietal cortex in anthropoid primates (Roumazeilles et al., 2022). It is thought that the ILF has expanded in great apes and that the dorsal component has a role in social cognition, allowing some of the temporal cortex machinery for visual processing to be adapted for social information processing (Pitcher and Ungerleider, 2020; Roumazeilles et al., 2020). The current results connect these two findings of TPJ's role in social cognition and ILF's prominent expansion by showing the TPJ is innervated by the ILF in the human.

Comparison of connectivity profiles across species based on a priori homologs

It is important to note that the above analyses select those vertices in the chimpanzee and macaque brain that have the least divergent connectivity profile with the chosen vertex in the human brain, independent of their location. This allows an unbiased assessment of divergence across the different species' brains. As has been shown previously, this analysis is capable of identifying homologous regions that are known to have similar connectivity profiles across species (Mars et al., 2018b) while not relying on priors. It is therefore more principled than comparing known homologs across species. For completion, however, we also present comparisons of the connectivity profiles of human areas with those of known homologs in the chimpanzee and macaque for all areas in Figure 1.

The left dorsal prefrontal region overlaps with anterior area 6, the inferior 6-8 transition area, and the frontal eye fields (Glasser et al., 2016). In humans, this area has much stronger connectivity to SLF2, compared to its best matching chimpanzee and macaque counterparts. We extracted the connectivity profiles of area FB in the chimpanzee (Bailey et al., 1950), which has been suggested to contain the frontal eye fields (Percheron et al., 2015), and macaque FEF (Petrides, 2005, p. 200). As with the best matching vertices, the human has much stronger SLF2 connectivity than the other species (Extended Data Fig. 1-2).

Anterior ventral frontal cortex in the human received innervations of the arcuate fascicle (AF), which was evident to a lesser extent in the chimpanzee and absent in the macaque. The human area of maximum divergence overlaps with area IFSa of Glasser and colleagues (Glasser et al., 2016). and area IFS of Neubert and co-workers (Neubert et al., 2014). The homolog of this area in the chimpanzee is difficult to establish. We extracted the connectivity profile of a vertex in area FCBm (Bailey et al., 1950) in the chimpanzee and on the posterior bank of the inferior branch of the arcuate sulcus in the macaque. In both cases, these locations are, if anything, quite posterior, and therefore more likely to detect AF connectivity than human IFS. Nevertheless, the pattern of most AF connectivity in the human, less in the chimpanzee, and very little in the macaque was replicated (Extended Data Fig. 1-3).

The human middle temporal gyrus shows strong AF connectivity, which is much lower even in the best matching areas in the other two species. When extracting the connectivity profile of middle temporal gyrus in the chimpanzee and macaque, this pattern of relatively reduced AF in the non-human primates is even stronger (Extended Data Figs. 1-4 and 1-5).

The right temporoparietal junction (TPJ) area in the human brain shows strong innervation of the ILF, which is not seen in the best matching vertices in the chimpanzee and macaque. The homolog of TPJ is difficult to establish. Although the area overlaps with area PGi of Glasser and colleagues (Glasser et al., 2016), it is uncertain whether it is homologous to area PG in the macaque (Pandya and Seltzer, 1982). Mars et al. (2012) identified two subregions of TPJ, which they labeled TPJp and TPJa, the posterior of which shows strong activation in social cognition tasks, as found in our decoding analysis. Connectivity profiles of regions in the macaque inferior parietal lobule do not show a prominent ILF, but rather the IFO and MdLF. In addition, the small macaque inferior parietal lobule shows strong connectivity with the AF, which does not extend ventrally as it does in the human, as discussed above (Extended Data Fig. 1-6).

Discussion

449

450 Comparing brain organization across species typically involves detailed analysis of small
451 parts of the brain using measures such as cytoarchitecture or transcriptomics on the one
452 hand or comparisons of large subdivisions using global measures such as relative brain
453 size on the other. In contrast, here we compared the organization of the human cortex
454 directly with that of two other species at a level of direct relevance to function: connectivity.
455 We exploit the availability of white matter atlases created using diffusion MRI to provide a
456 detailed comparison of cortical organization between the human brain and that of one of
457 its closest relatives, the chimpanzee, and the most often studied non-human primate, the
458 macaque monkey. We demonstrate uniquely human organization of large parts of
459 association cortex and relate them for the first time to the behavioral domains in which
460 they show functional activation.

461

462 Although most debates regarding what might be special about the human brain focus on
463 prefrontal cortex (Barton and Venditti, 2013; Donahue et al., 2018), the current results
464 demonstrate that major areas of difference between the human, chimpanzee, and
465 macaque are in other parts of association cortex. The most different region is in the middle
466 temporal gyrus. This region was previously identified in our human-macaque comparisons
467 (Mars et al., 2018b) and the current results extend this result to the human-chimpanzee
468 comparison. This change is primarily driven by the extension of the arcuate fascicle.
469 Arcuate expansion has been identified as a hall-mark for human language (Rilling et al.,
470 2008; Roelofs, 2014), but a focus solely on language might be a too narrow interpretation
471 of this major between-species difference. For instance, the arcuate expansion is bilateral
472 and, although right temporal cortex also has some language functions, our functional
473 decoding shows its involvement in other functions as well. Moreover, the arcuate extension
474 is partly driven by the short parietal-temporal aspect of the arcuate (Sierpowska et al.,
475 2022) integrating information processing between dorsal and ventral cortical pathways.

476

477 An important difference between the human-chimpanzee and the human-macaque
478 comparisons is in the ventral frontal cortex. Although the cortical territory termed 'Broca's
479 area' has been associated with uniquely human organization and function, the picture of
480 the precise pattern of evolutionary change is only now starting to become clear. When
481 comparing the human to the chimpanzee, there is no clear hotspot of change in ventral
482 prefrontal cortex, whereas this is clear in the human-macaque comparison. This result
483 extends earlier demonstrations of a differences in both area 44 and 45 between the adult
484 human and adult macaque brain, but only in area 45 between the adult macaque and the
485 human infant (Warrington et al., 2022). Another prominent frontal cortex difference
486 between the human and both non-human primates was in the strength of parietal-frontal
487 connections. Again, some of these differences had been identified in human-macaque
488 comparisons, but are now shown to be unique to the human lineage.

489

490 The between-species differences in temporal and temporo-parietal cortex are not solely
491 driven by the arcuate. It had previously been established that the temporal longitudinal

white matter pathways are more extensive and show more complex subdivisions in apes than in monkeys (Roumazeilles et al., 2020). Here, we demonstrate that the inferior longitudinal fascicle reaches part of the so-called temporoparietal junction area (TPJ) in the human. This area has previously been shown to share some anatomical and functional properties with face sensitive areas in the macaque middle superior temporal sulcus (Mars et al., 2013; Roumazeilles et al., 2021), but human TPJ seems to process the more complex information associated with human social cognition, by either entertaining others' belief states (Koster-Hale et al., 2017) or the difference between one's own and other's knowledge (Kolling et al., 2021).

Differences between the human and non-human primate are less prominent on the medial wall, but medial parietal cortex does show a hotspot of divergence between species. This dovetails with earlier reports comparing human and macaque (Mars et al., 2018b). Precuneus has previously been identified as a region of expansion in the brain of modern humans based on fossil endocasts (Bruner, 2018). Here, we show that such changes are accompanied by changes in connectivity profile, although it is unknown whether such changes have coincided.

Our approach of using white matter tracts as a common space in which to describe brain organization of the three species contrasts with that of a direct spatial registration of the brains based on sulcal morphology (Chaplin et al., 2013; Vickery et al., 2024). There are two reasons the common space approach is beneficial. First, the homology of sulci across the human, chimpanzee, and macaque is far from established. Major longitudinal sulci such as the macaque principal sulcus may not be homologous to any of the frontal sulci of the great apes (Petrides, 2005) and the pattern of smaller sulci is more complex in the human brain (Hathaway et al., 2023). Secondly, while sulcal-based registration might identify regional expansion and even relocation of certain cortical areas (Hill et al., 2010), these results do not speak to the different possible scenarios of evolutionary change that can accompany such changes, including whether a region has simply expanded or also changed its profile of connectivity with the rest of the brain (Eichert et al., 2020). In the latter case, the interaction of the region with other parts of the brain has changed, which likely results in different functional roles. Indeed, changes in connectivity of cortical areas have been proposed to be a prominent way in which brain organization changes throughout evolution (Krubitzer and Kaas, 2005).

Although our approach addresses problems of differences in brain size and morphology when comparing different species' brains, as any method it has some limitations that should be kept in mind when interpreting the results. Our definition of common tracts relies on the correct placement of seed, waypoint, stop, and exclusion masks that for the tractography recipe of each tract. Our approach has been to define masks based on explicit anatomical landmarks that can be recognized easily across species. Previous work has validated these recipes compared to known tracts in the human and the macaque (Mars et al., 2018b; Warrington et al., 2020) and definitions of new species are created in

as similar a way as possible. But we acknowledge that the tractography masks are the basis of the comparisons. The masks defined for the chimpanzee and their comparisons to the human and macaque have been the topic of a previous communication (Bryant et al., 2020). All recipes used in this approach and can be found on the website of the XTRACT tool; the modular organization of XTRACT means that researchers can easily substitute their own recipes and study the effects on the between-species comparisons.

Due to the limited availability of data from the chimpanzee, our sample only consisted of female subjects. Similarly, our age range is limited to young adults for all species. Although to our knowledge differences in connectivity across sex are limited to white matter volume and the strength of particular connections rather than the presence or absence of particular fiber bundles (Gong et al., 2011), subtle differences in connectivity across sexes and how these differences manifest themselves across species is an important avenue in research. Translational neuroscience has long been biased by inclusion of mostly single-sex data, while it is now known that sex differences occur even in rodent brains (Guma et al., 2024). The connectivity blueprint method has been used to compare young adult and infant humans (Warrington et al., 2022) and developmental changes in other species are the topic of ongoing research, where the data are available. However, for the current study, the single time point and sex bias in the data are a limitation of the scope.

Although comparison of the organization of the entire neocortex of the human to two other species of primate is unique, future work will strengthen and extend our results by inclusion of more species and direct comparisons across them. The current manuscript has focused on the human as the reference species, but a full understanding of primate phylogeny necessitates comparisons that are less human-centric. Using the same protocols as used in the present study, partial white matter atlases primate species are now available (Bryant et al., 2023, 2021) and work to extend these to include the same range of tracts as the present study are ongoing. Moreover, data-driven methods for identification of white matter tracts have also shown promise in comparative studies (Mars et al., 2019).

Overall, our results thus argue against a single explanatory factor or evolutionary event driving the uniquely human behavioral repertoire. While current theories on human brain uniqueness focus on changes to prefrontal areas, our findings support a two-step evolutionary process, in which changes in prefrontal cortex organization emerge prior to changes in temporal areas. Unlike global connectivity or gross anatomical approaches, anatomically informed comparative connectivity makes it possible to reveal major changes in multiple association fiber systems underlying a variety of cognitive functions that have changed in a stepwise manner in the great ape and human lineages.

References

577 Andersson, J.L.R., Skare, S., Ashburner, J., 2003. How to correct susceptibility distortions in
578 spin-echo echo-planar images: application to diffusion tensor imaging. *Neuroimage* 20,
579 870–888. [https://doi.org/10.1016/S1053-8119\(03\)00336-7](https://doi.org/10.1016/S1053-8119(03)00336-7)

580 Assem, M., Glasser, M.F., Van Essen, D.C., Duncan, J., 2020. A Domain-General Cognitive
581 Core Defined in Multimodally Parcellated Human Cortex. *Cereb Cortex* 30, 4361–4380.
582 <https://doi.org/10.1093/cercor/bhaa023>

583 Assimopoulos, S., Warrington, S., Bryant, K.L., Pszczolkowski, S., Jbabdi, S., Mars, R.B.,
584 Sotiropoulos, S.N., 2024. Generalising XTRACT tractography protocols across common
585 macaque brain templates. *Brain Struct Funct.* [https://doi.org/10.1007/s00429-024-02760-](https://doi.org/10.1007/s00429-024-02760-0)
586 0

587 Autrey, M.M., Reamer, L.A., Marengo, M.C., Sherwood, C.C., Herndon, J.G., Preuss, T.,
588 Schapiro, S.J., Hopkins, W.D., 2014. Age-related effects in the neocortical organization of
589 chimpanzees: Gray and white matter volume, cortical thickness, and gyrification.
590 *NeuroImage* 101, 59–67. <https://doi.org/10.1016/j.neuroimage.2014.06.053>

591 Bailey, P., Von Bonin, G., McCulloch, W., 1950. The isocortex of the chimpanzee. University of
592 Illinois Press, Urbana.

593 Barton, R.A., Venditti, C., 2014. Rapid evolution of the cerebellum in humans and other great
594 apes. *Curr Biol* 24, 2440–2444. <https://doi.org/10.1016/j.cub.2014.08.056>

595 Barton, R.A., Venditti, C., 2013. Human frontal lobes are not relatively large. *Proc Natl Acad Sci*
596 *USA* 110, 9001–9006. <https://doi.org/10.1073/pnas.1215723110>

597 Behrens, T.E.J., Berg, H.J., Jbabdi, S., Rushworth, M.F.S., Woolrich, M.W., 2007. Probabilistic
598 diffusion tractography with multiple fibre orientations: What can we gain? *Neuroimage* 34,
599 144–155. <https://doi.org/10.1016/j.neuroimage.2006.09.018>

600 Binney, R.J., Parker, G.J.M., Lambon Ralph, M.A., 2012. Convergent Connectivity and Graded
601 Specialization in the Rostral Human Temporal Lobe as Revealed by Diffusion-Weighted
602 Imaging Probabilistic Tractography. *Journal of Cognitive Neuroscience* 24, 1998–2014.
603 https://doi.org/10.1162/jocn_a_00263

604 Bruner, E., 2018. Human Paleoneurology and the Evolution of the Parietal Cortex. *Brain Behav*
605 *Evol* 91, 136–147. <https://doi.org/10.1159/000488889>

606 Bryant, K.L., Ardesch, D.J., Roumazeilles, L., Scholtens, L.H., Khrapitchev, A.A., Tendler, B.C.,
607 Wu, W., Miller, K.L., Sallet, J., van den Heuvel, M.P., Mars, R.B., 2021. Diffusion MRI
608 data, sulcal anatomy, and tractography for eight species from the Primate Brain Bank.
609 *Brain Struct Funct* 226, 2497–2509. <https://doi.org/10.1007/s00429-021-02268-x>

610 Bryant, K.L., Glasser, M.F., Li, L., Jae-Cheol Bae, J., Jacquez, N.J., Alarcón, L., Fields, A.,
611 Preuss, T.M., 2019. Organization of extrastriate and temporal cortex in chimpanzees
612 compared to humans and macaques. *Cortex, The Evolution of the Mind and the Brain*
613 118, 223–243. <https://doi.org/10.1016/j.cortex.2019.02.010>

614 Bryant, K.L., Li, L., Eichert, N., Mars, R.B., 2020. A comprehensive atlas of white matter tracts in
615 the chimpanzee. *PLoS Biol* 18, e3000971.

616 Bryant, K.L., Manger, P.R., Bertelsen, M.F., Khrapitchev, A.A., Sallet, J., Benn, R.A., Mars, R.B.,
617 2023. A map of white matter tracts in a lesser ape, the lar gibbon. *Brain Struct Funct.*
618 <https://doi.org/10.1007/s00429-023-02709-9>

Chaplin, T.A., Yu, H.-H., Soares, J.G.M., Gattass, R., Rosa, M.G.P., 2013. A Conserved Pattern of Differential Expansion of Cortical Areas in Simian Primates. *J Neurosci* 33, 15120–15125. <https://doi.org/10.1523/JNEUROSCI.2909-13.2013>

Donahue, C.J., Glasser, M.F., Preuss, T.M., Rilling, J.K., Van Essen, D.C., 2018. Quantitative assessment of prefrontal cortex in humans relative to nonhuman primates. *Proc Natl Acad Sci USA* 115, E5183–E5192. <https://doi.org/10.1073/pnas.1721653115>

Eichert, N., Robinson, E.C., Bryant, K.L., Jbabdi, S., Jenkinson, M., Li, L., Krug, K., Watkins, K.E., Mars, R.B., 2020. Cross-species cortical alignment identifies different types of anatomical reorganization in the primate temporal lobe. *eLife* 9, e53232. <https://doi.org/10.7554/eLife.53232>

Eickhoff, S.B., Bzdok, D., Laird, A.R., Roski, C., Caspers, S., Zilles, K., Fox, P.T., 2011. Co-activation patterns distinguish cortical modules, their connectivity and functional differentiation. *NeuroImage, Special Issue: Educational Neuroscience* 57, 938–949. <https://doi.org/10.1016/j.neuroimage.2011.05.021>

Folloni, D., Sallet, J., Khrapitchev, A.A., Sibson, N., Verhagen, L., Mars, R.B., 2019. Dichotomous organization of amygdala/temporal-prefrontal bundles in both humans and monkeys. *eLife* 8, e47175. <https://doi.org/10.7554/eLife.47175>

Fox, P.T., Laird, A.R., Fox, S.P., Fox, P.M., Uecker, A.M., Crank, M., Koenig, S.F., Lancaster, J.L., 2005. BrainMap taxonomy of experimental design: description and evaluation. *Hum Brain Mapp* 25, 185–198. <https://doi.org/10.1002/hbm.20141>

Fox, P.T., Lancaster, J.L., 2002. Mapping context and content: the BrainMap model. *Nat Rev Neurosci* 3, 319–321. <https://doi.org/10.1038/nrn789>

Genovese, C.R., Lazar, N.A., Nichols, T., 2002. Thresholding of Statistical Maps in Functional Neuroimaging Using the False Discovery Rate. *NeuroImage* 15, 870–878. <https://doi.org/10.1006/nimg.2001.1037>

Genovesio, A., Wise, S.P., Passingham, R.E., 2014. Prefrontal–parietal function: from foraging to foresight. *Trends Cogn Sci* 18, 72–81. <https://doi.org/10.1016/j.tics.2013.11.007>

Glasser, M.F., Coalson, T.S., Robinson, E.C., Hacker, C.D., Harwell, J., Yacoub, E., Ugurbil, K., Andersson, J., Beckmann, C.F., Jenkinson, M., Smith, S.M., Van Essen, D.C., 2016. A multi-modal parcellation of human cerebral cortex. *Nature* 536, 171–178.

Glasser, M.F., Sotiropoulos, S.N., Wilson, J.A., Coalson, T.S., Fischl, B., Andersson, J.L., Xu, J., Jbabdi, S., Webster, M., Polimeni, J.R., Van Essen, D.C., Jenkinson, M., 2013. The minimal preprocessing pipelines for the Human Connectome Project. *NeuroImage, Mapping the Connectome* 80, 105–124. <https://doi.org/10.1016/j.neuroimage.2013.04.127>

Gong, G., He, Y., Evans, A.C., 2011. Brain Connectivity: Gender Makes a Difference. *Neuroscientist* 17, 575–591. <https://doi.org/10.1177/1073858410386492>

Guma, E., Beauchamp, A., Liu, S., Levitis, E., Ellegood, J., Pham, L., Mars, R.B., Raznahan, A., Lerch, J.P., 2024. Comparative neuroimaging of sex differences in human and mouse brain anatomy. *eLife* 13, RP92200. <https://doi.org/10.7554/eLife.92200>

Hathaway, C.B., Voorhies, W.I., Sathishkumar, N., Mittal, C., Yao, J.K., Miller, J.A., Parker, B.J., Weiner, K.S., 2023. Defining putative tertiary sulci in lateral prefrontal cortex in

- chimpanzees using human predictions. *Brain Struct Funct*.
<https://doi.org/10.1007/s00429-023-02638-7>
- Healy, S.D., 2021. *Adaptation and the brain*. Oxford University Press, Oxford.
- Hecht, E.E., Gutman, D.A., Bradley, B.A., Preuss, T.M., Stout, D., 2015. Virtual dissection and comparative connectivity of the superior longitudinal fasciculus in chimpanzees and humans. *NeuroImage* 108, 124–137. <https://doi.org/10.1016/j.neuroimage.2014.12.039>
- Herculano-Houzel, S., 2012. The remarkable, yet not extraordinary, human brain as a scaled-up primate brain and its associated cost. *PNAS* 109, 10661–10668.
<https://doi.org/10.1073/pnas.1201895109>
- Hill, J., Inder, T., Neil, J., Dierker, D., Harwell, J., Van Essen, D., 2010. Similar patterns of cortical expansion during human development and evolution. *Proceedings of the National Academy of Sciences* 107, 13135–13140. <https://doi.org/10.1073/pnas.1001229107>
- Karadachka, K., Assem, M., Mitchell, D.J., Duncan, J., Medendorp, W.P., Mars, R.B., 2023. Structural connectivity of the multiple demand network in humans and comparison to the macaque brain. *Cereb Cortex* 33, 10959–10971. <https://doi.org/10.1093/cercor/bhad314>
- Kolling, N., Braunsdorf, M., Vijayakumar, S., Bekkering, H., Toni, I., Mars, R.B., 2021. Constructing others' beliefs from one's own using medial frontal cortex. *J. Neurosci*.
<https://doi.org/10.1523/JNEUROSCI.0011-21.2021>
- Koster-Hale, J., Richardson, H., Velez, N., Asaba, M., Young, L., Saxe, R., 2017. Mentalizing regions represent distributed, continuous, and abstract dimensions of others' beliefs. *Neuroimage* 161, 9–18. <https://doi.org/10.1016/j.neuroimage.2017.08.026>
- Krubitzer, L., Kaas, J., 2005. The evolution of the neocortex in mammals: how is phenotypic diversity generated? *Curr Opin Neurobiol* 15, 444–453.
<https://doi.org/10.1016/j.conb.2005.07.003>
- Li, L., Preuss, T.M., Rilling, J.K., Hopkins, W.D., Glasser, M.F., Kumar, B., Nana, R., Zhang, X., Hu, X., 2010. Chimpanzee (*Pan troglodytes*) Precentral Corticospinal System Asymmetry and Handedness: A Diffusion Magnetic Resonance Imaging Study. *PLOS ONE* 5, e12886. <https://doi.org/10.1371/journal.pone.0012886>
- Mars, R.B., Jbabdi, S., Rushworth, M.F.S., 2021. A Common Space Approach to Comparative Neuroscience. *Annu Rev Neurosci* 44, 69–86. <https://doi.org/10.1146/annurev-neuro-100220-025942>
- Mars, R.B., O'Muircheartaigh, J., Folloni, D., Li, L., Glasser, M.F., Jbabdi, S., Bryant, K.L., 2019. Concurrent analysis of white matter bundles and grey matter networks in the chimpanzee. *Brain Struct Funct* 224, 1021–1033. <https://doi.org/10.1007/s00429-018-1817-8>
- Mars, R.B., Passingham, R.E., Jbabdi, S., 2018a. Connectivity Fingerprints: From Areal Descriptions to Abstract Spaces. *Trends Cogn Sci* 22, 1026–1037.
<https://doi.org/10.1016/j.tics.2018.08.009>
- Mars, R.B., Sallet, J., Neubert, F.-X., Rushworth, M.F.S., 2013. Connectivity profiles reveal the relationship between brain areas for social cognition in human and monkey temporoparietal cortex. *Proc Natl Acad Sci USA* 110, 10806–10811.
<https://doi.org/10.1073/pnas.1302956110>

703 Mars, R.B., Sallet, J., Schüffelgen, U., Jbabdi, S., Toni, I., Rushworth, M.F.S., 2012.
704 Connectivity-based subdivisions of the human right “temporoparietal junction area”:
705 evidence for different areas participating in different cortical networks. *Cereb. Cortex* 22,
706 1894–1903. <https://doi.org/10.1093/cercor/bhr268>

707 Mars, R.B., Sotiropoulos, S.N., Passingham, R.E., Sallet, J., Verhagen, L., Khrapitchev, A.A.,
708 Sibson, N., Jbabdi, S., 2018b. Whole brain comparative anatomy using connectivity
709 blueprints. *eLife* 7, e35237. <https://doi.org/10.7554/eLife.35237>

710 Neubert, F.-X., Mars, R.B., Thomas, A.G., Sallet, J., Rushworth, M.F.S., 2014. Comparison of
711 Human Ventral Frontal Cortex Areas for Cognitive Control and Language with Areas in
712 Monkey Frontal Cortex. *Neuron* 81, 700–713.
713 <https://doi.org/10.1016/j.neuron.2013.11.012>

714 Pandya, D.N., Seltzer, B., 1982. Intrinsic connections and architectonics of posterior parietal
715 cortex in the rhesus monkey. *Journal of Comparative Neurology* 204, 196–210.
716 <https://doi.org/10.1002/cne.902040208>

717 Percheron, G., François, C., Pouget, P., 2015. What makes a frontal area of primate brain the
718 frontal eye field? *Front. Integr. Neurosci.* 9. <https://doi.org/10.3389/fnint.2015.00033>

719 Petrides, M., 2005. Lateral prefrontal cortex: architectonic and functional organization. *Phil.*
720 *Trans. R. Soc. B* 360, 781–795. <https://doi.org/10.1098/rstb.2005.1631>

721 Pitcher, D., Ungerleider, L.G., 2020. Evidence for a Third Visual Pathway Specialized for Social
722 Perception. *Trends Cogn Sci* 0. <https://doi.org/10.1016/j.tics.2020.11.006>

723 Rilling, J.K., Glasser, M.F., Preuss, T.M., Ma, X., Zhao, T., Hu, X., Behrens, T.E.J., 2008. The
724 evolution of the arcuate fasciculus revealed with comparative DTI. *Nat Neurosci* 11, 426–
725 428. <https://doi.org/10.1038/nn2072>

726 Roelofs, A., 2014. A dorsal-pathway account of aphasic language production: the
727 WEAVER++/ARC model. *Cortex* 59, 33–48. <https://doi.org/10.1016/j.cortex.2014.07.001>

728 Roumazeilles, L., Eichert, N., Bryant, K.L., Folloni, D., Sallet, J., Vijayakumar, S., Foxley, S.,
729 Tendler, B.C., Jbabdi, S., Reveley, C., Verhagen, L., Dershowitz, L.B., Guthrie, M., Flach,
730 E., Miller, K.L., Mars, R.B., 2020. Longitudinal connections and the organization of the
731 temporal cortex in macaques, great apes, and humans. *PLoS Biol* 18, e3000810.

732 Roumazeilles, L., Lange, F.J., Benn, R.A., Andersson, J.L.R., Bertelsen, M.F., Manger, P.R.,
733 Flach, E., Khrapitchev, A.A., Bryant, K.L., Sallet, J., Mars, R.B., 2022. Cortical
734 Morphology and White Matter Tractography of Three Phylogenetically Distant Primates:
735 Evidence for a Simian Elaboration. *Cereb Cortex* 32, 1608–1624.
736 <https://doi.org/10.1093/cercor/bhab285>

737 Roumazeilles, L., Schurz, M., Lojkiewicz, M., Verhagen, L., Schüffelgen, U., Marche, K.,
738 Mahmoodi, A., Emberton, A., Simpson, K., Joly, O., Khamassi, M., Rushworth, M.F.S.,
739 Mars, R.B., Sallet, J., 2021. Social prediction modulates activity of macaque superior
740 temporal cortex. *Science Advances* 7, eabh2392. <https://doi.org/10.1126/sciadv.abh2392>

741 Saygin, Z.M., Osher, D.E., Norton, E.S., Youssoufian, D.A., Beach, S.D., Feather, J., Gaab, N.,
742 Gabrieli, J.D.E., Kanwisher, N., 2016. Connectivity precedes function in the development
743 of the visual word form area. *Nat Neurosci* 19, 1250–1255.
744 <https://doi.org/10.1038/nn.4354>

745 Schurz, M., Tholen, M.G., Perner, J., Mars, R.B., Sallet, J., 2017. Specifying the brain anatomy
746 underlying temporo-parietal junction activations for theory of mind: A review using
747 probabilistic atlases from different imaging modalities. *Hum Brain Mapp* 38, 4788–4805.
748 <https://doi.org/10.1002/hbm.23675>

749 Sierpowska, J., Bryant, K.L., Janssen, N., Blazquez Freches, G., Römkens, M., Mangnus, M.,
750 Mars, R.B., Piai, V., 2022. Comparing human and chimpanzee temporal lobe
751 neuroanatomy reveals modifications to human language hubs beyond the frontotemporal
752 arcuate fasciculus. *Proc Natl Acad Sci U S A* 119, e2118295119.
753 <https://doi.org/10.1073/pnas.2118295119>

754 Sotiropoulos, S.N., Jbabdi, S., Xu, J., Andersson, J.L., Moeller, S., Auerbach, E.J., Glasser,
755 M.F., Hernandez, M., Sapiro, G., Jenkinson, M., Feinberg, D.A., Yacoub, E., Lenglet, C.,
756 Van Essen, D.C., Ugurbil, K., Behrens, T.E.J., 2013. Advances in diffusion MRI
757 acquisition and processing in the Human Connectome Project. *NeuroImage* 80, 125–143.
758 <https://doi.org/10.1016/j.neuroimage.2013.05.057>

759 Suddendorf, T., Redshaw, J., Bulley, A., 2022. The invention of tomorrow. A natural history of
760 foresight. Basic Books, New York.

761 Thiebaut de Schotten, M., Dell'Acqua, F., Forkel, S.J., Simmons, A., Vergani, F., Murphy,
762 D.G.M., Catani, M., 2011. A lateralized brain network for visuospatial attention. *Nat*
763 *Neurosci* 14, 1245–1246. <https://doi.org/10.1038/nn.2905>

764 Thiebaut de Schotten, M., Forkel, S.J., 2022. The emergent properties of the connected brain.
765 *Science* 378, 505–510. <https://doi.org/10.1126/science.abq2591>

766 Tomasello, M., Vaish, A., 2013. Origins of Human Cooperation and Morality. *Annu Rev Psychol*
767 64, 231–255. <https://doi.org/10.1146/annurev-psych-113011-143812>

768 Van ESSEN, D.C., 2002. Surface-Based Atlases of Cerebellar Cortex in the Human, Macaque,
769 and Mouse. *Annals of the New York Academy of Sciences* 978, 468–479.
770 <https://doi.org/10.1111/j.1749-6632.2002.tb07588.x>

771 Van Essen, D.C., Smith, S.M., Barch, D.M., Behrens, T.E.J., Yacoub, E., Ugurbil, K., 2013. The
772 WU-Minn Human Connectome Project: An overview. *NeuroImage* 80, 62–79.
773 <https://doi.org/10.1016/j.neuroimage.2013.05.041>

774 Vickery, S., Patil, K.R., Dahnke, R., Hopkins, W.D., Sherwood, C.C., Caspers, S., Eickhoff, S.B.,
775 Hoffstaedter, F., 2024. The uniqueness of human vulnerability to brain aging in great ape
776 evolution. *Sci Adv* 10, eado2733. <https://doi.org/10.1126/sciadv.ado2733>

777 Warrington, S., Bryant, K.L., Khrapitchev, A.A., Sallet, J., Charquero-Ballester, M., Douaud, G.,
778 Jbabdi, S., Mars, R.B., Sotiropoulos, S.N., 2020. XTRACT - Standardised protocols for
779 automated tractography in the human and macaque brain. *NeuroImage* 217, 116923.
780 <https://doi.org/10.1016/j.neuroimage.2020.116923>

781 Warrington, S., Thompson, E., Bastiani, M., Dubois, J., Baxter, L., Slater, R., Jbabdi, S., Mars,
782 R.B., Sotiropoulos, S.N., 2022. Concurrent mapping of brain ontogeny and phylogeny
783 within a common space: Standardized tractography and applications. *Sci Adv* 8,
784 eabq2022. <https://doi.org/10.1126/sciadv.abq2022>

785

786

787 **Figure captions**

788

789 **Figure 1. Mapping connectivity divergence between primates identifies multiple**
790 **hotspots of human specialization.** Divergence maps of the human brain showing
791 vertices with connectivity profiles that have a poor match in the chimpanzee (left) or in
792 either the chimpanzee or the macaque (right). Bar graphs show the normalized
793 connectivity ($\pm$ SEM) of the selected vertex with a tract driving these differences in the
794 human (red) and of its best matching vertices in the chimpanzee (dark blue) and macaque
795 (light blue). Tracts include SLF2 (superior longitudinal fascicle 2), ILF (inferior longitudinal
796 fascicle) and AF (arcuate fascicle). Histograms in the center show the distribution of KL
797 values comparing human and chimpanzee (blue) and human and macaque (red). The
798 complete connectivity profile of each human vertex and its best matches are displayed in
799 the other species are in Extended Data Figs. 1-1; the connectivity profile of anatomical
800 homologs are displayed in Figs. 1-2 to 1-6.


801

802


803 **Figure 2. Decoding areas of high divergence highlights multiple behavioral**
804 **domains.** Functional activations that correlate most with areas of high KL divergence for
805 the human : chimpanzee comparison (A) and the human : macaque comparison (B). Color
806 coding of areas according to the parcellation of Glasser and colleagues (Glasser et al.,
807 2016) is done by assigning each area the divergence value of the most divergent vertex
808 in that area. We note that the procedure of assigning a whole region with a single
809 divergence value accentuates the spatial representation of this value and emphasizes that
810 the actual vertex-wise presentation of Figure 1 presents the most spatially precise
811 representation of the data. Full decoding of the areas is listed in the Extended Data Tables
812 2-1 and 2-2.

813

814



Human : Chimpanzee



Human : Macaque

