

Article

Organic Rice Transition in a Changing Environment: Linking Farmers' Benefits to Adaptation and Mitigation

Jack O'Connor ¹, Joachim H. Spangenberg ^{2,*}, Ngan Ha Nguyen ¹, Gioia Emidi ¹, Arne Kappenberg ³, Linda Klamann ², Nick Kupfer ², Huynh Ky ⁴, Nguyen Thi Thu Nga ⁵, Chau Minh Khoi ⁶, Cao Dinh An Giang ⁶, Jürgen Ott ⁷, Björn Thiele ², Bei Wu ² and Lutz Weihermüller ²

¹ Institute for Environment and Human Security (UNU-EHS), United Nations University, 53113 Bonn, Germany; oconnor@ehs.unu.edu (J.O.); emidi@ehs.unu.edu (G.E.)

² Agrosphere Institute IBG-3, Forschungszentrum Jülich GmbH, 52426 Jülich, Germany; l.klamann@fz-juelich.de (L.K.); n.kupfer@fz-juelich.de (N.K.); b.thiele@fz-juelich.de (B.T.); b.wu@fz-juelich.de (B.W.); l.weihermueller@fz-juelich.de (L.W.)

³ Institute for Soil Sciences, University Bonn, 53113 Bonn, Germany; arne.kappenberg@uni-bonn.de

⁴ Department of Genetics and Plant Breeding, College of Agriculture, Can Tho University, Can Tho 900000, Vietnam; hky@ctu.edu.vn

⁵ Department of Plant Protection, College of Agriculture, Can Tho University, Can Tho 900000, Vietnam; nttnga@ctu.edu.vn

⁶ Department of Soil Science, College of Agriculture, Can Tho University, Can Tho 900000, Vietnam; cmkhoi@ctu.edu.vn (C.M.K.); cdagiang@ctu.edu.vn (C.D.A.G.)

⁷ L.U.P.O GmbH, 67705 Trippstadt, Germany; ott@lupogmbh.de

* Correspondence: joachim.spangenberg@gmail.com or j.spangenberg@fz-juelich.de; Tel.: +49-175-3868982

Abstract

Organic rice farming (ORF) can support both climate change mitigation and adaptation. However, a deeper understanding of its specific benefits and challenges is needed. This paper synthesises current knowledge on the potential of ORF to enhance resilience in regions exposed to natural hazards, with particular attention to the climate-vulnerable region of the Mekong Delta (MKD), Vietnam. ORF can deliver multiple benefits: reducing production costs, revitalising degraded and pesticide-contaminated soils, improving water and soil quality, enhancing biodiversity, and contributing to human health and sustainable livelihoods. In the context of MKD, where rice production intersects with acute vulnerability to salinity intrusion, storms, and drought, ORF also presents opportunities for long-term adaptation by improving ecosystem health and reducing socio-ecological vulnerability. Despite these benefits, ORF remains limited in scale and impact due to the lack of integrated, landscape-level implementation strategies. Challenges like chemical contamination, limited access to certified organic inputs, and insufficient institutional and technical support leave many existing ORF initiatives vulnerable and constrain further expansion. To fully realise ORF's resilience and sustainability potential, more targeted research and policy attention are needed. An integrated governance approach that considers both biophysical and socio-economic dimensions is essential to support a meaningful and scalable transition to organic rice farming in climate-sensitive regions like the MKD.

Keywords: rice organic agriculture; transition; soil quality; water; biodiversity; health; income; institutions



Academic Editor: Cezary Kabala

Received: 4 August 2025

Revised: 4 September 2025

Accepted: 9 September 2025

Published: 17 October 2025

Citation: O'Connor, J.; Spangenberg, J.H.; Nguyen, N.H.; Emidi, G.; Kappenberg, A.; Klamann, L.; Kupfer, N.; Ky, H.; Nga, N.T.T.; Khoi, C.M.; et al. Organic Rice Transition in a Changing Environment: Linking Farmers' Benefits to Adaptation and Mitigation. *Land* **2025**, *14*, 2074. <https://doi.org/10.3390/land14102074>

Copyright: © 2025 by the authors.

Licensee MDPI, Basel, Switzerland.

This article is an open access article

distributed under the terms and

conditions of the Creative Commons

Attribution (CC BY) license

(<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Climate change and biodiversity loss threaten global livelihoods and food production, prompting calls for sustainable methods that stabilize yields, enhance co-benefits, and

build resilience [1–3]. With food demand projected to rise by up to 56% by 2050 [4], climate models predict yield declines of up to 23% under severe scenarios [5]. Consequently, many countries promote transitioning from conventional to organic agriculture to strengthen food system resilience and community well-being [6–8]. Between 2000 and 2020, organic agricultural land expanded over 500%, and organic markets grew nearly eightfold [9]. Organic practices can boost biodiversity and mitigate greenhouse gas emissions, though effects vary by context and methods [10–12]. Understanding the benefits and trade-offs of organic transitions across diverse socio-ecological systems, especially vulnerable deltas, is crucial.

Despite growing adoption, scaling organic agriculture faces challenges including vested interests, weak policies, economic perceptions, social inertia and uneven input and market access [13,14]. These are compounded by limited awareness of organic agriculture's interconnected socio-ecological impacts, at both the policy and community level, particularly in regions like the MKD where large vulnerable populations reside. Conventional agrochemical-based agriculture remains dominant, supported by agribusiness, markets, technology, and policy [15]. However, decades of intensive agriculture in Asia have degraded soil fertility and yields, increasing reliance on synthetic fertilizers and pesticides [16]. While the Green Revolution improved productivity in some areas [17], it also caused negative environmental, social, and health consequences [18,19]. These concerns have driven some governments to promote organic agriculture as a sustainable development strategy to restore food production and foster resilient livelihoods amid climate change [20,21]. Organic farming's socio-economic and environmental resilience, especially in vulnerable rural areas, calls for holistic evaluation of its adaptive capacity.

In the ecologically sensitive MKD, rice cultivation profoundly impacts environment and health. Excessive fertilizer and pesticide use exceed national averages, contributing to nitrogen pollution in waterways and pesticide residues in food [22–32]. Intensive agriculture has also caused biodiversity loss, soil degradation, and significant greenhouse gas emissions, notably methane and nitrous oxide [33–35]. Agriculture, dominated by rice production, accounts for a large share of Vietnam's GHG emissions [36]. Biodiversity and natural habitats have declined due to intensified agricultural and water management [37]. Sustainable rice cultivation in Vietnam, and in the MKD in particular, requires balancing economic objectives with conservation and ensuring socio-ecological resilience to climate change, biodiversity loss, and pollution [22].

This paper has been motivated by discussion between Vietnamese and German scientists from different disciplines such as sociology, economy, soil fertility, soil pollution, crop protection, and hydrology on the topic of organic rice production within the *OrganoRice* project (<https://organorice.org>). It reviews existing evidence of the benefits and challenges of transitioning to organic agriculture and explores their relevance for climate adaptation and sustainability frameworks, drawing on a literature review and comparisons with our findings from the MKD. Discussions and interviews were held with local stakeholders, including administrators and business, between spring 2023 and summer 2025. They showed that there is need to provide a detailed overview of problems and solutions for the transition from conventional high inputs to organic rice production. Additionally, interviews with farmers were held aimed at identifying motivation, experienced/expected gains (monetary and non-monetary) and experienced/perceived obstacles during the transition. This information was used to embed the literature findings in a regional context and to answer our research question “What are necessary conditions, primary benefits and motivations, and what are key obstacles for a large-scale transition to organic rice agriculture in the MKD”. In the final section of this paper, we give some preliminary

answers, discuss transition challenges and propose recommendations to scale organic rice farming and integrate diverse benefits into policy frameworks.

2. Materials and Methods

To understand the state of the art in assessing the diverse benefits of transitioning to organic rice farming (ORF) at the global level, literature was gathered from SCOPUS and Web of Science (WoS) databases. The review focused on both environmental and socio-economic benefits to gain a holistic picture from different fields of research. The following search terms were used for articles written in English and published after 2000: “organic rice” AND “services” OR “benefits” OR “impacts” OR “health” OR “econom*” OR “soci*” OR “ecolog*” OR “environment*”.

The review involved multiple steps. Firstly, the results of the database searches were compared to filter duplicates. Between the papers found from SCOPUS ($n = 413$) and Web of Science (papers found $n = 69$), 46 duplicates were detected, leaving 436 papers for review (Figure 1). Titles were then screened for further review, keeping papers suggesting a context of changes in social or environmental metrics ($n = 245$). In the next step, papers were screened based on their abstract. Papers that mentioned organic rice contexts where specific aspects were assessed related to environmental or socio-economic changes were kept for in-depth review ($n = 147$). In a third step, papers were reviewed in depth. All articles with evidence of changes in ORF areas relating to environmental, ecological, social and socio-economic impacts and benefits were considered relevant ($n = 96$). Further papers were sourced through snowball sampling by reviewing the references of papers selected ($n = 22$). Publications in the Vietnamese context related were searched using similar search terms in both English and Vietnamese using Google Scholar. Grey literature on government standards, regulations and management policies were sourced from online databases and Vietnamese government websites ($n = 23$).




Figure 1. Methodological steps involved in the review.

In addition, the different authors or authors teams specialised in the different subjects and disciplines tackled in this paper, such as soil quality, water quality, biodiversity, rice quality, socio-economics and disaster risk reduction, gathered literature from various

sources. Each individual group was searching for its own literature using different, topic-specific keywords, contributing additional 22 references.

Literature was also compiled using ChatGPT-3.5 and DeepSeek-V-2 AI tools with the search terms “organic rice agriculture”, “organic rice farming”, “transition to organic rice”, “challenges of organic rice” and “ecosystem services in organic rice”, limited to 50 references each. The references suggested and not yet covered by the WoS search were checked for their existence with Google Scholar, resulting in 8 additional references.

Grey literature and technical documents on the study topic were sourced from expert consultations with German and Vietnamese researchers in the OrganoRice project. Vietnamese language publications were sourced with similar terms, using Google Scholar and the University of Can Tho literature database and translated into Vietnamese by project partners. All literature collected covers the time span from 1950 to 2022, when the project began to generate its own data.

They were completed by our own interview experience from five provinces in the MKD, representing a trajectory from the Cambodian border in the West to the sea shore in the East (provinces Dong Thap, An Giang, Vinh Long, Tra Vinh and Ca Mau). We interviewed farmers, cooperative managers, traders, extension workers, local and provincial agricultural and nature protection authority representatives, identified by the Can Tho University team. Due to the rather restrictive legal regulations, the group interviews with farmers (participating and non-participating in organic rice production) were held in the premises of local cooperatives in five provinces (Vinh Long, An Giang, Dong Thap, Tra Vinh, Ca Mau). Individual in depth interviews with officials were held in their respective offices and in Can Tho University. The results were stored in a database and are used for comparative analyses across provinces and over time (to be published).

In parts of the text, Chat GPT has been used for English language editing the manuscript, to improve grammar and wording. Their use was strictly limited to language correction without generating substantive content.

3. Results

3.1. Benefits of Organic Rice Farming

In this section, the different benefits we identified are introduced one by one, aware that they combine and interact. For an overview see Table 1.

3.1.1. Reducing Environmental Impact of Weed and Pest Management

For rice farmers, weeds are a major production barrier, particularly in organic rice farming where managing intense crop–weed competition is challenging [38]. Weeds reduce yields by competing for water, light, and nutrients and by increasing crop vulnerability to pests and diseases [39,40]. Without chemical herbicides, organic systems rely on hand weeding, biological control, optimised habitat and water management, and crop rotation to suppress weeds [41–43]. Though more labour-intensive, these methods can lower costs and minimize environmental impacts compared to conventional agriculture [41].

Unlike conventional rice farms (CRF), pests and diseases are generally less problematic in organic systems due to healthier plants grown in nutrient-balanced soils [44,45]. Organic fields often suffer fewer pest outbreaks [46,47]. Organic fertilizers support growth while reducing pests like brown plant hopper, stem borer, leaf folder, blast and sheath blight [48]. Selecting resistant varieties is a cost-effective strategy but requires knowledge and may affect yields [45,49].

Organic pesticides, often plant-derived and locally sourced, offer crop protection with minimised environmental impacts and can reduce cost, as seen in West Java [50]. While it is known that synthetic insecticides can disrupt natural biocontrol and enable new pest

outbreaks [51,52], the broader ecological impacts of organic treatments remain under study. Research in India's Godavari delta showed that organic mulch (*Sesbania* sp.) effectively managed some pests while maintaining yields [53].

Enhanced biological pest control by increased numbers or diversity of natural predators can be supported by intercropping, e.g., by flower strips [42,54–56]. Predators like spiders and dragonflies are effective antagonists of pest insects [57], which can significantly boost rice yields and hence ecosystem services [58]. Among insects, Odonata have been widely used as bioindicators of insecticide impact, reflecting effects at both lethal and sublethal levels [59–61]. Farmers who are aware of beneficial species, recognising and making use of them can reduce pesticide reliance [62].

Effective pest management must integrate socio-ecological principles—biodiversity, host resistance, landscape ecology, and social dynamics—since human knowledge, perceptions, and decision-making shape outcomes from individual farmers to policymakers [45].

Mekong Delta context

Farmers in the MKD report that ORF approaches can work well to suppress weeds and pests while simultaneously being beneficial for rice crops. In An Giang province, the use of biologically derived (as opposed to chemically derived) organic fertiliser was linked to increased rice plant size and health, while simultaneously fostering the proliferation of natural predators controlling the spread of pests. In Thoai Son district, in particular, the use of organic fertiliser led to thick, large, green leaves less susceptible to damage [63]. In Bac Lieu province, experiments using organic fertiliser resulted in an increased rice plant vigour reducing pest pressures (e.g., rice blast incidence) [64]. In Dong Thap province, the implementation of mechanical weeding and organic fertiliser use has decreased the necessity for chemical treatments while enriching nutrient absorption by rice plants [65]. Reducing chemical inputs including inorganic Nitrogen-Phosphorus-Potassium (NPK) fertiliser can also lower pest risks. This was demonstrated by a study performed in organic fields in Quang Tri province, which experienced minor (2–4%) leaf roller infestation versus 50–70% in conventional fields [66]. Alternatively, some farmers are using a combination of managing water levels, manual weeding and biocontrol organisms such as *Trichoderma* to effectively control weeds and diseases in ORF fields.

In line with the Vietnamese government's interest in organic rice expansion as a way to achieve more sustainable agriculture [67], the promotion of natural pest control is also outlined in Section 5.1.10 of the Vietnam National Standards for organic agriculture (Tiêu chuẩn Việt Nam) TCVN 11041-5:2018, which reflects the government's recognition of the viable benefits of organic approaches. Despite this, there is currently only a limited number of research projects on alternative approaches to herbicides and pesticides supporting farmers transitioning from conventional to organic farming [68].

3.1.2. Promoting Soil Health and Nutrient Management

Long-term use of organic residues on certified organic farms improves key physical, chemical, and biological soil health indicators by enhancing soil organic carbon (SOC), nutrient availability, and microbial activity [44,69]. However, outcomes vary. Amarasekara et al. [70] observed higher soil pH, organic matter, and cation exchange capacity under organic management, but lower available nitrogen and phosphorus. Thus, transitioning to organic rice farming can benefit soil health, though results are context-dependent, particularly on initial soil quality [46]. Organic practices significantly boost SOC compared to chemical fertilizers [47], with five-year trials in India showing SOC increases of 50–58%, along with rises in available N, P, and K stocks of 3–10%, 10–30%, and 8–25%, respectively [71].

Soil organic matter (SOM) is central to soil health, influencing both physicochemical properties and biological processes [72–74]. Intensive farming without organic inputs depletes SOM and nutrients, undermining sustainability [75]. Organic rice systems, through compost, manure, cover crops, and biomass additions, improve nitrogen supply and SOM levels. The higher SOM leads to improved soil structure, water retention, and reduced bulk density [69,76], and enhances fertility and resilience over time. In regions like Lao PDR, where soils are low in fertility, SOM and N-availability, organic farming can be particularly beneficial [77]. Given SOM's susceptibility to rising temperatures, elevated atmospheric CO₂, and changes in soil water content, its increase also supports climate resilience [78].

Organic practices enhance nutrient cycling by relying on natural nutrient sources such as compost, manure etc., and crop rotations. Gradual SOM decomposition releases nutrients gradually, reducing leaching and hence eutrophication of neighbouring streams and lakes, and improving plant uptake. Applying both macro and micronutrients through organic residue addition can provide optimum supply of biologically available nutrients in organic rice systems [38,69,79]. Studies in Bhutan and India reported increased total nitrogen and available phosphorus on all elevation levels [47,75]. Rice quality also benefitted—rice grain analysis showed a significant increase in Fe and Mn content when two or more organic amendments had been applied, and in Zn and Cu content with combined application of three or four of the organic amendments [47]. In Ghana, N uptake improved significantly under organic nutrient amendments [80]. Enhanced enzyme activity linked to SOM boosts nutrient cycling [69], while long-term organic input can increase micronutrient levels stimulating the mineralisation of organic matter, reducing redox potential, and improving the overall soil environment [69,72].

The measurement of soil pH provides valuable insights on various factors that influence plant growth, such as the availability of nutrients, the behaviour of added nutrients, the level of salt, the degree of soil aeration, the composition of soil minerals, and the prevailing meteorological conditions in the region [81]. Unlike chemical fertilizers, compost and manure stabilize or raise soil pH, improving nutrient uptake and creating favourable growth conditions [69,72,73].

Contamination with heavy metals such as As, Cd, and Pb—stemming from industrial waste, mining, pesticides, wastewater irrigation, sewage sludge application and natural processes—poses risks to soil fertility, food safety, and human health [82–86]. Organic farming mitigates metal accumulation by avoiding synthetic inputs and using amendments like compost, zeolites, and manure which enhance metal immobilization, improve soil structure, and promote microbial activity [87,88]. Crop selection and rotation with phytoremediation plants such as Brassica and sunflowers help manage contamination levels [89,90]. Microbial bioremediation with plant growth promoting rhizobacteria and mycorrhizae can further stabilize metals and reduce plant uptake [91].

However, these methods face challenges when applied to organic rice. Rice readily absorbs heavy metals such as Pb, As, and Cd, and crop rotation is rarely practised in rice cultivation, making phytoremediation or microbial bioremediation methods less applicable [90,92]. Silicon amendments offer promise by reducing heavy metal uptake and toxicity through forming silicate complexes, cell wall reinforcement, and altering metal bioavailability in the soil [93,94].

Mekong Delta context

ORF has been linked to benefits for soil health and quality in the MKD as well. The implementation of ORF in Ca Mau province in rice-shrimp agriculture has led to a rise in shrimp yields, underscoring its beneficial effects on both the environment and soil health [95]. In Dong Thap province, the utilisation of mechanical transplanting techniques and organic fertilisers has been found to enhance soil activation and structure [65,96]

while organic farms in An Giang province had higher organic matter, more available nutrients and lower levels of heavy metals compared to traditional farms [97]. The latter is of particular importance in the MKD as geogenic arsenic is found pervasively in deep aquifers, with nearly 900 wells at depths of 200–500 m now contaminated while the usual methods of excavation and disposal of contaminated soil do not apply when contamination is continuously replenished from the ground water. The benefits of implementing ORF and organic fertiliser management on soil health and quality are reflected in Section 5.1.7 and 5.1.9 of Vietnam's National Standard TCVN 11041-5:2018 on Organic agriculture—Part 5: Organic rice. It dictates that organic rice farmers comply with standards on heavy metals, pesticides and pH in soil, as well as taking measures to maintain soil biodiversity and avoid soil degradation.

3.1.3. Water Quality and Management

ORF shows promise as a sustainable approach to reducing water pollution, as it eliminates the use of synthetic pesticides and fertilisers and reduces the risk of nitrate pollution. Additionally, ORF reduces excess of nutrients such as phosphorus and potassium in soils, lowering the risk of ground and surface water pollution, and eutrophication of surface waters [98]. Long-term ORF practices are characterised by favourable water pH, electrical conductivity (EC), nitrate, residual sodium carbonate (RSC), and sodium absorption ratio (SAR) values compared to conventional systems. Organic fields, with lower levels of nitrate, RSC, and SAR, consistently have better water quality in drinking water sources such as wells compared to conventional fields, a reduction in the comparative risk of well water contamination [99]. For example, in Haryana, India, areas with 15 to 20 years of ORF practices implemented, saw a significant benefit to drinking water quality [99]. For irrigation purposes, well water quality parameters indicating salinity such as RSC and SAR follow a similar trend, with higher values in conventional agriculture indicating long-term soil risks due to salinisation [99].

The efficiency of water management is one of the aspects that have to be considered in benefit evaluation for ORF practices. In addition to overseeing the irrigation schedule of the rice field, water management encompasses the practices of evaporation control [50,100,101]. The increase in SOM and its capability to increase the water retention of the soil benefits the water supply to the rice plants, particularly during drought conditions [100,101]. Therefore, increasing SOM in the field is a straightforward approach to saving or retaining water. In order to regulate evaporation, the implementation of mulching techniques might also be employed [102] as the application of mulch to the topsoil layer can effectively mitigate evaporation and enhance water infiltration rates. This approach can, on the other hand, increase some GHG emissions in semi-arid zones and needs particular approaches in order to reduce the global warming potential in those regions. However, in humid regions such as Vietnam soil mulching can minimise GHG emissions by significantly decreasing methane emissions [50].

In general, improving the soil hydrology via organic farming can lead to numerous advantages in water use efficiency, hence reducing agricultural water demand [50]. For example, in Sindangkerta Village, Indonesia, interviewed stakeholders reported that ORF required a lower volume of water and led to a higher ability of soil to retain water compared to conventional methods [50]. Organic wastewater management approaches can also be used for cleaning drainage runoff and, as shown by Prikhodko et al. [103], wastewater successfully cleaned the drainage runoff from mechanical impurities, organic and biogenic elements if halophytes planted in phyto-sections located in the discharge channels of the rice irrigation system were introduced. By doing so, the irrigation rate was increased by

10%, the reclamation state of soils improved, and the cost of rice production were reduced by 7% [103].

Mekong Delta Context

Organic rice farming techniques in different parts of Vietnam have enhanced water quality and improved management, including in the MKD [97]. In Bac Lieu Province, switching to organic rice farming within a rice-shrimp cropping model—by avoiding chemical inputs—has improved water quality, enhanced environmental conditions for post-rice-harvest shrimp cultivation and crop diversification, and increased farmers' income by enabling wet-season rice production with freshwater irrigation and enabling shrimp cultivation during the dry season, when high water salinity prevents rice production in certain areas [64,104]. Similarly, farmers from an An Xuyen Commune in Ca Mau Province reported augmented shrimp yields complementing their organic rice cultivation due to improved water quality management, which resulted in both enhanced economic opportunities and water quality in the region [95]. The abundance of shrimp feed and healthy soil characteristics resulting from ORF comply with national regulations related to sustainable water management (QCVN 08-MT:2015/BTNMT on surface water quality and QCVN 09-MT:2015/BTNMT on groundwater quality).

3.1.4. Human Health

While agrochemicals enhance productivity, their overuse and improper application have been linked to acute and chronic health issues among farmers, workers, and consumers. Nitrates from fertilisers contribute to the formation of potentially carcinogenic nitrosamines [27,28,105]. Farmers are especially vulnerable to acute pesticide poisoning via skin contact, inhalation, or ingestion, while prolonged exposure has been associated with respiratory, dermatological, and neurological disorders. Pesticide residues may persist on harvested crops, entering the food chain and posing further risks, including cancer, hormonal disruption, and developmental issues [28,106,107].

In addition to reducing the health risks associated with rice grown in environments with higher chemical content, ORF has the potential for improving nutritional benefits for rice consumers. Studies assessing rice found that certain cultivars (e.g., IRGA 410) show higher yield, protein and lipid content in conventionally grown rice, while organic rice shows higher total carbohydrates, soluble protein, amylose content, and phenolic acids [108]. Especially phenolic compounds, as natural antioxidants, have gained prominence for their health-promoting effects. According to Bergman and Pandhi [109], conventional farming increases rice grain length, kernel width, and their ratios. Organic farming, on the other hand, lowers quality by increasing the amylose content and decreasing crude protein content. Furthermore, Sihi et al. [110] found that the use of organic farming methods increased the levels of micronutrients (namely, iron, manganese, and zinc) in rice grains, something often overlooked in nutritional comparisons of CRF and ORF [111]. And although organically produced rice is less likely to contain residues of pesticides (e.g., organochlorine) compared to rice grown conventionally, some evidence suggests that organically grown rice can be more likely to be contaminated with mycotoxin-producing fungi and some mycotoxins [109].

Mekong Delta context

In the Mekong Delta as well, the use of NPK fertilisers and pesticides in rice farming poses significant human health risks. Due to the intensification of the agricultural sector in Vietnam over the last decades, ground- and drinking water were found to be polluted with pesticides at high levels [112–114]. Lack of personal protective equipment (PPE) and pesticide-related knowledge have been identified as major drivers of pesticide exposure for local farmers [115]. In several provinces including Tra Vinh and Dong Thap, farmers

reported symptoms like exhaustion, respiratory and digestive problems and skin irritations. Local farming authorities in Dong Thap, however, while well aware of the potential impacts of CRF chemicals, denied that such problems existed in their province

The persistent exposure of the wider population resulted in detectable pesticide residues in human blood, breast milk, and urine, which can yield to various negative health effects [116–119]. Excluding synthetic fertilisers, chemical pesticides, plant growth regulators as needed, to fulfil EU organic standards—or at least strictly limiting pesticide use as needed to fulfil US organic standards—will eliminate a number of well-established sources of health risks, primarily through reduced exposure to chemical pesticides.

Beyond direct exposure, pesticide packaging and waste handling also present environmental and health hazards. Improper disposal—such as leaving containers exposed to heat, on the ground, in the fields or on the banks of the canals—can release airborne toxins that contaminate water sources and food supplies, causing digestive and respiratory illnesses [26,120]. Transitioning to ORF offers a pathway to reduce these risks by eliminating harmful chemical inputs, thereby improving health outcomes for local communities in the MKD, reducing sick leave days and improve productivity.

3.1.5. Biodiversity

Biodiversity plays a vital role in stabilising and enhancing the resilience of ecosystems, including rice-based agroecosystems, against climate change [121–123]. However, conventional agriculture poses global threats to biodiversity through habitat loss and land use changes [124,125]. The persistence of native flora and fauna is strongly influenced by farming practices [126,127]. Arable plants, field birds, and insects—key to functions such as pollination and pest control—depend on agricultural habitats [127]. Paddy fields also function as anthropogenic wetlands, supporting aquatic and semi-aquatic species such as waterfowl, amphibians, and invertebrates [128–130]. Biodiversity patterns in paddies are shaped by proximity to non-paddy habitats, climate, soil conditions, and water availability [131]. Cocultures in rice fields can enhance both biodiversity and agroecosystem functioning [132]. Yet, rice-associated ecosystems face stressors from fungicides, invasive species, and infrastructural changes like concrete bank protection [133]. Warming may further amplify insecticide effects on aquatic insect communities [134].

Unsustainable farming and land conversion contribute to ecosystem degradation, affecting over 3 billion people globally [135]. Agricultural intensification has diminished the ecological value of rice fields [124]. CRF, which relies on synthetic inputs, compromises habitat quality through pollution, erosion, pesticide resistance, and biodiversity loss [136,137]. In contrast, ORF aims to reduce environmental harm and support biodiversity, nutrient cycling, and soil health [138–140]. By minimizing chemical inputs, ORF fosters biodiversity across taxa—including endangered plants, spiders, dragonflies, frogs, and soil microbes [124,141]. Both taxonomic and functional diversity improve under landscape heterogeneity and decline with pesticide use [142]. Waterbird richness increases with the proportion of organic fields, likely due to greater prey availability [124]. Enhanced soil microbial activity, nutrient cycling, and structure are linked to organic inputs and the presence of beneficial organisms like mycorrhizal fungi [47,69,70,79].

A comparative study found 58% of 474 cases showed higher species richness or abundance in ORF, with only 4% indicating negative effects; median plant species richness was 95% higher under organic management [127]. Gains were observed across birds (+35%), insects (+36%), and spiders (+55%). While ORF may slow biodiversity loss, its potential trade-offs—such as possible land expansion due to lower yields—must be considered holistically [124,143]. A summary of benefits by socio-ecological categories is provided in Table 1.

Mekong Delta context

Rice fields in Vietnam, including those in the Mekong Delta, provide habitat for a range of aquatic and terrestrial species [37]. These fields serve as temporary wetlands, providing breeding and feeding grounds for many species of birds, insects, amphibians, and fish [37,144]. However, intensified agricultural practices, characterised by increased use of agrochemicals, drainage changes and monoculture, disrupt ecological functions and can lead to soil degradation, reduced natural pest control services and pest outbreaks, and negative impacts on biodiversity. Agricultural land use changes in the MKD are evident in the shrinking of natural grasslands and melaleuca forests, which are converted for agricultural production. Even in protected areas or reserves, such as Lung Ngoc Hoang Nature Reserve in Hau Giang province, land is still being reclaimed for the purpose of farming, with over 800 households in the area producing rice and sugar cane [145]. Due to the interconnected nature of the landscapes and habitats in the MKD via the maze of waterways, the expansion and intensification of agricultural production has negatively impacted biodiversity both within and outside of protected areas [145,146] as we could confirm in the Dong Thap crane nature reserve. While there is an intuitive expectation that wider transition to ORF will have positive effects on biodiversity, more research is needed in the region [147].

Table 1. Diversity of benefits attributed to ORF from Vietnamese sources.

Benefit Category	Benefits Reported in Vietnamese Context	Sources
<i>Environmental benefits</i>	Improved soil health via organic farming techniques and organic fertilisers	[65,96,97,148]
	Reduced pollution from chemical fertilisers, herbicides, and pesticides	[64,65,97,104,149,150]
	Enhanced biodiversity and encouragement of natural pest predators	[97,148,151]
	Improved crop resistance to diseases & pests	[64,66]
	Preservation of ecological condition	[65,152]
<i>Health benefits</i>	Soil activation condition leading to favourable habitat for beneficial animals	[95,96]
	The production of nutritious organic goods for the consumption of consumers	[65,68,148]
	Reduced health risks for farmers and consumers	[97,152]
	Strong plants with increased resilience to extreme events (e.g., storms)	[63]
<i>Agricultural and farming benefits</i>	Enhanced rice grain yield	[63,64,152]
	Reduction in pest & disease pressure on crops	[63,64,66]
	Enhancement of nutrient uptake efficiency in rice plants	[65]
	Improved livelihoods for farmers	[66]
<i>Social benefits</i>	Provide farmers with support, training and resources to grow organic rice	[148]
	Strengthening cooperatives and partnerships in agricultural production	[148,150]
	Creating sustainable production chains	[63,150,153]
	Increasing farmers' incomes	[65,148,150,152]

3.1.6. Socio-Economic Opportunities and Challenges

Higher consumer prices for organic produce are probably the best-known economic characteristic of ORF, but by far not the only and not necessarily the most important one. Here we distinguish three kinds of direct economic effects: increased income, reduced expenditure and the effects on the farm workload. Indirect monetary co-benefits include

those from improved human and animal health, and the economic losses avoided by ORF. So far, such positive economic side effects are usually not accounted for in cost–benefit analyses of ORF. Avoiding damages pays out, but it does not pay in; hence it is invisible in most economic analyses. In the social dimension, we focus on the impact on reputation and urbanisation.

Increasing income

Increased public awareness of a healthy lifestyle and environmental sustainability has spurred an uptake in organic rice consumption, despite its higher market price, a phenomenon not constrained to high-income countries but shared by the consumer class in many low- and middle-income countries. Grimm et al. [154] tested consumers' willingness to pay for organic rice in urban and suburban Indonesia and found that respondents were willing to pay an average price premium of up to 20% compared to conventional rice. This is a key element of what prompts farmers to transition to ORF, as they hope to increase their business income [155]; our MKD interviews confirmed this. Such local demand is an essential requirement for increased adoption of ORF among smallholders, who typically have only limited access to export markets [154], unless they organize in cooperatives and these strike deals with larger rice trading companies. Although in the 2024 global rice price crisis, the sales of rice decreased for both conventional and organic rice, and the cost crises reduced the overall demand for premium products, in Vietnam traders like LotusRice continued to pay a premium of 40–50%. Nonetheless a significant number of farmers gave up organic practices, arguing that the high rice prices at the time allowed them to maintain their income without the organic-specific workload, although the income difference to conventional or safe rice (2–3% premium) would still have been significant. Apparently, for some farmers, the quality of life was more important than income maximization.

Organic rice production is usually assumed to result in inevitably lower yields and higher costs compared to conventional farming methods. However, there are diverging results regarding yield trends in the literature [44,156,157]. Eyhorn et al. [158] found that in Northern India, organic farmers achieved the same yields of cereals and pulses as conventional farmers, with considerably lower external inputs. Similarly, the average productivity of organic rice production in Nepal was found to be consistently higher than national average [159]. This aligns with earlier results from Java, which found that although the first year's harvest during the transition phase was significantly reduced, organic yields caught up with and surpassed those of conventional farming by the third year [160]. According to the authors, this was partly attributed to the lower pest infestation levels in organic as compared to conventional systems. Moreover, farmers in Vietnam and India using a mixed approach of both organic and inorganic fertilization benefited from applying organic fertilizers, as the amount of inorganic fertilizer required decreased [161,162].

Reducing expenditure

A study carried out in Lao PDR found that the most promising inputs and strategies available to optimize yields in organic rice production systems were identified to be (1) optimizing use of locally available nutrients, mostly from manure, crop residues and weed biomass, (2) N addition through green manure and legumes growing in rotation and (3) additions of P through guano or rock-phosphate [77]. This is similar, though not identical, to the experience of Singh et al. [47], who reported that, in India, different treatments comprising organic amendments such as Blue Green Algae (15 kg/ha), Azolla (1 t/ha), Vermicompost and organic manure (5.0 t/ha), each applied alone or in combination, resulted in a significant enhancement in rice grain yield compared to rice crops that did not receive any fertilization. Moreover, rice grain yields matching the yield were achieved with the recommended dose of inorganic fertilizer application. In a study in Bangladesh,

organic fertilizer users required 42% less farm capital for rice production, despite achieving 17% higher yields [163].

In Vietnam, on the one hand, organic agriculture tends to require less input of disease combating and pest regulating inputs, but on the other hand certified organic substitutes are not easily available. Organic fertilizers and pest regulation substances can be produced from different mixtures of local plants, farm residues and fish waste, but as they are usually not certified, they cannot be used in certified organic fields. What has become known as “safe rice” in the last two years (no management rules but zero pesticide residues in the final product required) can and does use such cheaper, locally produced, uncertified organic inputs alongside reduced volumes of seed (about 1/3 less, like in organic rice, as our interviews in Dong Thap and An Giang confirmed). This makes it competitive with organic produce despite a much lower market price premium. However, also the sourcing of uncertified organic inputs is not always easy—the optimal composition varies with location and resource availability, and traders we interviewed do not routinely offer it as the demand is too small to justify investments into setting up a production and supply network. On-farm production, while economically attractive, requires additional skills and working capacity. Hence, the cost and availability of organic fertilizers, specifically animal manure, could be a bottleneck for large-scale transition to organic [164].

Increased workload

For the household economic balance, it is essential to consider not only market prices of sales and potential savings on inputs, plus salaries for remunerated farm workers, but the workload of (unpaid) family members as well.

Increasing workloads are frequent in organic agriculture. While in mountainous regions the natural slope limits the size of fields, and with it the level of mechanization, in flatland areas, typical for major deltas, fields tend to be large and the level of mechanization high. Direct seeding and fertilizing by drones (from a contractor, or for members from the cooperative) and with high-pressure “seed guns”, and the use of big harvesters go together well with chemical inputs using the same equipment. While reducing insecticide spraying tasks, the transition to organic usually enhanced the demand for hand work in the fields, in some cases for transplanting instead of direct seeding, and in most cases including hand weeding, hand picking of snails (in particular GAS), and regular control of the fields regarding the occurrence of pests or diseases and checking the traps for rodents (rats). However, while heavy work, transplanting has its benefits: it allows for better control of the distance between rice plants, which can reduce the vulnerability to disease outbreaks and pest infestation. It also reduces the damages from Golden Apple Snails as the time between rice seedlings being put in the field and becoming indigestible to GAS due to silicon incorporation of rice plants is shorter. Hence, although organic farmers benefit from reduced input costs, they usually must contend with an increased workload. If not undertaken by unpaid family members or as mutual support on a community level, the salaries of farm workers have to be taken into account. However, in Vietnam they tend to be a minor factor in overall expenditure of a farm, according to An Giang farmers’ estimates: even if all farm work would be carried out by paid workers, the total labour cost would be just 1/3 of the total profit a rice farm generates. is one solution.

Improved health for humans and animals

Besides the direct effects, there are also indirect economic co-benefits. Rice is the staple food of throughout South and East Asia—often in countries without a comprehensive system of free health care and sick leave payments. There ORF economic co-benefits include, for instance, less household spending on medical treatments, a health-induced higher working capability, or decreasing days of (usually uncompensated) sick leave.

Health benefits beyond humans occur when poultry and molluscs, indigestible or even dying in and around conventionally managed fields, as farmers reported in Tra Vinh province, become fit for consumption, or if honeybees survive and produce good honey. If collected and sold on the market, they provide monetary income—if used in the household diet, they provide a non-monetary economic benefit.

Economic losses avoided by ORF

Enhanced resilience against climate change and intensifying disasters such as storms and saltwater intrusions generates economic benefits from avoided losses. Hence, even additional monetary costs may appear justified as a kind of insurance premium, but this largely depends on the prevailing attitudes and the general level of knowledge. Benefits accrue not only to the farm itself, but also to the larger community, including, e.g., ground-water and biodiversity protection, or social stability due to higher resilience. Finally, loss avoidance is the key motivation for going organic when ORF is applied out of necessity to counteract collapsing soil fertility and reduce the significant losses of yield experienced, as our interviewees in An Giang province reported. Then ORF helps avoiding further deterioration and revitalizing the soils to first stabilize and then increase harvests again.

Reputation and Migration

In many countries, and in Vietnam in particular, becoming a farm worker is one of the least appreciated job choices—whoever can, chooses a different career path. The attractiveness of farming as a profession for the younger generation is currently too low to guarantee a sustained production of healthy food and export crops. Farmers' children are leaving for better paid jobs in town or abroad, benefiting from the higher-level education they enjoyed as compared to their parents. Due to the resulting negative selection process, those left over are usually of low qualification, and have problems following the rules of land management and, in particular, the documentation requirements which are part of certified organic agriculture.

Besides the hard work, bad reputation is another reason. Reputation may increase when producing safe and health food, generating more reputation and more respect from society. Many parent farmers are confident that some of their children will return to take over the land after their careers, in particular if the farm is organic, offering a healthy lifestyle and earning social respect from the community.

3.2. Challenges for the Transition to Organic Farming in a Changing Climate

Vietnam is one of the countries worldwide which is most susceptible to damages from the climate crisis [165]. However, storms, floods and rising sea water levels are but one of the sustainability challenges which together require an integrated compound response to safeguard a sustainable future for the MKD. For instance, overusing ground water and uncontrolled sand mining lead to falling ground water levels and accelerates saltwater intrusion beyond the effects of climate change [166]. Besides high chemicals intensity, the dearth of nature protection plans and areas enhances biodiversity loss, with potentially significant impacts on rice production, organic or not. Upstream poldering and dams reduce sediment loads and hence the fertilisation effects of flooding (locally known as 'poor floods'), while high dykes enhance riverbed erosion affecting agriculture and fisheries [167]. Lack of recognition, demographic change and urbanisation threaten to reduce the number of available farm workers. Infrastructure development, diets changing with increasing income and higher revenues from fruit, tree crops, orchards, sugarcane, etc., increase the pressure for converting rice land to other uses.

ORF is an important tool mitigating a wide variety of these challenges, but no silver bullet solving all of them. It can build resilience to various threats and contribute both directly and indirectly to the reduction in vulnerability of local people, communities and

ecosystems to different hazards faced in the MKD. For a future-proof MKD, however, ORF, while being crucial, has to be embedded into a larger framework of sustainability transition strategies.

3.2.1. Salinity Intrusion

Coastal soils and groundwater worldwide are increasingly affected by salinity intrusion, driven by climate change [168]. Rising sea levels are causing salinisation of marshy and previously non-saline rice-growing soils [169] and aquifers. Salinity-tolerant rice varieties are thus critical for food security [169]. In Kerala's Kaippad integrated organic rice–aquaculture system, five traditional varieties—*Chovverian*, *Kuthiru*, *Kuttusan*, *Orkazhama*, *Orthadian*—and two novel varieties, *Ezhome-1* and *Ezhome-2*, demonstrated distinct salinity tolerance and contributed to ecosystem diversity [169–171]. Alongside plant breeding approaches, organic amendments—such as rice hull, straw, and sawdust—have proven effective in mitigating salinity stress, enhancing plant growth, yield, and soil reclamation [172].

Mekong Delta context

In the MKD, salinity intrusion, driven by climate change (altered precipitation, sea level rise, storm surges) and human activities (groundwater over-extraction, altered river flows), threatens water supplies and ecosystems [166]. Future projections indicate worsening conditions, making not only mitigation, but also adaptation urgent [173]. Evidence suggests ORF can improve groundwater quality and reduce salinity vulnerability compared to conventional systems [99], while mulching with organic materials helps retain soil moisture and lower surface salt concentrations [174]. Farmers in An Giang province report that organic fertilisers strengthen root systems and enhance resilience to weather extremes [63]. Consequently, ORF offers a viable climate adaptation strategy by increasing salinity resilience [175]. Socio-economic vulnerability may also decline as organic rice commands higher market prices and facilitates alternative livelihoods, such as shrimp farming, in increasingly saline areas [95].

3.2.2. Flooding

Changes in flooding frequency and intensity are increasingly linked to climate-driven shifts in weather patterns, including heavier rainfall and rising sea levels [176]. Vietnam is particularly vulnerable due to its extensive low-lying delta regions [165]. To mitigate flooding impacts, farmers and policymakers should focus on enhancing soil health, increasing water retention, and reducing surface runoff during intense precipitation [176]. ORF practices such as year-round soil cover and diverse crop rotations improve soil structure and humus, contributing to flood risk reduction [176].

Mekong Delta context

The Mekong River Delta (MKD) experiences seasonal flooding that plays a vital role in delivering sediment and nutrients, supporting local agriculture and aquaculture [177]. However, extreme floods—such as the 2000 event causing 539 deaths and \$210 million in damages [178]—can cause severe soil erosion and loss of agricultural inputs, disrupting farming cycles [179]. Climate-smart crops, selected for resilience to drought, heat, or flooding, are therefore increasingly valuable [179]. Moreover, flood-tolerant rice cultivars offer adaptive solutions; in particular, organic varieties 'Ezhome-1' and 'Ezhome-4' demonstrate strong tolerance to prolonged flooding and tidal inundation in both saline and non-saline wetlands [170,180]. These varieties not only maintain high yields but also possess desirable grain quality traits, benefiting both farmers and consumers [170,180].

3.2.3. Drought

Organic rice farming enhances soil and water resilience, crucial for mitigating drought [69,72]. Practices such as organic manure application improve soil fertility and nutrient uptake, supporting robust crop growth under water stress [73,181]. Higher SOM in ORF reduces soil density, thereby increasing water retention [182–184]. In contrast, chemical inputs tend to increase bulk density, decreasing water-holding capacity (WHC) and pore space [182]. Organic nutrition management, including manure and crop residues, enhances WHC and soil stability by promoting polysaccharide production, which fosters stable soil aggregates and improves moisture retention [72]. Conservation tillage minimizes soil disturbance, preserving structure and enhancing infiltration and root growth. For instance, in the Philippines, crop residue recycling (3–4 t/ha) and animal manure application (1–2 t/ha) increased SOM, resulting in looser soils and deeper mud [185]. Similarly, soils under organic management generally show better granular structure and lower bulk density than conventionally managed soils [186].

Mekong Delta context

In the MKD, extreme droughts in 2016 and 2020 affected millions of hectares and caused economic losses of about \$500 to \$600 million USD [97,187]. Drought exacerbates salinity intrusion, creating multi-hazard challenges that reduce yields and threaten groundwater [188]. ORF improves soil porosity, permeability, and water retention, building drought resilience while ground cover—such as green manure, hedges, and flower strips—protects soil from erosion.

In the MKD, organic farms exhibited higher nutrient absorption and less susceptibility to erosion compared to traditional farms [97]. These soil retention services contribute to reducing drought and erosion impacts, increasing resilience to salinity intrusion, and decreasing sedimentation downstream, benefiting reservoirs and coastal zones [189]. Widespread adoption of ORF could also mitigate pollution and eutrophication in surface and coastal waters, though effective solutions require landscape-level management beyond individual farms [189], but solutions have to be found on the landscape level, not by individual farms.

3.2.4. Reducing GHG Emissions

In 2022, global greenhouse gas (GHG) emissions reached a record 53.8 Gt CO_{2eq}, with agriculture—including crop and livestock production—accounting for approximately 12% [190]. Rice cultivation is a major GHG emitter per unit area, largely due to fertiliser and water management practices that create conditions favouring anaerobic organic matter decomposition and excess nitrogen release, particularly methane [191]. Although emissions in rice production are partly inevitable, they can be mitigated. Conservation-based approaches such as sustainable intensification reduce soil disturbance and modify crop rotations, thereby lowering emissions [192]. Bacenetti et al. [193] found that in Italian organic systems, climate impacts were primarily driven by methane emissions from flooded fields (41%) and compost production (49%). Chen et al. [194] demonstrated through modelling and life-cycle assessments that integrated soil–crop system management can substantially reduce reactive nitrogen losses and GHG emissions. Organic fertilisers can produce up to five times lower GHG emissions than conventional fertilisers, especially when methane emissions from their production are minimised [195].

Mekong Delta context

In Vietnam, agriculture accounts for 25–30% of total GHG emissions, half of which originate from rice production [36]. Rice cultivation contributes 75% of agricultural methane and 46% of nitrous oxide emissions [68]. Local authorities continue to prioritise economic growth and food security over emission reductions. Despite a recent central government initiative, mechanisms to encourage low-carbon rice production remain insufficient, and

production costs under low-carbon agriculture are high [36]. New initiatives, such as the World Bank-funded “1 million hectare” project, promote reduced seeding and fertiliser use in the Mekong Delta to lower emissions and potentially enable carbon credit payments for farmers—a disputed option.

Organic agricultural practices can enhance climate resilience, particularly in vulnerable regions, by delivering multidimensional sustainability benefits (Table 2). Research with MKD communities and decision-makers shows that Sustainable Development Goal (SDG) targets on sustainable agriculture have strong positive linkages with other key sustainability goals [196]. Despite ORF’s potential to support climate resilience, further empirical research is needed on its effectiveness in reducing risks from floods, droughts, and salinity intrusion in Vietnam. Given the MKD’s vulnerability, assessing organic farming’s role within broader sustainable development strategies is urgent to improve resilience and support climate adaptation in this hotspot.

Table 2. Benefits of ORF linked to climate resilience via increase (+) or reduction (−) of socio-ecological factors compared to CRE.

	Benefit	Link to Climate Resilience	Challenge	Recommendation
	Chemical contamination (−)	Social and environmental vulnerability (−)		
Weed & pest management	Number of species (+)	Human health (+)	Farmer scepticism on effectiveness and resulting yield	Horizontal communication strategies, company compensation for yield deficits
	Cost-effectiveness (+)	Biodiversity support (+)		
	SOM/SOC (+)	Resilience to hazards (flood, drought, salinity intrusion, erosion) (+)		
Soil Health	Heavy metals (−)	Human health (+)	Managing cross-contamination	Integrated irrigation strategies at the farm and inter-provincial level
	Nutrient availability (+)	Biodiversity support (+)		
	Chemical contamination (−)			
	Chemical contamination (−)	Resilience to hazards (flood, drought, salinity intrusion) (+)		
Water Quality	Water-use efficiency (+)	Human health (+)	Managing chemical flows and extreme flood events	Identify point sources, change pollution management approaches
	Water retention (+)	Biodiversity support (+)		
Economic gain	Input costs (−)	Socio-economic vulnerability (−)	Certification, market prices and labour costs	Enhance market connections, develop local production of farm inputs
	Profit potential (+)			

Despite the wide-ranging potential benefits ORF can provide for climate change adaptation the increasing integration of organic agriculture targets in policy, the speed of transition has been slowed down by a number of challenges which must be comprehensively addressed. One of the reasons appears to be that the most effective incentive for transition is via market prices, where a higher market price for organic rice can be a strong (but not always sufficient) motivator for transition, while a nominally different price can lead to a reluctance for transition [184,197]. In many cases the social and environmental benefits are given lower priority in farmers’ decision-making [198]. Key challenges emerging from the literature and our observations are summarised below.

3.2.5. Managing Water and Soil Contamination

Over recent decades, concerns have grown globally about the impacts of agricultural activities on water and soil pollution due to cascading effects on environmental and human health. In Bangladesh, for example, farmers and consumers recognize the risks of chemical inputs but remain largely unaware of organic rice farming [199]. In rice paddy systems, water quality issues arise from both contaminated irrigation inflows and nutrient-laden outflows [200]. While the environmental impacts of CRF like water use inefficiencies, flooding, and nutrient leaching, and aerobic methane generation are well-documented [201–203], research on ORF—particularly during the transition phase—remains limited [149,204,205].

Furthermore, while pesticide pollution in CRF is widely acknowledged, its role in hindering organic transition via contaminant residues in irrigation and floodwaters remains underexplored. Agrochemicals from adjacent CRF areas can contaminate transitioning farms through shared irrigation networks. Floodwaters during the rainy season also carry pollutants from distant farms, villages, and upstream cultivation areas [201]. Despite this, the influence of regional crop landscapes on agrochemical transfer is poorly studied, partly due to limited high-resolution pesticide and crop data [206].

Land use has been shown to significantly influence water quality [207]. In the U.S. and Thailand, studies have linked pesticide use and human exposure to specific crop types, including rice, via land use classifications [208,209], indicating the need for landscape-scale solutions.

Mekong Delta context

In the MKD, rice—especially in triple-cropped areas—is the primary pesticide-exposed crop. Authorities also identify aquaculture and livestock as major pollution sources due to the discharge of pesticides, antibiotics, and hormones [32,203]. Dyke systems are additional hotspots for agrochemical accumulation [210,211].

Pesticide use is typically higher on large, intensively managed farms and is linked to reduced crop diversity, decreased natural pest control, the removal of non-crop habitat, though this varies by crop type [206,212]. Past use can play a role when accumulated pesticides contaminate the breeding or foraging areas of biocontrol species. Transitions in the MKD have at times failed due to inadequate monitoring of residual agrochemicals and the incidental import of pollutants via irrigation before certification [149]. However, no study has thoroughly examined the link between specific crops in the MKD (e.g., orchards, tree crops, sugarcane) and pesticide-related water pollution. This gap, likely due to data limitations and a focus on spatial intensity over crop type, complicates the identification of CRF areas suitable for conversion. The risk is significant: certification for organic production may be denied if chemical residues are detected in soil or crops.

3.2.6. Certification Processes, Costs and Markets

Farmers must undergo a minimum two-year transition period before obtaining organic certification, during which they are required to follow organic practices but cannot market their products as organic. This period poses economic risks, as yields may be lower while soil benefits remain limited, and prices for “pre-organic” rice remain similar to those of conventional rice [76]. Certification involves following stringent rules covering conversion, seed use, fertilisation, pest and disease management, crop rotation, labelling, and post-harvest handling [213]. The process is often costly and time-consuming, particularly for smallholders, due to high fees and labour-intensive documentation, and has to be regularly repeated to maintain the validity of the certificate [198,214]. In Nepal, additional barriers include limited organic markets, high input costs, lack of training, and insufficient government support [214]. In Iran, inspection costs and the absence of branding efforts

hinder organic adoption [215]. These challenges underscore the need for support from partners willing to share investment risks and promote branding and marketing [216].

Perceptions of higher labour requirements also deter adoption, though impacts vary by task. In the Philippines, family labour was essential during the transition, especially for composting, hand-weeding, and pest control, while reduced pesticide use and easier land preparation offset some burdens [187]. Similarly, in Bangladesh, organic fertiliser use required 12% less labour despite achieving 17% higher yields [163]. Nonetheless, labour demands and the limitations of smallholder capacity remain significant barriers to scaling organic rice production [217].

3.2.7. Knowledge Management—Farmer Education and Training

Organic farming is only economically viable when production factors are efficiently managed; efficiency—defined as the optimal use of inputs for maximum output at minimal cost—directly influences yields and income. Beyond physical inputs, knowledge and skills are essential. As Pawitri et al. [155] observed in Indonesia, failure to recognize their importance limits productivity. However, awareness alone is insufficient. Devi et al. [198] found that while Indonesian farmers understood organic principles, they often failed to meet the procedural standards set by the Indonesian National Standard (SNI) and international bodies like Organics International (IFOAM) the United States Department of Agriculture (USDA). This reflects a broader knowledge–action gap, where individuals do not always act on what they know to be beneficial [218]. Lal et al. [219] similarly reported that only half of the perceived barriers to organic transition in India were linked to knowledge, with other challenges including complexity of application, poor-quality inputs, and limited guidance.

Training has been identified as a key factor in organic adoption, particularly within sustainable agricultural networks. In Northern Italy, a participatory project showed that enhancing farmer skills led to improved yields and reduced variability, emphasizing that successful organic management depends on applying agroecological principles flexibly in response to local conditions [220]. For risk-averse, rice-dependent smallholders, lack of knowledge can be a significant barrier. However, strong social networks and information exchange—such as in Vietnam and Taiwan—can foster trust and investment in organic practices through peer influence and “neighbourhood effects” [161,221,222]. Nonetheless, informal exchanges must be supported by formal interventions, including training, education, and comparative studies in cultivation and marketing [223].

3.2.8. Values, Institutional and Policy Frameworks

Organic rice is primarily valued by farmers for its income, environmental and health benefits. However, although they clearly and frequently say they care about their health, they are often willing to trade it in for higher profits. This aligns well with the attitude of officials on all levels who agree that whatever transition is suggested, farm incomes must not decline (some farmers would prefer to switch from 3 to 2 seasons). Agricultural authorities appear focussed on yield levels, downplaying soil fertility loss and health impacts of conventional farming. Organic is then considered a risky experiment, with a number of farmers and officials “cherry picking”, adopting some immediately economically beneficial elements from the organic rulebook, but not all, and keeping in case of non-increasing yield conventional farming as the fall-back option. This is a major barrier to adopting any sustainable techniques in the Mekong Delta.

Demographic change, migration and urbanisation reduce the rural work force, resulting in a manifest shortage of agricultural labour. At the same time, organic farming usually requires more work in the fields—an emerging bottleneck, already felt in some

provinces. The use of high tech in rice production, such as drones, is intended to reduce labour demand. However, certain stages, such as re-transplanting or weeding, still require a large amount of manual work.

While already in June 2020, the Project on Organic Agriculture Development for the period 2020–2030 was approved, affirming Vietnam's determination to develop organic agriculture products certified according to organic standards locally and globally, the current picture is more mixed. There is support for safe/clean/ecological rice with reduced chemical inputs and no detectable pesticide contamination by the time of harvest, although the price premium is clearly below 5%. In an initiative supported by the World Bank, 1 million ha have been earmarked for low emission rice, however with so far unclear definitions and supports structures, and not necessarily organic. The rapidly changing and apparently uncoordinated prioritisation by authorities makes even medium-term planning difficult. Local implementation suffers from limited direct communication with active farmers, instead through hierarchies (village heads). Doing so, however, would require more extension staff at the local level.

4. Discussion and Recommendations

Our international literature compilation has highlighted the significant benefits of ORF compared to conventional methods, especially in times of climate change and biodiversity loss, but also some of the challenges. Vietnam, and, in particular, the MKD, is particularly susceptible to damages from the climate crisis [165]. Stabilising the harvests in this vital yet climate-vulnerable rice-producing region is of social, economic and political relevance, as the MKD is not only the “bread basket” of Vietnam, but also the source of significant export earnings.

Evidence, also from Vietnam, shows that ORF has the potential to deliver economic gains, improved soil and water quality, enhanced biodiversity, and better health outcomes for farmers and communities. These benefits also bolster climate resilience by improving soil properties and mitigating impacts from floods, drought, and salinity intrusion. Nonetheless, the expansion of organic rice agriculture is slow, and the share in total rice area still marginal. Hence, challenges such as the ones described in the previous section must be addressed to unlock ORF's full potential.

4.1. Values, Institutional and Policy Frameworks

While organic transition often requires areas with lower residual chemical pollution, in Vietnam it is usually recommended for regions with good soil qualities. However, regions of poorer soil quality can also benefit from ORF, maybe even more, as international examples illustrate. Successful organic conversions frequently occur in less-favoured agricultural areas with smaller fields and limited mechanisation, where yields have increased and stabilized [224]. Farmers in low-potential areas face lower opportunity costs and may be more willing to adopt organic methods [222]. Government institutes in several countries therefore recommend focusing organic agriculture on poor soil areas while seeking alternative strategies to reduce environmental impacts for high-input agriculture in fertile areas [111]. Declining soil quality under conventional farming may further motivate adoption of organic practices.

Given ORF's potential to enhance resilience to climate hazards like salinity, storms, and drought, mitigate biodiversity loss, and improve the social status of farmers as well as their income, expansion plans should incorporate broad assessments and climate modelling to prioritize benefits. A landscape-level, integrated approach is essential, combining farm-level organic practices that reduce chemical loads and promote biodiversity with inter-provincial sustainable water management. This approach should leverage existing

irrigation infrastructure to minimize chemical runoff, where possible restore sediment flow, and improve groundwater recharge.

To overcome the perception of transition to ORF as risky, an insurance policy could help, eliminating the transition period risk and building a bridge to the higher income level from certified organic farming. It should be backed by scientific analysis and government enforcement as so far, the major obstacle to such a solution is the mutual mistrust, if only transition-induced losses are claimed, or if the deficit is really fully compensated. On the policy level, the indirect economic benefits (see Section 3.1.6) should be valued and more clearly communicated, also between departments.

There is no easy solution to address the labour shortage challenge, as despite growing farm incomes, the discrepancy to non-farming work tends to be increasing. Using high tech like drones is an important step, but merging fields and upscaling machinery has its limits in rice paddy soil sensitivity.

4.2. Managing Water and Soil Contamination

Given the Mekong Delta's complex hydrology and intensive agriculture, landscape-level planning and transformation are essential to avoid cross-contamination of water and soil. Additional data on the sources and fate of chemical contaminants across the MKD are critical for managing chemical residues in irrigation and floodwaters during organic transition and ongoing organic production. Existing irrigation infrastructure may inadequately control chemical flows, especially for farmers who cannot afford or must share facilities with conventional producers. Therefore, diverse irrigation strategies tailored to maintaining organic standards should be explored to effectively manage chemical contamination. The trend to crop diversification, with co-production of shrimps and fish, or planting fruit, may contribute to reduce the overall pesticide contamination of soils and water.

4.3. Context-Specific Policy

It is essential to recognize that (1) ORF is not necessarily beneficial in comparison to CRF in all contexts and for all criteria, and (2) the drivers that reduce its ability to deliver benefits are multiple. Prescribing market-driven farming notions in different cultural and ecological settings can lead to contradictions, and certification requirements, resource constraints, and labour demands can exclude some farmers [225]. Therefore, the analysis of organic farming as a rural development strategy should consider not only economic returns but also the broader socio-political context and the influence of development agencies on poverty reduction potential [225]. This also applies to the suggestions made in this section: none of them is a 'silver bullet', none fits to all circumstances, but one each has to be adapted to the local natural, economic and social conditions.

Ideology and practices play a significant role in this context, and recommendations should be sensitive to local conditions [226]. The importance of environmental benefits for health and well-being of delta-dwellers should also not be understated and accounted for in policy. While this often includes non-monetary benefits such as increases in biodiversity, the indirect economic benefits through saved expenditure, lower health cost and increased opportunity for income diversification and crop resilience should be considered in long-term policy approaches. Clearly delineating when and where organic, low emission and other forms of less intensive rice production are encouraged, and—if so—there is a sequence of land management forms foreseen, would be a kind of expectation management helping farmers and local decision-makers to adapt their plans and investments.

4.4. Knowledge Management–Farmer Education, Training and Support

As income appears to be the most effective incentive for transition to ORF, part of any information dissemination should be that not only the market prices determine the profit, but also the reduced cost due to less expenditure for chemical inputs to avoid that a nominally too small price difference leads to a reluctance for transition [184,197]. Unfortunately, currently in Vietnam no domestically produced, internationally certified fertilisers are available, reducing the economic benefits due to the dependence on expensive imported inputs for ORF. This may be one of the reasons why more farmers adopted safe rice farming and similar options, which provide some of the ORF indirect benefits without the cost for imported certified fertilisers and pesticides (roughly speaking, both fertilisers and pesticides, and labour each account for a fifth of CRF farm expenditure).

A second ORF specific cost factor is the organic certification and its regular refreshment, in particular for export markets, i.e., following US, EU and Japanese standards. So far, in cooperatives certified as organic, the supervision and documentation are mainly undertaken by the cooperative, but training for non (or not yet) member farmers is missing: extension workers could contribute, but that would require that they themselves receive hands-on training, and that more staff is available on the local level. Traders buying the organic produce have a self-interest in organic quality, and could also offer training (some already do). However, the challenge of shortage in qualified staff limits these options.

Regarding the certification cost, which are a major obstacle in particular ahead of a transition, when an intermediate reduction in yield is to be expected (although it not always materialises), different options exist. A government-based solution would be state-backed transition insurance, while a market based one would be if traders planning to export the organic produce after certification could offer medium term contracts under which they cover the certification cost and receive a repayment from the higher prices per ton of rice afterwards. However, in both cases the lack of trust between agents, and limited trust in enforcement through the legal system, are limiting factors. A lack of clear organic-specific policies (distinguished from ‘safe’, ‘clean’ and ‘ecological’ rice, for instance) contributes to mixed and not necessarily coherent approaches to ORF in the MKD. Limited mandates and staff shortages in local extension workers’ offices shift the role of monitoring and evaluating protocols to companies. In particular, the system of training and information for farmers through agricultural extension workers can vary in effectiveness depending on how they themselves are trained and how far they are entitled to engage with individual farmers.

Without a standardised oversight of both ORF and CRF, there is scope for approaches that could compromise the ability to control chemical flows and realise diverse benefits of ORF. Direct exchange between farmers from different villages and provinces is a promising way to facilitate learning and problem solving for ORF, and should be seen as a way to promote success stories in organic farming while addressing reasons for scepticism in farmers who have little insight to the reality of organic farming. Administrators at the district and higher levels could provide platforms for farmers to share experience amongst each other, and to raise concerns and needs with government officials.

4.5. Limitations of the Work Presented

The MKD faces compounded threats from extreme weather, sea-level rise, and altered hydrology, biodiversity loss, demography and urbanisation. This complexity makes it virtually impossible to derive “one size fits all” solutions, as the combination of challenges varies even on relatively small scales. Scale is also an issue in data integration across disciplines, with soil science and plant chemical analysis generating results on the field level, while biodiversity analyses happen on landscape level, and socio-economic ones on province, district and commune level (farmers could be from different hamlets).

Comparison to international experiences helps only to a limited degree. While overall impacts of organic farming such as increasing soil carbon content and moderation of biodiversity loss are documented across the board, the different case studies are incomparable regarding soil and water conditions, and, in particular, the organic inputs which are usually generated from regionally available resources and differ from case to case. Hence the studies quoted in this paper should be considered not as blueprints for transition purposes, but as inspirations for locally adapted measures.

Conducting expert interviews as a survey method has its limitations. The long duration of the interviews and the fact that they were conducted as individual or small group interviews made it possible to gain very in-depth insights into the perceptions of the interviewees. Nevertheless, there is also a susceptibility to bias and a natural limitation of the horizon of knowledge due to the knowledge and perception limits of the respondents, and thus also a high degree of personal dependence. In particular, in interviews in which the respondents acted as representatives of an administrative unit (DARDs, extension officers, agricultural and nature protection administration) or a company (cooperatives, traders), significant bias due to positive self-reference is to be expected. In general, the respondents were willing to criticise their institutions and other actors to a very limited extent. It must therefore be assumed that, despite the comprehensive survey, individual inhibiting factors may have remained undetected.

Furthermore, the survey should by no means be considered exhaustive, as not all relevant actors could be interviewed. On the one hand, several specialist services declined the request for interviews; on the other hand, local and provincial politicians were explicitly excluded from the survey, as this would have led to an even greater normative bias.

It would have been particularly exciting and valuable in terms of a transdisciplinary research process to develop and discuss the results iteratively with the interviewees in several survey rounds in the process of transitioning to or establishing a permanent organic rice regime. It would also have allowed to implement the research results, including recommendations for action, in a joint praxis process. Unfortunately, the repeated changes in framework conditions (market prices, incentives and government regulations and recommendation, plus agents leaving the transition arena and others entering) made such a stepwise and continuous approach impossible – the research had to adapt to the changing circumstances in the framework conditions and agents' preferences.

Author Contributions: Conceptualization, J.O. (Jack O'Connor), J.H.S., and L.W.; methodology, J.O. (Jack O'Connor), J.H.S., and L.W.; validation, J.O. (Jack O'Connor), J.H.S., and L.W.; investigation, J.O. (Jack O'Connor), J.H.S., N.H.N., G.E., A.K., L.K., N.K., H.K., N.T.T.N., C.M.K., C.D.A.G., J.O. (Jürgen Ott), B.T., B.W., and L.W.; writing—original draft preparation, J.O. (Jack O'Connor), and J.H.S.; writing—review and editing, J.H.S.; supervision, L.W.; project administration, L.W., and C.M.K.; funding acquisition, L.W., C.M.K., A.K., J.O. (Jürgen Ott), and J.H.S. All authors have read and agreed to the published version of the manuscript.

Funding: The bilateral OrganoRice project, from which this paper emerged, is funded by the German Federal Ministry of Research, Technology and Space (BMFTR) within the “CLIENT II—International Partnerships for Sustainable Innovations” funding initiative, funding code 01LZ1806A and by the Vietnamese Ministry of Science and Technology (MOST), funding code NDT/DE/22/29.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by the Ethics Committee of the Juelich Research Centre.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The original contributions presented in this study are included in the article. Further inquiries can be directed to the corresponding author. An anonymised publication

of the interview results mentioned is in preparation; the original data are not accessible due to data protection regulations and ethics committee standards.

Acknowledgments: We are grateful to Nguyen Thi Thuy Ngan and Nguyen Thi Quynh Anh who supported the interviews in the early phase of the project. Bùi Chúc Ly was an essential contributor to data gathering and interpretation in the last two years of the project.

Conflicts of Interest: Authors Joachim H. Spangenberg, Linda Klamann, Nick Kupfer, Björn Thiele, Bei Wu and Lutz Weihermüller were employed by the Agrosphere Institute IBG-3, Forschungszentrum Jülich GmbH. Author Jürgen Ott was employed by the company L.U.P.O. GmbH. The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Abbreviations

The following abbreviations are used in this manuscript:

MKD	Mekong Delta
ORF	Organic rice farming
CRF	Conventional rice farming
GHG emissions	Greenhouse gas emissions
SOM	Soil organic matter
SOC	Soil organic carbon
RSC	Residual sodium carbonate
Pb, As, Cd, Zn, Cu	Lead, arsenic, cadmium, zinc, copper
NPK	Nitrogen, phosphorus, potassium

References

1. Borron, S. *Building Resilience for an Unpredictable Future: How Organic Agriculture Can Help Farmers Adapt to Climate Change*; Food and Agriculture Organization of the United Nations: Rome, Italy, 2011. Available online: <https://www.fao.org/3/ah617e/ah617e.pdf> (accessed on 24 November 2024).
2. Gonzalez, C.G. Climate change, food security, and agrobiodiversity: Toward a just, resilient, and sustainable food system. *Fordham Environ. Law Rev.* **2011**, *22*, 493–522.
3. Smith, P.; Calvin, K.; Nkem, J.; Campbell, D.; Cherubini, F.; Grassi, G.; Korotkov, V.; Le Hoang, A.; Lwasa, S.; McElwee, P.; et al. Which practices co-deliver food security, climate change mitigation and adaptation, and combat land degradation and desertification? *Glob. Change Biol.* **2020**, *26*, 1532–1575. [\[CrossRef\]](#) [\[PubMed\]](#)
4. van Dijk, M.; Morley, T.; Rau, M.L.; Saghai, Y. A meta-analysis of projected global food demand and population at risk of hunger for the period 2010–2050. *Nat. Food* **2021**, *2*, 494–501. [\[CrossRef\]](#) [\[PubMed\]](#)
5. Rezaei, E.E.; Webber, H.; Asseng, S.; Boote, K.; Durand, J.L.; Ewert, F.; Martre, P.; MacCarthy, D.S. Climate change impacts on crop yields. *Nat. Rev. Earth Environ.* **2023**, *4*, 831–846. [\[CrossRef\]](#)
6. Criveanu, R.C.; Sperdea, N.M. Organic agriculture, climate change, and food security. *Econ. Manag. Financ. Mark.* **2014**, *9*, 118–123.
7. Muller, A.; Olesen, J.; Smith, L.; Davis, J.; Dyrtrtová, K.; Gattinger, A.; Lampkin, N.; Niggli, U. *Reducing Global Warming and Adapting to Climate Change: The Potential of Organic Agriculture. Working Papers in Economics: No. 526*; University of Gothenburg, Department of Economics: Gothenburg, Sweden, 2012.
8. Ziervogel, G.; Ericksen, P.J. Adapting to climate change to sustain food security. *Wiley Interdiscip. Rev.* **2010**, *1*, 525–540. [\[CrossRef\]](#)
9. Willer, H.; Trávníček, J.; Schlatter, S. *The World of Organic Agriculture. Statistics and Emerging Trends 2024*; Research Institute of Organic Agriculture FiBL & IFOAM—Organics International: Bonn, Germany, 2024.
10. Scialabba, N.E.-H.; Müller-Lindenlauf, M. Organic agriculture and climate change. *Renew. Agric. Food Syst.* **2010**, *25*, 158–169. [\[CrossRef\]](#)
11. Goh, K.M. Greater Mitigation of Climate Change by Organic than Conventional Agriculture: A Review. *Biol. Agric. Hortic.* **2011**, *27*, 205–229. [\[CrossRef\]](#)
12. Patle, G.T.; Badyopadhyay, K.K.; Kumar, M. An overview of organic agriculture: A potential strategy for climate change mitigation. *J. Appl. Nat. Sci.* **2014**, *6*, 872–879. [\[CrossRef\]](#)
13. Ferdous, Z.; Zulfiqar, F.; Datta, A.; Hasan, A.K.; Sarker, A. Potential and challenges of organic agriculture in Bangladesh: A review. *J. Crop Improv.* **2021**, *35*, 403–426. [\[CrossRef\]](#)

14. Reganold, J.; Wachter, J. Organic agriculture in the twenty-first century. *Nat. Plants* **2016**, *2*, 15221. [CrossRef]
15. Kerdnoi, T.; Prabudhanitisarn, S.; Sangawongse, S.; Prapamontol, T.; Santasup, C. The struggle of organic rice in Thailand: A multi-level perspective of barriers and opportunities for up scaling. *Environ. Nat. Resour. J.* **2014**, *12*, 95–115.
16. Charyulu, D.K. Technical, allocative and economic efficiency of organic input units in India. *Indian J. Agric. Econ.* **2010**, *65*, 722–738.
17. Evenson, R.E.; Gollin, D. Assessing the impact of the Green Revolution, 1960 to 2000. *Science* **2003**, *300*, 758–762. [CrossRef] [PubMed]
18. Singh, R.B. Environmental consequences of agricultural development: A case study from the Green Revolution state of Haryana, India. *Agric. Ecosyst. Environ.* **2000**, *82*, 97–103. [CrossRef]
19. John, D.A.; Babu, G.R. Lessons from the aftermaths of green revolution on food system and health. *Front. Sustain. Food Syst.* **2021**, *5*, 644559. [CrossRef]
20. Noppol, A.; Sereenonchai, S.; Wang, C. Carbon footprint and predicting the impact of climate change on carbon sequestration ecosystem services of organic rice farming and conventional rice. *J. Environ. Manag.* **2021**, *289*, 112458. [CrossRef]
21. Ditzler, L.; Breland, T.A.; Francis, C.; Chakraborty, M.; Singh, D.K.; Srivastava, A.; Eyhorn, F.; Groot, J.C.J.; Six, J.; Decock, C. Identifying viable nutrient management interventions at the farm level: The case of smallholder organic Basmati rice production in Uttarakhand, India. *Agric. Syst.* **2018**, *161*, 61–71. [CrossRef]
22. Le, T. Human Behavior to the Ecological Environment in the Mekong Delta. *Communist Review*, 2008. Available online: <https://tapchiconsan.org.vn/tin-tieu-diem-10-06/-/2018/3734/ung-xu-cua-con-nguoi-voi-moi-truong-sinh-thai-o-dong-bang-song-cuu-long.aspx> (accessed on 1 April 2025). (In Vietnamese)
23. Galli, A.; Winkler, M.S.; Doanthu, T.; Fuhrmann, S.; Huynh, T.; Rahn, E.; Stamm, C.; Staudacher, P.; Van Huynh, T.; Loss, G. Assessment of pesticide safety knowledge and practices in Vietnam: A cross-sectional study of smallholder farmers in the Mekong Delta. *J. Occup. Environ. Hyg.* **2022**, *19*, 509–523. [CrossRef] [PubMed]
24. Phung, D.; Connell, D.; Miller, G.; Rutherford, S.; Chu, C. Pesticide regulations and farm worker safety: The need to improve pesticide regulations in Viet Nam. *Bull. World Health Organ.* **2012**, *90*, 468–473. [CrossRef]
25. Hou, X.; Han, M.; Dai, X.; Yang, X.; Yi, S. A multi-residue method for the determination of 124 pesticides in rice by modified QuEChERS extraction and gas chromatography—Tandem mass spectrometry. *Food Chem.* **2013**, *138*, 1198–1205. [CrossRef]
26. Truong, A. The Risk of Environmental Pollution from Field Waste. *Quang Nam Online*. Available online: <https://baoquangnam.vn/moi-truong/nguy-co-o-nhiem-moi-truong-tu-rac-thai-dong-ruong-140758.html> (accessed on 3 November 2023). (In Vietnamese)
27. Dhankhar, N.; Kumar, J. Impact of increasing pesticides and fertilisers on human health: A review. *Mater. Today Proc.* **2023**, *3*, 766.
28. Le, D.H. *Pesticide Abuse: A Threat to Health and the Environment*; Tap Chi Viet Nam Huong Sac. Available online: <https://tapchivietnamhuongsac.vn/lam-dung-thuoc-bao-ve-thuc-vat-moi-de-doa-den-suc-khoe-va-moi-truong-1800.html> (accessed on 26 April 2025). (In Vietnamese)
29. Nguyen, H.L.; Ngo, Q.D.; Nguyen, V.C.; Ngo, K.D.; Lam, V.N.; Dang, T.N.; Tran, Q.H.; Phung, T.D.; Nguyen, K.T.; Nguyen, T.V.; et al. Organophosphate Pesticide Exposure: Effect on Farmers’ Sperm Quality in the Mekong Delta, Vietnam. *J. Agromedicine* **2024**, *29*, 404–414. [CrossRef]
30. Do, H. *To Ensure that Agricultural Activities Do not Affect the Ecosystem and Biodiversity*; Government News; 27 August 2021. Available online: <https://baochinhphu.vn/de-hoat-dong-nong-nghiep-khong-tac-dong-toi-he-sinh-thai-da-dang-sinh-hoc-102299273.htm> (accessed on 17 November 2024). (In Vietnamese)
31. Nguyen, T.H. *An Overview of Agricultural Pollution in Vietnam: The Crops Sector*; World Bank: Washington, DC, USA, 2017.
32. Sebesvari, Z.; Le, H.T.T.; Van Toan, P.; Arnold, U.; Renaud, F. Agriculture and Water Quality in the Vietnamese Mekong Delta. In *The Mekong Delta System*; Renaud, F., Kuenzer, C., Eds.; Springer Environmental Science and Engineering: Dordrecht, The Netherlands, 2012; pp. 331–361.
33. Minh, G. *Abuse of Chemical Fertilisers and Pesticides in Agricultural Production*; Dien Bien Radio and Television Station: 2012. Available online: <https://dienbientv.vn/tin-tuc-su-kien/kinh-te/201704/dien-bien-lam-dung-phan-bon-hoa-hoc-va-thuoc-bao-ve-thuc-vat-trong-san-xuat-nong-nghiep-5529127> (accessed on 16 November 2024). (In Vietnamese)
34. Doan, T.T. *It is Necessary to Limit Environmental Pollution from Agricultural Production*. Tap San Khuyen Nong; 3 February 2023. Available online: <https://khuyennongnghean.com.vn/index.php/trong-trot/can-han-che-gay-o-nhiem-moi-truong-tu-san-xuat-nong-nghiep-1388.html> (accessed on 16 November 2024). (In Vietnamese)
35. Wassmann, R. Fertiliser use and GHG Emissions in Agriculture/Paddy Field. *Clim. Change Crop Prod.* **2010**, 419–443.
36. Canh, K. *25–30% of Total Emissions in Vietnam from Agricultural Production*. Tien Phong. Available online: <https://tienphong.vn/25-30-tong-luong-khi-thai-o-viet-nam-tu-san-xuat-nong-nghiep-post1456582.tpo> (accessed on 16 November 2024). (In Vietnamese)
37. Nguyen, D.T.; Pham, T.T.; Tang, T.K.H. *The State of Biodiversity in the Mekong Delta, Vietnam*; Technical Report 30; CIFOR: Bogor, Indonesia; ICRAF: Nairobi, Kenya, 2024. Available online: https://www.cifor-icraf.org/publications/pdf_files/WPapers/CIFOR-ICRAF-WP-30.pdf (accessed on 8 September 2025). (In Vietnamese)

38. Hazra, K.K.; Swain, D.K.; Bohra, A.; Singh, S.S.; Kumar, N.; Nath, C.P. Organic rice: Potential production strategies, challenges and prospects. *Org. Agric.* **2018**, *8*, 39–56. [[CrossRef](#)]
39. Dass, A.; Shekhawat, K.; Choudhary, A.K.; Sepat, S.; Rathore, S.S.; Mahajan, G.; Chauhan, B.S. Weed management in rice using crop competition—A review. *Crop Prot.* **2017**, *95*, 45–52. [[CrossRef](#)]
40. Kaur, S.; Singh, S. Impact of crop establishment methods and weed control on weeds, insect-pest and disease infestation in rice in north-western indo-gangetic plains. *IJ Agric. Sci.* **2015**, *7*, 487–491.
41. Singh, A.; Pazhanisamy, S.; Devi, R.C.; Singh, A.K.; Mehta, C.M. Weed Management Strategies in Organic Rice Production System—A Review. *IJ Environ. Clim. Change* **2020**, *10*, 519–528. [[CrossRef](#)]
42. Muneret, L.; Mitchell, M.; Seufert, V.; Aviron, S.; Djoudi, E.A.; Pétilion, J.; Plantegenest, M.; Thiéry, D.; Rusch, A. Evidence that organic farming promotes pest control. *Nat. Sustain.* **2018**, *1*, 361–368. [[CrossRef](#)]
43. Scherner, A.; Schreiber, F.; Andres, A.; Concenço, G.; Martins, M.B.; Pitol, A. Rice crop rotation: A solution for weed management. In *Rice Crop-Current Developments*; Shah, F., Khan, Z.H., Iqbal, A., Eds.; InTechOpen: London, UK, 2018; pp. 83–98.
44. Ram, M.; Davari, M.; Sharma, S. Organic farming of rice (*Oryza sativa* L.)—Wheat (*Triticum aestivum* L.) cropping system: A review. *IJ Agron. Plant Prod.* **2011**, *2*, 114–134.
45. Savary, S.; Horgan, F.; Willocquet, L.; Heong, K.L. A review of principles for sustainable pest management in rice. *Crop Prot.* **2012**, *32*, 54–63. [[CrossRef](#)]
46. Sukristiyonubowo, R.; Wiwik, H.; Sofyan, A.; Benito, H.; Neve, S.D. Change from conventional to organic rice farming system: Biophysical and socioeconomic reasons. *Int. Res. J. Agric. Sci. Soil Sci.* **2011**, *1*, 172–182.
47. Singh, Y.V.; Singh, B.V.; Pabbi, S.; Singh, P.K. Impact of Organic Farming on Yield and Quality of BASMATI Rice and Soil Properties. In *Zwischen Tradition und Globalisierung. 9. Wissenschaftstagung Ökologischer Landbau*; Zikeli, S., Claupein, W., Dabbert, S., Kaufmann, B., Müller, T., Valle Zárate, A., Eds.; Verlag Dr. Köster: Berlin, Germany, 2007; p. 10742.
48. Chau, L.M.; Heong, K.L. Effects of organic fertilisers on insect pest and diseases of rice. *Omonrice* **2005**, *13*, 26–33.
49. Fujita, D.; Kohli, A.; Horgan, F.G. Rice Resistance to Planthoppers and Leafhoppers. *Crit. Rev. Plant Sci.* **2013**, *32*, 162–191. [[CrossRef](#)]
50. Johannes, H.P.; Priadi, C.R.; Herdiansyah, H. Organic rice farming: An alternative to sustainable agriculture. *IOP Conf. Ser.* **2019**, *546*, 022008. [[CrossRef](#)]
51. Heong, K.L.; Wong, L.; Delos Reyes, J.H. Addressing Planthopper Threats to Asian Rice Farming and Food Security: Fixing Insecticide Misuse. In *Rice Planthoppers*; Heong, K., Cheng, J., Escalada, M., Eds.; Springer: Dordrecht, The Netherlands, 2015; pp. 65–76.
52. Spangenberg, J.H.; Douguet, J.-M.; Settele, J.; Heong, K.L. Escaping the lock-in of continuous insecticide spraying in rice: Developing an integrated ecological and socio-political DPSIR analysis. *Ecol. Model.* **2015**, *295*, 188–195. [[CrossRef](#)]
53. Srinivas, M.; Anusha, B.; Reddy, C.V.; Satyanarayana, P.; Munirathnam, P. Effect of Weed Management Practices on Productivity and Economics of Rice-Rice Crop sequence under Organic Production System. *Int. J. Bio-Resour. Stress Manag.* **2019**, *10*, 124–127. [[CrossRef](#)]
54. Horgan, F.G.; Vu, Q.; Mundaca, E.A.; Crisol-Martínez, E. Restoration of Rice Ecosystem Services: Ecological Engineering for Pest Management; Incentives and Practices in the Mekong Delta Region of Vietnam. *Agronomy* **2022**, *12*, 1042. [[CrossRef](#)]
55. Tschumi, M.; Albrecht, M.; Entling, M.H.; Jacot, K. High effectiveness of tailored flower strips in reducing pests and crop plant damage. *Proc. Biol. Sci.* **2015**, *282*, 20151369. [[CrossRef](#)]
56. Westphal, C.; Vidal, S.; Horgan, F.G.; Gurr, G.M.; Escalada, M.; Van Chien, H.; Tscharnkte, T.; Heong, K.L.; Settele, J. Promoting multiple ecosystem services with flower strips and participatory approaches in rice production landscapes. *Basic Appl. Ecol.* **2015**, *16*, 681–689. [[CrossRef](#)]
57. Saksongmuang, V.; Michalko, R.; Petcharad, B.; Bumrungsri, S. Changes in community composition and prey capture of web-building spiders during rice field development. *Basic Appl. Ecol.* **2024**, *79*, 29–37. [[CrossRef](#)]
58. De Andrade França, J.; Latini, A.O.; Stein, K.; Barbosa, M.A.; Araújo, G.S.S.; Pimenta Pereira, A.C. Enhancing rice yield in paddy fields through beneficial organisms. *J. Nat. Conserv.* **2024**, *77*, 126544. [[CrossRef](#)]
59. Ohtsu, K. Study of acute toxicity test method to Fipronil and Imidacloprid for the larvae of Japanese common meadowhawk (*Sympetrum frequens*). *Jpn. J. Environ. Toxicol.* **2019**, *22*, 31–40.
60. Hayasaka, D.; Korenaga, T.; Suzuki, K.; Saito, F.; Sánchez-Bayo, F.; Goka, K. Cumulative ecological impacts of two successive annual treatments of imidacloprid and fipronil on aquatic communities of paddy mesocosms. *Ecotoxicol. Environ. Saf.* **2012**, *80*, 355–362. [[CrossRef](#)]
61. Jingui, H.; Ohtsu, K.; Ueda, T.; Goka, K. Effects of short-term, sublethal fipronil and its metabolite on dragonfly feeding activity. *PLoS ONE* **2018**, *13*, e0200299. [[CrossRef](#)]
62. Horgan, F.G.; Mundaca, E.A.; Hadi, B.A.; Crisol-Martínez, E. Diversified rice farms with vegetable plots and flower strips are associated with fewer pesticide applications in the Philippines. *Insects* **2023**, *14*, 778. [[CrossRef](#)] [[PubMed](#)]

63. Le, B.Y. *WEHG Biological Products Accompany Organic Rice*. Vietnam Farmer's Union: Agriculture and Environment, Hanoi. 2021. Available online: <https://nongnghiep.vn/che-pham-sinh-hoc-wehg-dong-hanh-cung-lua-huu-co-d302519.html> (accessed on 16 November 2024). (In Vietnamese)
64. Tuan, K. *Bac Lieu Expands Agricultural Production in an Organic Direction*; Dan Toc & Mien Nui, Hanoi; 17 May 2022. Available online: <https://dantocmiennui.vn/bac-lieu-nhan-rong-san-xuat-nong-nghiep-theo-huong-huu-co/320154.html> (accessed on 17 November 2024). (In Vietnamese)
65. Nguyen, C.D. *Initial Effectiveness of Some Rice Production Models in an Organic Direction*; Thanh Tuu Khoa Hoc—Cong Nghe: 2021. Available online: <https://vinhlong.gov.vn/Portals/1038/Khoa%20h%E1%BB%8Dc%20-%20C%C3%B4ng%20ng%E1%BB%87/50.pdf?ver=FF7NtYSurOBIXWVeTqm2w%3D%3D> (accessed on 17 November 2024). (In Vietnamese)
66. Lan, A.; Tran, T. *Quang Tri-Double Profits Through Organic Rice Growing Linked To Product Consumption*; Agriculture Extension Center, Quang Tri Province: 7 June 2023. Available online: <https://khuyennong.quangtri.gov.vn/mo-hinh-kinh-nghiem/quang-tri-loi-nhuan-gap-doi-nho-trong-lua-huu-co-lien-ket-tieu-thu-san-pham-703.html> (accessed on 16 November 2024). (In Vietnamese)
67. Ngan, N.T.; Ngoc, T.H. Development of organic agriculture in Vietnam: Some theoretical and practical issues. *Am. Res. J. Humanit. Soc. Sci.* **2022**, *5*, 36–42.
68. Nguyen, V.K. Perception of Challenges in Opportunities for Organic Food Research and Development in Vietnam. In *Regulatory Issues in Organic Food Safety in the Asia Pacific*; Goh, B., Price, R., Eds.; Springer: Singapore, 2020; pp. 199–216.
69. Sihi, D.; Dari, B.; Sharma, D.K.; Pathak, H.; Nain, L.; Sharma, O.P. Evaluation of soil health in organic vs. conventional farming of basmati rice in North India. *J. Plant Nutr. Soil Sci.* **2017**, *180*, 389–406.
70. Amarasekara, M.G.T.S.; Waduge, K.; Bandara, A.M.K.R. Characteristics of Rice Growing Soils under Conventional and Organic Input Systems in Sri Lanka. *Int. J. Adv. Sci. Res. Manag.* **2019**, *5*, 10–18.
71. Surekha, K.; Satishkumar, Y.S. Productivity, Nutrient Balance, Soil Quality, and Sustainability of Rice (*Oryza sativa* L.) under Organic and Conventional Production Systems. *Commun. Soil Sci. Plant Anal.* **2014**, *45*, 415–428. [CrossRef]
72. Babu, S.; Singh, R.; Avasthe, R.K.; Yadav, G.S.; Das, A.; Singh, V.K.; Mohapatra, K.P.; Rathore, S.S.; Chandra, P.; Kumar, A. Impact of land configuration and organic nutrient management on productivity, quality and soil properties under baby corn in Eastern Himalayas. *Sci. Rep.* **2020**, *10*, 16129. [CrossRef]
73. Kumawat, A.; Kumar, D.; Shivay, Y.S.; Bhatia, A.; Rashmi, I.; Yadav, D.; Kumar, A. Long-term impact of biofertilization on soil health and nutritional quality of organic basmati rice in a typic ustchrept soil of India. *Front. Environ. Sci.* **2023**, *1*, 1031844. [CrossRef]
74. Wang, H.; Xu, J.; Liu, X.; Di, Z.; Li, L.; Li, W.; Sheng, L. Effects of long-term application of organic fertiliser on improving organic matter content and retarding acidity in red soil from China. *Soil Tillage Res.* **2019**, *195*, 104382. [CrossRef]
75. Singh, Y.; Singh, B.; Ladha, J.K.; Khind, C.S.; Gupta, R.K.; Meelu, O.P.; Pasuquin, E. Long-Term Effects of Organic Inputs on Yield and Soil Fertility in the Rice—Wheat Rotation. *Soil Sci. Soc. Am. J.* **2004**, *68*, 845–853. [CrossRef]
76. Tashi, S.; Wangchuk, K. Organic vs. conventional rice production: Comparative assessment under farmers' condition in Bhutan. *Org. Agric.* **2016**, *6*, 255–265. [CrossRef]
77. Roder, W.; Schürmann, S.; Chittanavanh, P.; Sipaseuth, K.; Fernandez, M. Soil fertility management for organic rice production in the Lao PDR. *Renew. Agric. Food Syst.* **2006**, *21*, 253–260. [CrossRef]
78. Dalal, R.C.; Allen, D.E.; Chan, K.Y.; Singh, B.P. Soil organic matter, soil health and climate change. In *Soil Health and Climate Change—Soil Biology*; Singh, B., Cowie, A., Chan, K., Eds.; Springer: Berlin/Heidelberg, Germany, 2011; Volume 29, pp. 87–106.
79. Surekha, K.; Latha, P.C.; Rao, K.V.; Kumar, R.M. Grain Yield, Yield Components, Soil Fertility, and Biological Activity under Organic and Conventional Rice Production Systems. *Commun. Soil Sci. Plant Anal.* **2010**, *41*, 2279–2292. [CrossRef]
80. Ofori, J.; Kamidouzono, A.; Masunaga, T.; Wakatsuki, T. Organic Amendment and Soil Type Effects on Dry Matter Accumulation, Grain Yield, and Nitrogen Use Efficiency of Rice. *J. Plant Nutr.* **2005**, *28*, 1311–1322. [CrossRef]
81. Cedeño, K. Soil Quality of a Rice Organic Farm in Langkong, M'lang, Cotabato. *Southeast Philipp. J. Res. Dev.* **2019**, *24*, 35–53. [CrossRef]
82. Min, B. Identifying an influential spreader from a single seed in complex networks via a message-passing approach. *Eur. Phys. J. B* **2018**, *91*, 18. [CrossRef]
83. Ghori, N.-H.; Ghori, T.; Hayat, M.Q.; Imadi, S.R.; Gul, A.; Altay, V.; Ozturk, M. Heavy metal stress and responses in plants. *Int. J. Environ. Sci. Technol.* **2019**, *16*, 1807–1828. [CrossRef]
84. Zakaria, Z.; Zulkafflee, N.S.; Mohd Redzuan, N.A.; Selamat, J.; Ismail, M.R.; Praveena, S.M.; Tóth, G.; Abdull Razis, A.F. Understanding Potential Heavy Metal Contamination, Absorption, Translocation and Accumulation in Rice and Human Health Risks. *Plants* **2021**, *10*, 1070. [CrossRef] [PubMed]
85. Duan, G.; Zhang, H.; Liu, Y.; Jia, Y.; Hu, Y.; Cheng, W. Long-term fertilization with pig-biogas residues results in heavy metal accumulation in paddy field and rice grains in Jiaxing of China. *Soil Sci. Plant Nutr.* **2012**, *58*, 637–646.
86. Angon, P.B.; Islam, M.; Shafiul, K.; Shreejana, D.; Arpan, A.; Nafisa, P.; Amrit, S.; Shaharia, A. Sources, effects and present perspectives of heavy metals contamination: Soil, plants and human food chain. *Heliyon* **2024**, *10*, e28357. [CrossRef]

87. Azhar, U.; Ahmad, H.; Shafqat, H.; Babar, M.; Shahzad, M.; Hafiz, M.; Sagir, M. Remediation techniques for elimination of heavy metal pollutants from soil: A review. *Environ. Res.* **2022**, *214 Pt 4*, 113918. [CrossRef]
88. Nazia, T.; Ullah, A.; Tahir, A.; Rashid, H.U.; Rehman, T.U.; Danish, S. Strategies for reducing Cd concentration in paddy soil for rice safety. *J. Clean. Prod.* **2022**, *316*, 128116.
89. Cao, C.; Yang, Y.; Kwan, M.-P.; Ma, Z.-B.; Karthikeyan, R.; Wang, J.-J.; Chen, H. Crop selection reduces potential heavy metal(loid)s health risk in wastewater contaminated agricultural soils. *Sci. Total Environ.* **2022**, *819*, 152502. [CrossRef]
90. Puppe, D.; Kaczorek, D.; Stein, M.; Schaller, J. Silicon in Plants: Alleviation of Metal(loid) Toxicity and Consequential Perspectives for Phytoremediation. *Plants* **2023**, *12*, 2407. [CrossRef]
91. Hnini, M.; Rabeh, K.; Oubohssaine, M. Interactions between beneficial soil microorganisms (PGPR and AMF) and host plants for environmental restoration: A systematic review. *Plant Stress* **2024**, *11*, 100391. [CrossRef]
92. Ngo, H.T.T.; Hang, N.T.T.; Nguyen, X.C.; Nguyen, N.T.M.; Truong, H.B.; Liu, C. Toxic metals in rice among Asian countries: A review of occurrence and potential human health risks. *Food Chem.* **2024**, *460 Pt 1*, 140479. [CrossRef]
93. Cai, Y.; Zhang, S.; Cai, K.; Huang, F.; Pan, B.; Wang, W. Cd accumulation, biomass and yield of rice are varied with silicon application at different growth phases under high concentration cadmium-contaminated soil. *Chemosphere* **2020**, *242*, 125128. [CrossRef]
94. Yang, X.; Ni, Y.; Li, Z.; Yue, K.; Wang, J.; Li, Z. Silicon in paddy fields: Benefits for rice production and the potential of rice phytoliths for biogeochemical carbon sequestration. *Sci. Total Environ.* **2024**, *929*, 172497. [CrossRef]
95. Ca Mau People Committee. *Dual Benefits from the Organic Rice Model*. Doanh Nghiep Hoi Nhap 27 April 2021. Ca Mau province, Vietnam. Available online: <https://doanhnghiephoinhap.vn/ca-mau-loi-ich-kep-tu-mo-hinh-lua-huu-co-55904.html> (accessed on 17 November 2024). (In Vietnamese)
96. Huu, N. *Dong Thap Aims to Produce Organic Rice*; Nhan Dan Online, Official newspaper of the Communist Party of Vietnam, Hanoi; 26 October 2022. Available online: <https://nhandan.vn/dong-thap-huong-toi-san-xuat-lua-huu-co-post721282.html> (accessed on 18 November 2024). (In Vietnamese)
97. Tran, D.D.; Park, E.; Van, C.T.; Nguyen, T.D.; Nguyen, A.H.; Linh, T.C.; Quyen, P.H.; Tran, D.A.; Nguyen, H.Q. Advancing sustainable rice production in the Vietnamese Mekong Delta insights from ecological farming systems in An Giang Province. *Heliyon* **2024**, *10*, e37142. [CrossRef] [PubMed]
98. Lisenbee, W.; Saha, A.; Mohammadpour, P.; Cibin, R.; Kaye, J.; Grady, C.; Chaubey, I. Water quality impacts of recycling nutrients using organic fertilizers in circular agricultural scenarios. *Agric. Syst.* **2024**, *219*, 104041. [CrossRef]
99. Sihi, D.; Dar, B.; Yan, Z.; Sharma, D.K.; Pathak, H.; Sharma, O.P.; Nain, L. Assessment of Water Quality in Indo-Gangetic Plain of South-Eastern Asia under Organic vs. Conventional Rice Farming. *Water* **2020**, *12*, 960. [CrossRef]
100. Gheewala, S.H.; Silalertruksa, T.; Nilsalab, P.; Mungkung, R.; Perret, S.R.; Chaiyawannakarn, N. Water footprint and impact of water consumption for food, feed, fuel crops production in Thailand. *Water* **2014**, *6*, 1698–1718. [CrossRef]
101. Gattinger, A.; Muller, A.; Haeni, M.; Skinner, C.; Fliessbach, A.; Buchmann, N.; Niggli, U. Enhanced top soil carbon stocks under organic farming. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, 18226–18231. [CrossRef]
102. Junge, S.M.; Storch, J.; Finckh, M.R.; Schmidt, J.H. Developing Organic Minimum Tillage Farming Systems for Central and Northern European Conditions. In *No-Till Farming Systems for Sustainable Agriculture*; Dang, Y., Dalal, R., Menzies, N., Eds.; Springer: Cham, Switzerland, 2020; pp. 173–192.
103. Prikhodko, I.A.; Vladimirov, S.A.; Alexandrov, D.A. Improving the elements of organic farming in rice cultivation. *IOP Conf. Ser.* **2021**, *659*, 012062. [CrossRef]
104. Le, T.; Chau, P. *Growing Organic Rice Combined with Raising Clean Shrimp in Bac Lieu, Farmers are Healthy and Have Much Better Income*; Danviet. Vn 21 March 2023. Available online: <https://danviet.vn/trong-lua-huu-co-o-bac-lieu-kieu-gi-ma-chua-gat-da-co-nguoi-doi-mua-lai-con-nuoi-tom-su-to-bu-20230321124620828.htm> (accessed on 16 November 2024). (In Vietnamese)
105. UNEP United Nations Environment Programme. *Synthesis Report on the Environmental and Health Impacts of Pesticides and Fertilisers and Ways to Minimize Them*; UNEP: Geneva, Switzerland, 2022.
106. EEA European Environment Agency. *How Pesticides Impact Human Health and Ecosystems in Europe*; Office for the Publications of the European Union: Luxembourg, 2023.
107. Shah, R. Pesticides and Human Health. In *Emerging Contaminants*; Nuro, A., Ed.; IntechOpen: London, UK, 2020; pp. 57–78.
108. Alves, G.H.; Paraginski, R.T.; Lamas, N.D.; Hoffmann, J.F.; Vanier, N.L.; de Oliveira, M. Effects of Organic and Conventional Cropping Systems on Technological Properties and Phenolic Compounds of Freshly Harvested and Stored Rice. *J. Food Sci.* **2017**, *82*, 2276–2285. [CrossRef]
109. Bergman, C.; Pandhi, M. Organic rice production practices: Effects on grain end-use quality, healthfulness, and safety. *Foods* **2022**, *12*, 73. [CrossRef]
110. Sihi, O.; Sharma, O.; Pathak, H.; Singh, V.; Sharma, C.; Chaudhary, A.; Oari, B. Effect of organic farming on productivity and quality of basmati rice ORYZA-An. *Int. J. Rice* **2012**, *49*, 24–29.

111. Vanavichit, A. *The Next Green Revolution of Organic Rice*. Open Access Government. 6 January 2023. Available online: <https://www.openaccessgovernment.org/article/green-revolution-organic-rice-yield-environment/149446/> (accessed on 17 November 2024).
112. Lamers, M.; Anyusheva, M.; La, N.; Nguyen, V.V.; and Streck, T. Pesticide pollution in surface- and groundwater by paddy rice cultivation: A case study from Northern Vietnam. *Clean Soil Air Water* **2011**, *39*, 356–361. [\[CrossRef\]](#)
113. Toan, P.V.; Sebesvari, Z.; Blasing, M.; Rosendahl, I.; Renaud, F.G. Pesticide management and their residues in sediments and surface and drinking water in the Mekong Delta, Vietnam. *Sci. Total Environ.* **2013**, *452/453*, 28–39. [\[CrossRef\]](#)
114. Chau, N.D.G.; Sebesvari, Z.; Amelung, W.; Renaud, F.G. Pesticide pollution of multiple drinking water sources in the Mekong Delta, Vietnam: Evidence from two provinces. *Environ. Sci. Pollut. Res. Int.* **2015**, *22*, 9042–9058. [\[CrossRef\]](#)
115. Phung, D.T.; Connell, D.; Miller, G.; Rutherford, S.; Chu, C. Needs Assessment for Reducing Pesticide Risk: A Case Study With Farmers in Vietnam. *J. Agromed.* **2013**, *18*, 293–303. [\[CrossRef\]](#)
116. Dasgupta, S.; Meisner, C.; Wheeler, D.; Xuyen, K.; Lam, N.T. Pesticide poisoning of farm workers—Implications of blood test results from Vietnam. *Int. J. Hyg. Environ. Health* **2007**, *210*, 121–132. [\[CrossRef\]](#)
117. Nguyen, M.T.; Sudaryanto, A.; Tu, B.M.; Bui, H.N.; Isobe, T.; Takahashi, S.; Pham, H.V.; Tanabe, S. Kinetic differences of legacy organochlorine pesticides and polychlorinated biphenyls In Vietnamese human breast milk. *Chemosphere* **2010**, *81*, 1006–1011. [\[CrossRef\]](#)
118. Phung, D.T.; Connell, D.; Miller, G.; Hodge, M.; Patel, R.; Cheng, R.; Abeyewardene, M.; Chu, C. Biological monitoring of chlorpyrifos exposure to rice farmers in Vietnam. *Chemosphere* **2012**, *87*, 294–300. [\[CrossRef\]](#) [\[PubMed\]](#)
119. Lam, S.; Pham, G.; Nguyen-Viet, H. Emerging health risks from agricultural intensification in Southeast Asia: A systematic review. *Int. J. Occup. Environ. Health* **2017**, *23*, 250–260. [\[CrossRef\]](#)
120. Hien, M. *Agricultural Waste Treatment*. Nhan Dan Online, Official newspaper of the Communist Party of Vietnam 23 August 2022. Available online: <https://nhandan.vn/xu-ly-rac-thai-nong-nghiep-post711677.html> (accessed on 18 November 2024). (In Vietnamese)
121. Liebman, M.; Schulte, L.A. Enhancing agroecosystem performance and resilience through increased diversification of landscapes and cropping systems. *Elem. Sci. Anthr.* **2015**, *3*, 000041. [\[CrossRef\]](#)
122. Horgan, F.G.; Crisol-Martínez, E.; Stuart, A.M.; Villegas, J.M.; Peñalver-Cruz, A.; Mundaca, E.A.; Perez, M.O.; Bernal, C.C.; Almazan, M.L.P.; Ramal, A.F. Direct and Indirect Effects of Planting Density, Nitrogenous Fertiliser and Host Plant Resistance on Rice Herbivores and Their Natural Enemies. *Agriculture* **2022**, *12*, 2053. [\[CrossRef\]](#)
123. Dardonville, M.; Bockstaller, C.; Villerd, J.; Therond, O. Resilience of agricultural systems: Biodiversity-based systems are stable, while intensified ones are resistant and high-yielding. *Agric. Syst.* **2022**, *197*, 103365. [\[CrossRef\]](#)
124. Katayama, N.; Osada, Y.; Mashiko, M.; Baba, Y.G.; Tanaka, K.; Kusumoto, Y.; Okubo, S.; Ikeda, H.; Natuhara, Y. Organic farming and associated management practices benefit multiple wildlife taxa: A large-scale field study in rice paddy landscapes. *J. Appl. Ecol.* **2019**, *56*, 1970–1981. [\[CrossRef\]](#)
125. Newbold, T.; Hudson, L.N.; Hill, S.L.L.; Contu, S.; Lysenko, I.; Senior, R.A.; Börger, L.; Bennett, D.J.; Choimes, A.; Collen, B.; et al. Global effects of land use on local terrestrial biodiversity. *Nature* **2015**, *520*, 45–50. [\[CrossRef\]](#)
126. Firbank, L.G. Striking a new balance between agricultural production and biodiversity. *Ann. Appl. Biol.* **2005**, *146*, 163–175. [\[CrossRef\]](#)
127. Stein-Bachinger, K.; Gottwald, F.; Haub, A.; Schmidt, E. To what extent does organic farming promote species richness and abundance in temperate climates? A review. *Org. Agric.* **2021**, *11*, 1–12. [\[CrossRef\]](#)
128. Holzer, K.A.; Bayers, R.P.; Nguyen, T.T.; Lawler, S.P. Habitat value of cities and rice paddies for amphibians in rapidly urbanizing Vietnam. *J. Urban. Ecol.* **2017**, *3*, 3. [\[CrossRef\]](#)
129. Miyashita, T.; Chishiki, Y.; Takagi, S.R. Landscape heterogeneity at multiple spatial scales enhances spider species richness in an agricultural landscape. *Popul. Ecol.* **2012**, *54*, 573–581. [\[CrossRef\]](#)
130. Watanabe, K.; Koji, S.; Hidaka, K.; Nakamura, K. Abundance, diversity, and seasonal population dynamics of aquatic Coleoptera and Heteroptera in rice fields: Effects of direct seeding management. *Environ. Entomol.* **2013**, *42*, 841–850. [\[CrossRef\]](#)
131. Fried, O.; Westphal, C.; Schellenberg, J.; Grescho, V.; Kühn, I.; Nguyen, V.S.; Settele, J.; Bergmeier, E. Vascular plant species diversity in Southeast Asian rice ecosystems is determined by climate and soil conditions as well as the proximity of non-paddy habitats. *Agric. Ecosyst. Environ.* **2021**, *314*, 107346. [\[CrossRef\]](#)
132. Guo, L.; Zhao, L.; Ye, J.; Ji, Z.; Tang, J.-J.; Bai, K.; Zheng, S.; Hu, L.; Chen, X. Using aquatic animals as partners to increase yield and maintain soil nitrogen in the paddy ecosystems. *eLife* **2022**, *11*, 73869. [\[CrossRef\]](#) [\[PubMed\]](#)
133. Ito, H.C.; Shiraishi, H.; Nakagawa, M.; Takamura, N. Combined impact of pesticides and other environmental stressors on animal diversity in irrigation ponds. *PLoS ONE* **2020**, *15*, e0229052. [\[CrossRef\]](#)
134. Ishiwaka, N.; Hashimoto, K.; Masayoshi, K.; Hiraiwa, M.; Sanchez-Bayo, F.; Kadoya, T.; Hayasaka, D. Can warming accelerate the decline of Odonata species in experimental paddies due to insecticide fipronil exposure? *Environ. Pollut.* **2024**, *341*, 122831.

135. Scholes, R.; Montanarella, L.; Brainich, A.; Barger, N.; ten Brink, B.; Cantele, M.; Erasmus, B.; Fisher, J.; Gardner, T.; Holland, T.G.; et al. *Summary for Policymakers of the Assessment Report on Land Degradation and Restoration of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*; IPBES secretariat: Bonn, Germany, 2018.
136. Gao, P.; Wang, H.; Deng, S.; Dong, E.; Dai, Q. Influence of organic rice production mode on weed composition in the soil seed bank of paddy fields. *Front. Plant Sci.* **2022**, *13*, 1056975. [CrossRef]
137. Shaner, D.L. Lessons Learned From the History of Herbicide Resistance. *Weed Sci.* **2014**, *62*, 427–431. [CrossRef]
138. Baba, Y.G.; Kusumoto, Y.; Tanaka, K. Effects of agricultural practices and fine-scale landscape factors on spiders and a pest insect in Japanese rice paddy ecosystems. *BioControl* **2018**, *63*, 265–275. [CrossRef]
139. Ibáñez, C.; Curcó, A.; Riera, X.; Ripoll, I.; Sánchez, C. Influence on Birds of Rice Field Management Practices during the Growing Season: A Review and an Experiment. *Waterbirds* **2010**, *33*, 167–180. [CrossRef]
140. Lori, M.; Symnack, S.; Mäder, P.; de Deyn, G.; Gattinger, A. Organic farming enhances soil microbial abundance and activity—A meta-analysis and meta-regression. *PLoS ONE* **2017**, *12*, e0180442. [CrossRef]
141. Gong, S.; Zhou, X.; Zhu, X.; Huo, J.; Faghihinia, M.; Li, B.; Zou, Y. Organic rice cultivation enhances the diversity of above-ground arthropods but not below-ground soil eukaryotes. *Agric. Ecosyst. Environ.* **2023**, *347*, 108390.
142. Sattler, C.; Gianuca, A.T.; Schweiger, O.; Franzen, M.; Settele, J. Pesticides and land cover heterogeneity affect functional group and taxonomic diversity of arthropods in rice agroecosystems. *Agric. Ecosyst. Environ.* **2020**, *297*, 106927. [CrossRef]
143. Elphick, C.S.; Baicich, P.; Parsons, K.C.; Fasola, M.; Mugica, L. The Future for Research on Waterbirds in Rice Fields. *Waterbirds* **2010**, *33*, 231–243. [CrossRef]
144. Vietnam Academy of Agricultural Sciences. *Building a Sustainable Ecosystem for the Mekong Delta Region*. 11 January 2023. Available online: <https://vaas.vn/vi/nong-nghiep-trong-nuoc/xay-dung-he-sinh-thai-ben-vung-cho-vung-dbscl> (accessed on 18 November 2024). (In Vietnamese)
145. Tuyet, C. *Mekong Delta: Challenges in Nature Conservation and Biodiversity*. Resources and Environment 2 July 2019. Available online: <https://baotainguyenmoitruong.vn/dbscl-thach-thuc-trong-bao-ton-thien-nhien-va-da-dang-sinh-hoc-241920.html> (accessed on 17 November 2024). (In Vietnamese)
146. Mai, C. *Mekong Delta: Agricultural Wastewater Alarm*. Resources and Environment. 27 August 2019. Available online: <https://baotainguyenmoitruong.vn/dong-bang-song-cuu-long-bao-dong-nuoc-thai-nong-nghiep-294481.html> (accessed on 16 November 2024). (In Vietnamese)
147. WBCSD World Business Council for Sustainable Development. Deep dive: Rice production in the Mekong Delta, Vietnam. In *Roadmap to Nature Positive: Foundations for the Agri-Food System*; WBCSD: Geneva, Switzerland, 2024.
148. Nguyen, C.T.; Nguyen, V.M.; Nguyen, V.A.; Phan, T.P.T.; Doan, T.H.C.; Nguyen, T.H.; Nguyen, T.H. Some Initial Results on Research and Modeling of Organic Rice Production in the Mekong Delta, Vietnam. *Int. J. Energy Environ. Sci.* **2017**, *1*, 29–36.
149. Nguyen, C.T.; Vo, T.T.; Nguyen, T.X.; Nguyen, H.P.; Bui, D.D.; Nguyen, V.M.; Duong, V.H. Assessment of Pesticide Residues in Organic Rice Production in The Mekong Delta, Vietnam. *Eur. J. Dev. Stud.* **2022**, *2*, 1–11. [CrossRef]
150. Nguyen, C.T. Linking Farmers and Businesses in Integrated Organic Rice and Shrimp Farming—The Best Way for Enhancing Farmer’s Income and Sustainable Agriculture Development. *Agric. Ext. J.* **2019**, *3*, 58–66.
151. STAMEQ Directorate for Standards, Metrology and Quality. *Vietnam National Standard on Organic Agriculture—Part 2: Organic Crops (TCVN 11041-2:2017)*. Ministry of Science and Technology. 2017. Available online: <https://caselaw.vn/van-ban-phap-luat/343712-tieu-chuan-quoc-gia-tcvn-11041-2-2017-ve-nong-nghiep-huu-co-phan-2-trong-trot-huu-co-nam-2017> (accessed on 17 November 2024). (In Vietnamese)
152. Thai, H.; Vietnam’s Agriculture Is Making Steady Progress with Organic Rice. Vietnam Organic Agriculture Association, Hanoi. 23 December 2024. Available online: <https://nongnghiephuuc.vn/nen-nong-nghiep-viet-nam-tien-vung-chac-tu-lua-huu-co-3713.html>. (accessed on 8 September 2025). (In Vietnamese)
153. Nguyen, C.T.; Van, T.T.T. Development of Organic Agriculture in the Mekong Delta—Opportunities and Challenges. *Eur. J. Dev. Stud.* **2021**, *1*, 29–35.
154. Grimm, M.; Luck, N.; Steinhübel, F. Consumers’ willingness to pay for organic rice: Insights from a non-hypothetical experiment in Indonesia. *Aust. J. Agric. Res. Econ.* **2023**, *67*, 83–103.
155. Pawitri, G.; Budiraharjo, K.; Setiawan, B.M. The Production Efficiency in Organic Rice Farming. *SOCA J. Sos. Ekon. Pertan.* **2021**, *15*, 450–457.
156. Reddy, B.S. Organic Farming: Status, Issues and Prospects—A Review. *Agric. Econ. Res. Rev.* **2010**, *23*, 343–358.
157. Ashraf, I.; Ahmad, I.; Nafees, M.; Yousaf, M.M.; Ahmad, B. A review on organic farming for sustainable agricultural production. *Pure Appl. Biol.* **2021**, *5*, 277–286. [CrossRef]
158. Eyhorn, F.; van den Berg, M.; Decock, C.; Maat, H.; Srivastava, A. Does Organic Farming Provide a Viable Alternative for Smallholder Rice Farmers in India? *Sustainability* **2018**, *10*, 4424. [CrossRef]
159. Adhikari, R.K. Economics of organic rice production. *J. Agric. Environ.* **2011**, *12*, 97–103. [CrossRef]

160. Mungara, E.; Indradewa, D.; dan Rogomulyo, R. Analysis of Growth and Rice Yields (*Oryza sativa* L.) Conventional, Organic Transitional, and Organic Farming System. *Vegetalika* **2013**, *2*, 1–12.
161. Tuan, C.M.; Lee, S.H. Factors Affecting Organic Fertiliser Adoption in Rice Production in Vietnam. *J. Korean Soc. Int. Agric.* **2021**, *33*, 130–138. [\[CrossRef\]](#)
162. Rahman, F.H.; Mukherjee, S.; Das, S.; Mukhopadhyay, K.; Bera, R.R.; Seal, A. Improvement of Soil and Plant Health through Adoption of an Organic Package of Practice for Rice Cultivation in New Alluvial Soil of West Bengal. *Curr. J. Appl. Sci. Technol.* **2020**, *39*, 99–108. [\[CrossRef\]](#)
163. Salam, M.A.; Sarker, M.N.I.; Sharmin, S. Do organic fertiliser impact on yield and efficiency of rice farms? Empirical evidence from Bangladesh. *Heliyon* **2021**, *7*, e07731.
164. Aulakh, C.S.; Ravisankar, N. Organic farming in Indian context: A perspective. *Agric. Res. J.* **2017**, *54*, 149–164. [\[CrossRef\]](#)
165. Rentschler, J.; Salhab, M.; Jafino, B.A. Flood exposure and poverty in 188 countries. *Nat. Commun.* **2022**, *13*, 3527. [\[CrossRef\]](#) [\[PubMed\]](#)
166. Xiao, H.; Tang, Y.; Li, H.; Zhang, L.; Ngo-Duc, T.; Chen, D.; Tang, Q. Saltwater intrusion into groundwater systems in the Mekong Delta and links to global change. *Adv. Clim. Change Res.* **2021**, *12*, 342–352. [\[CrossRef\]](#)
167. Thai, D.V.H.; Cong, V.T.; Nestmann, F.; Oberle, P.E.; Trung, N.N. Land use based flood hazards analysis for the Mekong Delta. In Proceedings of the 19th IAHR-APD Congress, Hanoi, Vietnam, 21–24 September 2014.
168. Hopmans, J.W.; Qureshi, A.S.; Kisekka, I.; Munns, R.; Grattan, S.R.; Rengasamy, P.; Ben-Gal, A.; Assouline, S.; Javaux, M.; Minhas, P.S.; et al. Critical knowledge gaps and research priorities in global soil salinity. *Adv. Agron.* **2021**, *16*, 1–191.
169. Chandramohan, K.T.; Sreeja, P. An indigenous saline resistant organic rice tract in north Kerala—Practices and challenges. In Proceedings of the Lake 2016: Conference on Conservation and Sustainable Management of Ecologically Sensitive Regions in Western Ghats, 10th Biennial Lake Conference, Karnataka, India, 28–30 December 2016.
170. Vanaja, T.; Neema, V.P.; Mammooty, K.P.; Balakrishnan, P.C.; Jayaprakash, N.B. The first high yielding saline tolerant rice variety suited to the Kaipad tidal farming ecosystem of Kerala, India, and suited for flood prone and water scarce environments: ‘Ezhome-1’. *J. Org.* **2015**, *2*, 21–32.
171. Vanaja, T.; Neema, V.P.; Mammooty, K.P.; Balakrishnan, P.C.; Jayaprakash, N.B. A high yielding organic rice variety suited for coastal saline and non-saline fields: ‘Ezhome-2’. *J. Org.* **2017**, *4*, 21–28.
172. Hossain, M.S.; Khan, H.R.; Akter, S. Consequences of Indigenous Organic Amendments and Moisture Conditions on the Growth and Yield of Rice Grown on Saline Soil. *Int. J. Sci. Eng. Investig.* **2017**, *6*, 108–114.
173. Eslami, S.; Hoekstra, P.; Minderhoud, P.S.; Trung, N.N.; Hoch, J.M.; Sutanudjaja, E.H.; Dung, D.D.; Tho, T.Q.; Voepel, H.E.; Woillez, M.N.; et al. Projections of salt intrusion in a mega-delta under climatic and anthropogenic stressors. *Commun. Earth Environ.* **2021**, *2*, 142. [\[CrossRef\]](#)
174. Kaveney, B.; Barrett-Lennard, E.; Minh, K.C.; Duy, M.D.; Thi, K.P.N.; Kristiansen, P.; Orgill, S.; Stewart-Koster, B.; Condon, J. Inland dry season saline intrusion in the Vietnamese Mekong River Delta is driving the identification and implementation of alternative crops to rice. *Agric. Syst.* **2023**, *207*, 103632. [\[CrossRef\]](#)
175. Poelma, T.; Bayrak, M.M.; Van Nha, D.; Tran, T.A. Climate change and livelihood resilience capacities in the Mekong Delta: A case study on the transition to rice–shrimp farming in Vietnam’s Kien Giang Province. *Clim. Change* **2021**, *164*, 9. [\[CrossRef\]](#)
176. Kolsky, M. *Flood Resilience in Agriculture: Safeguarding Food Supply*. Agritask. 26 July 2023. Available online: <https://start.agritask.com/blog/flood-resilience-in-agriculture/> (accessed on 22 November 2024).
177. Nguyen, H.N.; Vu, K.T.; Nguyen, X.N. Flooding in Mekong River Delta, Viet Nam. In *Human Development Report 2007/2008*; Human Development Report Office, Ed.; UNDP United Nations Development Programme: Nairobi, Kenya, 2007; pp. 1–23.
178. Van, C.T.; Thuy, H.T.T.; Viet, C.T.; Anh, L.N.; Van Anh, V.T.; Tran, D.D. Unveiling flood vulnerability in the Vietnamese Mekong Delta: A case study of An Giang province. *Int. J. Disaster Risk Reduct.* **2024**, *106*, 104429. [\[CrossRef\]](#)
179. Jacob, S.; Subramannian, S. Pokkali: An Exquisite Climate Resilient and Organic Rice-Prawn Production System for Future. *Kerala Karshakan* **2022**, *10*, 4–8.
180. Vanaja, T.; Neema, V.P. Saline and flood tolerant organic red rice variety (*Oryza sativa* L.) from Kerala. *Int. J. Trop. Agric.* **2021**, *39*, 113–118.
181. Setyorini, D.; Hartatik, W. Nutrients balance under organic rice farming system in Central Java, Indonesia. *IOP Conf. Ser. Earth Environ. Sci.* **2021**, *648*, 012154. [\[CrossRef\]](#)
182. Alagappan, S.; Venkitaswamy, R.; Mariappan, G. Impacts of different sources of organic manures on soil physico-chemical properties, nutrient balance and yield of rice-green gram cropping sequence under organic farming. *Agric. Updat.* **2017**, *12*, 2044–2066. [\[CrossRef\]](#)
183. Gaiind, S. Effect of organic farming on quality of soil under rice cultivation. *J. Agroecol. Nat. Resour.* **2015**, *2*, 391–393.
184. Sujianto; Gunawan, E.; Saptana; Syahyuti; Darwis, V.; Ashari; Syukur, M.; Ariningsih, E.; Saliem, H.P.; Mardianto, S.; et al. Farmers’ perception, awareness, and constraints of organic rice farming in Indonesia. *Open Agric.* **2022**, *7*, 284–299. [\[CrossRef\]](#)

185. Mendoza, T.C. Evaluating the Benefits of Organic Farming in Rice Agroecosystems in the Philippines. *J. Sustain. Agric.* **2004**, *24*, 93–115. [\[CrossRef\]](#)
186. Hossain, M.; Elahi, S.; Khondaker, M. A comparison between ecological and conventional rice farming systems in Bangladesh. In Proceedings of the 7th International Specialized Conference on Diffuse Pollution and Basin Management, Dublin, Ireland, 18–22 August 2003.
187. Loc, H.H.; Van Binh, D.; Park, E.; Shrestha, S.; Dung, T.D.; Son, V.H.; Truc, N.H.T.; Mai, N.P.; Seijger, C. Intensifying saline water intrusion and drought in the Mekong Delta: From physical evidence to policy outlooks. *Sci. Total Environ.* **2021**, *757*, 143919. [\[CrossRef\]](#) [\[PubMed\]](#)
188. Lavane, K.; Kumar, P.; Meraj, G.; Han, T.G.; Ngan, L.H.B.; Lien, B.T.B.; Van Ty, T.; Thanh, N.T.; Downes, N.K.; Nam, N.D.G.; et al. Assessing the effects of drought on rice yields in the Mekong Delta. *Climate* **2023**, *11*, 13. [\[CrossRef\]](#)
189. François, M.; Moreau, R.; Sylvander, B. Question 3. How can organic farming contribute to environmental conservation? In *Agriculture Biologique En Martinique*; François, M., Moreau, R., Sylvander, B., Eds.; IRD Éditions: Marseille, France, 2005; pp. 250–260.
190. Crippa, M.; Guizzardi, D.; Pagani, F.; Banja, M.; Muntean, M.; Schaaf, E.; Becker, W.; Monforti-Ferrario, F.; Quadrelli, R.; Risquez Martin, A.; et al. *GHG Emissions of All World Countries*; Publications Office of the European Union: Luxembourg, 2023.
191. Qian, H.; Zhu, X.; Huang, S.; Linqvist, B.; Kuzyakov, Y.; Wassmann, R.; Minamikawa, K.; Martinez-Eixarch, M.; Yan, X.; Zhou, F.; et al. Greenhouse gas emissions and mitigation in rice agriculture. *Nat. Rev. Earth Environ.* **2023**, *4*, 716–732. [\[CrossRef\]](#)
192. Sapkota, T.B.; Shankar, V.; Rai, M.; Jat, M.L.; Stirling, C.M.; Singh, L.K.; Jat, H.S.; Grewal, M.S. Reducing global warming potential through sustainable intensification of basmati rice-wheat systems in India. *Sustainability* **2017**, *9*, 1044. [\[CrossRef\]](#)
193. Bacenetti, J.; Fusi, A.; Negri, M.; Bocchi, S.; Fiala, M. Organic production systems: Sustainability assessment of rice in Italy. *Agric. Ecosyst. Environ.* **2016**, *225*, 33–44. [\[CrossRef\]](#)
194. Chen, X.; Cui, Z.; Fan, M.; Vitousek, P.; Zhao, M.; Ma, W.; Wang, Z.; Zhang, W.; Yan, X.; Yang, J.; et al. Producing more grain with lower environmental costs. *Nature* **2014**, *514*, 486–489. [\[CrossRef\]](#) [\[PubMed\]](#)
195. Yodkhum, S.; Gheewala, S.H.; Sampattagul, S. Life cycle GHG evaluation of organic rice production in northern Thailand. *J. Environ. Manag.* **2017**, *196*, 217–223. [\[CrossRef\]](#)
196. Tran, M.; Lam, O.; Long, G. *Unlocking SDG Interlinkages Through Integrated Governance for SDG6 in Vietnam's Mekong Delta*. Research Brief; Living Deltas Research Hub: 2023. Available online: <https://livingdeltas.org/resources/unlocking-sdg-interlinkages?rq=unlocking%20sdg> (accessed on 18 November 2024).
197. Setboonsarng, S.; Acharya, B.N. What Motivates Farmers to Adopt Organic Agriculture? A Case of Rainfed Organic Rice in Thailand. In *Organic Agriculture and Post-2015 Development Goals*; Setboonsarng, S., Markandya, A., Eds.; Asian Development Bank: Mandaluyong City, Philippines, 2015; pp. 133–154.
198. Devi, L.Y.; Irham, S.; Anatasari, E.; Nurhayati, A.; Wahyu Widada, A. Key drivers of organic rice productivity in Sleman and Magelang Regencies. *IOP Conf. Ser. Earth Environ. Sci.* **2021**, *746*, 012005. [\[CrossRef\]](#)
199. Hossain, S.T.; Sugimoto, H.; Ueno, H.; Huque, S.M.R. Adoption of organic rice for sustainable development in Bangladesh. *J. Org. Syst.* **2007**, *2*, 27–37.
200. Liu, J.; Liu, H.; Liu, R.; Amin, M.G.M.; Zhai, L.; Lu, H.; Wang, H.; Zhang, X.; Zhang, Y.; Zhao, Y.; et al. Water Quality in Irrigated Paddy Systems. In *Irrigation in Agroecosystems*; Ondrašek, G., Ed.; IntechOpen: London, UK, 2018; p. 77339.
201. Trinh, H.T.; Marcussen, H.; Hansen, H.C.B.; Le, G.T.; Duong, H.T.; Ta, N.T.; Nguyen, T.Q.; Hansen, S.; Strobel, B.W. Screening of inorganic and organic contaminants in floodwater in paddy fields of Hue and Thanh Hoa in Vietnam. *Environ. Sci. Pollut. Res.* **2017**, *24*, 7348–7358. [\[CrossRef\]](#) [\[PubMed\]](#)
202. Anyusheva, M.; Lamers, M.; La, N.; Nguyen, V.V.; Streck, T. Fate of Pesticides in Combined Paddy Rice—Fish Pond Farming Systems in Northern Vietnam. *J. Environ. Qual.* **2012**, *41*, 515–525. [\[CrossRef\]](#)
203. Braun, G.; Braun, M.; Kruse, J.; Amelung, W.; Renaud, F.G.; Khoi, C.M.; Duong, M.V.; Sebesvari, Z. Pesticides and antibiotics in permanent rice, alternating rice-shrimp and permanent shrimp systems of the coastal Mekong Delta, Vietnam. *Environ. Int.* **2019**, *127*, 442–451. [\[CrossRef\]](#) [\[PubMed\]](#)
204. Duong, H.V.; Nguyen, T.C.; Nguyen, X.T.; Nguyen, M.Q.; Nguyen, P.H.; Vo, T.T. Evaluating the Presence of Pesticide Residues in Organic Rice Production in An Giang Province, Vietnam. *J. Sustain. Dev.* **2022**, *15*, 49–64. [\[CrossRef\]](#)
205. Wahyudi, D.; Ardiansyah, A.; Asiah, N.; Madonna, S. Comparative study on agrochemical residue on rice cultivation in Tasikmalaya, Indonesia: Organic versus conventional. *Future Food* **2020**, *8*, 1–12.
206. Larsen, A.E.; Noack, F. Identifying the landscape drivers of agricultural insecticide use leveraging evidence from 100,000 fields. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 5473–5478.
207. Cheng, C.; Zhang, F.; Shi, J.; Kung, H. What is the relationship between land use and surface water quality? A review and prospects from remote sensing perspective. *Environ. Sci. Pollut. Res.* **2022**, *29*, 56887–56907. [\[CrossRef\]](#)

208. VoPham, T.; Wilson, J.P.; Ruddell, D.; Rashed, T.; Brooks, M.M.; Yuan, J.; Talbott, E.; Chang, C.H.; Weissfeld, J.L. Linking pesticides and human health: A geographic information system (GIS) and Landsat remote sensing method to estimate agricultural pesticide exposure. *Appl. Geogr.* **2015**, *62*, 171–181. [\[CrossRef\]](#)
209. Pinichka, C.; Issarapan, P.; Siriwong, W.; Kongtip, P.; Chotpantarat, S.; Bundhamcharoen, K. Application of Geographic Information Systems and Remote Sensing for Pesticide Exposure and Health Risk Assessment in Thailand. *Outbreak Surveill. Investig. Response J.* **2019**, *12*, 75–83.
210. Braun, G.; Sebesvari, Z.; Braun, M.; Kruse, J.; Amelung, W.; An, N.; Renaud, F. Does sea-dyke construction affect the spatial distribution of pesticides in agricultural soils?—A case study from the Red River Delta, Vietnam. *Environ. Pollut.* **2018**, *243*, 890–899. [\[CrossRef\]](#)
211. Livsey, J.; Da, C.T.; Scaini, A.; Lan, T.H.P.; Long, T.X.; Berg, H.; Manzoni, S. Floods, soil and food—Interactions between water management and rice production within An Giang province, Vietnam. *Agric. Ecosyst. Environ.* **2021**, *320*, 107589. [\[CrossRef\]](#)
212. Hou, L.; Liu, P.; Huang, J.; Deng, X. The influence of risk preferences, knowledge, land consolidation, and landscape diversification on pesticide use. *Agric. Econ.* **2020**, *5*, 759–776. [\[CrossRef\]](#)
213. Piadozo, M.E.S.; Lantican, F.A.; Pabuayon, I.M.; Quicoy, A.R.; Suyat, A.M.; Maghirang, P.K.B. Rice farmers' concept and awareness of organic agriculture: Implications for sustainability of Philippine organic agriculture program. *J. Int. Soc. Southeast Asian Agric. Sci.* **2014**, *20*, 142–156.
214. Sapkota, B.K.; Subedi, A.P.; Tripathi, K.M.; Dhakal, S.C. Economics of organic vs inorganic rice production: A Case of Chitwan district of Nepal. *J. Nepal Agric. Res. Council.* **2021**, *7*, 109–121. [\[CrossRef\]](#)
215. Mahdavi, H.; Allahyari, M.S.; Damalas, C.A.; Dunn, E.S. Drivers and barriers for organic rice (*Oryza sativa* L.) production in northern Iran: Experts' consensus using the Delphi method. *Biol. Agric. Hortic.* **2020**, *36*, 96–106. [\[CrossRef\]](#)
216. Hasanah, Y.; Hanum, H.; Rusmarilin, H. Organic Rice Cultivation System to Support Food Security. *J. St. Transf.* **2020**, *2*, 47–54.
217. Woranoot, I. Implications of Organic Farming in Development: Experiences from Organic Rice Farms in Northeastern Thailand. Master's Thesis, Erasmus University, Rotterdam, The Netherlands, 2009.
218. Colombo, S.L.; Chiarella, S.G.; Lefrançois, C.; Fradin, J.; Raffone, A.; Simone, L. Why Knowing about Climate Change Is Not Enough to Change: A Perspective Paper on the Factors Explaining the Environmental Knowledge-Action Gap. *Sustainability* **2023**, *15*, 14859. [\[CrossRef\]](#)
219. Lal, A.E.; Mazhar, S.H.; Jahanara, J. Study of Socio-Economic Condition and Constraints faced by the Rice Growing Farmers in Adoption of Organic and Bio-Fertiliser in Bilaspur District of Chhattisgarh. *Int. J. Adv. Agric. Sci. Technol.* **2021**, *8*, 45–47.
220. Orlando, F.; Alali, S.; Vaglia, V.; Pagliarino, E.; Baig, J.; Bocchi, S. Participatory approach for developing knowledge on organic rice farming: Management strategies and productive performance. *Agric. Syst.* **2020**, *178*, 102739. [\[CrossRef\]](#)
221. Boun My, K.; Nguyen-Van, P.; Kim Cuong Pham, T.; Stenger, A.; Tiet, T.; To-The, N. Drivers of organic farming: Lab-in-the-field evidence of the role of social comparison and information nudge in networks in Vietnam. *Ecol. Econ.* **2022**, *196*, 107401. [\[CrossRef\]](#)
222. Lu, C.-F.; Cheng, C.-Y. Exploring the distribution of organic farming: Findings from certified rice in Taiwan. *Ecol. Econ.* **2023**, *212*, 107915. [\[CrossRef\]](#)
223. Suminah, S.; Sundari, M.T.; Wijianto, A.; Rusdiyana, E. Farmers attitudes to organic rice cultivation at the industrial revolution era. *IOP Conf. Ser. Earth Environ. Sci.* **2022**, *1001*, 012031.
224. Parajuli, S.; Shrestha, J.; Ghimire, S. Organic farming in Nepal: A viable option for food security and agricultural sustainability. *Arch. Agric. Environ. Sci.* **2020**, *5*, 223–230. [\[CrossRef\]](#)
225. Beban, A. Is Organic Agriculture a Viable Strategy in Contexts of Rapid Agrarian Transition? Evidence from Cambodia. *J. Agric. Food Syst. Community Dev.* **2014**, *4*, 131–147. [\[CrossRef\]](#)
226. Rubinos, R.; Jalipa, A.T.; Bayacag, P. Comparative economic study of organic and conventional rice farming in Magsaysay, Davao Del Sur. In Proceedings of the Philippine Statistical Association 10th National Convention on Statistics, Manila, Philippines, 1 October 2007.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.