



1 **Detecting the resilience of soil moisture dynamics to drought** 2 **periods as function of soil type and climatic region**

3 Nedal Aqel¹, Jannis Groh^{2,3}, Lutz Weihermüller², Ralf Gründling⁴, Andrea Carminati¹, Peter
4 Lehmann¹

5 ¹Physics of Soils and Terrestrial Ecosystems, ETH Zurich, Zurich, Switzerland

6 ²Institute of Bio- and Geoscience IBG-3: Agrosphere, Forschungszentrum Jülich GmbH, Jülich, Germany

7 ³Biogeochemistry and Gas Fluxes, Leibniz Institute for Agricultural and Landscape Research (ZALF),
8 Müncheberg, Germany

9 ⁴Department of Soil System Science, Helmholtz-Zentrum für Umweltforschung GmbH – UFZ, Halle, Germany

10 *Corresponding author, nedal.aqel@usys.ethz.ch, Universitätstrasse 16, 8092 Zurich, Switzerland

11 **Abstract**

12 Abrupt changes in climatic conditions and land management can cause permanent shifts in soil
13 hydraulic response to climatic inputs, impacting soil functions and established soil–climate interactions.
14 To quantify the resilience of soil water content dynamics after abrupt changes in environmental
15 conditions, we present a model framework combining a neural network with seasonal trend analysis
16 (STL). Using data from a series of lysimeters from the TERrestrial ENvironmental Observatories
17 (TERENO) - SOILCan lysimeter network, we identified changes in soil water content responses after
18 an extremely hot and dry summer in Germany in 2018. The model incorporates meteorological variables
19 decomposed into seasonal and long-term components along with a categorical indicator of current
20 moisture conditions. It is trained on data from a reference site with stable soil water content response
21 and applied to lysimeters from multiple origins exposed to contrasting climates. By analysing annual
22 residual patterns—particularly mean bias over time—soil water content state dynamics is classified as
23 ‘stable’, ‘resilient’, or ‘changed’, reflecting whether the system maintains, recovers, or diverges from
24 its original state. We found that soils preserve the response function to environmental forcing under
25 typical conditions but exhibit structural change when relocated to new environments, even when soil
26 texture remains constant. The proposed method offers a scalable and non-invasive tool for tracking
27 changes in the response of soil water content to climatic change and provides early indicators of changes
28 in essential soil functions and soil health status.



1. Introduction

Soil water content plays a fundamental role in hydrological processes and land–atmosphere interactions, governing the exchange of water and energy at the Earth’s surface (Seneviratne et al., 2010; Sun et al., 2025). It regulates key hydrological functions, including infiltration, runoff generation, evapotranspiration, and groundwater recharge. Through these processes, soil water content influences water availability, ecosystem productivity, and climatic conditions across local to global scales (Bogena et al., 2015; Fatichi et al., 2020). Soil water content status and related soil environmental conditions change with short- and long-term atmospheric processes. This response of soil water content to atmospheric conditions, which we define as ‘soil water content response function’, determines, for example, if anaerobic conditions are inhibited after heavy rainfall (by fast percolation to deeper soil layers) and if enough water remains available after dry periods for plant growth, temperature regulation, and chemical reactions. In short, the soil water content response function is a dominant factor of the soil health status. This response function is shaped by soil formation processes and reflects adaptation to the dominant climatic conditions (Kuzakov and Zamanian, 2019; Sainju et al., 2022; Al-Shammari et al., 2025). Accordingly, a change in the response function after extreme climatic events is likely to imply a change in soil health.

In the core of this study is the question how changes in the soil water content response function, and thus in soil properties and health, can be detected. The standard approach to determine the response function is to apply physically based models, for example by inverse modelling of soil water content dynamics under varying boundary conditions (Šimůnek et al., 2016). These models require detailed knowledge of soil hydraulic properties and extensive calibration, limiting their generalization beyond the spatial scale and the local conditions used in the calibration process (Or, 2020; Lehmann et al., 2020; O. & Orth, 2021). Additionally, such approaches are typically destructive, limiting the options of conducting a time series analysis. Moreover, these models typically assume static soil characteristics, failing to adequately represent structural changes in soil properties — such as compaction, degradation, or organic matter loss — that can substantially alter hydraulic behaviour over time (Fatichi et al., 2020; Melsen & Guse, 2021; Wankmüller et al., 2024). Most of the models also neglect or oversimplify the



56 hysteretic nature of the soil water retention curve, as well as seasonal changes in soil hydraulic
57 properties that can substantially alter infiltration, drainage, and plant water availability (Aqel et al.,
58 2024; Hannes et al., 2016; Herbrich & Gerke, 2017). Therefore, we used a non-invasive approach based
59 on neural networks as discussed below.


60 In recent years, artificial intelligence (AI), particularly neural networks, has emerged as a promising
61 alternative for modelling complex hydrological processes (Reichstein et al., 2019). These data-driven
62 models have demonstrated the capacity to learn nonlinear relationships directly from observational
63 datasets without relying heavily on explicit physical equations (Kratzert et al., 2019; Mosavi et al.,
64 2018; Shen et al., 2018). Within soil hydrology, neural networks have been used to characterise
65 hysteretic soil–water behaviour from training data, improving the representation of wetting–drying
66 cycles without explicit hysteresis parameterisation (Aqel et al., 2024). They have also been applied to
67 soil-moisture time-series modelling using Long Short-Term Memory (LSTM) networks with recurrent
68 architectures suited to capture long-range temporal dependencies (Liu et al., 2023; O. & Orth, 2021).
69 Across diverse hydro-climatic regimes, LSTM have been shown to effectively learn nonlinear
70 relationships between climatic inputs and soil water content, often matching or surpassing traditional
71 physically based approaches and demonstrating strong generalisation (Kratzert et al., 2019; J. Liu et al.,
72 2022, 2023; O. & Orth, 2021).

73 Independent of the chosen modelling approach, these models ignore that the soil water content response
74 function (i.e., the variations in soil water content following changes in atmospheric conditions) can
75 change at larger time scale. Experimental studies have shown that extreme events such as drought can
76 induce persistent shifts in soil water content dynamics, potentially leading to alternative stable states
77 (Robinson et al., 2016). Moreover, changes in land use — such as forest conversion to agriculture or
78 bare land — alter soil hydraulic properties, with effects on infiltration, water retention, and saturated
79 hydraulic conductivity (Fu et al., 2021; Robinson et al., 2022). Considering that texture remains
80 constant at time scales of decades, the changes in the response function are likely to be related to
81 changes in soil structure primarily.



82 Soil systems are subject to temporal change, yet models are often trained on historical datasets without
 83 evaluating whether the system dynamics remain stable throughout the training period (Montanari et al.,
 84 2013; Vaze et al., 2010). As a result, models may be applied to prediction settings where underlying
 85 soil–climate interactions differ from those on which the model was trained. For example, a recent study
 86 comparing different crop models using the TERENO-SOILCan set-up showed that predicting
 87 agronomic and environmental variables under different climatic conditions to those represented in the
 88 training datasets resulted in significant discrepancies between simulations and observations (Groh et
 89 al., 2022). This underscores the critical need to assess whether site-specific representations of soil–
 90 water behaviour remain valid over time (Hrachowitz et al., 2013) and highlight the need for more
 91 adaptable modelling approaches under evolving environmental conditions (Blöschl et al., 2019; Milly
 92 et al., 2008). Considering these limitations, none of the discussed approaches can capture the change in
 93 the soil water content response function. A recent modelling framework by Jarvis et al. (2024) takes a
 94 major conceptual step forward by explicitly representing soil-structure dynamics and their feedback on
 95 hydraulic behaviour. However, as the authors emphasize, such process-based models still depend on
 96 detailed observational data to constrain temporal changes in structure and hydraulic properties.

97
 98 To address this gap, the present study introduces a framework based on neural networks and seasonal
 99 trend decomposition. Specifically, we quantify the change in the response function after the 2018
 100 drought in summer, which was a Europe-wide event, but in Germany was particularly characterized by
 101 an extreme combination of high temperatures and low precipitation (Xoplaki et al., 2025). The response
 102 of soil water content on this drought will be analysed for a set of lysimeters. As shown in Fig. 1 for one
 103 of the study sites, the monthly water deficit (potential evapotranspiration minus precipitation) peaked
 104 in summer 2018 indicating the strong drought in this period.



105
 106 **Figure 1** Variations in climatic conditions at Selhausen (SE) expressed as difference between potential
 107 evapotranspiration (PET) and precipitation (P) cumulated over the precedent 30 days (one month). The extreme
 108 summer 2018 is manifested by a maximum monthly deficit of ~150 mm. Details on the calculation of PET are
 109 provided in section 2.1.1.

110 The general objective of this study is thus to introduce a model framework to quantitatively detect
 111 changes in the soil water content response function and to classify the response as ‘stable’, ‘resilient’,
 112 or ‘changed’.

113 In principle, it would be possible to develop a quantitative framework to detect changes in the response
 114 function exclusively based on experimental data of time series (without modelling) as for example by
 115 the application of wavelet analysis (Ehrhardt et al., 2025). In this study, however, we pursue a different
 116 approach based on predictive modelling, in which the temporal evolution of differences between
 117 measurements and predictions serves as an indicator of changes in the response function, yields both,
 118 accurate predictions of soil water content dynamics (not in the focus of this study) and detection of
 119 changes in the relationship between climate and soil water content dynamics.

120 2. Material and Methods

121 We developed a data-driven modelling framework that combines time-series decomposition of climatic
 122 inputs with a feed-forward neural network to predict the daily soil water content (Fig. 2).




Figure 2 Schematic overview of the modelling framework for daily soil water content prediction. Input features include the analysis of a climatic variable (top left), its long-term trend component (bottom left), and the categorical soil water content state ('wet', 'moderate', 'dry'; top middle). These features, derived from observed data, are used to train a feed-forward neural network (centre), which outputs daily predictions of volumetric water content (right). The model thus captures temporal soil water content dynamics based on structured climate signals and categorical conditions.

The approach explicitly incorporates precipitation, potential evapotranspiration, and their difference (climatic water balance) as primary inputs. Each of these climate drivers was decomposed into seasonal variations and long-term trend components using Seasonal-Trend decomposition (STL) and was included as a separate feature in the model (Boergens et al., 2024; Cleveland et al., 1990). The input features also included a categorical moisture class (type) that reflects the expected current soil water condition ('wet', 'moderate', 'dry'). This design reflects the understanding that changes in climate - such as shifts in rainfall and evaporative demand - substantially affect soil water availability and fluxes (Vereecken et al., 2022). The methodology is detailed in the following sections, which consists of five key steps: selecting study sites and datasets collected in contrasting hydro-climatic conditions (subsection 2.1), preprocessing the data and extracting meaningful signals features including STL (subsection 2.2), constructing and training a neural network model on a reference dataset (subsection 2.3), generating soil water content predictions for independent (non-training) sites using the trained



142 model (subsection 2.4), evaluating the model's performance with statistical metrics (subsection 2.5),
 143 and physical consistency checks (subsection 2.6).


144 2.1 Study Sites and Data Selection

145 2.1.1 Lysimeter Network TERENO SOILCan

146 The study was conducted using lysimeter data from the TERENO-SOILCan lysimeter network in
 147 Germany (Pütz et al., 2016) with a focus on two locations: Bad Lauchstädt (BL) and Selhausen (SE)
 148 (Fig. 3). These sites were selected for their contrasting climatic regimes and the specific set-up of
 149 lysimeters, providing a natural experiment on how climate variability influences soil hydrological
 150 behaviour for a variety of soils. The TERENO-SOILCan lysimeters were moved between and within
 151 observatories according to a modified space-for-time approach, to expose them to different climates
 152 (Groh et al. 2020). This allows us to compare the ecosystem response of the same soil, but under
 153 different climatic conditions. Selhausen is characterized by a humid, Atlantic-influenced climate
 154 (annual precipitation around 720 mm and mean air temperature around 10 °C), whereas Bad Lauchstädt
 155 represents a drier, more continental climate (annual precipitation roughly 487 mm and mean air
 156 temperature approximately 8.8 °C); both climate descriptions are based on Pütz et al. (2016). Long-
 157 term observations confirm that Bad Lauchstädt experiences significantly lower rainfall and higher
 158 evaporative demand than Selhausen, yielding a higher aridity index (ratio of potential
 159 evapotranspiration to precipitation) and more pronounced dry spells in the growing season. By
 160 including both, a wetter site Selhausen and a drier site Bad Lauchstädt, the model is evaluated under
 161 distinctly different moisture regimes, which is critical for testing the generality of the approach and to
 162 separate between climatic and soil type effects. For each lysimeter station (Bad Lauchstädt and
 163 Selhausen), 12 lysimeters (1 m² surface area, 1.5 m depth) arranged in hexagons with 6 lysimeters
 164 around a service well were included in the analysis to monitor soil water content along with
 165 meteorological variables. In this study, lysimeters are not used for drainage or storage estimates, but
 166 rather as instruments providing long-term, high-resolution time series of soil water content and matric
 167 potential under field conditions. The lysimeters contain undisturbed soil columns collected at four
 168 different locations (see Fig. 3a), each with three replicates (Pütz et al., 2016) and were managed as



169 arable land under crop rotation. While fertilization practices differed regionally until spring 2019, the
 170 overall management concept was comparable, ensuring that differences in water dynamics can be
 171 attributed to changes in the climate and soil rather than the management (Pütz et al., 2016).



172 **Figure 3** Overview of the study area with site locations, topsoil texture, and soil origin. (a) The TERENO-
 173 SOILCan network contains lysimeters from four climatic regions (different symbols and colours in map). Our
 174 analysis focuses on the TERENO–SOILCan sites Selhausen (SE) and Bad Lauchstädt (BL), located in the
 175 Eifel/lower rhine valley and Harz/central German lowland observatory of TERENO, respectively, because at both
 176 sites lysimeter clusters were built (represented by shaded areas and hexagons), collecting large soil columns from
 177 four distinct source regions (i.e., Dedelow (DD), Bad Lauchstädt (BL), Sauerbach (SB), and Selhausen (SE)). (b)
 178 The analysed soil horizons (10 cm depth) cover two textural classes, shown in the USDA soil texture triangle,
 179 assigned to four different soil types and a range of soil organic carbon contents (SOC) (numbers in the legend).
 180

181 To investigate the effects of climate on soil water dynamics, daily time series of precipitation (P),
 182 potential evapotranspiration (PET), matric potential, and volumetric soil water content (used as the
 183 target variable) were compiled. P at Selhausen was measured at the on-site SOILCan weather station,
 184 while for Bad Lauchstädt it was taken from the nearest long-term monitoring station operated by the
 185 Deutscher Wetterdienst (Leipzig/Halle, ID 2932; DWD Climate Data Center, 2025). PET was
 186 calculated with the FAO-56 Penman–Monteith model (Allen et al., 2006), using measured
 187 meteorological variables (air temperature, air pressure, relative humidity, radiation, and wind speed)
 188 according to SOILCan protocols (Pütz et al., 2016; Groh et al., 2020). In this context, the reference
 189 evapotranspiration (ET_0) calculated with the Penman–Monteith model for a clipped grass surface (FAO-



56) is used as a proxy for potential evapotranspiration (PET), representing the site-independent atmospheric evaporative demand. Soil matric potential was measured using MPS-1 sensors (Decagon Devices Inc., Pullman, WA, USA), and volumetric soil water content was measured with time-domain reflectometry probes (CS610, Campbell Scientific, North Logan, UT, USA). The observational record spans the period from 2015 to 2023 and includes measurements taken at a depth of 10 cm (deeper soil layers were not analysed; Pütz et al., 2016).

2.1.2 Definition of Reference Site for Model Framework

For model development, a single lysimeter moved from Dedelow to the Bad Lauchstädt lysimeter station (see Fig. 3a and 3b) was selected as the training dataset. This lysimeter was chosen due to its stable soil water content dynamics and minimal temporal drift in water retention properties over the observation period (see Fig. 4a). This lysimeter served as the reference dataset for developing the predictive model because it allows the definition of soil water content response function for the seasonal climatic conditions. The 23 remaining lysimeters at the Bad Lauchstädt and Selhausen site were used as independent test datasets (a contrasting example is shown in Fig. 4b) to evaluate model generalization and detect potential shifts in soil hydraulic behaviour across sites and years.




Figure 4 Soil water retention curves using data collected between 2015 and 2023 at 10 cm depth. Matric potential is plotted against volumetric water content, with data colour-coded by period: 2015–2016 (blue), 2017–2020 (green), and 2021–2023 (red). (a) Training lysimeter (moved from Dedelow to Bad Lauchstädt). (b) Test lysimeter (original soil from Selhausen in lysimeter station at Selhausen).



210 2.2 Data Preprocessing and Feature Engineering

211 All raw data were aggregated or resampled to a daily time step to support time-series analysis and
212 modelling. Any misaligned or duplicated timestamps were corrected to ensure consistency.

213 2.2.1 Seasonal-Trend decomposition

214 After cleaning and aligning the data, the climatic variables used as inputs in the modelling were selected.
215 In addition to raw P and PET data, the daily climatic water balance (WB) was included as an explicit
216 input. This variable reflects the net difference between P and PET, serving as a proxy for wetting or
217 drying conditions. Positive values indicate potential moisture accumulation (e.g., during rainfall-
218 dominated periods), while negative values reflect high evaporative demand and drying conditions (e.g.,
219 during hot, dry days). Including the WB helps the model to distinguish humid periods from dry ones.
220 By providing WB alongside P and PET, the model can learn both the individual and combined effects
221 of P and evaporative demand on soil water content dynamics (Brocca et al., 2010; Uber et al., 2018).
222 For example, it can infer that 10 mm of P during a high-PET summer day (low positive or negative WB)
223 is less likely to increase soil water content than the same P on a cool, low-PET Day (high positive WB).

224 To provide the model with structured representations of climate variability, each climatic time series
225 was decomposed into additive components using STL based on LOESS with LOESS as acronym for
226 ‘Locally Estimated Scatterplot Smoothing’ (Cleveland et al., 1990). STL is a non-parametric method
227 that separates a time series into three interpretable parts: a seasonal component representing repeating
228 seasonal patterns (such as wetting and drying cycles), a trend component capturing gradual long-term
229 changes (such as climate shifts), and a residual component containing short-term irregularities and high-
230 frequency noise (Cleveland et al., 1990). This decomposition was applied independently to the P, PET,
231 and WB time series. Only the seasonal and trend components were retained as input features, as they
232 contain meaningful patterns relevant to soil water content dynamics. The residual component, which
233 lacks systematic structure, was excluded from further analysis. STL was configured with a cycle length
234 of 180 days, representing the semi-annual wet–dry phases at the study sites. A LOESS smoother with
235 a 90-day window was then applied to the de-seasonalized series to extract the trend component. This



236 configuration was chosen to capture gradual, long-term changes in the climatic variables while reducing
237 short-term fluctuations. Note, that near the ends of the time series the absence of future values causes
238 the smoothing window to become asymmetric. As a result, the estimated trend becomes more sensitive
239 to recent variability. This limitation does not affect the outcome of the analysis, as both the input
240 features and the target variable (water content) are equally influenced by it. Each of the extracted
241 seasonal and trend components from P, PET, and WB was included as input to the neural network
242 alongside the original raw values. This allowed the model to learn structured seasonal behaviour—such
243 as distinguishing the rising phase of spring wetting up of the soil profile from the declining phase of a
244 summer dry-down—and to account for long-term shifts, such as gradual drying or changes in mean
245 climate conditions.

246 *2.2.2 Wetness classification*

247 In addition to the continuous climate-related features, a categorical input was included to describe the
248 soil's moisture condition as either 'dry', 'moderate', or 'wet'. These categories were defined using the
249 soil water content time series from the training site, with thresholds based on quantiles of the full
250 distribution. Specifically, values below the 30th percentile were labelled as 'dry', between the 30th and
251 70th percentiles as 'moderate', and above the 70th percentile as 'wet'. These categories were then
252 encoded numerically prior to modelling, using values of 30 for 'dry', 20 for 'moderate', and 1 for 'wet'.
253 This encoding allowed the categorical feature to be treated as ordinal variable and integrated into the
254 neural network input layer alongside the other features. There are two reasons to include this feature.
255 Firstly, the soil's current moisture condition can strongly influence its response to P and PET (Western
256 & Grayson, 1998). For example, under dry conditions, more water can be absorbed by the soil due to
257 its high storage capacity. In contrast, when soils are already wet or near saturation, infiltration capacity
258 is reduced, and additional rainfall is more likely to result in runoff (Tromp-van Meerveld & McDonnell,
259 2006; Zehe & Blöschl, 2004). The second reason is the motivation to use remote sensing data in similar
260 follow up studies, which are not yet accurate enough for modelling purposes but allow a general
261 classification of the wetness status. Because (i) corresponding information on soil matric potential



cannot be deduced at larger scale from remote sensing data and (ii) hysteresis in the soil water retention curve may lead to ambiguous thresholds, we focus here on soil water content measurements.

Another choice for the model framework that must be discussed is the choice of percentile thresholds. From a soil hydrological point of view, it would make sense that the thresholds defining the three classes ‘wet’, ‘moderate’ and ‘dry’ are chosen individually for each lysimeter (a wet clay soil may have very different water content values than a sandy soil). However, from a methodological point of view, we prefer to ensure that the model does not require a long time series to determine quantiles of soil water content data and that the model can be run solely based on the training site’s distribution. Accordingly, the same percentile thresholds, derived from the training site, were applied to label daily water content values at the prediction sites. Note, that the application of the same percentile thresholds for all sites is not relevant for the detection of changes in the soil water content response function. Very similar results will be obtained for a site-specific percentile definition as shown in the supplementary material (section S1). After constructing all the above features, each daily input to the model consisted of (i) raw climate variables (P, PET, and WB), (ii) the STL-derived seasonal and trend components for each of those variables (six variables), and (iii) the categorical moisture label. All ten features were aligned by date to ensure consistency across inputs. This combination of raw values, decomposed temporal signals, and qualitative soil condition provides the model with a detailed daily representation of both external climatic forcing and internal system state.

2.3 Neural Network Architecture and Training

To model daily volumetric soil water contents, a feed-forward neural network was implemented. The architecture consisted of three hidden layers: two dense layers with 12 neurons each using ReLU activation functions (Rectified Linear Unit), followed by a batch normalization layer, and a third dense layer with 6 neurons. ReLU was chosen for its ability to introduce non-linearity while maintaining computational efficiency and avoiding vanishing gradient problems during training (Lu et al., 2020; Montesinos López et al., 2022). Batch normalization was applied to stabilize learning by reducing internal covariate shift, which improves convergence speed and training stability (Montesinos López et



288 al., 2022). The output layer consisted of a single neuron with a linear activation function, which is
289 standard for continuous regression tasks such as predicting soil water content.

290 The network was trained using input features derived from daily observations at the reference lysimeter
291 at the Bad Lauchstädt site, covering the period 2015–2023. Prior to training, all continuous input
292 features were standardized to have a mean of zero and a standard deviation of one using z-score
293 normalization. The standardization parameters (mean and standard deviation) were computed solely
294 from the training dataset and applied unchanged to the validation sets at the training site, as well as to
295 the prediction sites, ensuring consistency across all data splits. The target variable, volumetric soil water
296 content, was preserved in its original physical units ($\text{m}^3 \text{m}^{-3}$), allowing for direct interpretation of the
297 model outputs and associated errors in hydrologically meaningful terms. The model was compiled with
298 the Adam optimizer, which adaptively adjusts learning rates and is widely used for its computational
299 efficiency and stable convergence. Mean squared error (MSE) was used as the loss function due to its
300 sensitivity to large deviations, making it suitable for continuous regression tasks. To monitor
301 generalization, 30% of the data was withheld as a validation set and excluded from updating the weights
302 between the nodes during training. The training procedure was initially set to proceed for a maximum
303 of 1000 epochs. To prevent overfitting, an early stopping criterion was implemented based on validation
304 loss. Specifically, training was terminated if no improvement in validation performance was observed
305 over a predefined number of consecutive epochs (patience threshold). The model parameters from the
306 epoch exhibiting the lowest validation loss were retained for final evaluation.

307 2.4 Testing the Neural Network

308 After training, the model was applied to the remaining 23 lysimeters across both Selhausen and Bad
309 Lauchstädt, none of which were included in the training phase. All test inputs were processed using the
310 same structure and normalization parameters derived from the training data. As outlined in Section 2.1,
311 the experimental setup includes four soil types, each installed with three replicates at both sites (see Fig.
312 3). While the lysimeters at the Bad Lauchstädt lysimeter station share the same climatic setting as the
313 training site, the lysimeters at Selhausen represent a more humid region. Accordingly, the raw data and
314 seasonal trend data of the Selhausen climate were used as input for the prediction of soil water content



in lysimeters located at Selhausen. This configuration allows to evaluate (i) whether the soil water content response function determined for the training remains valid for different climates and soil types, and (ii) to detect potential temporal or structural changes in soil hydraulic behaviour. The evaluation and classification procedures are described in the following two subsections.

2.5 Detection of Change in Soil Water Content Response Function Based on Error Metrics

As explained in the introduction, we use error metrics to detect changes in the soil water content response function. While we use the Nash-Sutcliffe -Efficiency (NSE, see eq. 5) as a general descriptor of model error, we investigate temporal changes in model performance based on the Mean Bias (MB) that was calculated on an annual basis from 2015 to 2023. This year-by-year assessment does not rely on predefined change points and enables the detection of gradual or abrupt shifts in model performance directly from the data. MB measures the average signed difference between predicted and observed values, providing an estimate of systematic overestimation or underestimation over time (Moriassi et al., 2007; Liu et al., 2011) and is defined as:

$$MB = \frac{1}{N} \sum_{i=1}^N (\hat{\theta}_i - \theta_i) \quad (1)$$

where $\hat{\theta}_i$ is the predicted volumetric water content at day i , θ_i is the corresponding observation and N defines the number of available observations–prediction pairs. Although, the calculation uses daily values, MB is aggregated over yearly intervals to produce a single value per year, capturing annual patterns in prediction bias. Volumetric water contents ($\text{m}^3 \text{ m}^{-3}$) were multiplied by 100 prior to calculation, and MB is therefore reported in percentage (%). Positive MB values indicate systematic overestimation by the model, while negative values reflect underestimation. The annual assessment of MB allowed us to evaluate whether the soil water content response function remains consistent across time or shows temporal dynamics.

To classify the soil water content dynamics with respect to the resilience after the extreme summer 2018, we check if the deviation of the predictions based on a stable response function (developed with the training data) changes over the years. When the deviation in the first year (2015; i.e., before the



drought) is different from the deviation in the year 2023, we consider that the soil water content response function has changed (it is still possible that the response function may recover in the future) and the soil water content dynamics is classified accordingly as ‘changed’. When the deviations at beginning and end are similar, but there was a period between 2018 and 2022 with a different deviation level, we conclude that the soil water content response function changed reversibly over time but recovered within the observation window and the lysimeter is classified as ‘resilient’. The soil water content response is considered as ‘stable’ when the deviations remain similar during the entire observation period. As threshold we chose 1.52%, that equals the 3-fold of the standard deviation of the nine yearly MB values computed for the training site. The classification of the time series was thus expressed formally as:

$$\text{‘changed’}: |MB_{2023} - MB_{2015}| > 1.52\% \quad (2)$$

$$\text{‘resilient’}: |MB_{2023} - MB_{2015}| \leq 1.52\% \wedge |MB_{20xx} - MB_{2015}| > 1.52\% \quad (3)$$

$$\text{‘stable’}: |MB_{2023} - MB_{2015}| \leq 1.52\% \wedge |MB_{20xx} - MB_{2015}| \leq 1.52\% \quad (4)$$

with the logical operator $\wedge$ and the mean bias of a specific year with MB_{20xx} , that shows the largest difference $|MB_{\text{year}} - MB_{2015}|$ for the time period between 2018 and 2022 (starting with dry year 2018).

As general information on the different response function, we calculated the NSE coefficient (Moriassi et al., 2007; Nash & Sutcliffe, 1970). The NSE is a standard metric for hydrological model skill, with $NSE = 1$ indicating perfect agreement and $NSE \leq 0$ stating that the model predictions are no better than using the mean of the observations. Mathematically, it is defined as:

$$NSE = 1 - \frac{\sum_{i=1}^N (\theta_i - \hat{\theta}_i)^2}{\sum_{i=1}^N (\theta_i - \bar{\theta})^2} \quad (5)$$

where $\hat{\theta}_i$ is the predicted volumetric water content at day i , θ_i is the corresponding observed value, $\bar{\theta}$ is the mean observed volumetric water content over the evaluation period, and N denotes the total number of valid data points used in the calculation. Following Moriassi et al. (2015), model performance was classified as very good for $NSE > 0.80$, good for $0.70 < NSE \leq 0.80$, satisfactory for $0.50 < NSE \leq 0.70$, and unsatisfactory for $NSE \leq 0.50$. Lower NSE values were interpreted as indicators of deviations from



366 the soil hydraulic behaviour represented in the training data, potentially due to differences in soil
367 properties or climate-induced structural changes.

368 2.6 Interpretation of Change in Response Function in Soil Physical Terms

369 The dynamics of MB (see above) was also used to assess changes in soil water retention curves
370 (SWRCs), which were plotted for each test lysimeter on a yearly basis. As stated in eq. (1), a positive
371 MB value corresponds to measured values that are smaller than the predictions. Because the predictions
372 are based on the model trained for a specific lysimeter, we expect for a positive MB that the water
373 content for the same environmental conditions (as manifested in the matric potential) is smaller in the
374 test lysimeter compared to the lysimeter used for training (SWRC is shifted to the left). Analogously,
375 for a consistently negative MB we expect that the test site retained more water at a given matric potential
376 than the training site and the SWRC is shifted to the right. For a 'resilient' soil, the soil water retention
377 curve will be shifted over time and will shift back close to the original position at the end of the
378 observation period. Finally, for a soil with 'changed' response function, the water retention curve is
379 drifting over time as well but without returning to its original position. In some cases, the temporal
380 evolution of MB may not exactly follow the apparent shift of the SWRC, as additional vertical or slope
381 changes could occur due to variations in porosity or pore-size distribution. These effects cannot be
382 identified within the current framework but may contribute to deviations between MB dynamics and
383 the apparent SWRC shift.

384 3. Results


385 Following the methodological framework described in Section 2.3-2.6, we present the results of model
386 predictions across the test lysimeters to assess the resilience of the soil water content response function
387 for the different lysimeters. We organize the section in four subsections according to the four different
388 origins of the soil in the lysimeters (see Fig.3b) to discuss effects of soil origin and climatic conditions
389 on the response function. In the last subsection (3.5) the results are summarized to allow direct
390 comparison of all 24 lysimeters. Note, that all model results presented below are based on the soil water
391 content classification ('wet', 'moderate', 'dry') as deduced from the lysimeter used for model training.



392 The corresponding figures using specific classification for each lysimeter are shown in supplementary
 393 information Fig. S4-S7.

394 3.1 Lysimeters with Same Soil as Used in Model Training (Dedelow Soils)

395 The neural network was trained to capture the soil water content response function of one lysimeter
 396 with sandy loam topsoil (Luvisol) extracted from Dedelow and translocated to the dry climatic region
 397 in Bad Lauchstädt (see Fig. S1 in supplementary material). The NSE of the training and validation of
 398 that specific lysimeter was very high with 0.91 indicating good model performance. The application of
 399 this response function to the other two lysimeters from Dedelow that were translocated to Bad
 400 Lauchstädt resulted in relatively high NSE values (0.79 and 0.84). However, the lower soil water
 401 contents observed during the summer of 2018 were not adequately captured as shown in Fig. 5a. More
 402 specifically, the time series show that predictions and observations matched closely in 2015, while after
 403 the dry summer of 2018 the model systematically overestimated water content in 2019 and 2020, before
 404 the agreement improved again towards the end of the period.



405



Figure 5 Analysis of soil water content dynamics (2015–2023) for a Dedelow-origin lysimeter tested at Bad Lauchstädt. Panel (a) shows the time series of observed (blue) and predicted (orange) water content, with close agreement in 2015, clear overestimation in 2019–2020 (predictions above observations), and improved agreement again towards the end of the period. Panel (b) presents the temporal evolution of mean bias (MB), remaining near zero until 2017, increasing to about 2–3 % in 2019–2020, and decreasing again to approximately zero in 2022. Such soil water content response was classified as ‘resilient’. Panel (c) displays soil water retention curves from the training site (grey) and from selected years representing different MB conditions, with low-MB years (2015, 2022) and high-MB years (2019, 2020). The curves are close to the training site in 2015, show a shift to lower water contents in 2019–2020, and in 2022 return to the training site data.

These changes are reflected in the MB development (Fig. 5b), with values increasing from near zero in 2015 to about 2–3% in 2019–2020 and then decreasing again towards 2022. The retention curves confirm this interpretation (Fig. 5c). The year with low MB (2015) produced a SWRC close to the measured curve of the training site, the years with high MB (2019–2020) were shifted to lower water contents, and the later year with reduced MB (2022) returned to the measured SWRC of the training site. Taken together, the time series, MB trend, and SWRCs show that the soil response was disturbed after 2018 but later recovered, defining this lysimeter as ‘resilient’. The same finding holds for the simulations for the lysimeters translocated to Selhausen (less dry climate) with high NSE between 0.80–0.82. This indicates, that for this coarse soil (i) the effect of changing climatic conditions was rather small (very good NSE classification for both sites) but (ii) that also these coarse textured topsoils do not show identical response to the extreme year but each lysimeter reacts slightly different, indicating slightly different structural properties.


3.2 Lysimeters with Soils Adapted to Climatic Conditions Similar with Those of the Model Training Site (Bad Lauchstädt soil)

The lysimeters filled with soil from Bad Lauchstädt (Chernozem) represent soils adapted to the climatic conditions under which the response function was calibrated. In case of dominant effect of climate on the soil water content response function, we could expect similar results as for the training lysimeter. For the soil remaining at the original site (Bad Lauchstädt), the model performance was very good (0.88–0.89). As shown for an example in Fig. 6a, the fit between observed and predicted water content was consistently close, with a tendency to slightly underestimate in the early years and to mildly overestimate after 2018, particularly in 2019–2020, before the agreement improved again in later years.



436 This is also manifested in the MB values that increased from slightly negative values in 2015 to about
437 +1.5% in 2019, before decreasing again towards zero (Fig. 6b). The plotted SWRCs support this
438 interpretation (Fig. 6c), with low MB years (2015 and 2016) showing a slight shift to higher water
439 contents relative to the measured SWRC of the training site, and high-MB years (2019 and 2020)
440 displaying a modest shift to lower water contents. Accordingly, the soil water content dynamics was
441 classified as ‘resilient’.

442 For the lysimeters transported from Bad Lauchstädt to Selhausen, the performance was more variable
443 (NSE ranging from 0.50 to 0.84) corresponding to satisfactory to very good classifications, reflecting
444 the stronger effect of the wetter climate. None of the three lysimeters who stayed in Bad Lauchstädt
445 were classified as ‘changed’ but two out of three showed a systematic shift and were classified as
446 ‘changed’ when translocated to Selhausen (see Table 1). In short, those examples show that the Bad
447 Lauchstädt soil remained resilient under unchanged climate at Bad Lauchstädt but changed under the
448 wetter climate at Selhausen.





450 **Figure 6** Analysis of soil water content dynamics (2015–2023) for a Bad Lauchstädt-origin soil lysimeter tested
 451 at Bad Lauchstädt. (a) Comparison of measured (blue) and simulated (orange) daily water content values,
 452 showing high agreement in the early years and temporary overestimation in 2019–2020. (b) Mean Bias (MB)
 453 started slightly negative in 2015, increased to about +1.5 % in 2019, and then decreased again towards 2022. (c)
 454 Soil water retention curves (SWRCs) from the training site (grey) and from the same replicate for selected years
 455 with low MB (2015, 2016) and high MB (2019, 2020) show close agreement in the early years and a shift to lower
 456 water contents in 2019–2020.

457 3.3 Lysimeters with Soils Adapted to Climatic Conditions comparable with Those 458 of the Model Training Site (Sauerbach soil)

459 The findings are similar for the silt loam (Cambisol) from Sauerbach, representing soils adapted to
 460 climatic conditions comparable to those in Bad Lauchstädt. As in case of the soil from Bad Lauchstädt,
 461 soils from Sauerbach show higher NSE values when translocated to Bad Lauchstädt (0.81–0.88, very
 462 good) compared to those transferred to the wetter climate in Selhausen (0.74–0.79, good). This reflects,
 463 that the soil water content response function in the drier climate is not the same as in the wetter climate.
 464 In one illustrative case, the observed water content initially showed wetter dynamics than predicted, but
 465 gradually converged toward the model predictions by 2023, indicating a possible structural adjustment
 466 over time (Fig. 7a).




Figure 7 Analysis of soil water content dynamics (2015–2023) for Sauerbach-origin lysimeter relocated to Selhausen (a) Comparison of observed and predicted daily volumetric water content ($NSE = 0.74$) After initial underestimation by the model, the observed and predicted values gradually converged, indicating a possible structural adjustment. (b) Temporal evolution of the mean bias (MB), which increased from about -5% in 2015 to values close to zero by 2019–2023, consistent with the improved match between observed and predicted values shown in panel (a). (c) Soil water retention curves (SWRCs) from the training site (grey) and from selected years with low MB (2015, 2016) and high MB (2020, 2022) illustrate the same trend, with early years showing higher water contents at a given matric potential and later years shifting towards to the training curve.

This development is also evident in the MB values (Fig. 7b), which started strongly negative (-5%) in 2015–2016 and steadily increased toward values close to zero by 2023, indicating a progressive reduction of underestimation. The corresponding SWRCs (Fig. 7c) confirm this trend, with curves from early years (2015, 2016) showing higher water contents at a given matric potential compared to the measured SWRC of the training site and later years (2021, 2023) shifting closer to the reference, suggesting a gradual adjustment of hydraulic behaviour. In the case of soils from Sauerbach, there was a difference in quantification of resilience with respect to the classification of the soil water content used as input variable: with the classification based on the training lysimeter (with sandy loam in the



484 topsoil), the soil water content dynamics was classified as ‘changed’ for all six lysimeters. But using
 485 the classification based on the soil water content statistics obtained for each lysimeter individually (see
 486 Fig. S6), the large water contents at the beginning were captured and only one lysimeter out of three
 487 was classified ‘changed’. Independent of the water content classification, all lysimeters translocated to
 488 Selhausen were classified as ‘changed’, exhibiting the strongest response to relocation of all soils.

489 3.4 Lysimeters with Soils Adapted to Climatic Conditions Different from Those of 490 the Model Training Site (Selhausen)

491 At last, we discuss the Selhausen silt loam (Luvisol), representing soils adapted to climatic conditions
 492 that were not included in the neural network training. The model performance was better for the
 493 replicates translocated to the drier Bad Lauchstädt climate ($NSE = 0.86\text{--}0.92$, very good), compared to
 494 slightly lower performance at their site of origin under humid Atlantic conditions ($0.76\text{--}0.86$, good to
 495 very good). The classification with respect to resilience helps to explain this, since Selhausen soils at
 496 their origin were mainly assigned to ‘stable’ or ‘resilient’ categories (see Table I), while the same soils
 497 translocated to Bad Lauchstädt showed a more variable pattern. This indicates, that the lower NSE at
 498 Selhausen does not represent a misfit of the model but reflects that the soils follow their own stable soil
 499 water content response function. One replicate at Selhausen ($NSE = 0.85$) reproduced the seasonal
 500 dynamics well, although differences between observed and predicted values remained visible in the wet
 501 season across several years (Fig. 8a). The MB shifted from negative values in the first years toward
 502 zero after 2019. Note, that an MB value of 0 does not mean that deviations disappeared, but that errors
 503 in wetter and drier phases compensated each other (Fig. 8b). The SWRCs were generally close to the
 504 training reference, but in later years small deviations appeared mainly at the saturated end (Fig. 8c).
 505 Overall, these changes remained below the assumed threshold, supporting a classification of the soil
 506 water content dynamics as ‘stable’.




Figure 8 Soil water content dynamics (2015–2023) for a Selhausen-origin lysimeter tested at Selhausen. (a) Observed (blue) and predicted (orange) water content show fair agreement, with underestimation of water contents in the wet season. (b) Mean Bias (MB) fluctuated from negative values in the early years to values close to zero after 2019, but these variations remained below the threshold for change. (c) Soil water retention curves (SWRCs) from the training site (grey) and from selected years with low MB (2015, 2016) and higher MB (2019, 2020) reflect these minor variations, with the 2019 curve showing the strongest deviation yet remaining close to the training reference, consistent with the stable classification. The apparent cutoff at the wet end in (a) arises from the use of absolute rather than normalized values during training, as discussed in the Supplementary Material (Text S1).

3.5 Comparison of All Lysimeters

The comparison of the soil water content dynamics of all lysimeters indicate, that climatic shifts between sites - particularly between the continental Bad Lauchstädt and Atlantic-influenced Selhausen - can significantly alter the hydraulic response of the soil, even when texture remains constant. In general, prediction performance at Selhausen was lower, likely because the model was trained under the drier climate of Bad Lauchstädt, and therefore, failed to fully capture the soil–water interactions



emerging under wetter conditions (Fig. 9). The broader NSE range observed at Selhausen location further suggests increased structural variability among replicates.





Figure 9 Spread of Nash–Sutcliffe Efficiency (NSE) values across different soil origins and test locations. Each symbol represents one lysimeter from a given origin (x-axis) evaluated at Bad Lauchstädt or Selhausen (indicated by colour). The results highlight the influence of climate–soil interactions on model performance. Notably, Bad Lauchstädt-origin soils exhibited strong performance at their origin but a wider and lower range when tested at Selhausen, reflecting increased structural variability or climate-induced divergence in hydraulic response.

With respect to resilience of the soil water content response function, we show the temporal evolution of the mean bias for all lysimeters in Fig. 10 and summarize the results in Table 1. In Table 1 we add the general classification type (‘stable’, resilient’ and ‘changed’) and calculate the average of 3 lysimeters (same material and same location) for the drift in mean bias value between the year 2015 and 2023 and the maximum deviation from year 2015 for the years between 2018 and 2022. The table shows that deviations from a ‘stable’ or ‘resilient’ response function mainly occur when soils from Dedelow, Bad Lauchstädt, and Sauerbach were translocated to Selhausen. Only in case of the soil from Selhausen, the response function remains ‘stable’. It seems, that the soil material ‘trained over decades’ to the wetter climate in Selhausen adapts better to the extreme summer 2018.



540
541 **Figure 10** Temporal evolution of Mean Bias (MB) for three representative lysimeter replicates, each classified
542 into one of three structural response categories: (a) ‘stable’, (b) ‘resilient’, and (c) ‘changed’. Thick dashed lines
543 indicate the mean of the MB trend across all lysimeters within each classification group, with sample size (n)
544 specified in the legend. Shaded areas represent ± 1 standard deviation. Thin grey lines show individual MB
545 trajectories of the remaining lysimeters in each group. Highlighted blue lines depict selected replicates
546 originating from and/or tested at distinct sites: (a) BL $\rightarrow$ BL (soil material from Bad Lauchstädt tested at its
547 origin), (b) DD $\rightarrow$ BL (soil material from Dedelow tested at Bad Lauchstädt), and (c) BL $\rightarrow$ SE (soil material
548 from Bad Lauchstädt tested at Selhausen). These examples illustrate contrasting temporal patterns in structural
549 response, ranging from sustained stability to progressive divergence from the trained site dynamics.

550 **Table 1:** Resilience of soil water content response function for the four soil materials translocated to Bad
551 Lauchstädt and Selhausen. The ‘type’ describes the class of response function of the individual lysimeters (S for
552 ‘stable’, R for ‘resilient’ and C for ‘changed’). The ‘drift’ is the average value $|\text{MB}_{2023} - \text{MB}_{2015}|$ of the three
553 lysimeters with the difference in Mean Bias (MB) between years 2023 and 2015. The ‘amplitude’ is the maximum



554 difference of the Mean Bias between the first year (2015) and the years between 2018 and 2022 (denoted as year
 555 20xx).

	Located at Bad Lauchstädt			Located at Selhausen		
	Type	Drift	Amplitude	Type	Drift	Amplitude
Dedelow	S,R,C	1.09	1.76	R,C,C	1.72	2.11
Bad Lauchstädt	S,S,R	0.94	1.36	R,C,C	2.15	2.99
Sauerbach	C,C,C	2.17	3.73	C,C,C	3.38	4.24
Selhausen	S,R,C	0.96	2.24	S,S,R	0.47	1.78

556

557 4. Discussion

558 The results presented in Section 3 demonstrate that the model can reproduce soil water content dynamics
 559 reliably under stable conditions (as indicated by high NSE-values), but it exhibits limitations when soils
 560 undergo structural changes or are exposed to a different climate. Several soils showed a shift in the wet
 561 range, indicating that differences in soil water content response cannot be explained by texture alone
 562 but reflect the combined effects of climatic conditions and structural evolution. Based on these findings,
 563 the following discussion evaluates how assumptions of static hydraulic behaviour and response function
 564 affect model performance, examines the role of NSE and MB in identifying evolving system dynamics,
 565 and reflects on the broader implications for long-term modelling and soil water content monitoring.

566 4.1 Soil–Climate Interactions as Drivers of Hydraulic Response Function

567 The predictive success of data-driven models depends not only on the physical properties of soils but
 568 also on the climatic context in which those properties developed and continue to function. The present
 569 study shows that soils exhibit the most consistent replicate behaviour when evaluated under climate
 570 conditions similar to those of their origin, where gradual climatic changes over time have allowed their
 571 structure to adjust naturally. When exposed to faster or contrasting climatic shifts, as in translocated
 572 settings, the soil response becomes less predictable and less stable. This can be shown using the table
 573 S1 in the supplementary material file, which lists the average trends (difference in MB between year
 574 2023 and 2015) and amplitudes (difference in MB between 2015 and the dry years) for three lysimeter
 575 replicates: only for the three lysimeters at the original locations Bad Lauchstädt or Selhausen), both



576 drift and amplitude were below the stability threshold of 1.52% and can be classified as ‘stable’ as group
 577 of lysimeters.

578 This suggests that the structure and function of the soil system cannot be meaningfully decoupled from
 579 its climatic history. Soils may develop pore arrangements, aggregation patterns, and, as a consequence,
 580 moisture retention characteristics that reflect long-term adaptation to local hydrological regimes. When
 581 these soils are translocated to environments with contrasting P and atmospheric demand (PET), their
 582 hydraulic response can shift in ways that are not captured by static texture-based estimates of soil
 583 hydraulic properties.

584 Such context-dependent behaviour highlights the limitation of the common assumption that soils with
 585 the same texture will show comparable retention across regions, an assumption often made in the
 586 absence of better descriptors. Experimental evidence collected under natural conditions also indicates
 587 that this description is oversimplified (Hannes et al., 2016; Robinson et al., 2016; Aqel et al., 2024). In
 588 our case, even soils with similar textural composition exhibited different levels of model agreement
 589 depending on climate, highlighting that physical similarity (e.g. soil texture) does not guarantee
 590 functional equivalence in retention. For example, Selhausen-origin soils achieved higher NSE values
 591 when translocated to Bad Lauchstädt, likely because the model was trained under similar dry climatic
 592 conditions. However, classification results showed, that these soils retained greater structural stability
 593 at their origin, suggesting that predictive success under familiar climatic forcing does not necessarily
 594 imply hydraulic consistency. After the 2018 drought, the Selhausen soils translocated to Bad Lauchstädt
 595 converged toward similar dynamics across replicates, with MB stabilizing close to zero, indicating that
 596 their response functions adjusted consistently to the drier climate (see Fig. S2 in the supplementary
 597 material). However, a clear carry-over effect was observed: soil water in the upper 10 cm was not fully
 598 replenished during the wet phase of autumn and winter 2019 and only reached comparable, though
 599 slightly lower, values in winter 2020. This persistent deficit points to a structural legacy of the drought,
 600 where reduced pore connectivity and altered aggregation limited subsequent rewetting. A comparable
 601 multi-year legacy across the full soil column was reported in the TERENO-SOILCan lysimeter network
 602 by Groh et al. (2020).



603 All mentioned points underscore the importance of including very broad range of climatic forcing in
 604 the assessment of soil model transferability, as demonstrated by Groh et al. 2022. Our results also
 605 suggest that future efforts to generalize hydrological models should consider training under a range of
 606 climatic conditions to capture the full expression of soil–climate interactions, rather than relying on a
 607 single static representation. From a process-based perspective, these findings reflect that climate does
 608 not simply modulate soil water content inputs but actively shapes the retention and release behaviour
 609 of the soil pore network through structural evolution or breakdown. While management practices across
 610 sites were similar, minor differences in tillage and fertilization cannot be completely excluded and may
 611 have influenced soil structure and water retention. Nonetheless, the dominant control remains climatic
 612 forcing, which makes this consideration particularly relevant for climate-change experiments: models
 613 calibrated under past climatic conditions may not remain valid under the rapid climatic shifts projected
 614 for the coming decades. Neglecting this evolving soil–climate feedback could lead to substantial
 615 underestimation of future changes in soil hydraulic behaviour and associated ecosystem responses

616 4.2 High Predictive Performance Can Mask System Evolution

617 Although, several lysimeters achieved high predictive performance as expressed by high NSE values
 618 (Fig. 9), systematic trends in MB over time suggest that the underlying retention behaviour and soil
 619 water content response function may have shifted (Fig. 10 and Table 1). This was most apparent in
 620 Dedelow soils translocated to Bad Lauchstädt, where the model maintained high NSE values, but the
 621 MB increased across years (see Fig. S3). The corresponding shifts in the soil water retention curves
 622 confirmed a gradual change in how the soil retained water, despite the model continuing to predict
 623 moisture levels accurately.

624 This suggests that local structural changes can occur without immediate deterioration in model fit. The
 625 predictive framework remained effective in capturing the general moisture dynamics, but the
 626 relationship between matric potential and water content was no longer consistent with that observed
 627 during training. These findings highlight that high model accuracy does not guarantee stability in the
 628 hydraulic characteristic, particularly under changing environmental conditions. Identifying such
 629 divergence early is critical for maintaining reliable predictions in long-term monitoring.



4.3 Implications for Monitoring, Remote Sensing, and Soil Health

The classification outcomes across all lysimeters highlight the role of site memory and structural resilience in maintaining hydraulic behaviour under climatic stress. Soils assessed at their origin were more frequently classified as ‘stable’ or ‘resilient’ (e.g., Selhausen at Selhausen), while those translocated to different locations were more likely to be classified as ‘changed’ (e.g., Sauerbach at Bad Lauchstädt). These patterns indicate that soil structure, once adapted to specific climate regimes, may lose its functional integrity when exposed to new conditions. The presented methods allow us to detect emerging structural shifts that may be relevant for soil health assessment and could be used as indicator for deteriorated soil health status.

This has direct implications for long-term monitoring and remote sensing. Our model framework - by avoiding reliance on matric potential data and instead using moisture state categories and decomposed climatic features - is compatible with satellite-derived products. As remote sensing missions increasingly provide continuous global soil water content estimates, the proposed framework could be adapted for large-scale assessment of soil system stability. Furthermore, under scenarios of future climate change, where shifts in precipitation patterns and evaporative demand are expected, models trained on historical data may become progressively outdated. The presented residual-based approach (quantifying MB) enables early detection of such divergence, offering a method for identifying when re-training or reparameterization is needed to maintain predictive reliability under non-stationary conditions.

5. Summary and conclusions

The temporal variations in the water content of the topsoil define the amount of plant available water and oxygen supply, affecting ecosystem functions and soil health status. Reliable information on soil water content dynamics in response to atmospheric conditions is thus essential to detect and mitigate critical conditions. This response depends on soil hydraulic properties that are traditionally characterized by a time-invariant and unambiguous relationship between matric potential, water content, and hydraulic conductivity as deduced from small-scale lab experiments. In this study, we



656 developed and applied a feed-forward neural network combined with seasonal trend analysis of climatic
657 time series to quantify the soil water content response function after an extreme drought in summer
658 2018 in Germany. By analysing the time series of topsoil water content measured at two lysimeter
659 stations of the TERENO SOILCan network, we summarize the conclusion on the soil water content
660 response function as follows:

- 661 • 50% of the lysimeters showed changes in soil water content dynamics after the dry summer
662 2018.
- 663 • The other half showed a resilient behaviour, and the soil water content response function was
664 not permanently changed.
- 665 • The changes in soil water content response function were manifested as (i) temporal trends in
666 prediction error (mean bias) and (ii) shifts in the soil water characteristics function.
- 667 • The soil water content response function is adapted to climatic conditions as manifested by (i)
668 smallest changes in lysimeters that were not translocated and (ii) decreased model performance
669 for applications of a response function that was determined for another climate
- 670 • Good model performance as expressed by high Nash-Shutcliff-Efficiency values does not
671 correspond to stable soil water content response function that was only detected by temporal
672 trends in error metrics

673 The study revealed that extreme climatic events can permanently change the soil hydraulic properties,
674 but the lack of resilience depends on the soil and the climatic conditions. We argue that the response
675 depends on the range of climatic conditions experienced in the past that allowed adaptation of soil
676 structural properties. Because the presented model framework (i) does not aim to predict successfully
677 time series in water content and (ii) does only require categorical water content information ('stable',
678 'resilient', 'changed'), it can be applied to larger scale using remote sensing data that do not provide
679 accurate soil water content values but reliable trends, enabling to detect changes in hydraulic behaviour
680 at the ecosystem scale.



681 Author contributions

682 NA, AC, and PL designed the study. NA and PL conducted the research. NA developed the model and
683 wrote the code. NA and PL prepared the manuscript with contributions of all co-authors. JG and RG
684 provided and quality-controlled the lysimeter data.

685 Acknowledgements

686 We acknowledge the support of the TERENO–SOILCan network that were funded by the Helmholtz
687 Association (HGF) and the Federal Ministry of Education and Research (BMBF). We thank Ferdinand
688 Engels, Robert Lüdtke, Werner Küpper, Ines Merbach, Philipp Meulendick, and Syliva Schmögner for
689 the instrument operation and data processing at both sites. We also thank Hans-Jörg Vogel for his
690 constructive feedback and insightful comments on the manuscript. Nedal Aqel acknowledges the
691 utilization of ChatGPT to enhance coherence within certain sections of the paper.

692 Financial support

693 This research is part of the project AI4SoilHealth of the European Union’s Horizon research and
694 innovation programme (grant agreement No. 101086179). This work has received funding from the
695 Swiss State Secretariat for Education, Research and Innovation (SERI).

696



References

- Allen, R. G., Pereira, L. S., Raes, D., and Smith, M.: Crop evapotranspiration – Guidelines for computing crop water requirements, FAO Irrigation and Drainage Paper 56, Food and Agriculture Organization of the United Nations, Rome, 300 pp., 2006.
- Aqel, N., Reusser, L., Margreth, S., Carminati, A., and Lehmann, P.: Prediction of hysteretic matric potential dynamics using artificial intelligence: application of autoencoder neural networks, *Geosci. Model Dev.*, 17, 6949–6966, <https://doi.org/10.5194/gmd-17-6949-2024>, 2024.
- Blöschl, G., Bierkens, M. F. P., Chambel, A., Cudennec, C., Destouni, G., Fiori, A., Kirchner, J. W., McDonnell, J. J., Savenije, H. H. G., Sivapalan, M., Stumpp, C., Toth, E., Volpi, E., Carr, G., Lupton, C., Salinas, J., Széles, B., Viglione, A., Aksoy, H., and Zhang, Y.: Twenty-three unsolved problems in hydrology (UPH) – a community perspective, *Hydrol. Sci. J.*, 64, 1141–1158, <https://doi.org/10.1080/02626667.2019.1620507>, 2019.
- Boergens, E., Güntner, A., Sips, M., Schwatke, C., and Dobsław, H.: Interannual variations of terrestrial water storage in the East African Rift region, *Hydrol. Earth Syst. Sci.*, 28, 4733–4754, <https://doi.org/10.5194/hess-28-4733-2024>, 2024.
- Bogena, H. R., Huisman, J. A., Güntner, A., Hübner, C., Kusche, J., Jonard, F., Vey, S., and Vereecken, H.: Emerging methods for noninvasive sensing of soil moisture dynamics from field to catchment scale: a review, *WIREs Water*, 2, 635–647, <https://doi.org/10.1002/wat2.1097>, 2015.
- Brocca, L., Melone, F., Moramarco, T., Wagner, W., Naeimi, V., Bartalis, Z., and Hasenauer, S.: Improving runoff prediction through the assimilation of the ASCAT soil moisture product, *Hydrol. Earth Syst. Sci.*, 14, 1881–1893, <https://doi.org/10.5194/hess-14-1881-2010>, 2010.
- Cleveland, R. B., Cleveland, W. S., McRae, J. E., and Terpenning, I.: STL: A seasonal-trend decomposition procedure based on Loess, *J. Off. Stat.*, 6, 3–73, 1990.
- Detty, J. M. and McGuire, K. J.: Threshold changes in storm runoff generation at a till-mantled headwater catchment, *Water Resour. Res.*, 46, W07525, <https://doi.org/10.1029/2009WR008102>, 2010.
- Deutscher Wetterdienst (DWD) Climate Data Center (CDC): Historical daily precipitation data, station Leipzig/Halle (ID 2932), available at: <https://www.dwd.de/cdc> (last access: 28 August 2025), 2025.
- Ehrhardt, A., Groh, J., and Gerke, H. H.: Effects of different climatic conditions on soil water storage patterns, *Hydrol. Earth Syst. Sci.*, 29, 313–334, <https://doi.org/10.5194/hess-29-313-2025>, 2025.
- Fatichi, S., Or, D., Walko, R., Vereecken, H., Young, M. H., Ghezzehei, T. A., Hengl, T., Kollet, S., Agam, N., and Avisar, R.: Soil structure is an important omission in Earth system models, *Nat. Commun.*, 11, 522, <https://doi.org/10.1038/s41467-020-14411-z>, 2020.
- Fu, Z., Hu, W., Beare, M., Thomas, S., Carrick, S., Dando, J., Langer, S., Müller, K., Baird, D., and Lilburne, L.: Land use effects on soil hydraulic properties and the contribution of soil organic carbon, *J. Hydrol.*, 602, 126741, <https://doi.org/10.1016/j.jhydrol.2021.126741>, 2021.



- 737 Groh, J., Diamantopoulos, E., Duan, X., Ewert, F., Heinlein, F., Herbst, M., Holbak, M., Kamali, B.,
738 Kersebaum, K.-C., Kuhnert, M., Nendel, C., Priesack, E., Steidl, J., Sommer, M., Pütz, T.,
739 Vanderborght, J., Vereecken, H., Wallor, E., Weber, T. K. D., and Gerke, H. H.: Same soil,
740 different climate: crop model intercomparison on translocated lysimeters, *Vadose Zone J.*, 21,
741 e20202, <https://doi.org/10.1002/vzj2.20202>, 2022.
- 742 Groh, J., Vanderborght, J., Pütz, T., Vogel, H.-J., Gründling, R., Rupp, H., Rahmati, M., Sommer,
743 M., Vereecken, H., and Gerke, H. H.: Responses of soil water storage and crop water use
744 efficiency to changing climatic conditions: a lysimeter-based space-for-time approach, *Hydrol.*
745 *Earth Syst. Sci.*, 24, 1211–1225, <https://doi.org/10.5194/hess-24-1211-2020>, 2020.
- 746 Hannes, M., Wollschläger, U., Wöhling, T., and Vogel, H.-J.: Revisiting hydraulic hysteresis based
747 on long-term monitoring of hydraulic states in lysimeters, *Water Resour. Res.*, 52, 3847–3865,
748 <https://doi.org/10.1002/2015WR018319>, 2016.
- 749 Hari, V., Rakovec, O., Markonis, Y., Hanel, M., and Kumar, R.: Increased future occurrences of
750 the exceptional 2018–2019 Central European drought under global warming, *Sci. Rep.*, 10,
751 12207, <https://doi.org/10.1038/s41598-020-68872-9>, 2020.
- 752 Herbrich, M. and Gerke, H. H.: Scales of water retention dynamics observed in eroded Luvisols
753 from an arable postglacial soil landscape, *Vadose Zone J.*, 16, 1–12,
754 <https://doi.org/10.2136/vzj2017.01.0003>, 2017.
- 755 Hrachowitz, M., Savenije, H. H. G., Blöschl, G., McDonnell, J. J., Sivapalan, M., Pomeroy, J. W.,
756 Arheimer, B., Blume, T., Clark, M. P., Ehret, U., Fenicia, F., Freer, J. E., Gelfan, A., Gupta, H. V.,
757 Hughes, D. A., Hut, R. W., Montanari, A., Pande, S., Tetzlaff, D., and Cudennec, C.: A decade of
758 Predictions in Ungauged Basins (PUB) – a review, *Hydrol. Sci. J.*, 58, 1198–1255,
759 <https://doi.org/10.1080/02626667.2013.803183>, 2013.
- 760 Jarvis, N. J., Beven, K. J., Larsbo, M., van Genuchten, M. T., Vereecken, H., and Vogel, H.-J.:
761 Interactions between soil structure dynamics, hydrological processes and organic matter
762 cycling: a new soil–crop model, *Eur. J. Soil Sci.*, 75, e13455, <https://doi.org/10.1111/ejss.13455>,
763 2024.
- 764 Kratzert, F., Klotz, D., Shalev, G., Klambauer, G., Hochreiter, S., and Nearing, G. S.: Towards
765 learning universal, regional, and local hydrological behaviours via machine learning applied to
766 large-sample datasets, *Hydrol. Earth Syst. Sci.*, 23, 5089–5110, [https://doi.org/10.5194/hess-](https://doi.org/10.5194/hess-23-5089-2019)
767 23-5089-2019, 2019.
- 768 Lehmann, P., Bickel, S., Wei, Z., and Or, D.: Physical constraints for improved soil hydraulic
769 parameter estimation by pedotransfer functions, *Water Resour. Res.*, 56, e2019WR025963,
770 <https://doi.org/10.1029/2019WR025963>, 2020.
- 771 Liu, J., Hughes, D., Rahmani, F., Lawson, K., and Shen, C.: Evaluating a global soil moisture
772 dataset from a multitask model (GSM3 v1.0) with potential applications for crop threats,
773 *Geosci. Model Dev.*, 16, 1553–1567, <https://doi.org/10.5194/gmd-16-1553-2023>, 2023.
- 774 Liu, J., Rahmani, F., Lawson, K., and Shen, C.: A multiscale deep learning model for soil moisture
775 integrating satellite and in situ data, *Geophys. Res. Lett.*, 49, e2021GL096847,
776 <https://doi.org/10.1029/2021GL096847>, 2022.
- 777 Liu, Y. Y., Parinussa, R. M., Dorigo, W. A., De Jeu, R. A. M., Wagner, W., van Dijk, A. I. J. M.,
778 McCabe, M. F., and Evans, J. P.: Developing an improved soil moisture dataset by blending



- 779 passive and active microwave satellite-based retrievals, *Hydrol. Earth Syst. Sci.*, 15, 425–436,
 780 <https://doi.org/10.5194/hess-15-425-2011>, 2011.
- 781 Lu, L., Shin, Y., Su, Y., and Karniadakis, G. E.: Dying ReLU and initialization: theory and numerical
 782 examples, *Commun. Comput. Phys.*, 28, 1671–1706, [https://doi.org/10.4208/cicp.OA-2020-](https://doi.org/10.4208/cicp.OA-2020-0165)
 783 0165, 2020.
- 784 Melsen, L. A. and Guse, B.: Climate change impacts model parameter sensitivity – implications
 785 for calibration strategy and model diagnostic evaluation, *Hydrol. Earth Syst. Sci.*, 25, 1307–
 786 1332, <https://doi.org/10.5194/hess-25-1307-2021>, 2021.
- 787 Milly, P. C. D., Betancourt, J., Falkenmark, M., Hirsch, R. M., Kundzewicz, Z. W., Lettenmaier, D.
 788 P., and Stouffer, R. J.: Stationarity is dead: whither water management?, *Science*, 319, 573–
 789 574, <https://doi.org/10.1126/science.1151915>, 2008.
- 790 Montanari, A., Young, G., Savenije, H. H. G., Hughes, D., Wagener, T., Ren, L.-L., Koutsoyiannis,
 791 D., Cudennec, C., Toth, E., Grimaldi, S., Blöschl, G., Sivapalan, M., Beven, K., Gupta, H., Hipsey,
 792 B., Schaefli, B., Arheimer, B., Boegh, E., Schymanski, S. J., and Belyaev, V.: “Panta Rhei—
 793 Everything Flows”: change in hydrology and society—The IAHS Scientific Decade 2013–2022,
 794 *Hydrol. Sci. J.*, 58, 1256–1275, <https://doi.org/10.1080/02626667.2013.809088>, 2013.
- 795 Montesinos López, O. A., Montesinos López, A., and Crossa, J.: Fundamentals of artificial neural
 796 networks and deep learning, in: *Multivariate Statistical Machine Learning Methods for*
 797 *Genomic Prediction*, Springer International Publishing, Cham, 379–425,
 798 https://doi.org/10.1007/978-3-030-89010-0_10, 2022.
- 799 Moriasi, D. N., Arnold, J. G., Van Liew, M. W., Bingner, R. L., Harmel, R. D., and Veith, T. L.:
 800 Model evaluation guidelines for systematic quantification of accuracy in watershed
 801 simulations, *Trans. ASABE*, 50, 885–900, <https://doi.org/10.13031/2013.23153>, 2007.
- 802 Moriasi, D. N., Gitau, M. W., Pai, N., and Daggupati, P.: Hydrologic and water quality models:
 803 Performance measures and evaluation criteria, *Trans. ASABE*, 58, 1763–1785, 2015.
 804 <https://web.ics.purdue.edu/~mggitau/pdf/Moriasi%20et%20al%202015.pdf>
- 805 Mosavi, A., Ozturk, P., and Chau, K.-W.: Flood prediction using machine learning models:
 806 literature review, *Water*, 10, 1536, <https://doi.org/10.3390/w10111536>, 2018.
- 807 Nash, J. E. and Sutcliffe, J. V.: River flow forecasting through conceptual models part I — A
 808 discussion of principles, *J. Hydrol.*, 10, 282–290, [https://doi.org/10.1016/0022-1694\(70\)90255-](https://doi.org/10.1016/0022-1694(70)90255-6)
 809 6, 1970.
- 810 Nearing, G. S., Kratzert, F., Sampson, A. K., Pelissier, C. S., Klotz, D., Frame, J. M., Prieto, C., and
 811 Gupta, H. V.: What role does hydrological science play in the age of machine learning?, *Water*
 812 *Resour. Res.*, 57, e2020WR028091, <https://doi.org/10.1029/2020WR028091>, 2021.
- 813 O., S. and Orth, R.: Global soil moisture data derived through machine learning trained with in-
 814 situ measurements, *Sci. Data*, 8, 170, <https://doi.org/10.1038/s41597-021-00964-1>, 2021.
- 815 Pütz, T., Kiese, R., Wollschläger, U., Groh, J., Rupp, H., Zacharias, S., Priesack, E., Gerke, H. H.,
 816 Gasche, R., Bens, O., Borg, E., Baessler, C., Kaiser, K., Herbrich, M., Munch, J.-C., Sommer, M.,
 817 Vogel, H.-J., Vanderborght, J., and Vereecken, H.: TERENO-SOILCan: a lysimeter-network in
 818 Germany observing soil processes and plant diversity influenced by climate change, *Environ.*
 819 *Earth Sci.*, 75, 1242, <https://doi.org/10.1007/s12665-016-6031-5>, 2016.



- 820 Reichstein, M., Camps-Valls, G., Stevens, B., Jung, M., Denzler, J., Carvalhais, N., and Prabhat:
821 Deep learning and process understanding for data-driven Earth system science, *Nature*, 566,
822 195–204, <https://doi.org/10.1038/s41586-019-0912-1>, 2019.
- 823 Robinson, D. A., Jones, S. B., Lebron, I., Reinsch, S., Domínguez, M. T., Smith, A. R., Jones, D. L.,
824 Marshall, M. R., and Emmett, B. A.: Experimental evidence for drought induced alternative
825 stable states of soil moisture, *Sci. Rep.*, 6, 20018, <https://doi.org/10.1038/srep20018>, 2016.
- 826 Robinson, D. A., Nemes, A., Reinsch, S., Raddbourne, A., Bentley, L., and Keith, A. M.: Global
827 meta-analysis of soil hydraulic properties on the same soils with differing land use, *Sci. Total*
828 *Environ.*, 852, 158506, <https://doi.org/10.1016/j.scitotenv.2022.158506>, 2022.
- 829 Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., Orlowsky, B., and
830 Teuling, A. J.: Investigating soil moisture–climate interactions in a changing climate: a review,
831 *Earth-Sci. Rev.*, 99, 125–161, <https://doi.org/10.1016/j.earscirev.2010.02.004>, 2010.
- 832 Shen, C., Laloy, E., Elshorbagy, A., Albert, A., Bales, J., Chang, F.-J., Ganguly, S., Hsu, K.-L., Kifer,
833 D., Fang, Z., Fang, K., Li, D., Li, X., and Tsai, W.-P.: HESS Opinions: Incubating deep-learning-
834 powered hydrologic science advances as a community, *Hydrol. Earth Syst. Sci.*, 22, 5639–5656,
835 <https://doi.org/10.5194/hess-22-5639-2018>, 2018.
- 836 Sun, W., Zhou, S., Yu, B., Zhang, Y., Keenan, T., and Fu, B.: Soil moisture–atmosphere
837 interactions drive terrestrial carbon–water trade-offs, *Commun. Earth Environ.*, 6, 169,
838 <https://doi.org/10.1038/s43247-025-02145-z>, 2025.
- 839 Tromp-van Meerveld, H. J. and McDonnell, J. J.: Threshold relations in subsurface stormflow: 1.
840 A 147-storm analysis of the Panola hillslope, *Water Resour. Res.*, 42, W02410,
841 <https://doi.org/10.1029/2004WR003778>, 2006.
- 842 Uber, M., Vandervaere, J.-P., Zin, I., Braud, I., Heistermann, M., Legoût, C., Molinié, G., and
843 Nord, G.: How does initial soil moisture influence the hydrological response? A case study from
844 southern France, *Hydrol. Earth Syst. Sci.*, 22, 6127–6146, <https://doi.org/10.5194/hess-22-6127-2018>, 2018.
- 846 Vaze, J., Post, D. A., Chiew, F. H. S., Perraud, J.-M., Viney, N. R., and Teng, J.: Climate non-
847 stationarity – validity of calibrated rainfall–runoff models for use in climate change studies, *J.*
848 *Hydrol.*, 394, 447–457, <https://doi.org/10.1016/j.jhydrol.2010.09.018>, 2010.
- 849 Vereecken, H., Amelung, W., Bauke, S. L., Bogen, H., Brüggemann, N., Montzka, C.,
850 Vanderborght, J., Bechtold, M., Blöschl, G., Carminati, A., Javaux, M., Konings, A. G., Kusche, J.,
851 Neuweiler, I., Or, D., Steele-Dunne, S., Verhoef, A., Young, M., and Zhang, Y.: Soil hydrology in
852 the Earth system, *Nat. Rev. Earth Environ.*, 3, 573–587, <https://doi.org/10.1038/s43017-022-00324-6>, 2022.
- 854 Vereecken, H., Huisman, J. A., Bogen, H., Vanderborght, J., Vrugt, J. A., and Hopmans, J. W.:
855 On the value of soil moisture measurements in vadose zone hydrology: a review, *Water*
856 *Resour. Res.*, 44, W00D06, <https://doi.org/10.1029/2008WR006829>, 2008.
- 857 Wankmüller, F. J. P., Delval, L., Lehmann, P., Baur, M. J., Cecere, A., Wolf, S., Or, D., Javaux, M.,
858 and Carminati, A.: Global influence of soil texture on ecosystem water limitation, *Nature*, 635,
859 631–638, <https://doi.org/10.1038/s41586-024-08089-2>, 2024.



860 Western, A. W. and Grayson, R. B.: The Tarrawarra data set: soil moisture patterns, soil
861 characteristics, and hydrological flux measurements, *Water Resour. Res.*, 34, 2765–2768,
862 <https://doi.org/10.1029/98WR01833>, 1998.

863 Xoplaki, E., Ellsäßer, F., Grieger, J., Nissen, K. M., Pinto, J. G., Augenstein, M., Chen, T.-C., and
864 Wolf, F.: Compound events in Germany in 2018: drivers and case studies, *Nat. Hazards Earth*
865 *Syst. Sci.*, 25, 541–564, <https://doi.org/10.5194/nhess-25-541-2025>, 2025.

866 Zehe, E. and Blöschl, G.: Predictability of hydrologic response at the plot and catchment scales:
867 role of initial conditions, *Water Resour. Res.*, 40, W10240,
868 <https://doi.org/10.1029/2003WR002869>, 2004.

869 Šimůnek, J., van Genuchten, M. T., and Šejna, M.: Recent developments and applications of the
870 HYDRUS computer software packages, *Vadose Zone J.*, 15, 1–25,
871 <https://doi.org/10.2136/vzj2016.04.0033>, 2016.

872