

Discussion Paper | Discussion Paper | Discussion Paper

11, 28797–28849, 2011

D. Zyryanov et al.

D. Zyryanov¹, G. Foret¹, M. Eremenko¹, M. Beekmann¹, J.-P. Cammas²,
M. D'Isidoro^{3,12}, H. Elbern⁴, J. Flemming⁵, E. Friese⁴, I. Kioutsoutis⁹,
A. Maurizi³, D. Melas⁹, F. Meleux⁶, L. Menut⁷, P. Moinat⁸, V.-H. Peuch⁸,
A. Poupkou⁹, M. Razingar⁵, M. Schultz¹⁰, O. Stein¹⁰, A. M. Suttie⁵,
A. Valdebenito¹¹, C. Zerefos⁹, G. Dufour¹, G. Bergametti¹, and J.-M. Flaud¹

⁷Laboratoire de Météorologie Dynamique, IPSL, Ecole Polytechnique, Palaiseau, France

Interactive Discussion



**3-D evaluation of
tropospheric ozone
simulations**

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

⁸ Météo-France, Centre National de Recherche Météorologique, Toulouse, France⁹ Laboratory of Atmospheric Physics, Aristotle University of Thessaloniki, Greece¹⁰ FZ Jülich, Institute for chemistry and dynamics of the Geosphere-2: Troposphere, Jülich, Germany¹¹ Norwegian Meteorological Institute, Norway¹² Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy

Received: 10 August 2011 – Accepted: 12 October 2011 – Published: 26 October 2011

Correspondence to: G. Foret (foret@lisa.u-pec.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Abstract

A detailed 3-D evaluation of an ensemble of five regional CTM's and one global CTM with focus on free tropospheric ozone over Europe is presented. It is performed over a summer period (June to August 2008) in the context of the GEMS-RAQ project. A data set of about 400 vertical ozone profiles from balloon soundings and commercial aircraft at 11 different locations is used for model evaluation, in addition to satellite measurements with the infrared nadir IASI sounder showing largest sensitivity to free tropospheric ozone. In the free troposphere, models using the same top and boundary conditions from MOZART-IFS exhibit a systematic negative bias with respect to observed profiles of about -20% . RMSE values are constantly growing with altitude, from 22% to 32% to 53% , respectively for 0–2 km, 2–8 km and 8–10 km height ranges. Lowest correlation is found in the free troposphere, with minimum coefficients (R) between 0.2 to 0.45 near 8 km, as compared to 0.7 near the surface and similar values around 10 km. Use of hourly instead of monthly chemical boundary conditions generally improves the model skill. Lower tropospheric 0–6 km partial ozone columns derived from IASI show a clear North-South gradient over Europe, which is qualitatively reproduced by the models. Also the temporal variability showing decreasing ozone concentrations in the lower troposphere (0–6 km columns) during summer is well caught by models even if systematic bias remains (the value of the bias being also controlled by the type of BC used). A multi-day case study of a through with low tropopause was conducted and showed that both IASI and models were able to resolve strong horizontal gradients of middle and upper tropospheric ozone occurring in the vicinity of an upper tropospheric frontal zone.

1 Introduction

Regional Chemical Transport Models (RCTM) are now central tools of air quality policy. In the case of ozone, their operational use for short-term forecast and monitoring

ACPD

11, 28797–28849, 2011

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Rouil et al., 2009; www.airnow.gov) implies to identify and reduce the remaining uncertainties. Classically, RCTM are evaluated against surface observations (Honoré et al., 2008; Van Loon et al., 2007; Vautard et al., 2007) since their primary goal is to simulate pollutants to which humans and more generally the biosphere, are directly exposed. On the contrary, performances of such models to simulate free tropospheric ozone have been less evaluated (in contrast to global scale models, Johnson et al., 2010a). Nevertheless, precise simulation of tropospheric ozone fields is crucial from the point of view of air quality. Since ozone is known to be harmful for humans (West et al., 2007) and vegetation development (Felzer et al., 2007), it is important to evaluate its long-range transport from source regions (Liang et al., 2004; Jonson et al., 2010) and the downward exchange between free troposphere and the boundary layer, which is poorly documented at the moment, but which is thought to be significant (Fiore et al., 2002; Foret et al., 2009; Parrington et al., 2009). In addition, the correct simulation of regional scale tropospheric ozone is important to assess its impact on regional climate change: ozone is the third most important greenhouse gas of the atmosphere (Forster et al., 2007) and, as an oxidant, it controls concentrations of other important greenhouse gases (mostly methane via OH production, Forster et al., 2007).

Vertical profiles of free tropospheric ozone provided by balloon borne ozone sondes and performed on board of commercial aircrafts (MOZAIC program) are very precious, because of their high vertical resolution. For summer 2008, ozone vertical profiles made by sondes have been obtained at 9 sites over Europe, among which five sites with a frequency of one or more soundings per week. MOZAIC vertical ozone profiles have been also obtained near 2 airports (Frankfurt, London) with sometimes more than one profile per day. In addition, the new generation of nadir viewing infrared sounders (IASI, Clerbaux et al., 2009; TES, Worden et al., 2007) is now operational and it opens new perspectives to study free tropospheric ozone. Thanks to its twice daily coverage of Europe (under cloud free conditions), IASI is a particularly good candidate due to a higher sensitivity to the free tropospheric (comparing to older instruments like GOME and/or SCIAMACHY) and in some cases also boundary layer ozone

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



concentrations, these observations offer the possibility to evaluate/constrain pollution models (Eremenko et al., 2008; Foret et al., 2009).

The FP6 European project “Global and regional Earth-system (atmosphere) Monitoring using Satellite and in-situ data” (GEMS) aimed at developing a pre-operational system for forecasting the chemical composition of the atmosphere at the global scale and more specifically at the regional scale for Europe by using an ensemble of RAQ (where RAQ stands for Regional Air Quality) models (Hollingsworth et al., 2008). In the framework of the RAQ-GEMS subproject, ten European RCTM have been set up (since June 2008) to forecast pollutants concentrations (ozone, NO₂, SO₂, CO and particles) over Europe (<http://gems.ecmwf.int/d/products/raq/>). The IFS-MOZART (Global CTM) forecast (Flemming et al., 2009) is used as boundary conditions (for top and lateral boundaries) for most of the RAQ models, but it produces also forecast over the regional domain. Model skills scores (such as bias, RMSE ...) have been calculated on-line for pollutants surface concentrations using measurements made by European air quality networks (<http://acm.eionet.europa.eu/databases/airbase/>). On the contrary, few efforts have been paid to evaluate the model abilities to reproduce free tropospheric concentrations. One reason for this is the lack of suitable (near real time) observations. It is proposed here is to conduct such an evaluation for tropospheric ozone in a hindcast mode. To do so, a specific exercise has been set-up where 5 of the GEMS-RAQ RCTM have re-simulated the summer 2008 period, with, for some of them, new configurations allowing simulating the whole troposphere. These models are state-of-the-art models in Europe and together they are a representative sample of European RCTM's. They will be confronted to an extended set of tropospheric ozone measurements from sondes, commercial aircraft (MOZAIC), and thermal-infrared measurements onboard satellite (IASI). To our knowledge, this is the first study that uses IASI ozone observations to evaluate RCTM's. The frequency of observations (especially the daily coverage for IASI observations) allows performing comparisons between observations and models from the seasonal to the day-to-day temporal scale. More specifically, we discuss uncertainties induced by the different representation in models

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of some of the processes controlling tropospheric ozone concentrations. Especially, for boundary conditions, we compare the impact of climatological and daily resolved boundary conditions but also differences in model transport between regional scale and global scale CTM.

In Sect. 2 in situ and satellite observations are described. Section 3 presents the models participating to the exercise. The results of the systematic comparison between observations and models (over a whole summer period) are shown in Sect. 4 including also a case study that illustrates the synergy between models and satellite data to analyse specific events.

2 Description of observations

2.1 In situ observations: ozone sondes and aircraft

2.1.1 Tropospheric ozone measurements by sondes

Vertical ozone soundings are obtained from electrochemical sensors lifted by hydrogen filled rubber balloons up to 30 km altitude. The vertical resolution of the stored measurements is about 100 m. The accuracy of such measurements is estimated to be better than 5 % in the troposphere (Smit et al., 2007). Over the “GEMS” European domain (covering part of European Russia, see model domain in Sect. 3), we have gathered data from 9 sounding sites for summer 2008 (June to August, Fig. 1). Table 1 indicates coordinates and altitude for each site as well as the number of profiles available and the databases from which they are available.

2.1.2 Tropospheric ozone measurements by commercial aircraft

Since 1994, ozone is measured onboard commercial airliners in the framework of the MOZAIC program (Marenco et al., 1998). The principle of ozone measurements is a dual beam UV absorption with an accuracy estimated at ± 2 ppb or +2 % (Thouret et

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



al., 1998). Vertical resolution of profiles taken during the take-off and landing phases is about few tens of meters. For summer 2008, a large number of profiles were available at Frankfurt (162 profiles) and London (58 profiles) airports. For Frankfurt, this corresponds to a daily frequency of nearly two (1.76 day^{-1}).

2.2 Tropospheric ozone measurements by satellite

The IASI (Infrared Atmospheric Sounding Interferometer) instruments (Clerbaux et al., 2009) are nadir viewing Fourier-transform spectrometers designed for operation on the meteorological Metop satellites (ESA/EUMETSAT). The first instrument was launched aboard the satellite Metop-A on 19 October 2006, and started operational measurements in June 2007. IASI is a Michelson-type Fourier-transform spectrometer providing infrared atmospheric emission-absorption spectra with a large spectral coverage ($645\text{--}2760 \text{ cm}^{-1}$), high radiometric sensitivity and accuracy, and the rather fine spectral resolution (the apodized spectral resolution is 0.5 cm^{-1}). This allows deriving global distributions of several important atmospheric trace gases (among which ozone (e.g. Boynard et al., 2009), CO (e.g. George et al., 2009), ammoniac (Clarisse et al., 2009)). The vertical Nadir field of view for one IASI pixel has the diameter of 12 km at the surface. The maximum scan angle of 48.3° from Nadir corresponds to coverage of about 2200 km across-track for one swath.

The retrieval of ozone profiles from IASI spectra used in the present study is performed with the radiative transfer model KOPRA (Karlsruhe Optimised and Precise Radiative transfer Algorithm, Stiller et al., 2000) and its numerical inversion module KOPRAFIT. The inversion method was set-up and first applied by Eremenko et al. (2008). To achieve maximal information content in the troposphere a constrained least squares fit method with an analytical altitude-dependent regularization is used. The regularization matrix is a combination of zero, first and second order Tikhonov constraints with altitude-dependent coefficients that were optimised to both maximise the Degrees of Freedom (DOF) of the retrieval in the troposphere and to minimise the total error of the retrieved profile. A validation exercise performed over the first one-year-and-a-half

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of IASI operation for the northern midlatitudes showed a bias of less than 5 % in the retrieved ozone. Calculated instrumental and retrieval errors (in total about 18 % for 0–6 km partial columns for mid-latitudes) are consistent with the standard deviation of the differences between sonde measurements and IASI observations (Keim et al., 2009).

Due to the limited number of degrees of freedom in the troposphere and considering the GEMS-RAQ focus on lower tropospheric ozone, partial 0–6 km ozone columns have been chosen as the basis of comparison between IASI observations and RAQ model simulations in this paper, as in Eremenko et al. (2008). In order to make simulations comparable to the retrieval, the simulated ozone profile vector \mathbf{x}_s needs to be transformed into a pseudo-retrieved profile \mathbf{x}_r by applying Eq. (1):

$$\mathbf{x}_r = \mathbf{x}_a + \mathbf{AVK}(\mathbf{x}_s - \mathbf{x}_a) \quad (1)$$

Here **AVK** denotes the Averaging Kernel Matrix, which expresses the sensitivity of the retrieved profile to the true profile and, by extension, to the a priori information (\mathbf{x}_a). A row of the AVK indicates the sensitivity of retrieved ozone, at a given layer, to changes in ozone at the same and other layers. This matrix is calculated during the retrieval process for each individual retrieved profile. An example of a typical AVK is shown in Fig. 2. The left panel shows the rows of AVK for different altitudes (black curves correspond to levels between 0 and 6 km, red curves to levels between 7 to 12 km). The right panel shows the integrated AVK over these two height ranges. This figure indicates that due to the measurement set-up and as a result of the retrieval method: (1) it is impossible to separate information originating from nearby vertical levels; (2) the sensitivity to the lower levels of the ozone profile (below 3 km) is relatively small. Nevertheless the lower and the upper parts of the troposphere (0–6 km and 7–12 km) are almost independent and can be separated when thermal condition (surface temperature, thermal contrast) are favorable, i.e. mainly during summer (Dufour et al., 2010). In Eq. (1), \mathbf{x}_a represents the a priori ozone profile used in the retrieval. Application of Eq. (1) to the “high resolution” vertical profile \mathbf{x}_s ensures that \mathbf{x}_a (the a priori) has no impact on the IASI-simulation comparison.

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In the following, we name “raw” columns the vertical columns integrating the simulated profiles x_s , and “smoothed” columns those calculated from the x_r profile (Eq. 1). For comparison purposes, individual IASI measurements (pixels) are regridded over the CHIMERE model grid cells with 0.5° resolution for the comparison exercise.

3 Model set-up and processing

Five models from the RAQ activity within the FP6/GEMS project participate to this exercise (BOLCHEM, CAMx, CHIMERE, EURAD, and MOCAGE). In addition, MOZART global fields provided by IFS (Fleming et al., 2009) are also included in the comparison. Model runs are all performed over the common European GEMS-RAQ domain (Fig. 1) for three summer months 2008 (June to August). Note that all models during this period have also been active for real time air quality forecast over Europe within the FP6 GEMS project (Hollingsworth et al., 2008). However, in this work hindcast simulations are used, because the whole set of tropospheric ozone simulations has not been stored during real time forecast. Moreover, some models have modified their operational version for this exercise; especially the BOLCHEM and CHIMERE model have moved the top of their domain from 500 to 200 hPa.

Global meteorological analysis and previsions with a spectral resolution of T799 and with 91 vertical levels up to 1 hPa are provided by ECMWF to the GEMS project (called IFS meteorology). All models use it either as direct meteorological input (CHIMERE, MOCAGE, MOZART) or as large scale fields for separate mesoscale simulations, on-line in BOLCHEM based on BOLAM dynamics (Buzzi et al., 1994), and off-line with the MM5 meteorological regional model for EURAD and CAMx. For all models, anthropogenic emissions are taken from the high resolution emission data base provided by TNO for GEMS (Visschedijk et al., 2007). For biogenic emissions, two models (BOLCHEM, CAMx) use the grid-based Biogenic Emission model (BEM; Poupkou et al., 2010) that allows calculating NMVOC (Non-Methane Volatil Organic Compound) emissions from vegetation at high spatial ($30 \text{ km} \times 30 \text{ km}$) and temporal (hour) scale.

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Similar approaches are also used by other models: EURAD (Guenther et al., 1993); MOCAGE (Guenther et al., 1995; Dentener et al., 2004) and the MEGAN model (Guenther et al., 2006) is used for CHIMERE. Hourly varying boundary conditions (BC) for ozone (but also for CO, NO, NO₂, HNO₃, peroxyacetyl nitrate, C₂H₆, isoprene, toluene and some others) are taken from MOZART-IFS global fields for most of the models except for MOCAGE that is using MOCAGE global simulations as hourly boundary conditions. As we will see later, the choice of boundary conditions can be a crucial parameter in model's behaviour. Models also include various formulations for atmospheric chemistry using well characterised reduced chemical schemes (Table 2). Dry deposition schemes are based on the classical “resistance” approach (Wesely, 1989). Another important model feature is the representation of pollutant transport. Table 2 indicates choices made in each RCTM to describe horizontal and vertical advection, turbulent transport in the planetary boundary layer, and convection by clouds. The impact of using different formulations for some of these processes will be analysed in Sect. 4.

The horizontal resolution of models varies between 0.2° and 0.5°, the model top between 200 (approximate tropopause height over Europe) and 10 hPa (only the CAMx model has a top above 200 hPa). The number of vertical tropospheric levels used to discretise the troposphere is about 20 between surface and 200 hPa (Table 2). Daily concentrations calculated by each model have been horizontally transposed on a 0.5° × 0.5° lat/lon grid (Fig. 1) and a 1 km steps vertical grid (from 0 to 12 km), in order to have a common reference frame for comparison. This should reduce partially the impact of having different horizontal and vertical resolution used by the models (cf. Table 2). For the comparisons with satellite data, “raw” and “smoothed” partial 0–6 km columns have been calculated as explained in the previous section.

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Systematic model evaluation over the summer period

We first present here the systematic comparison of ozone tropospheric profiles simulated with RAQ models (Table 2) against in-situ observations available from sondes and aircraft (Table 1). Results are analysed in terms of bias, RMSE (Root Mean Square Error) and correlation as a function of altitude and integrated over the whole summer period (June to August 2008). Also the impact of the chemical boundary conditions (for lateral and top limits of the modelling domain) is investigated. Next, this evaluation is completed by the confrontation of models to satellite observations (IASI partial tropospheric columns of ozone) at different time scales (from seasonal to daily time scale). This section is also completed by a case study that illustrates the models and IASI ability to reproduce strong ozone gradient in the troposphere associated to the tropopause height variability.

4.1 Comparisons between models and in situ vertical profiles

The comparison between simulations and in situ vertical ozone profiles is performed in the following way. In the horizontal plane, the model grid point closest to the observations is used. For MOZAIC measurements, we take into account the horizontal displacement of aircraft during take-off and landing (up to 500 km until the flight level is reached). In the vertical, we interpolate observations and simulations to a uniform grid, stretching from 0 to 10 km (a.s.l.) with 1 km steps. We apply linear interpolation. Second order interpolation was also tested, but differences in error statistics revealed negligible. With respect to time, the closest hourly model output with respect to the mean observation time is taken. Figure 3 shows the results of this comparison: vertical profiles of mean bias (Model-Observation), RMSE (Root Mean Square Error) and Pearson's correlation coefficient. Values are averaged over 1 km-height layers (plotted here in the middle of each layer) and over all available data from soundings and aircrafts (about 400 vertical profiles). To analyse these results, we have chosen to first consider models using the MOZART-IFS hourly boundary conditions that constitute by

ACPD

11, 28797–28849, 2011

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



construction a coherent sub-ensemble (i.e. BOLCHEM, CAMX, CHIMERE, EURAD and MOZART-IFS itself). For the CAMX model, we have restricted the comparison to below 6km height due to unsatisfying results above probably associated to problems in the coupling with MOZART-IFS.

4.1.1 Results in the planetary boundary layer (PBL; 0–2 km height)

In the PBL, mean ozone concentration is about 46 ppb with a 12.4 ppb standard deviation (Table 3). The mean model bias is -7% with values varying, as a function of site, in a range between -24% and $+8\%$ (Table S1). Largest negative biases are observed at Valentia (-24%), Lerwick (-21%) and Sodankyla (-19%), stations more directly under the influence of air masses from northern Atlantic and polar origin. This reflects probably a bias in the MOZART-IFS ozone fields for these regions. From Fig. 3, we observe a median bias of -2% between 0 and 1 km height, which increases to about -12% between 1 and 2km. RMSE is almost constant in the PBL with a median value of about 10 ppb (22%) (Fig. 3; Table. 3). This value is fairly similar at all sites (Table S1). We can note that these RMSE are similar to those obtained from previous evaluation studies using operational surface ozone measurements (Honoré et al., 2008; Vautard et al., 2007).

Concerning the Pearson's coefficient of correlation, the median value in the PBL is 0.71 but with significant variations from site to site (from 0.24 at Barajas to 0.86 at De-Bilt). These discrepancies between sites are difficult to understand because they do not follow a clear pattern related to their geographical situation (i.e. coastal sites, mountainous sites which would be more complex to model do not show lower correlations in a systematic way). The general good correlation in PBL indicates that ozone build-up in the boundary layer is fairly well represented in the models as it is generally confirmed also by comparisons with ground stations (GEMS Final report, 2010). The variability ratio (model standard deviation divided by observation standard deviation) shows that models reproduce well the observations variability ($\pm 10\%$) except for Barajas (0.68) and Valentia (0.58) (Table 3 and Supplement Table S1).

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.1.2 Results in the free troposphere (FT; >2–8 km height)

In the FT, mean ozone concentrations are about 66 ppb with a 16.7 ppb standard deviation (Table 3). Mean model bias is negative (−16 %, Table 3). Largest negative values of the model median (about −20 %) are reached at 5 km height (Fig. 3). This behaviour is fairly systematic at all sites with a more or less pronounced minimum (Fig. S1). This type of negative model bias in free tropospheric ozone over Europe has already been observed for several global models in earlier studies. Law et al. (2000) already pointed out a negative model bias of ozone in the troposphere for European sites presenting a summer maximum. They postulated that it could be due to “a lack of chemistry, deficiencies in transport schemes, as well as inadequate resolution”. Tarasick et al. (2007) also observed such a feature and postulated inaccuracies in the representation of stratosphere to troposphere exchange. More recently, Jonson et al. (2010) showed this negative bias for the Uccle station, especially in the middle and upper troposphere during a summer period (cf. Fig. 3 of their work). Finally, Ordonez et al. (2010) came also to this conclusion after comparing GEMS-GRG (GRG stands for Global Reactive Gases) global models, including MOZART, to aircraft data, and concluded that a combination of uncertainties affecting model simulations (coarse horizontal resolution, uncertainties in long-range transport of pollution, limitations of the chemistry scheme, under-estimated emissions) were responsible of this underestimation in FT ozone. Since the RCTM's evaluated here use boundary conditions derived from the global MOZART model, it is plausible that free tropospheric boundary condition at the edge of Europe are also biased negatively, causing the negative bias in FT – ozone in the RCTMs studied here.

RMSE (of the mean) increases almost linearly from about 24 % at 3 km height to 35 % at 8 km height (Fig. 3). This feature is observed at almost all sites except Barajas and Valencia (located near the western edge of the domain) for which values stays almost constant. In parallel, correlations are decreasing from 0.66 at 3 km height to a minimum of about 0.41 at 8 km height. Such patterns are observed for the three sites, Frankfurt,

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



London and, to a lesser extent, Payerne, that are dominating the statistics (273 profiles of 395). At most other locations a more or less broad minimum centred at 5–6 km height is observed (Supplement Fig. 1). Apparently models better reproduce ozone variability due to photochemical build-up in the PBL and due to the tropopause height variability in the upper troposphere (see below) than the more “diffuse” forcing in the middle troposphere due to long range transport, slow photochemistry, and exchange with lower and upper layers. The fact that RMSE increases with altitude, even when correlation increases again beyond its minimum, is due to increasing variability in ozone profiles at higher altitudes.

The ratio between the modelled and observed standard deviation is close to 1 on average (0.96; Table 3) but a certain spread is observed from one site to another ranging from 0.65 at London to 1.40 at Södankyla.

4.1.3 Results in the Upper Troposphere (UT; >8–10 km height)

In the UT, as expected, much higher and variable ozone concentrations are observed ranging from about 84 ppb (Barajas) to 148 ppb (Lerwick) with associated standard deviations representing 27 % to 65 % of these values (Supplement; Table S1). Spatial gradients and the large temporal variability in observed ozone levels are induced by the vicinity of the tropopause and its spatial-temporal variability, which determine the degree of stratospheric and ozone enhanced character of air masses. Vertical transport across the tropopause is an additional process affecting ozone fields (e.g. Stohl et al., 2003). As a consequence, ozone fields simulated by RAQ models are highly influenced by meteorological forcing and model transport (i.e. tropospheric height, advection by winds) as well as by top and lateral boundary chemical forcing. Chemistry plays a minor role due to the residence time of air masses over the domain less than a few days, but it impacts boundary conditions. Mean model bias is weak (below 1 %, Table 3), with a large variability for individual sites ranging from –18 % at Lerwick to +30 % at Payerne. A rapid increase of relative RMSE with height is observed reaching about 55 % at 10 km height associated to a significant improvement of correlations

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(~0.55 at 10 km height). These features are similar at almost all sites (Supplement; Fig. S1; Table S1). This increase in RMSE with altitude despite of an increase in correlation is explained by the increase in variability in ozone profiles reaching unity (1σ standard deviation at 10 km height). The average variability ratio between simulations and observations is close to one (0.88) as for other height ranges (Table 3) but again with a large spread for individual sites (Supplement; Table S1).

These results indicate that models perform quite well in the PBL region for which they had been initially designed. Considering the vertical structure of model errors, it clearly shows a C-shape form of the bias with a minimum in the free troposphere (~5–6 km height) of about –20 %. RMSE exhibits increasing values from about 20 % in the PBL to about 55 % at 10 km height that correlate to the vertical gradient of ozone variability. Correlation also follows a kind of C-shape but with a minimum (0.4) at about 8–9 km height.

4.1.4 Analysis of differences between models

Figure 3 also illustrates that discrepancies exist between models themselves even if global meteorological forcing, anthropogenic emissions (at least for RAQ models) and chemical boundary conditions are similar as it is the case for the subset of models using MOZART-IFS as boundary conditions (black curve and associated bars on Fig. 3). Nevertheless various formulations (chemistry, transport ...), forcings (natural emissions ...) and numerical set-up (horizontal and vertical resolutions) remain different between the models.

A weak dispersion in models results for biases and RMSE is observed in the PBL where ozone concentrations are strongly controlled by emissions (anthropogenic and natural), turbulent and horizontal transport and photochemistry. This indicates that differences in these processes likely do not induce large differences between models. This idea is reinforced by the fact that differences between models are increasing with height when the influence of these “PBL” processes decreases (Fig. 3). As net ozone production due to photochemistry in the middle and upper troposphere is expected

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to be weak during the residence time of air masses in the regional model domain of several days, we are suspecting that discrepancies between models are induced by the (horizontal and vertical) advection scheme, and horizontal and vertical resolution. The way the top boundary is handled can also be an issue. Sensitivity tests for short periods (ten days) are performed with the CHIMERE model have shown that discrepancies can occur when using different advection schemes (a simple first-order upwind scheme; the Van Leer second-order scheme (Van Leer et al., 1979) used in the reference run; the PPM (Piecewise Parabolic Method) third-order scheme, Colella and Woodward, 1984). Differences are bigger when winds are stronger (i.e. at high altitudes and latitudes) but remain generally weak (a few ppb). Thus this error source does not explain the observed model-to-model differences. Differences due to horizontal resolution will be discussed below.

Differences could be due also to vertical transport due to differences in vertical advection schemes (different for each model), to differences in the treatment of top boundary conditions and in vertical resolution. Also the way vertical velocities are computed from the continuity equation, either as a diagnostic (CAMx, CHIMERE) either as a direct output of meteorological models (BOLCHEM, EURAD, MOZART-IFS), could play a role. All models except CAMx (monthly mean from IFS-MOZART) use hourly IFS-MOZART as top conditions but with different top levels, 300 hPa for CAMx, 200 hPa for BOLCHEM and CHIMERE and 100 hPa for EURAD. For example, changing the top of the CHIMERE model from 200 hPa to 150 hPa (i.e. here we use 18 vertical levels instead of 17) induces differences (that grow with altitude) of +10 ppb for the 9–12 km layer with 95 % of the values included between 0 and 20 ppb. Thus the choice of the level of the model top boundary could have some impact on model errors in the upper troposphere.

We can note that differences in correlations are more constant throughout the troposphere with especially weaker differences in the upper troposphere (contrary to bias) where all models probably follows IFS-MOZART. Last, it is interesting to discuss, whether RCTM simulations with horizontal resolutions between 0.2° and 0.5°

show larger correlation coefficients than the global MOZART-IFS model with nearly 2° horizontal resolution. Indeed, near the surface, the global model shows lower correlations coefficients (0.67) than the regional models (0.69–0.78), making evident the benefit of higher resolution to improve simulation of PBL photochemical ozone build-up. However, from 2 km height on, the MOZART-IFS correlation coefficient is close to the median one (not shown), so in the free troposphere improved resolution does not necessarily result in better ozone simulations. As expected, the MOZART-IFS model exhibits (not shown) a lower variability than observations and than the RCTM's over the whole troposphere.

In conclusion, model-to model differences are most pronounced in the upper troposphere. A large variety of model settings could be responsible for errors, in particular related to transport processes. Some of them could be tested within CHIMERE, for instance the impact of the horizontal advection scheme (minor) or of the choice of top boundary (potentially contributing to part of the errors for the case of CHIMERE), but a final explanation for the model to model differences could not be achieved in this work.

4.1.5 Impact of chemical boundary conditions

By construction, limited-area models need to be provided at their boundaries with concentrations of long-lived (CO, O₃ ...) and shorter-lived pollutants (as NO_x ...). The impact of use of different boundary conditions on regional model results is analysed here.

As described previously, it is common to use large-scale climatologies to prescribe top and boundary conditions of RCTM's to avoid the set-up of more complicated combined global-regional modelling chains. One of the achievements of the FP6 GEMS project was to set up this type of systems in which global models provide hourly chemical boundary conditions (BC) to regional models. Szopa et al. (2009) have shown that the impact of improving BC variability (use of daily instead of monthly BC) on surface ozone concentrations remains limited (less than 5 %) in the centre of the regional European modelling domain. Nevertheless, the authors did not evaluate the impact on free tropospheric ozone concentrations. Here, we use our ensemble of different RCTM's

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with different forcings from GCTM output or from climatologies. Two different types of boundary forcings are evaluated: (1) hourly forcing from another GCTM than MOZART-IFS (namely MOCAGE); (2) the MOZART climatology in comparison to hourly forcing for one of the RAQ model (namely the CHIMERE model). Their impact is evaluated over the whole tropospheric height range using the in situ measurements presented earlier.

First, we have evaluated the impact of using climatological boundary conditions instead of hourly ones. To do so, we have simulated the whole period with the CHIMERE model using the monthly averaged values of the MOZART model instead of the hourly values and compared both model configurations to observations. As expected, both produce quite similar results in terms of biases with differences never exceeding 4 % (Fig. 3). For RMSE, differences increase but remain quite small reaching about 5 % at 9 km height. This height dependence is explained by a smaller influence of BC in the PBL due to local forcings, and by a temporal ozone variability increasing with height in the middle and upper troposphere. For correlations, differences are more systematic and the version with hourly IFS-MOZART BC is always better. Differences in the correlation coefficient are more significant above 3 km height, increasing from 0.03 to more than 0.2 at 10 km height. This indicates that temporal variations are better reproduced by the hourly BC than the monthly ones.

Second, we compared results of the MOCAGE model that uses its own BC in a nested global – regional simulation to those obtained with RAQ models forced by MOZART-IFS fields. For MOCAGE a positive mean surface bias (up to 20 % at the surface) is observed (Fig. 3). This result is in agreement with a parallel work of Ordóñez et al. (2010). In the free troposphere, bias remains positive until about 4 km height, and then becomes neutral or negative above this altitude. MOCAGE RMSE is larger than that for other models in the PBL (by about ~40 % at the surface) and becomes lower than for other models between 3 to 9 kilometres. Except in the PBL, correlations (Fig. 3) are similar to those of other models. Differences between MOCAGE and the median of IFS-MOZART driven models are indeed due to different boundary conditions. This can

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

be deduced from the fact that simulations are different for sites at western edge of the boundary (i.e Barajas, Valentia; figure not shown) and which are strongly influenced by boundary conditions. In addition, other differences in the model set-up (Table 2) can add to differences.

As a conclusion of these comparisons, we state an improvement of free tropospheric ozone simulations when passing from climatological ozone boundary conditions to hourly ones, although the benefit for boundary layer ozone predictions is rather small. This is an important finding of the GEMS project. It justifies the systematic coupling of global and regional models, if the aim is a consistent description of regional scale tropospheric ozone. However, if the aim is restricted to a prediction of boundary layer ozone only, this coupling is not mandatory and use of climatological boundary conditions for ozone seems sufficient (at least for the case of the CHIMERE model). BC from different global models can impact significantly vertical profiles at regional scales from the ground to the UT.

4.2 Comparisons between models and IASI 0–6 km columns

As a complementary data source for model evaluation, we use satellite observations obtained with the IASI instrument. As previously mentioned (Sect. 2.2), it is possible to derive 0–6 km tropospheric ozone partial columns from these measurements with good accuracy. Even though such observations still give limited vertical information (especially compared to those from sondes and aircraft), they are attractive because of their large spatial coverage (two complete ozone fields per day under cloud free conditions). It should be noted that results of the CAMx model are not included in the comparison presented here because the values simulated above 6 km height were not realistic as mentioned earlier. Given the observation averaging kernel, ozone values above 6 km height contribute to the retrieved 0–6 km columns, which makes it necessary to dispose of simulations also above 6 km.

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.2.1 Geographical distribution of summer averages

Figure 4 shows average IASI 0–6 km ozone partial columns for summer 2008 (June to August) interpolated on the model grid. The average is calculated using the more sensitive morning (by comparison with evening) observations. All IASI pixels available (up to five) within one model grid with 0.5° horizontal resolution are averaged to obtain a daily value. Individual profiles are smoothed using the averaging kernels to remove the a priori information (see Sect. 2.2 and Eremenko et al., 2008). The number of available “days” per grid cells (for the whole summer) is shown in Fig. 4. Indeed, pixels that do not fulfil the quality check (cloudy for example) are systematically discarded in the retrieval procedure (Eremenko et al., 2008). The number of available pixels is often less than 2/3 above 55° N, and generally less than 50 % over the Scandinavian Peninsula (Fig. 4). For the southern part of the domain, areas with low surface emissivity like desert areas (Maghreb, Southern Spain or even Turkey) are also poorly covered since. For such regions, strong aerosol loading (dust) as well as the presence of cirro-stratus along the subtropical jet-stream can also alter the measured radiances and then reduce the number of sampled pixels retrieved.

A clear north/northwest-south/southeast gradient in lower tropospheric 0–6 km ozone columns is observed by IASI over the European domain with largest values to the southeast especially over the Mediterranean basin (about 23–26 DU). Indeed, during summer, persistent anticyclonic (and subsident) conditions associated with strong photochemistry and low deposition rates are observed over the Mediterranean basin (Lelieveld et al., 2002; Foret et al., 2009). Such conditions favour the persistence of high ozone levels throughout the troposphere over this region. It should be noted that due to higher surface temperatures (and then higher thermal contrast between ground and surface air masses) in the southern part of the European domain, partial columns observed over this area are probably more sensitive to ozone concentrations at lower tropospheric altitudes. Strong horizontal gradients are often observed between land and marine surface for which surface temperature (and thus the observations sensitivity

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to ozone), but also orography is significantly different. The potential impact of these features on the gradient is not yet clear and should be further investigated. Over elevated or mountainous areas, ozone values are smaller since the thickness of atmospheric partial columns taken into account is reduced. Thus signature of Western European mountains an/or plateau (Meseta plateau, Pyrenées, Massif central, Alps, Scandinavian and Dinaric alps, Carpathian and Balkan mountains, Anatolian plateau) are visible on Fig. 4. We also note high ozone values over the Black Sea, Bulgaria, Romania, Moldavia and Ukraine with maxima of about 25 DU.

Corresponding smoothed columns calculated from models for hours with available observations. Models driven by MOZART-IFS BC qualitatively exhibit a similar, albeit less pronounced, north/northwest-south/southeast gradient as IASI (Fig. 4). Minimum values over Scandinavia in IASI observations, and maximum values over the eastern Mediterranean basin are reproduced by most of the models. Differences between the model median and IASI partial columns (Fig. 6a) exhibit a latitude dependence with a global model underestimation south of 60° N of about 2 to 4 DU (~10 to 20 %) and little bias (<1 DU) north of 60°. These results (i.e. negative bias) are well in line with the negative bias observed in the comparisons between models and vertical profiles (cf. Sect. 4.1). Discrepancies are more important over Spain and especially the Maghreb, regions with a weaker data coverage due to soil particularities (i.e. low emissivity) and, potentially, to the presence of airborne mineral dust. Also, over the northern coast of the Black sea and more generally over the south eastern part of the domain (near Romania), models underestimate the ozone maxima observed by IASI by about 6 DU. This value is still within the range of uncertainty of models (about 2 DU as seen from model dispersion in Fig. 4) and observations (10 to 20 %, about 2.5 DU).

Note that the 0–6 km partial columns of models without vertical smoothing (hereafter called “raw” columns) show a clear north-south gradient (Fig. 5). In the case of smoothed columns, differences between models themselves and/or IASI are less representative of the surface (due to the weak sensitivity of satellite observations to the surface and the use of a common a priori that dominates lowest levels) but integrate to

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



some extent information of the upper troposphere as seen from the averaging kernels (Fig. 2).

As expected from comparisons of models to sondes and aircraft, models using different chemical boundary conditions exhibit different behaviour. Figure 5 showing “raw” ozone columns confirms the positive bias of the MOCAGE model against other models below 6 km height as already shown by comparisons with in situ measurements. Comparisons with IASI (of the smoothed columns, Fig. 6b) show that MOCAGE performs well over the southern area of Europe but exhibits a positive bias over northern Europe, of more than 4 DU ($\sim 20\%$).

It is interesting to notice that comparisons between models and in situ measurements are fully consistent with comparisons between models and IASI: the median of models shows a negative bias with free tropospheric ozone from in-situ vertical profiles; this is confirmed by the comparison with 0–6 km IASI columns which indeed are most sensitive to free tropospheric ozone.

4.2.2 Summer ozone variability

As IASI inversions are available once per day from morning observations (under cloud free conditions), it is interesting to compare its temporal evolution for a summer season (here summer 2008) to the modelled one's. Figure 7 shows this variability expressed again as the smoothed 0–6 km partial columns and averaged over four model sub-domains that correspond to the four NW, NE, SW and SE model domain quarters). IASI daily (morning) observations are compared to the median of the models using MOZART-IFS as BC and the MOCAGE model. Both IASI and the models reproduce quite well the seasonal variability. This feature seems well in line with the expected slow decrease of ozone during the summer that follows the spring maximum (Monks, 2000) as observed at some remote stations in Europe (Chevalier et al., 2007; Gilge et al., 2010). Considering the median, as expected, a higher negative bias is observed for the southern part of the domain (-16%) instead of -8% (NW) and -5% (NE) in the north (Table 4), in line with the latitude of biases discussed before. It should be noted here that this bias is quite systematic (Fig. 7). Time correlations are relatively

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



high, between 0.74 for the NE to 0.63 (Table 4) for the SW sector indicating a good model ability to reproduce processes controlling regional scale ozone variability (either from BC or inside the domain itself). Also, we notice that correlations are systematically better in the eastern part compared to the western part at the same latitude when BC have less influence on the simulated concentrations. The dispersion of the ensemble is also plotted (Fig. 7) as the difference between the max and the min value of the ensemble for each day. In the northern part, the IASI observations are close or inside the model's variability while in the southern part of the domain where biases are more important they are almost systematically larger than the maximum model values. We notice that the mean dispersion of the ensemble is less important in the western part of the domain compared to the eastern part at the same latitude (2.7 DU in the NW against 3.5 DU in the NE and 1.8 DU in the SW against 3 DU in the SE). This is probably related to the use of common BC that have decreasing influence on simulated concentrations toward the east.

As expected from previous sections, the MOCAGE model (with its own boundary conditions) exhibits higher values in the lower free troposphere leading to a positive bias in the north (up to 10 %) and a weaker negative biases in the south (less than 5 %). The correlation remains good but is slightly lower than that of the IFS-MOZART driven model's median. Also, from Table 4, it is confirmed that the use of hourly BC compared to monthly averages largely improves the correlations for the case of the CHIMERE model across the whole domain.

In conclusion, the comparison between models and IASI shows that models qualitatively reproduce the observed lower tropospheric continental scale N/NW-S/SE gradient. Also the temporal variability of the columns at large geographical scales (1500–2000 km) is well reproduced (correlations between 0.63 and 0.8). These values are quite larger than those in the free troposphere for the comparison between simulations and in situ ozone profiles with large horizontal and vertical resolution (0.4 to 0.7). This nicely illustrates the scale dependence of the agreement between simulations and observations.

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.3 Case study of large ozone gradients in relation with an upper tropospheric wave

Since the IASI instrument is on board the METOP platform that samples the European domain at daily scale, it is conceptually possible to track specific ozone events. Especially, it is interesting to evaluate if models as well as IASI can reproduce large ozone gradients. To illustrate this point, we have focused our analysis on a case study of an upper tropospheric wave inducing a large variability in tropopause height and upper tropospheric ozone values.

From 8 to 11 June 2008, the median of raw simulated 0–6 km columns fields shows, to a varying degree, very prominent spatial features (Fig. 8, lower panel). A zone with enhanced ozone columns extends from Southern Norway to Northern Spain (also observed in the time series presented in the Fig. 7). Especially, spatial gradients at the western edge of this zone are very pronounced. Corresponding smoothed column fields show similar features although the spatial structure is less apparent, because only cloud free pixels for which also IASI observations are available are presented (Fig. 8, middle panel). For 8 June, spatial structures for smoothed simulated models and IASI observations (Fig. 8, upper panel) coincide rather well, the region of strong spatial gradients is only slightly shifted towards south in IASI observations with respect to models (from North Sea to the North sea coast). Observed and simulated spatial gradients coincide even better for 9 June, the steepest gradients being located at the German North Sea and the French channel coast. For 10 June, the correspondence is again very good, the steepest gradient zone being shifted about 100 km to south.

We now need to seek for an explanation why models (here represented by their median) show such similar spatial structures during this period, and moreover correspond very well with IASI observations. The potential vorticity (PV) contour map (figure not shown) at the 330 K potential temperature level (corresponding to about 12 km height) for 8 June shows a pronounced wave structure over Europe with a ridge over the British Islands (with low tropospheric PV values, below 1 PVU), and a trough covering a

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



large part of Western, Southern and Central Europe (with large stratospheric PV values, above 3 PVU). The region with strongest PV gradients follows the channel and North Sea coast from France to Denmark. Its NE-SW orientation and location correspond to the strong gradients in the IASI ozone column fields observed for this day (Fig. 9). Also the day to day evolution of IASI partial ozone columns and 330 K PV maps in the following days is correlated. This perfect coincidence of spatial structures suggests that variations in IASI and modelled partial ozone columns are caused by the upper tropospheric wave structure. It is well known that upper tropospheric ozone and PV are well correlated (for example, Danielsen, 1968, Beekmann et al., 1994). A vertical cross section through the upper tropospheric front along 51° N (Fig. 10) shows enhanced ozone values in the 4–10 km height region in the through region (>60 ppb), compared to ridge region (<40–50 ppb). Note that IASI observations are shown for specific altitudes (in km steps), but their implicit vertical resolution is of several kilometres. The picture in Fig. 10 is consistent with the spatial distribution in Fig. 8 when considering that due to the vertical sensitivity of IASI measurements (cf. Averaging Kernel in Fig. 2), the large ozone values in the 4–10 km region have a strong impact on the smoothed 0–6 km partial columns. Enhanced ozone values in the 4 to 8 km height range (between 60 ppb and 100 ppb) are also observed in a MOZAIC profile recorded from Frankfurt airport within the through region on 8 June at 06:45 UTC. The coincidence of enhanced ozone region with low CO, and low relative humidity indicates subsident motion from the tropopause region to the free troposphere. This is confirmed by Lagrangian particle simulations with the FLEXPART model (Stohl et al., 2005). For air masses arriving at Frankfurt, on 8 June, between 7 and 8 km altitude, they show subsiding anticyclonic motion of the retro-plume during the last three days, and indicate a significant fraction of air with stratospheric origin (from PV analysis). Nearly all models show strongly enhanced ozone values in the 4–10 km height region in the 51° N cross section east of –10° W (Fig. 10). For most of them ozone values in this region are somewhat stronger than those observed by IASI. Differences induced by the use of a monthly mean climatology (CHIMERE2) instead of hourly values (CHIMERE1) are

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



small for this case. Note that simulated fields in Fig. 10 are again smoothed in order to be comparable to IASI observations. Thus, in conclusion, both the agreement in the vertical and in the horizontal distribution between observed and simulated 0–6 km partial ozone columns is striking (Fig. 8), especially the gradient zone between the ridge and the through regions. Apparently, the deep through associated with low tropopause and high ozone values is well represented in IFS meteorological fields which are used by all models as input (either directly for the CTM or as large scale or boundary values for the mesoscale meteorological simulations). This case study illustrates the possibility to use IASI observations to evaluate the CTM model behaviour for cases of strong ozone gradients related to upper tropospheric wave structures.

5 Conclusions

The 3-D evaluation of an ensemble of RCTM to simulate tropospheric ozone concentrations over Europe is presented here. Several models have simulated ozone concentrations over an entire summer period (June to August 2008) in the context of the GEMS-RAQ project. Among those, five state of the art RCTM and the MOZART-IFS system have participated to this evaluation exercise. A large set of observed vertical ozone profiles, either from sondes or commercial aircraft have been used for this evaluation purpose, in addition to satellite derived partial columns. The data set used comprises about 400 vertical profiles at 11 different locations. The model skill of representing PBL ozone concentrations appears to be satisfying (relative bias of 4 %, RMSE of 24 %, correlation 0.77) and in the range of values observed in earlier studies using surface ozone measurements. In the free troposphere height region (>2–8 km), models using the same hourly top and boundary conditions from MOZART-IFS exhibit a systematic negative bias of about –20 %. This feature is commonly observed in global scale CTM's and not yet fully understood. RMSE values are constantly growing with altitude, both in an absolute and relative sense (from 32 % to 53 %, respectively in the >2–8 km and in the >8–10 km height range). Largest values in the UT are thought

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to be associated with the difficulty for models to catch to a full extent the variability of troposphere-stratosphere exchange processes or simply the height variation of the tropopause, although large correlation in the UT indicated that the basic processes governing ozone variability are taken into account. Correlation in the free troposphere revealed to be low, with minimum values of 0.2 to 0.45 near 8 km. Apparently, forcing processes for the ozone variability are not well captured in models in this height range. If long range transport of ozone contributes significantly to this variability, it is understandable that plume positions could not be easily predicted at several thousand kilometres distance from the sources. But misrepresentation of ozone chemistry as well as stratospheric intrusions upwind of Europe could also explain models errors. We also note that bias and RMSE are the lowest in the BL (as well as satisfying correlations) showing a better model capacity to reproduce ozone concentrations in the part of atmosphere for which these RAQ models have been originally designed. We also can add, that differences between models inside the domain are observed (generally increasing with altitude) especially for bias in spite of common meteorology and chemical boundary conditions. In this part of the troposphere, where surface processes like emissions and fast chemistry have a weak influence, transport processes most likely are responsible for differences. However, due to the multitudes of different settings within the models tested, the exact sources for model to model discrepancies could not be determined. During this exercise, the impact of using different chemical BC has also been investigated. Two ways of prescribing BC have been tested: variable BC using hourly forecast from either the IFS-MOZART or the global CTM MOCAGE and, using a monthly climatology calculated from IFS-MOZART instead of hourly values. It has been shown that the use of hourly (forecast) instead of monthly (climatology) BC generally improves the skill of one model to a certain extent (for example, the correlation in the 5 – 8 km height region increases from 0.2–0.3 to 0.4 when hourly BC are used with CHIMERE). Larger differences between models are observed when different CTM are used to produce BC (case for IFS-MOZART and MOCAGE, even if other settings are also different for MOCAGE).

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Another goal of the paper was to confront models to satellite data, especially to partial ozone columns (0–6 km) calculated from IASI observations. The IASI sounder is a thermal infrared instrument that allows estimating tropospheric ozone concentrations (mainly in the free troposphere) at twice daily frequency over Europe. It thus allows the identification of geographical pattern of the tropospheric ozone distribution and their temporal variations. The agreement between both IASI and models over the summer 2008 period is generally satisfying. Especially, IASI observations of minimum values over Scandinavia, and maximum values over the eastern Mediterranean basin are reproduced by most of the models. Below 60° north, a negative bias of models is observed well in line with comparisons between vertical profiles and models. Temporal variability in lower tropospheric ozone values during summer 2008 is also well reproduced by models (result obtained for IASI model comparisons averaged model sub-domains).

Finally, a case of a multiday upper tropospheric wave generating strong ozone gradients was observed by these satellite data, confirmed by a MOZAIC profile and meteorological analysis, and well reproduced by models. In particular, both IASI and models were able to resolve strong horizontal gradients in middle and upper tropospheric ozone occurring in the vicinity of the upper tropospheric frontal zone. This shows the potential of IASI observations for investigating the upper tropospheric ozone distribution. Ideally, these features should not only be analysed in 0–6 km partial ozone columns, which were the basis of this study, but also in 0–12 km or 6–12 km partial columns. During the summer 2008 period studied, no major photochemical ozone pollution event suitable for a case study occurred.

As a final general conclusion, it is shown in this paper that a combination of high resolution vertical ozone profiles at a limited number of sites and satellite observations with good spatial coverage, but low vertical resolution, allow for a thorough evaluation of tropospheric ozone simulations at various temporal scales (seasonal, case study). Within the framework of the GMES program (and its FP6/GEMS and FP7/MACC projects), this work also shows the ability of a combined system of vertical profile observations,

satellite observations, and model simulations to represent the free tropospheric vertical ozone distribution with a defined uncertainty, and to make evident key processes affecting its variability.

Supplementary material related to this article is available online at:

**[http://www.atmos-chem-phys-discuss.net/11/28797/2011/
acpd-11-28797-2011-supplement.pdf](http://www.atmos-chem-phys-discuss.net/11/28797/2011/acpd-11-28797-2011-supplement.pdf).**

Acknowledgements. This work has been funded by the FP6 GEMS (Global and regional Earth-system Monitoring using Satellite and in-situ data) and the FP7 MACC (Monitoring Atmospheric Composition Change) projects. The ozone sonde data have been taken from the following archives: theWorld Ozone and Ultraviolet Data Center (WOUDC) (<http://www.woudc.org>) and the NILU's Atmospheric Database for Interactive Retrieval (NADIR) at Norsk Institutt for Luftforskning (NILU) (<http://www.nilu.no/nadir/>). IASI has been developed and built under the responsibility of the Centre National d'Etudes Spatiales (CNES, France). It is flown aboard the Metop satellites as part of the EUMETSAT Polar System. The IASI L1C and L2 data have been received through the EUMETCast near real time data distribution service and provided by the Ether French atmospheric database (<http://ether.ipsl.jussieu.fr>). The presented study has been partly supported by CNES in the frame of the IASI-TOSCA project. The authors acknowledge the strong support of the European Commission, Airbus and airlines – Lufthansa, Air France, Austrian and former Sabena – to the MOZAIC program. This work is also part of the SMOQA project funded by the French LEFE program (CHAT/ASSIM). This work makes use of results produced by the Enabling Grids for E-science E project, a project co-funded by the European Commission (Under contract number INFOS-RI-22667). Full information is available at <http://www.eu-egee.org> This work has been developed with resources of the Earth Science Research virtual organization. The authors wish to acknowledge David Weissenbach (IPGP) for technical support of this research We thank the Institut für Meteorologie und Klimaforschung (IMK), Karlsruhe Institute of Technology (KIT), Germany, for a licence to use the KOPRA radiative transfer model, and especially M. Höpfner for his help with use of the code.

We are indebted to the initiator and coordinator of the GEMS project, Anthony Hollingsworth, who sadly passed away on 29 July 2007.

The authors are grateful to CNRS-INSU for publication support.

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The publication of this article is financed by CNRS-INSU.

References

- 5 Bechtold, P., Bazile, E., Guichard, F., Mascart, P., and Richard, E.: A mass flux convection scheme for mesoscale and global models, *Q. J. Roy. Meteor. Soc.*, 126, 865–889, 2001.
- Beekmann, M., Ancellet, G., and Mégie, G.: Climatology of tropospheric ozone in Southern Europe and its relation to potential vorticity, *J. Geophys. Res.*, 99, 12841–12853, doi:10.1029/94JD00228, 1994.
- 10 Bessagnet, B., Menut, L., Aymoz, G., Chepfer, H., and Vautard, R.: Modeling dust emissions and transport within Europe: the Ukraine March 2007 event, *J. Geophys. Res.*, 113, D15202, doi:10.1029/2007JD009541, 2008.
- Billet, S. J. and Toro, E. F.: On WAF-Type Schemes for Multidimensional Hyperbolic Conservation Laws, *J. Comput. Phys.*, 130, 1-24, 1997.
- 15 Blackadar, A. K.: Modelling pollutant transfer during daytime convection, Preprints, Fourth Symposium on Atmospheric Turbulence, Diffusion, and Air Quality, Reno, Am. Meteor. Soc., 443–447, 1978.
- Bott, A.: A positive definite advection scheme obtained by nonlinear renormalization of the advective fluxes, *Mon. Weather Rev.*, 117, 1006–1016, 1989.
- 20 Bousserez, N., Attié, J.-L., Peuch, V.-H., Michou, M., Pfister, G., Edwards, D., Avery, M., Sachse, G., Browell, E., and Ferrare, E.: Evaluation of MOCAGE chemistry and transport model during the ICARTT/ITOP experiment, *J. Geophys. Res.*, 112, D10S42, doi:10.1029/2006JD007595, 2007.
- Boynard, A., Clerbaux, C., Coheur, P.-F., Hurtmans, D., Turquety, S., George, M., Hadji-Lazaro, J., Keim, C., and Meyer-Arne, J.: Measurements of total and tropospheric ozone from IASI:

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- comparison with correlative satellite, ground-based and ozonesonde observations, *Atmos. Chem. Phys.*, 9, 6255–6271, doi:10.5194/acp-9-6255-2009, 2009.
- Buzzi, A., Fantini, M., Malguzi, P., and Nerozzi, F.: Validation of a limited area model in cases of Mediterranean cyclogenesis: Surface fields and precipitation scores, *Meteorol. Atmos. Phys.*, 53, 137–153, 1994.
- Carter, W.: A detailed mechanism for the gas-Phase atmospheric reactions of organic compounds, *Atmos. Environ.*, 27, 481–518, 1990.
- Carter, W.: Condensed atmospheric photooxidation mechanisms for isoprene, *Atmos. Environ.*, 30, 4275–4290, 1996.
- Chevalier, A., Gheusi, F., Delmas, R., Ordóñez, C., Sarrat, C., Zbinden, R., Thouret, V., Athier, G., and Cousin, J.-M.: Influence of altitude on ozone levels and variability in the lower troposphere: a ground-based study for western Europe over the period 2001–2004, *Atmos. Chem. Phys.*, 7, 4311–4326, doi:10.5194/acp-7-4311-2007, 2007.
- Clarisse, L., Clerbaux, C., Dentener, F., Hurtmans, D., and Coheur, P.-F.: Global ammonia distribution derived from infrared satellite observations, *Nature Geosci.*, 2, 479–483, 2009.
- Clerbaux, C., Boynard, A., Clarisse, L., George, M., Hadji-Lazaro, J., Herbin, H., Hurtmans, D., Pommier, M., Razavi, A., Turquety, S., Wespes, C., and Coheur, P.-F.: Monitoring of atmospheric composition using the thermal infrared IASI/MetOp sounder, *Atmos. Chem. Phys.*, 9, 6041–6054, doi:10.5194/acp-9-6041-2009, 2009.
- Colella, P. and Woodward, P. R.: The piecewise parabolic method (PPM) for gas-dynamical simulations, *J. Comput. Phys.*, 11, 38–39, 1984.
- Danielsen, E. F.: Stratospheric-tropospheric exchange based on radioactivity, ozone and potential vorticity, *J. Atmos. Sci.*, 25, 502–518, 1968.
- Dentener, F., Stevenson, D., Cofala, J., Mechler, R., Amann, M., Bergamaschi, P., Raes, F., and Derwent, R.: The impact of air pollutant and methane emission controls on tropospheric ozone and radiative forcing: CTM calculations for the period 1990–2030, *Atmos. Chem. Phys.*, 5, 1731–1755, doi:10.5194/acp-5-1731-2005, 2005.
- Dufour, G., Eremenko, M., Orphal, J., and Flaud, J.-M.: IASI observations of seasonal and day-to-day variations of tropospheric ozone over three highly populated areas of China: Beijing, Shanghai, and Hong Kong, *Atmos. Chem. Phys.*, 10, 3787–3801, doi:10.5194/acp-10-3787-2010, 2010.
- Elbern, H., Strunk, A., Schmidt, H., and Talagrand, O.: Emission rate and chemical state estimation by 4-dimensional variational inversion, *Atmos. Chem. Phys.*, 7, 3749–3769,

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



doi:10.5194/acp-7-3749-2007, 2007.

Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, 3, 43–67, doi:10.5194/gmd-3-43-2010, 2010.

Eremenko, M., Dufour, G., Foret, G., Keim, C., Orphal, J., Beekmann, M., Bergametti, G., and Flaud, J.-M.: Tropospheric ozone distributions over Europe during the heat wave in July 2007 observed from infrared nadir spectra recorded by IASI, *Geophys. Res. Lett.*, 35, L18805, doi:10.1029/2008GL034803, 2008.

Felzer, B., Cronin, T., Reilly, J. M., Melillo, J. M., and Wang, X.: Impacts of ozone on trees and crops, *C.R. Geosci.*, 339, 2007.

Fiore, A. M., Jacob, D. J., Bey, I., Yantosca, R. M., Field, B. D., and Fusco, A. C.: Background ozone over the United States in summer: Origin, trend, and contribution to pollution episodes, *J. Geophys. Res.*, 107, 4275, doi:10.1029/2001JD000982, 2002.

Fischer, H., Birk, M., Blom, C., Carli, B., Carlotti, M., von Clarmann, T., Delbouille, L., Dudhia, A., Ehnhalt, D., Endemann, M., Flaud, J. M., Gessner, R., Kleinert, A., Koopman, R., Langen, J., López-Puertas, M., Mosner, P., Nett, H., Oelhaf, H., Perron, G., Remedios, J., Ridolfi, M., Stiller, G., and Zander, R.: MIPAS: an instrument for atmospheric and climate research, *Atmos. Chem. Phys.*, 8, 2151–2188, doi:10.5194/acp-8-2151-2008, 2008.

Flemming, J., Inness, A., Flentje, H., Huijnen, V., Moinat, P., Schultz, M. G., and Stein, O.: Coupling global chemistry transport models to ECMWF's integrated forecast system, *Geosci. Model Dev.*, 2, 253–265, doi:10.5194/gmd-2-253-2009, 2009.

Foret, G., Hamaoui, L., Schmechtig, C., Eremenko, M., Keim, C., Dufour, G., Boynard, A., Coman, A., Ung, A., and Beekmann, M.: Evaluating the potential of IASI ozone observations to constrain simulated surface ozone concentrations, *Atmos. Chem. Phys.*, 9, 8479–8491, doi:10.5194/acp-9-8479-2009, 2009.

Forster, P., Ramaswamy, V., Artaxo, P., Berntsen, T., Betts, R., Fahey, D. W., Haywood, J., Lean, J., Lowe, D. C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., and Van Dorland, R.: Changes in Atmospheric Constituents and in Radiative Forcing. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L.,

ACPD

11, 28797–28849, 2011

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2007.
- Geiger, H., Barnes, I., Bejan, I., Benter, T., and Spittler, M.: The tropospheric degradation of isoprene: An updated module for the regional atmospheric chemistry mechanism, *Atmos. Environ.*, 37, 1503–1519, 2003.
- 5 George, M., Clerbaux, C., Hurtmans, D., Turquety, S., Coheur, P.-F., Pommier, M., Hadji-Lazaro, J., Edwards, D. P., Worden, H., Luo, M., Rinsland, C., and McMillan, W.: Carbon monoxide distributions from the IASI/METOP mission: evaluation with other space-borne remote sensors, *Atmos. Chem. Phys.*, 9, 8317–8330, doi:10.5194/acp-9-8317-2009, 2009.
- GEMS Final report: A monitoring and forecasting system for atmospheric composition, January 10 2010.
- Gery, M. W., Whitten, G., Killus, J., and Dodge, M.: A photochemical kinetics mechanism for urban and regional scale computer modeling, *J. Geophys. Res.*, 94, 12925–12956, 1989.
- Gilge, S., Plass-Duelmer, C., Fricke, W., Kaiser, A., Ries, L., Buchmann, B., and Steinbacher, M.: Ozone, carbon monoxide and nitrogen oxides time series at four alpine GAW mountain stations in central Europe, *Atmos. Chem. Phys.*, 10, 12295–12316, doi:10.5194/acp-10-12295-2010, 2010.
- 15 Guenther, A. B., Zimmermann, P. R., Harley, P. C., Monson, R. K., and Fall, R.: Isoprene and monoterpene emission rate variability: Model evaluations and sensitivity analyses, *J. Geophys. Res.*, 98, 12609–12617, 1993.
- Guenther, A. B., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Harley, L., Klinger, L., Lerdau, M., McKay, W. A., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.: A global model of natural volatile compound emissions, *J. Geophys. Res.*, 100, 8873–8892, 1995.
- 20 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, doi:10.5194/acp-6-3181-2006, 2006.
- 25 Hack, J. J.: Parameterization of moist convection in the NCAR community climate model (CCM2), *J. Geophys. Res.*, 99, 5551–5568, 1994.
- Hollingsworth, A., Engelen, R. J., Benedetti, A., Dethof, A., Flemming, J., Kaiser, J. W., Morcrette, J.-J., Simmons, A. J., Textor, C., Boucher, O., Chevallier, F., Rayner, P., Elbern, H., Eskes, H., Granier, C., Peuch, V.-H., Rouil, L., and Schultz, M. G.: Toward a monitoring and forecasting system for atmospheric composition: The GEMS project, *B. Am. Meteorol. Soc.*,
- 30

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



89, 1147–1164, 2008.

Holtstlag, A. and Boville, B.: Local versus nonlocal boundary-layer diffusion in a global climate model, *J. Clim.*, 6, 1825–1842, 1993.

Hong, S.-Y. and Pan, H.-L.: Nonlocal boundary layer vertical diffusion in a medium-range forecast model, *Mon. Weather Rev.*, 124, 2322–2339, 1996.

Honoré, C., Rouil, L., Vautard, R., Beekmann, M., Bessagnet, B., Dufour, A., Elichegaray, C., Flaud, J.-M., Malherbe, L., Meleux, F., Menut, L., Martin, D., Peuch, A., Peuch, V.-H., and Poisson, N.: Predictability of European air quality: Assessment of 3 years of operational forecasts and analyses by the PREVAIR system, *J. Geophys. Res.*, 113, D04301, doi:10.1029/2007JD008761, 2008.

Jonson, J. E., Stohl, A., Fiore, A. M., Hess, P., Szopa, S., Wild, O., Zeng, G., Dentener, F. J., Lupu, A., Schultz, M. G., Duncan, B. N., Sudo, K., Wind, P., Schulz, M., Marmer, E., Cuvelier, C., Keating, T., Zuber, A., Valdebenito, A., Dorokhov, V., De Backer, H., Davies, J., Chen, G. H., Johnson, B., Tarasick, D. W., Stübi, R., Newchurch, M. J., von der Gathen, P., Steinbrecht, W., and Claude, H.: A multi-model analysis of vertical ozone profiles, *Atmos. Chem. Phys.*, 10, 5759–5783, doi:10.5194/acp-10-5759-2010, 2010.

Josse, B., Simon, P., and Peuch, V.-H.: Rn-222 global simulations with the multiscale CTM MOCAGE, *Tellus B*, 56, 339–356, 2004.

Kain, J. S.: The Kain-Fristch convective parameterization: An update, *J. Appl. Meteorol.*, 43, 170–181, 2002.

Kain, J. S. and Fritsch, J. M. A.: One-dimensional entraining/detraining plume model and its application in convective parameterization, *J. Atmos. Sci.*, 47, 2784–2802, 1990.

Kinnison, D., Brasseur, G., Walters, S., Garcia, R., Marsh, D., Sassi, F., Harvey, V., Randall, C., Emmons, L., Lamarque, J., Hess, P., Orlando, J., Tie, X., Randel, W., Pan, L., Gettelman, A., Granier, C., Diehl, T., Niemeier, U., and Simmons, A.: Sensitivity of chemical tracers to meteorological parameters in the MOZART-3 chemical transport model, *J. Geophys. Res.*, 112, D20302, doi:10.1029/2006JD007879, 2007.

Keim, C., Eremenko, M., Orphal, J., Dufour, G., Flaud, J.-M., Höpfner, M., Boynard, A., Clerbaux, C., Payan, S., Coheur, P.-F., Hurtmans, D., Claude, H., Dier, H., Johnson, B., Kelder, H., Kivi, R., Koide, T., López Bartolomé, M., Lambkin, K., Moore, D., Schmidlin, F. J., and Stübi, R.: Tropospheric ozone from IASI: comparison of different inversion algorithms and validation with ozone sondes in the northern middle latitudes, *Atmos. Chem. Phys.*, 9, 9329–9347, doi:10.5194/acp-9-9329-2009, 2009.

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Law, K. S., Plantevin, P.-H., Thouret, V., Marengo, A., Asman, W. A. H., Lawrence, M., Crutzen, P. J., Muller, J.-F., Hauglustaine, D. A., and Kanakidou, M.: Comparison between global chemistry transport model results and Measurement of Ozone and Water Vapor by Airbus In-Service Aircraft (MOZAIC) data, *J. Geophys. Res.*, 105, 1503–1525, 2000.
- 5 Lefèvre, F., Brasseur, G. P., Folkins, I., Smith, A. K., and Simon, P.: Chemistry of the 1991-1992 stratospheric winter: three-dimensional model simulations, *J. Geophys. Res.*, 99, 8183–8195, 1994.
- Lelieveld, J., Berresheim, H., Borrmann, S., Crutzen, P. J., Dentener, F. J., Fischer, H., Feichter, J., Flatau, P. J., Heland, J., Holzinger, R., Korrman, R., Lawrence, M. G., Levin, Z.,
10 Markowicz, K. M., Mihalopoulos, N., Minikin, A., Ramanathan, V., de Reus, M., Roelofs, G. J., and Scheeren, H. A.: Global Air Pollution Crossroads over the Mediterranean, *Science*, 298, 794–799, doi:10.1126/science.1075457, 2002.
- Liang, Q., Jaegle, L., Jaffe, D., Weiss-Penzias, P., and Heckman, A.: Long-range transport of Asian pollution to the northeast Pacific: Seasonal variations and transport pathways of carbon monoxide, *J. Geophys. Res.*, 109, D23S07, doi:10.1029/2003JD004402, 2004.
- 15 Lin, S. J. and Rood, R. B.: A fast flux form semi-Lagrangian transport scheme on the sphere, *Mon. Weather Rev.*, 124, 2046–2070, 1996.
- Louis, J. F., A parametric model of vertical eddy fluxes in the atmosphere, *Bound.-Layer Meteor.*, 17, 197–202, 1979.
- 20 Marengo, A., Thouret, V., Nédélec, P., Smit, H., Helten, M., Kley, D., Karcher, F., Simon, P., Law, K., Pyle, J., Poschmann, G., von Wrede, R., Hume, C., and Cook, T.: Measurement of ozone and water vapor by Airbus in-service aircraft: The MOZAIC airborne program, An overview, *J. Geophys. Res.*, 103, 25631–25642, 1998.
- Michou, M. and Peuch, V.-H.: Surface exchanges in the multi-scale chemistry and transport model MOCAGE, *Water Sci. Rev.*, 15, 173–203, 2002.
- 25 Monks, P.: A review of the observations and origins of the spring ozone maximum, *Atmos. Environ.*, 34, 3545–3561, 2000.
- Mircea, M., D'Isidoro, M., Maurizi, A., Vitali, L., Monforti, F., Zanini, G., and Tampieri, F.: A comprehensive performance evaluation of the air quality model BOLCHEM to reproduce the ozone concentrations over Italy, *Atmos. Environ.*, 42, 1169–1185, 2008.
- 30 Morris, R., Lau, S., and Yarwood, G.: Development and Application of an Advanced Air Toxics Hybrid Photochemical Grid Modeling System, Presented at 96th Annual Conference and Exhibition of the A&WMA, San Diego, California, 2003.

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Ordóñez, C., Elguindi, N., Stein, O., Huijnen, V., Flemming, J., Inness, A., Flentje, H., Katragkou, E., Moinat, P., Peuch, V.-H., Segers, A., Thouret, V., Athier, G., van Weele, M., Zerefos, C. S., Cammas, J.-P., and Schultz, M. G.: Global model simulations of air pollution during the 2003 European heat wave, *Atmos. Chem. Phys.*, 10, 789–815, doi:10.5194/acp-10-789-2010, 2010.
- Parrington, M., Jones, D. B. A., Bowman, K. W., Thompson, A. M., Tarasick, D. W., Merrill, J., Oltmans, S. J., Leblanc, T., Witte, J. C., and Millet, D. B.: Impact of the assimilation of ozone from the Tropospheric Emission Spectrometer on surface ozone across North America, *Geophys. Res. Lett.*, 36, L04802, doi:10.1029/2008GL036935, 2009.
- Poupkou, A., Giannaros, T., Markakis, K., Kioutsioukis, I., Curci, G., Melas, D., and Zerefos, C.: A model for European Biogenic Volatile Organic compound emissions: Software development and first validation, *Environ. Modell. Softw.*, 25, 1845–1856, 2010.
- Randerson, J. T., van der Werf, G. R., Giglio, L., Collatz, G. J., and Kasibhatla, P. S.: Global Fire Emissions Database, Version 2 (GFEDv2), Data set, available at: <http://daac.ornl.gov/> from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, USA, doi:10.3334/ORNLDAAAC/834, 2006.
- Rouil, L., Honoré, C., Vautard, R., Beekmann, M., Bessagnet, B., Malherbe, L., Meleux, F., Dufour, A., Elichegaray, C., Flaud, J.-M., Menut, L., Martin, D., Peuch, A., Peuch, V.-H., and Poisson, N.: Prev'air: An Operational Forecasting and Mapping System for Air Quality in Europe, *B. Am. Meteorol. Soc.*, 90, 73–83, 2009.
- Schultz, M. G., Pulles, T., Brand, R., van het Bolscher, M., and Dalsøren, S. B.: A global data set of anthropogenic CO, NO_x, and NMVOC emissions for 1960–2000, available at: <http://retro.enes.org/dataemissions.shtml>, 2009.
- Simpson, D., Fagerli, H., Jonson, J., Tsyro, S., Wind, P., and Tuovinen, J.-P.: Transboundary acidification and eutrophication and ground level ozone in Europe: Unified EMEP Model Description, EMEP/MS-CW Report EMEP Status Report 1/2003 Part I, The Norwegian Meteorological Institute, Oslo, Norway, 2003.
- Slinn, S. A. and Slinn, W. G. N.: Predictions for particle deposition on natural waters, *Atmos. Environ.*, 24, 1013–1016, 1980.
- Smit, H. G. J., Straeter, W., Johnson, B., Oltmans, S., Davies, J., Tarasick, D. W., Hoegger, B., Stubi, R., Schmidlin, F., Northam, T., Thompson, A., Witte, J., Boyd, I., and Posny, F.: Assessment of the performance of ECC-ozone sondes under quasi-flight conditions in the environmental simulation chamber: Insights from the Juelich Ozone Sonde Intercomparison

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Experiment (JOSIE), J. Geophys. Res., 112, D19306, doi:10.1029/2006JD007308, 2007.
- Stiller, G. P. (Ed.) with contributions from v. Clarmann, T., Dudhia, A., Echle, G., Funke, B., Glatthor, N., Hase, F., Höpfner, M., Kellmann, S., Kemnitzer, H., Kuntz, M., Linden, A., Linder, M., Stiller, G. P., and Zorn, S.: The Karlsruhe Optimized and Precise Radiative Transfer Algorithm (KOPRA), vol. FZKA 6487 of Wissenschaftliche Berichte, Forschungszentrum Karlsruhe, Germany, 11448, 2000.
- Stockwell, W. R., Kirchner, F., Khun, M., and Seefeld, S.: A new mechanism for regional atmospheric chemistry modelling, J. Geophys. Res., 102, 25847–25879, 1997.
- Stohl, A., Wernli, H., James, P., Bourqui, M., Forster, C., Liniger, M. A., Seibert, P., and Sprenger, M.: A new perspective of stratosphere-troposphere exchange, B. Am. Meteorol. Soc., 84, 1565–1576, 2003.
- Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2, Atmos. Chem. Phys., 5, 2461–2474, doi:10.5194/acp-5-2461-2005, 2005.
- Szopa, S., Foret, G., Menut, L., and Cozic, A.: Impact of large scale circulation on European summer surface ozone: consequences for modelling forecast, Atmos. Environ., 43, 1189–1195, 2009.
- Tarasick, D. W., Moran, M. D., Thompson, A. M., Carey-Smith, T., Rochon, Y., Bouchet, V. S., Gong, W., Makar, P. A., Stroud, C., Ménard, S., Crevier, L.-P., Cousineau, S., Pudykiewicz, J. A., Kallaur, A., Moffet, R., Ménard, R., Robichaud, A., Cooper, O. R., Oltmans, S. J., Witte, J. C., Forbes, G., Johnson, B. J., Merrill, J., Moody, J. L., Morris, G., Newchurch, M. J., Schmidlin, F. J., and Joseph, E.: Comparison of Canadian air quality forecast models with tropospheric ozone profile measurements above midlatitude North America during the IONS/ICARTT campaign: Evidence for stratospheric input, J. Geophys. Res., 112, D12S22, doi:10.1029/2006JD007782, 2007.
- Thouret, V., Marenco, A., Logan, J., Nédélec, P., and Grouhel, C.: Comparisons of ozone measurements from the MOZAIC airborne program and the ozone sounding network at eight locations, J. Geophys. Res., 103, 25695–25720, 1998.
- Troen, I. and Mahrt, L.: A simple model of the atmospheric boundary layer: Sensitivity to surface evaporation, Bound.-Layer Meteorol., 37, 129–148, 1986.
- Van Leer, B.: Towards the ultimate conservative difference scheme. A second-order sequel to Godunov's method, J. Comput. Phys., 32, 101–136, 1979.
- Van Loon, M., Vautard, R., Schaap, M., Bergström, R., Bessagnet, B., Brandt, J., Builtjes, P. J.

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- H., Christensen, J. H., Cuvelier, C., Graff, A., Jonson, J. E., Krol, M., Langner, J., Roberts, P., Rouil, L., Stern, R., Tarrasón, L., Thunis, P., Vignati, E., and White, L.: Evaluation of long-term ozone simulations from seven regional air quality models and their ensemble, *Atmos. Environ.*, 41, 2083–2097, 2007.
- 5 Vautard, R., Builtjes, P. H. J., Thunis, P., Bessagnet, B., Brandt, J., Builtjes, P. J. H., Christensen, J. H., Cuvelier, C., Foltescu, V., Graff, A., Kerschbaumer, A., Krol, M., Roberts, P., Rouil, L., Stern, R., Tarrason, L., Thunis, P., Vignati, E., and Wind, P.: Evaluation and intercomparison of Ozone and PM₁₀ simulations by several chemistry transport models over four European cities within the CityDelta project, *Atmos. Environ.*, 41, 173–188, 2007.
- 10 Visschedijk, A. J. H., Zandveld, P. Y. J., and Denier van der Gon, H. A. C. A.: High resolution gridded European database for the EU Integrate Project GEMS, TNO-report 2007-A-R0233/B, 2007.
- Walcek, C. J. and Taylor, G. R.: A Theoretical Method for Computing Vertical Distributions of Acidity and Sulfate Production within Cumulus Clouds, *J. Atmos. Sci.*, 43, 339–355, 1986.
- 15 Wesely, M. L.: Parameterization of Surface Resistances to Gaseous Dry Deposition in Regional-Scale Numerical Models, *Atmos. Environ.*, 23, 1293–1304, 1989.
- West, J., Szopa, S., and Hauglustaine, D. A.: Human mortality effects of future concentrations of tropospheric ozone, *C.R. Geosci.*, 339, 775–783, 2007.
- Williamson, D. L. and Rash, P. J.: Two-dimensional semi-lagrangian transport with shape-preserving interpolation, *Mon. Weather Rev.*, 117, 102–129, 1989.
- 20 Worden, H. M., Logan, J. A., Worden, J. R., Beer, R., Bowman, K., Clough, S. A., Eldering, A., Fisher, B. M., Gunson, M. R., Herman, R. L., Kulawik, S. S., Lampel, M. C., Luo, M., Megretskaia, I. A., Osterman, G. B., and Shephard, M. W.: Comparisons of Tropospheric Emission Spectrometer (TES) ozone profiles to ozonesondes: Methods and initial results, *J. Geophys. Res.*, 112, D03309, doi:10.1029/2006JD007258, 2007.
- 25 Zhang, G. J. and McFarlane, N. A.: Sensitivity of climate simulations to the parameterization of cumulus convection in the Canadian climate centre general circulation model, *Atmos. Ocean*, 33, 407–446, 1995.
- Zhang, L., Brook, J. R., and Vet, R.: A revised parameterization for gaseous dry deposition in air-quality models, *Atmos. Chem. Phys.*, 3, 2067–2082, doi:10.5194/acp-3-2067-2003, 2003.
- 30

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Geographical characteristics of the sounding sites as well as the number of profiles available for the study. The sondes used in this paper are taken from two archives, namely (1) the World Ozone and Ultraviolet Data Center (WOUDC) (<http://www.woudc.org>) and (2) NILU's Atmospheric Database for Interactive Retrieval (NADIR) at Norsk Institutt for Luftforskning (NILU) (<http://www.nilu.no/nadir/>).

Sounding site	Country	Geographic coordinates	Altitude (meters)	# of profiles	Archive
DeBilt	The Netherlands	52.1° N, 5.18° E	4.0	15	NADIR
Legionowo	Poland	52.4° N, 20.97° E	96.0	17	WOUDC
Payerne	Switzerland	46.8° N, 6.95° E	491.0	53	NADIR
Uccle	Belgium	50.8° N, 4.35° E	100.0	13	NADIR
Hohenpeissenberg	Germany	47.8° N, 11° E	976.0	25	NADIR
Lerwick	Shetland	60.14° N, 1.19° W	82.0	13	NADIR
Sodankylä	Finland	67.37° N, 26.63° E	179.0	15	NADIR
Valentia	Ireland	51.93° N, 10.25° W	14.0	12	WOUDC
Barajas	Spain	40.47° N, 3.58° W	631.0	12	WOUDC

Table 2. Main characteristics of the RAQ and MOZART-IFS models as used for this study.

	MOZART-3 Kinnison et al. (2007)	MOCAGE Josse et al. (2004) Bousserze et al. (2007)	EURAD-IM Elbern et al. (2007)	CHIMERE Bessagnet et al. (2008)	CAMx Morris et al. (2003)	BOLCHEM Mircea et al. (2008)
Resol. (lon/lat) nlev (<200 hPa) toplev	1.9°×1.9° 24 0.1 hPa	0.2°×0.2° 23 5 hPa	0.4°×0.4° 23(20) 100 hPa	0.5°×0.5° 17 200 hPa	0.3°×0.3° 15 300 hPa	0.4°×0.4° 33 200 hPa
Meteorology Chem. BC	IFS(1 h) –	IFS(3 h) MOCAGE-CTM	MM5/IFS(3 h) MOZART-IFS	IFS(3 h) MOZART-IFS	MM5/IFS MOZART-IFS	On-line (IFS) MOZART-IFS
Emis. anthrop. natural	RETRO Schültz (2009) GFED Randerson (2006) Emmons et al. (2009)	TNO* Guenther et al. (1995) Dentener et al. (2004)	TNO* Guenther et al. (1993)	TNO* Guenther et al. (2006)	TNO* Poupkou et al. (2010)	TNO* Poupkou et al. (2010)
Chemistry	JPL-03/06 Kinnison et al. (2007) SO _x /NH ₃ /NH ₄	RACM Stockwell et al. (1997) + REP>ROBUS Lefèvre et al. (1994)	RACM-MIM Geiger et al. (2003)	MELCHIORII Schmidt et al. (2001)	CBM-IV+ updates Gery et al. (1989) Carter (1996)	SAPRC90 Carter (1990)
Advection	Lin and Rood (1996) Zhang and McFarlane (1995)	semi-lagrangian Williamson and Rasch (1989)	Bott (1989)	Collela and Woodward (1984) Van Leer et al. (1979)	Collela and Woodward (1984)	WAF Billet and Toro (1997)
Convection	Hack (1994)	Bechtold et al. (2001)	Walcek and Taylor (1986)	Tiedtke et al. (1989)	Kain-Fritsch 2 Kain (2002)	Kain and Fritsch (1990)
Diffusion	Holstlag and Boville (1993)	Louis (1979)	Blackadar (1978)	Troen and Mahrt (1986)	K-theory, coeff. MM5 Hong and Pan (1996)	Louis (1979)
Dry deposition	Wesely (1989)	Michou and Peuch (2002)	Zhang (2003)	Zhang (2003)	Wesely (1989) Slinn and Slinn (1980)	Simpson (2003)

* Visschedijk et al. (2007)

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**3-D evaluation of
tropospheric ozone
simulations**

D. Zyryanov et al.

Table 3. Mean ozone concentrations (ppb) and standard deviation have been calculated (over the 3 month period of summer 2008) for observations and models (forced by MOZART-IFS) for all profiles and for 3 altitude ranges: (1) 0–2 km; (2) >2–8 km; >8–10 km. Corresponding, mean bias, RMSE and Pearson's correlation coefficient are also displayed.

Altitude range	0–2 km		>2–8 km		>8–10 km	
# profiles: 395	Obs.	Model	Obs.	Model	Obs.	Model
Mean (ppb)	46.5	43.1	66.1	55.5	105.6	106.3
Std deviation	12.4	11.3	16.7	16.0	57.2	50.4
Bias (ppb)	–3.4 (–7 %)		–10.6 (–16 %)		0.74 (<1 %)	
Rmse (ppb)	10.1 (22 %)		21.3 (32 %)		55.8 (53 %)	
Correlation	0.71		0.47		0.45	
Variability ratio	0.91		0.96		0.88	

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Table 4. The simulation has been divided in 4 quarters NW (North-West), NE (North-East), SW (South-West) and SE (South-East). The temporal evolution of 0–6 km ozone partial columns from IASI and models are compared (for each quarter) in terms of their relative bias and Pearson's correlation. MEDIAN-IFS stand for the median of models using MOZART-IFS as boundary condition. CHIMERE-IFS is one of these models and is compared with CHIMERE-CLIM that is using the monthly mean of the MOZART-IFS hourly values as BC. Biases are express in DU.

	MEDIAN-IFS		MOCAGE		CHIMERE-IFS		CHIMERE-CLIM	
	bias	corr	bias	corr	bias	corr	bias	corr
NW	−1.5(−8 %)	0.68	2.6(11 %)	0.65	0.64(3 %)	0.68	0.4(2 %)	0.54
NE	−0.95(−5 %)	0.74	3.3(14 %)	0.67	1.8(8 %)	0.81	1.5(7 %)	0.66
SW	−2.9(−16 %)	0.63	−0.5(−2 %)	0.5	−1.6(−8 %)	0.62	−1.3(−7 %)	0.42
SE	−3.1(−16 %)	0.67	−0.6(−3 %)	0.58	−1.1(−5 %)	0.69	−0.5(−3 %)	0.54

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.




Fig. 1. Measurement sites of ozone profile over Europe. Red circles denote the location of balloon sounding sites. Green triangles indicate airports used in the frame of the MOZAIC program for summer 2008. Also indicated, centres of the horizontal grid (with 0.5° resolution) on which output from different models is projected.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.




Fig. 2. Typical averaging kernels on the vertical grid used for retrieval (and model evaluation) and spaced by 1 km height vertical layers (left panel) and in partial column space (right panel), the black curve displays the averaging kernel corresponding to the 0–6 km ozone partial columns and the red one shows the same for the 6–12 km ozone partial column.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 3. Vertical profiles (y-axis in km) of normalized **(a)** relative bias (Model minus Observations), **(b)** relative RMSE and **(c)** Pearson's correlation coefficients. Values are averaged over 1km vertical layers using all 395 available profiles of ozone. Black curves indicate the medians of the six models using IFS-MOZART at their boundaries (i.e. BOLCHEM, CAMX, CHIMERE, EURAD) and IFS-MOZART itself. Each value is also associated to the minimum and maximum value of the ensemble (bars). Results for MOCAGE (green), CHIMERE-IFS (full red) and CHIMERE.Clim (dashed red) (i.e. CHIMERE using a monthly climatology derived from IFS-MOZART as BC) are also plotted.

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.




Fig. 4. 0–6 km smoothed ozone partial columns (in Dobson Unit) averaged over the summer 2008 (JJA). The IASI columns are calculated with observations corresponding to the morning passage of the satellite. The “IASI counts” map indicates the number of (non cloudy) measurement days available during the period.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.




Fig. 5. 0–6 km raw ozone partial columns (in Dobson Unit) averaged over the summer 2008 (JJA).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 6a. Summer 2008 average differences (in DU) between smoothed 0–6 km ozone columns: model median minus IASI.

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



summer.2008
MOCAGE – IASI




Fig. 6b. The same as Fig. 6a: MOCAGE minus IASI.

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 7. Time evolution of 0–6 km partial columns (in Dobson Unit) for IASI (red), the smoothed median model (black) and the MOCAGE model. The Max and Min of the ensemble are also plotted as error cloud around the median. In each case, daily values (i.e. value at about 10:00 a.m.) are averaged over the considered geographical domain: North-Western Europe (upper left quarter of the model domain), South-Western Europe (bottom left), North-Eastern quarter of the domain (upper right), South-Eastern quarter of the domain (bottom right).

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.




Fig. 8. Time evolution of 0–6 km ozone partial columns (in Dobson Units) as observed by IASI (upper panel), and simulated by models for the 8 to 11 June 2008 period. The median value over all participating models is taken. For models both smoothed (middle panel) and raw (lower panel) columns are displayed.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 9. Contour map of Potential vorticity (at 330 K) obtained from ECMWF analysis for: (1) the 8 June 2008 at 12:00 UTC (upper panel); (2) the 9 June 2008 at 12:00 UTC (middle panel); (3) the 10 June 2008 at 12:00 UTC (lower panel).

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3-D evaluation of tropospheric ozone simulations

D. Zyryanov et al.




Fig. 10. Vertical cross section through IASI ozone concentrations (in ppb) and different model simulations at 48° N for 8 June, 10:00 GMT. The black box on the IASI cross section (upper panel) indicates roughly the location (at this latitude) of the corresponding MOZAIC flight for which measurements are presented in Fig. 10.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)