



Corrigendum to

“Identification of humic-like substances (HULIS) in oxygenated organic aerosols using NMR and AMS factor analyses and liquid chromatographic techniques” published in Atmos. Chem. Phys., 14, 25–45, 2014

M. Paglione¹, A. Kiendler-Scharr², A. A. Mensah^{2,*}, E. Finessi^{1,**}, L. Giulianelli¹, S. Sandrinelli¹, M. C. Facchini¹, S. Fuzzi¹, P. Schlag², A. Piazzalunga³, E. Tagliavini⁴, J. S. Henzing⁵, and S. Decesari¹

¹National Research Council (CNR), Institute of Atmospheric Sciences and Climate (ISAC), Bologna, Italy

²Forschungszentrum Jülich, IEK-8: Troposphere, Jülich, Germany

³University of Milano Bicocca, Department of Environmental, and territorial Science and University of Milan, Department of Chemistry, Milan, Italy

⁴Centro Interdipartimentale di Ricerca per le Scienze Ambientali (CIRSA), University of Bologna, Ravenna, Italy

⁵TNO, Netherlands Applied Research Organisation, Utrecht, the Netherlands

* now at: ETH Zurich, Institute for Atmospheric and Climate Science, 8092 Zurich, Switzerland

** now at: University of York, Department of Chemistry, Heslington, York, YO10 5DD, UK

Correspondence to: M. Paglione (m.paglione@isac.cnr.it)

In the paper “Identification of humic-like substances (HULIS) in oxygenated organic aerosols using NMR and AMS factor analyses and liquid chromatographic techniques” by M. Paglione et al., a wrong version of Fig. 1 was published. Please find the correct figure below.




Fig. 1. Average PM₁-filters' chemical composition and back-trajectory analysis for the four periods of the campaign. Organics are reported as total organic matter (OM) calculated by the relationship OM = OC_{PM₁} filters · 1.6.