



**The ACCENT model
evaluation protocol**

V. Grewe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[I◀](#)

[▶I](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



The ACCENT-protocol: a framework for benchmarking and model evaluation

V. Grewe¹, N. Moussiopoulos², P. Builtjes^{3,4}, C. Borrego⁵, I. S. A. Isaksen⁶, and A. Volz-Thomas⁷

¹Deutsches Zentrum für Luft- und Raumfahrt, Institut für Physik der Atmosphäre, Oberpfaffenhofen, Germany

²Department of Mechanical Engineering of the Aristotle University Thessaloniki, Thessaloniki, Greece

³TNO Environment and Geosciences, Utrecht, The Netherlands

⁴Institut für Meteorologie, Freie Universität Berlin, Germany

⁵Department of Environment an Planning, University of Aveiro, Portugal

⁶Center for International Climate and Environmental Research (CICERO), Oslo, Norway

⁷Institut für Energie- und Klimaforschung: Troposphäre, Forschungszentrum Jülich, Germany

Received: 28 November 2011 – Accepted: 7 December 2011 – Published: 13 December 2011

Correspondence to: V. Grewe (volker.grewe@dlr.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Abstract

We summarise results from a workshop on “Model Benchmarking and Quality Assurance” of the EU-Network of Excellence ACCENT, including results from other activities (e.g. COST Action 732) and publications. A formalised evaluation protocol is presented, i.e. a generic formalism describing the procedure how to perform a model evaluation. This includes eight steps and examples from global model applications are given for illustration. The first and important step is concerning the purpose of the model application, i.e. the addressed underlying scientific or political question. We give examples to demonstrate that there is no model evaluation per se, i.e. without a focused purpose. Model evaluation is testing, whether a model is fit for its purpose. The following steps are deduced from the purpose and include model requirements, input data, key processes and quantities, benchmark data, quality indicators, sensitivities, as well as benchmarking and grading. We define “benchmarking” as the process of comparing the model output against either observational data or high fidelity model data, i.e. benchmark data. Special focus is given to the uncertainties, e.g. in observational data, which have the potential to lead to wrong conclusions in the model evaluation if not considered carefully.

1 Introduction

The European Network of Excellence (NoE) ACCENT (Atmospheric Composition Change: the European Network; <http://www.accent-network.org>) had the goals “to promote a common European strategy for research on atmospheric composition change, to develop and maintain lasting means of communication and collaboration within the European scientific community, as well as to facilitate this research and optimise the interactions with policy-makers and the general public.” (ACCENT, 2009). In the past, there were many efforts from the scientific community to answer questions relevant to society, which focus on different topics and also scales. Examples (among many

GMDD

4, 3599–3621, 2011

The ACCENT model evaluation protocol

V. Grewe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**The ACCENT model
evaluation protocol**V. Grewe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

others) for questions with a global perspective are: “When will the ozone layer re-
cover?” (WMO, 2011) or “How large is the climate impact from air traffic?” (Penner
et al., 2009; Lee et al., 2010). For more regional aspects the question “How much
and for what reasons have background ozone levels in Europe changed during the last
5 decades?” (ACCENT, 2006; Parrish et al., 2009) may serve as an illustrative exam-
ple, which was one of many topics of the ACCENT project. Here, we concentrate on
global aspects, only, for simplicity reasons. These questions have been addressed us-
ing observational data in combination with modelling data, which both are affected by
uncertainties and errors. The assessment of the reliability of the answers relies very
10 much on the data quality and hence quality assurance, both for the benchmark data
and the models involved. In this paper we focus on a model evaluation protocol in a
generic form, which comprises previous model evaluations and summarises them as a
framework and general strategy for future efforts. This protocol has been emerged from
an ACCENT-workshop on “Model Benchmarking and Quality Assurance”, held in Thes-
15 saloniki in 2006 (Moussiopoulos and Isaksen, 2007), and is largely based on previous
activities (e.g. COST Action 723 “Quality Assurance of Microscale Meteorological Mod-
els”; <http://www.mi.uni-hamburg.de/Home.484.0.html>; Britter and Schatzmann, 2007a)
and publications (Schlesinger et al., 1979; AIAA, 1998). Although the described pro-
20 tocol is generic and applicable to all scales, we focus on examples from global models
for highlighting the different aspects.

The starting point of the protocol and every model evaluation is the formulation of its
purpose, that is the overall scientific or political question which is aimed to be answered
with the help of model simulations. Once this question has been clearly formulated, a
number of implications for the evaluation follow, concerning the model itself, as well as
25 the observational data required and the comparative analysis. This is formulated as
a framework, i.e. an evaluation protocol, which is outlined in Fig. 1 and discussed in
more detail in Sect. 2.

2 The ACCENT model evaluation protocol

A brief overview on the ACCENT model evaluation protocol is given in Fig. 1, including 8 topics, which are briefly laid out with a question and more deeply described in the following sections.

2.1 The purpose of the evaluation

The most important issue and first step is to be clear about the purpose of the evaluation. What is the overall question, which is aimed to be answered? The purpose of the evaluation has significant implications on the details of the model evaluation. In the Introduction we have raised two questions “Q1: When will the ozone layer recover?” and “Q2: How large is the climate impact from air traffic?” (Table 1). Both topics are addressed by chemistry-climate models, although, the model requirements differ significantly. For example, the inclusion of stratospheric polar clouds is a model requirement for question Q1, since they play an important role for the ozone hole (Crutzen and Arnold, 1986), whereas the formation of contrail cirrus is a model requirement for question Q2 (Burkhardt and Kärcher, 2011). Noteworthy, the opposite is not true. Contrail cirrus are not regarded to be an important process for stratospheric ozone and polar stratospheric clouds are not relevant for subsonic air traffic. Hence, the set-up of a flexible or modular model will be chosen differently for these two purposes.

Evidently, these examples show that a model evaluation necessarily requires a purpose, which implies that there is no general model evaluation per se.

2.2 The model requirements to fulfill the purpose

As discussed in the last section, the purpose leads to a number of model requirements for implemented processes, i.e. those related to the formation of polar stratospheric clouds for Q1 and the microphysics of contrail cirrus for Q2. The implemented processes are, however, only part of the model requirements, which can be summarized by

GMDD

4, 3599–3621, 2011

The ACCENT model evaluation protocol

V. Grewe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



be evaluated. A discrepancy between model data and observational data hints to a model deficiency only if this discrepancy exceeds the model's sensitivity to input data uncertainties. In many cases it is difficult to actually distinguish a model deficiency from a deficiency in the input data. Therefore, an evaluation protocol should include:

1. list of input data;
2. uncertainties of input data.

The sensitivities are important and hence covered separately in Sect. 2.7.

2.4 Key processes and quantities with respect to the purpose

A key part of the protocol is the description of the parameters, quantities or processes, which are important with respect to the purpose of the evaluation. In many cases this part of the protocol is best documented and reviewed. Concerning the evolution of the ozone layer (Q1) a SPARC-activity (Stratospheric Processes And their Role in Climate) was set-up to evaluate climate-chemistry models SPARC CCMVal (2010). A large number of quantities were identified and summarised in topics like radiation, dynamics, transport, stratospheric chemistry and microphysics.

There is a huge variety of possible parameters, e.g. temperatures, wind, ozone, nitrogen oxides, etc. and representations of, e.g. long-term monthly means or medians, both for a region and for a certain altitude, seasonal cycles, variabilities, etc. This representation of the data is important, since the quality indicators and the benchmark depend on them (Sects. 2.6 and 2.8).

A further key aspect concerns the counterpart to the model data, i.e. the benchmark data used in the evaluation (Sect. 2.5). Generally the most favourable are observational data. However, high fidelity models are also frequently used, e.g. high resolution line-by-line radiation codes to evaluate radiative fluxes (SPARC CCMVal, 2010). In many cases observational data are not available in the required representation. It is important to indicate (a) the need of these observational data and (b) to show the range

The ACCENT model evaluation protocol

V. Grewe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of simulated data. Even without benchmark data, a model intercomparison still reveals the range of uncertainty and might foster research in this direction.

Therefore the key parameters should be characterised by:

1. list of key parameters, quantities and processes;
2. representation;
3. description of benchmark data.

2.5 Benchmark data

As discussed in Sect. 2.4 the benchmark data, i.e. the data against which the model data are compared, are either observational data or high fidelity model data. It is essential for the benchmarking (Sect. 2.8) to include all available quality information for the observational data, in order to provide a quantitative estimate of their overall uncertainty, including

1. measurement techniques (accuracy and precision);
2. methodology;
3. representativity;
4. natural variability.

The uncertainties from the measurement techniques themselves are in most cases well documented. More difficult is to derive an estimate of uncertainties arising from the applied methodologies, such as retrieval algorithms, which convert a directly measured quantity into the targeted physical or chemical quantity. Van Noije et al. (2006) have presented results from a model intercomparison of tropospheric NO₂ columns, in comparison to three different retrieval algorithms. Their results show that the spread between the model results is basically as large as the spread between the observational data. The inclusion of different retrieval algorithms provided an indication of the

The ACCENT model evaluation protocol

V. Grewe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**The ACCENT model
evaluation protocol**

V. Grewe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

uncertainties associated with the methodology applied to the satellite data used as benchmark. Without these uncertainties, the assessment of the model results would have been very different: Taking only the Bremen satellite data into account suggests a model underestimate of a factor of two of the wintertime European NO₂ columns.

5 Whereas taking all observational data into account leads to the conclusion that observational data and model data do not show any statistical significant difference.

The representativity of the benchmark data is also a crucial point. For example in-situ measurements are often located at prominent locations, like mountain tops or at the sea side and are affected by the local environment. The local scale of such observations makes it difficult to assure the representativity of the measurement for the grid box volume. Other aspects of representativity are associated with sampling methods (e.g. Jöckel et al., 2010) and again retrievals. How accurate are height specifications of vertically resolved satellite data, i.e. how representative are satellite measurements for a certain height when the information is deduced from a column value and represents a certain height region? How representative are measurements of, e.g. tropospheric NO₂ columns, when clouds shield a large fraction of lower tropospheric NO₂. Richter and Burrows (2002) have investigated these effects and showed that they can lead to potentially large uncertainties.

Another example are HALOE (Halogen Occultation Experiment; Russel III et al., 1993) satellite measurements of e.g. HCl. Lary and Aulov (2008) showed a comparison of frequency distributions of satellite HCl measurements for a height region in isentropic levels and equivalent latitude bands representative for January conditions in the early 2000s. For some regions the difference between the individual measurement platforms was larger (up to 20%) than the variability within one individual measurement platform, which shows that a bias can be significantly larger than the variability. Aghedo et al. (2011) compared chemistry-climate model output with satellite observations. This comparison was performed in two ways: (a) by comparing the model output directly with the satellite data and (b) by processing the model output in a way the satellite would have observed the model's atmosphere (sampling, satellite processing). By

**The ACCENT model
evaluation protocol**

V. Grewe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the latter procedure they derived data which were based on observational operators representing the satellite retrieval algorithms. They concluded that for most species the sampling would have a low error, but neglecting observational operators have an impact for, e.g. ozone and water vapour on the order of 30% and 100%, respectively.

5 These examples show that uncertainties directly associated with the compilation of these examples show that uncertainties directly associated with the compilation of observational data, i.e. points data, i.e. points (1) to (3) (see beginning of this section), severely impact on the quality of the data. While this does not devalue the data, it places great demands on the way the data are to be compared, i.e. the quality indicators used and their statistical interpretation (see Sect. 2.6). The determination of these uncertainties is indeed often challenging. However, to disregard the uncertainty rather than to use an educated guess or even a rough estimate is misjudging the consequences. Neglecting the uncertainties in the data implies the assumption of zero uncertainty, which is in general worse than using a rough estimate. For this reason, it is recommended to better rely on expert judgment than to disregard uncertainty.

15 The last point (4) is associated with natural variability and representativity of the data. If the targeted quantity is, e.g. a climatological mean mixing ratio for winter-time northern polar ozone at 50 hPa, a large interannual variability will lead to a large confidence interval for the climatological mean mixing ratio derived from the measurements, depending on sampling statistics. On the other hand, significant deviations between model and reality can only be claimed if the differences between model and observational data are smaller than this confidence interval. Again this is a matter of the quality indicators (see Sect. 2.6) which, if chosen in a more appropriate way, might have reduced the statistical limitations.

25 2.6 Quality indicators

Quality indicators, i.e. the way model data and benchmark data are compared, have a large impact on the outcome of the evaluation. The list of possibilities is endless and ranges from a simple intercomparison of climatological mean values to RMSE (root

mean square error), NMSE (normalized mean square error), correlation coefficients, or so-called Taylor diagrams (Taylor, 2001), which compare the normalised standard deviation versus the correlation coefficient. However, there are requirements to the quality indicators, which mainly arise from the previous sections: quality indicators should

1. resemble the representation of the key parameters (point 2 in Sect. 2.4);
2. include the uncertainties from benchmark data (Sect. 2.5);
3. include statistical tests on the significance of differences or
4. include confidence intervals for the quality indicators.

Therefore, although there are many possibilities for quality indicators, the definition of the key parameters, given in Sect. 2.4 largely constrains them. For example, if in Sect. 2.4 a seasonal cycle is identified as a key parameter, then a parametric approach (Eq. 1) or the RMSE between deviations from the annual means (Eq. 2) can be taken into account.

$$f_1(t) = a \sin(bt - c) \quad (1)$$

$$f_2^2 = \sum_{i=1}^{12} [(x_{\text{mod}}(i) - x_{\text{mod}}^{\text{ann}}) - (x_{\text{obs}}(i) - x_{\text{obs}}^{\text{ann}})]^2, \quad (2)$$

where a , b , c are parameters, which have to be fitted to either data set, $x_{\text{mod}}(i)$ and $x_{\text{obs}}(i)$ monthly mean values and $x_{\text{mod}}^{\text{ann}}$ and $x_{\text{obs}}^{\text{ann}}$ the respective annual means. In the first case differences in the fitted parameters a , b , and c between model and observational data would be the quality indicator and in the second case f_2 is directly the quality indicator. The parametric approach f_1 might provide more information, i.e. on amplitude, period and phase.

From the previous Sects. 2.3, 2.5 it is clear that the use of the data, as they are, will very likely lead to misinterpretations, because the differences might arise from

The ACCENT model evaluation protocol

V. Grewe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



uncertainties in the data rather than from model deficiencies. Britter and Schatzmann (2007b) defined the total uncertainty of the model evaluation as a combination of uncertainties in the input data, benchmark data and natural variability (Fig. 3). This requires a calculation of confidence intervals or a statistical testing of the quality indicators. In the example above, this implies a test whether the fitted parameters a , b , and c differ statistically significantly for the model and observational data. In the second example, a Monte-Carlo simulation could provide an uncertainty range and a confidence interval for f_2 . This Monte-Carlo simulation replaces a complicated analytical calculation of the confidence interval and converts uncertainty ranges in the observational data into confidence interval for, e.g. f_2 .

2.7 Sensitivities

The application of the quality indicators including statistical tests leads to the conclusion, which parameters are simulated with deficiencies. This has an impact on the quality with which the overall question, i.e. the purpose of the model (Sect. 2.1), can be answered. Hence it is important to further understand these deficiencies and to provide a better understanding of the simulated processes. This can be achieved by additional simulations or diagnostics, which provide information on the sensitivity of the results to e.g. input data. Again a wide range of examples exists in the literature. For example concerning the impact of air traffic on climate (Q2), the question of the contribution of other NO_x sources (e.g. lightning, Grewe, 2007) is of interest as well as the impact of model resolution and domain, feedbacks, altitude of aircraft emissions (Grewe et al., 2002). A prominent difference between the climate-chemistry models is the ratio between the predicted response to air traffic emissions for methane and ozone (Lee et al., 2010). Stevenson et al. (2006) showed that although the methane lifetime was simulated very differently by individual models, the sensitivity to different scenarios was very similar among the models, which provides more confidence in the sensitivities than the absolute numbers. Within the “Model and Measurement Intercomparison II” (Park et al., 1999) idealised air traffic emission simulations were performed to investigate the

The ACCENT model evaluation protocol

V. Grewe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[I◀](#)

[▶I](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



impact of mid-latitude air traffic emissions in comparison to high latitude emissions, in order to provide a basis for the robustness in simulating the effect of mid-latitude air traffic and to pin down differences in model results. Hence, a sensitivity analysis is a step beyond an uncertainty assessment (Saltelli et al., 2000).

2.8 Benchmarking and grading

The evaluation protocol is a mixture of subjective choices and objective analysis. The choice of the key parameters and the corresponding quality indicator is based on expert knowledge and hence subjective. The analysis of the quality indicators itself is a mathematical procedure and hence objective. A statistical test which includes the uncertainties in the data shows whether the model contradicts reality. A statistical test which excludes uncertainties in the benchmark data only shows, whether or not the model output contradicts the benchmark data (e.g. Grewe and Sausen, 2009).

Benchmarking (Fig. 4), i.e. the quantitative comparative analysis with respect to a reference value, is again highly subjective, since it uses the quality indicators and quantifies and rates its outcome with respect to a reference value and sets a margin. If the quality indicator is within this margin, the model output is, based on current knowledge, within an acceptable range. The choice of this margin is again highly subjective and, in general, cannot be based on objective measures.

Grading goes one step further by translating the outcome of the quality indicators and the margin into standardised units, e.g. a mark between 0 (bad) to 1 (excellent). Again the choice of the key parameter, the quality indicator, the margin and the grading formula are purely subjective, thus leading to pseudo-objective metrics. This is not devaluating grades, but has to be taken into account when comparing grades from, e.g. different model evaluations.

Moreover, grades can be and have been combined to multi-model grades (mean grade of many models for one key parameter) or overall model grades (mean grade of all key parameter for one model). Multi model grades have the same basis, i.e. quality indicator and margin, and can be combined, whereas overall model grades are lacking

The ACCENT model evaluation protocol

V. Grewe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a common basis, i.e. an ozone grade and a temperature grade is not comparable, which limits the interpretation of this metric.

We have already stressed the importance to include a statistical test to the quality indicators in order to identify statistically significant differences between model prediction and reality deduced from benchmark data and based on uncertainties in the model input data, measurement data and natural variability. It is a challenge to transfer this into a reliable margin and grade. There is a substantial methodological difference between the mentioned test for quality indicators and for grades. The outcome of the test for quality indicators shows how much the model is differing from the available benchmark data, whereas grades include a much more quantitative aspect. The outcome of a test for grades includes, depending on the grading formula, that the model differs from the real atmosphere by a certain distance to the margin. The test for the quality indicator is directly linked to the test whether a grade is 1, whereas for the grades the test for lower values than 1 has to be performed in addition. Grewe and Sausen (2009) have provided an example, which transfers a widely used grading formula (Douglass et al., 1999; Kawa et al., 1999; Waugh and Eyring, 2008; SPARC CCMVal, 2010) into a statistically tested grading formula by calculating confidence intervals for the quality indicators and using its lower bound as a new margin for grading. Their results show that without such a transformation the grading results cannot be interpreted in a reasonable manner. To summarise, benchmarking and grading requires:

1. definition of a margin, within which quality indicators are indicating acceptable model results;
2. definition of a grading formula, which represents a statistical significant difference of model results from the margin and takes into account an estimate of the reality deduced from observations.

The ACCENT model evaluation protocol

V. Grewe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Summary

The aim of this study is to provide a framework for model evaluation, i.e. a model evaluation protocol. It was prepared during an ACCENT workshop held in Thessaloniki in May 2006 (Moussiopoulos and Isaksen, 2007) and further refined afterwards. Hence this protocol is based on input from a broad community including experts from observational and modelling communities, and covering expertise from local, regional and global aspects.

The protocol has 8 topics (Fig. 1), of which the first one is the most important, that is the purpose, the scientific or political question, which is aimed to be answered by the modelling activity. Other issues, like model requirements, model input data, key parameters and processes, etc., can then be deduced from the purpose, though subjectively based on expert knowledge.

It is important to stress the role of uncertainties in the evaluation. Britter and Schatzmann (2007b) defined the total uncertainty of quality indicators as a combination of uncertainties in the input data, benchmark data and natural variability. Especially the uncertainties in the benchmark data or observational data are crucial and include uncertainties of the measurements, the uncertainty in the methodology used for conversion of a measured physical quantity into the targeted atmospheric quantity, and the representativity of the observational data for the modelling counterpart (spatial/temporal representativity). Disregarding any of these uncertainties implies the assumption of a zero uncertainty and, hence, an expert judgment of the uncertainties should be used if independent information is not available.

The benchmarking itself, i.e. the quantitative comparative analysis with respect to a reference value, has to include statistical tests to ensure that the uncertainties of the quality indicator are not leading to misjudgment of the model performance.

As a result, two main conclusions can be formulated. A model evaluation demands the purpose of the model application and uncertainties in the observational data have to be included to be able to deduce a statistical significant deviation of the model from reality.

The ACCENT model evaluation protocol

V. Grewe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acknowledgements. This work was supported by the Network of Excellence ACCENT and the DLR-project ESMVal.

References

- 5 ACCENT (Atmospheric Composition Change the European Network of Excellence): Answers to the Urbino Questions, ACCENTs first policy-driven synthesis, edited by: Raes F. and Hjorth J., published by: ACCENT Secretariat, University Urbino, Italy, ISBN 92-79-02413-2, <http://www4.nilu.no/portal/publications/accent-series-reports>, 2006. 3601
- 10 ACCENT (Atmospheric Composition Change the European Network of Excellence): Editorial, edited by: Fuzzi, S. and Maione, M., *Atmos. Environm.*, 43, 5136–5137, 2009. 3600
- 10 Aghedo, A. M., Bowman, K. W., Shindell, D. T., and Faluvegi, G.: The impact of orbital sampling, monthly averaging and vertical resolution on climate chemistry model evaluation with satellite observations, *Atmos. Chem. Phys.*, 11, 6493–6514, doi:10.5194/acp-11-6493-2011, 2011. 3606
- 15 AIAA (American Institute of Aeronautics and Astronautics): Guide for the Verification and Validation of Computational Fluid Dynamics Simulations, American Institute of Aeronautics and Astronautics, AIAA-G-077-1998, Reston, VA, USA, 1998. 3601
- Braesicke, P. and Pyle, J.: Sensitivity of dynamics and ozone to different representations of SSTs in the Unified Model, *Q. J. Roy. Meteor. Soc.*, 130, 2033–2045, 2004. 3603
- Britter, R. and Schatzmann, M.: Model evaluation guidance and protocol document, COST Action 732, p. 28, ISBN 3-00-018312-4, Hamburg, Germany, 2007a.
- 20 Britter, R. and Schatzmann, M.: Background and justification document to support the model evaluation guidance and protocol, COST Action 732, p. 88, ISBN 3-00-018312-4, Hamburg, Germany, 2007b. 3609, 3612
- Burkhardt, U. and Kärcher, B.: Global radiative forcing from contrail cirrus, *Nature Climate Change*, 1, 54–58, 2011. 3602
- 25 Crutzen, P. J. and Arnold, F.: Nitric acid cloud formation in the cold Antarctic stratosphere: a major cause for the springtime “ozone hole”, *Nature*, 324, 651–655, 1986. 3602
- Douglass, A. R., Prather, M. J., Hall, T. M., Strahan, S. E., Rasch, P. J., Sparling, L. C., Coy, L., and Rodriguez, J. M.: Choosing meteorological input for the global modeling initiative assessment of high-speed aircraft, *J. Geophys. Res.*, 104, 27545–27564, 1999. 3611
- 30 Garny, H., Dameris, M., and Stenke, A.: Impact of prescribed SSTs on climatologies and

The ACCENT model evaluation protocol

V. Grewe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- long-term trends in CCM simulations, *Atmos. Chem. Phys.*, 9, 6017–6031, doi:10.5194/acp-9-6017-2009, 2009. 3603
- Grewe, V.: Impact of climate variability on tropospheric ozone, *Sci. Total Environ.*, 374, 167–181, 2007.
- 5 Grewe, V. and Sausen, R.: Comment on “Quantitative performance metrics for stratospheric-resolving chemistry-climate models” by Waugh and Eyring (2008), *Atmos. Chem. Phys.*, 9, 9101–9110, doi:10.5194/acp-9-9101-2009, 2009. 3611
- Grewe, V., Dameris, M., Fichter, C., and Sausen, R.: Impact of aircraft NO_x emissions. Part 1: Interactively coupled climate-chemistry simulations and sensitivities to climate-chemistry feedback, lightning and model resolution, *Meteorol. Z.*, 3, 177–186, 2002. 3603, 3609
- 10 Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., Tost, H., Riede, H., Baumgaertner, A., Gromov, S., and Kern, B.: Development cycle 2 of the Modular Earth Submodel System (MESSy2), *Geosci. Model Dev.*, 3, 717–752, doi:10.5194/gmd-3-717-2010, 2010.
- Kawa, S. R., Anderson, J. G., Baughcum, S. L., Brock, C. A., Brune, W. H., Cohen, R. C., Kinnison, D. E., Newman, P. A., Rodriguez, J. M., Stolarski, R. S., Waugh, D., and Wofsy, S. C.: Assessment of the Effects of High-Speed Aircraft in the Stratosphere: 1998, NASA-Report, NASA/TP-1999-209237, 1999. 3611
- 15 Lary, D. J. and Aulov, O.: Space-based measurements of HCl: Intercomparison and historical context, *J. Geophys. Res.*, 113, D15S04, doi:10.1029/2007JD008715, 2008. 3606
- 20 Lee, D. S., Pitari, G., Grewe, V., Gierens, K., Penner, J. E., Petzold, A., Prather, M. J., Schumann, U., Bais, A., Bernsten, T., Iachetti, D., Lim, L. L., and Sausen, R.: Transport impacts on atmosphere and climate: Aviation, *Atmos. Environm.* 44, 4678–4734, 2010. 3601, 3609
- Moussiopoulos, N. and Isaksen, I.: Proceedings of the Workshop on Model Benchmarking and Quality Assurance, ACCENT Series Reports 2.2007, 60 p., <http://www4.nilu.no/farcry-accent/download.cfm?DownloadFile=3DCD0697-17A4-8DB2-8C63CCDF7478BCB6>, 2007. 3601, 3612
- 25 Park, J. H., Ko, M. K. W., Jackman, C. H., Plumb, R. A., Kaye, J. A., and Sage, K. H.: M&M-2, NASA: Models and Measurements Intercomparison II. TM-1999-209554, 502 pp., 1999. 3609
- 30 Parrish, D. D., Millet, D. B., and Goldstein, A. H.: Increasing ozone in marine boundary layer inflow at the west coasts of North America and Europe, *Atmos. Chem. Phys.*, 9, 1303–1323, doi:10.5194/acp-9-1303-2009, 2009. 3601
- Penner, J. E., Lister, D. H., Griggs, D. J., Dokken, D. J., and McFarland, M.: Aviation and the

**The ACCENT model
evaluation protocol**

V. Grewe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**The ACCENT model
evaluation protocol**

V. Grewe et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- Global Atmosphere. A special report of IPCC Working Groups I and III, 373, Cambridge Univ. Press, Cambridge, UK, 1999. 3601
- Richter, A. and Burrows, J. P.: Tropospheric NO₂ from GOME measurements, *Adv. Space Res.*, 29, 1673–1683, 2002. 3606
- 5 Russell III, J. M., Gordley, L. L., Park, J. H., Drayson, S. R., Hesketh, D. H., Cicerone, R. J., Tuck, A. F., Frederick, J. E., Harries, J. E., and Crutzen, P.: The Halogen Occultation Experiment, *J. Geophys. Res.*, 98(D6), 10777–10797, 1993.
- Saltelli, A., Chan, K., and Scott, E. M.: *Sensitivity Analysis*, John Wiley & Sons, Chichester, New York, 2000. 3610
- 10 Schlesinger, S., Crosbie, R. E., Gagné, R. E., Innis, G. S., Lalwani, C. S., Loch, J., Sylvester, R. J., Wright, R. D., Kheir, N., and Bartos, D.: Terminology for Model Credibility, *Simulation*, 32(3), 103–104, 1979. 3601
- Stevenson, D. S., Dentener, F. J., Schultz, M. G., Ellingsen, K., van Noije, T. P. C., Wild, O., Zeng, G., Amann, M., Atherton, C. S., Bell, N., Bergmann, D. J., Bey, I., Butler, T., Cofala, J., Collins, W. J., Derwent, R. G., Doherty, R. M., Drevet, J., Eskes, H. J., Fiore, A. M., Gauss, M., Hauglustaine, D. A., Horowitz, L. W., Isaksen, I. S. A., Krol, M. C., Lamarque, J.-F., Lawrence, M. G., Montanaro, V., Müller, J.-F., Pitari, G., Prather, M. J., Pyle, J. A., Rast, S., Rodriguez, J. M., Sanderson, M. G., Savage, N. H., Shindell, D. T., Strahan, S. E., Sudo, K., and Szopa, S.: Multimodel ensemble simulations of present-day and near-future tropospheric ozone, *J. Geophys. Res.*, 111, D08301, doi:10.1029/2005JD006338, 2006. 3609
- 20 SPARC Report on the Evaluation of Chemistry-Climate Models: SPARC Report No. 5, edited by: Eyring, V., Shepherd, T. G., and Waugh, D. W., WCRP-132, WMO/TD-No. 1526, <http://www.atmosp.physics.utoronto.ca/SPARC>, 2010. 3604, 3611
- Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, *J. Geophys. Res.*, 106, 7183–7192, 2001. 3608
- 25 van Noije, T. P. C., Eskes, H. J., Dentener, F. J., Stevenson, D. S., Ellingsen, K., Schultz, M. G., Wild, O., Amann, M., Atherton, C. S., Bergmann, D. J., Bey, I., Boersma, K. F., Butler, T., Cofala, J., Drevet, J., Fiore, A. M., Gauss, M., Hauglustaine, D. A., Horowitz, L. W., Isaksen, I. S. A., Krol, M. C., Lamarque, J.-F., Lawrence, M. G., Martin, R. V., Montanaro, V., Müller, J.-F., Pitari, G., Prather, M. J., Pyle, J. A., Richter, A., Rodriguez, J. M., Savage, N. H., Strahan, S. E., Sudo, K., Szopa, S., and van Roozendaal, M.: Multi-model ensemble simulations of tropospheric NO₂ compared with GOME retrievals for the year 2000, *Atmos. Chem. Phys.*, 6, 2943–2979, doi:10.5194/acp-6-2943-2006, 2006. 3605
- 30

Waugh, D. W. and Eyring, V.: Quantitative performance metrics for stratospheric-resolving chemistry-climate models, *Atmos. Chem. Phys.*, 8, 5699–5713, doi:10.5194/acp-8-5699-2008, 2008. 3611

WMO: Scientific assessment of ozone depletion: 2010, World Meteorological Organisation, Geneva, 2011. 3601

5

GMDD

4, 3599–3621, 2011

The ACCENT model evaluation protocol

V. Grewe et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[I◀](#)

[▶I](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



GMDD

4, 3599–3621, 2011

**The ACCENT model
evaluation protocol**

V. Grewe et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)**Table 1.** Two questions serving as examples for the purpose of a model evaluation.

Abbreviation	Questions characterising the purpose of the evaluation
Q1	When will the ozone layer recover?
Q2	How large is the climate impact from air traffic?

Overview on the ACCENT model evaluation protocol

- **Purpose**
What is the scientific or political question ?
- **Model requirements**
What are the model requirements to answer the question ?
- **Input data**
What data are necessary to run the model ?
- **Key processes / quantities**
What are the key processes and quantities to be evaluated ?
- **Benchmark data**
What observations or high fidelity model data are required for the validation ?
- **Quality indicators**
How are model and observational data compared ?
- **Sensitivities**
What model sensitivities have to be investigated to understand the robustness of the answers to the question ?
- **Benchmarking & Grading**
What are the conclusions on the accuracy and robustness of the answer to the posed question, implied by the comparative analysis ?

Fig. 1. Overview on the ACCENT model evaluation protocol.

The ACCENT model evaluation protocol

V. Grewe et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 2. Sketch of an intercomparison of modelling data to observational data, illustrating the difficulties in attributing discrepancies to model deficiencies rather than input or boundary data uncertainties.

The ACCENT model evaluation protocol

V. Grewe et al.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[I◀](#) [▶I](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Fig. 3. Sketch, showing the components of model and observational data and the consequences for the uncertainties of the quality indicators.

The ACCENT model evaluation protocol

V. Grewe et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
I◀	▶I
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



The ACCENT model evaluation protocol

V. Grewe et al.




Fig. 4. Overview on the whole benchmarking procedure.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

