

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Canopy-scale flux measurements and bottom-up emission estimates of volatile organic compounds from a mixed oak and hornbeam forest in northern Italy

W. J. F. Acton¹, S. Schallhart², B. Langford³, A. Valach^{1,3}, P. Rantala², S. Fares⁴, G. Carriero⁵, R. Tillmann⁶, S. J. Tomlinson³, U. Dragosits³, D. Gianelle^{7,8}, C. N. Hewitt¹, and E. Nemitz³

¹Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK

²Division of Atmospheric Sciences, Department of Physics, University of Helsinki, Gustaf Hållströmin katu 2, 00560 Helsinki, Finland

³Centre for Ecology & Hydrology, Bush Estate, Penicuik, Midlothian, EH26 0QB, UK

⁴Research Centre for Soil-Plant System, Council for Agricultural Research and Economics, Rome, Italy

⁵Institute for Plant Protection CNR Research Area – Building E, Via Madonna del Piano 10, 50019 Sesto Fiorentino, Italy

⁶Institute of Energy and Climate Research, IEK-8: Troposphere, Research Centre Jülich, 52425 Jülich, Germany

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



⁷Sustainable Agro-ecosystems and Bioresources Department, Research and Innovation Centre, Fondazione Edmund Mach, 38010 S. Michele all'Adige, Italy

⁸Foxlab Joint CNR-FEM Initiative, Via E. Mach 1, 38010 San Michele all'Adige, Italy

Received: 24 September 2015 – Accepted: 3 October 2015 – Published: 27 October 2015

Correspondence to: E. Nemitz (en@ceh.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

This paper reports the fluxes and mixing ratios of biogenically emitted volatile organic compounds (BVOCs) 4 m above a mixed oak and hornbeam forest in northern Italy. Fluxes of methanol, acetaldehyde, isoprene, methyl vinyl ketone + methacrolein, methyl ethyl ketone and monoterpenes were obtained using both a proton transfer reaction-mass spectrometer (PTR-MS) and a proton transfer reaction-time of flight-mass spectrometer (PTR-ToF-MS) together with the methods of virtual disjunct eddy covariance (PTR-MS) and eddy covariance (PTR-ToF-MS). Isoprene was the dominant emitted compound with a mean day-time flux of $1.9 \text{ mg m}^{-2} \text{ h}^{-1}$. Mixing ratios, recorded 4 m above the canopy, were dominated by methanol with a mean value of 6.2 ppbv over the 28 day measurement period. Comparison of isoprene fluxes calculated using the PTR-MS and PTR-ToF-MS showed very good agreement while comparison of the monoterpene fluxes suggested a slight over estimation of the flux by the PTR-MS. A basal isoprene emission rate for the forest of $1.7 \text{ mg m}^{-2} \text{ h}^{-1}$ was calculated using the MEGAN isoprene emissions algorithms (Guenther et al., 2006). A detailed tree species distribution map for the site enabled the leaf-level emissions of isoprene and monoterpenes recorded using GC-MS to be scaled up to produce a “bottom-up” canopy-scale flux. This was compared with the “top-down” canopy-scale flux obtained by measurements. For monoterpenes, the two estimates were closely correlated and this correlation improved when the plant species composition in the individual flux footprint was taken into account. However, the bottom-up approach significantly underestimated the isoprene flux, compared with the top-down measurements, suggesting that the leaf-level measurements were not representative of actual emission rates.

ACPD

15, 29213–29264, 2015

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

The term volatile organic compound (VOC) describes a broad range of chemical species emitted from natural and anthropogenic sources into the atmosphere. VOCs emitted from the biosphere are commonly termed biogenic VOCs (BVOCs). Of the BVOCs isoprene is almost certainly the dominant species globally with an estimated annual emission of $535\text{--}578 \times 10^{12}$ g C (Arneth et al., 2008; Guenther et al., 2012). Isoprene, along with larger terpenoids, are the BVOCs that have received the most attention in the literature to date. Although isoprene is the most commonly measured BVOC global emission estimates continue to differ and there are still large uncertainties associated with the emission estimates of many other compounds. A better understanding of how emissions change with land cover, temperature, soil moisture and solar radiation is required to constrain model descriptions of the effects of BVOCs on atmospheric chemistry in the past, present and future (Monks et al., 2009).

BVOCs are a major source of reactive carbon into the atmosphere and as such exert an influence on both climate and local air quality. BVOCs are oxidised primarily by the hydroxyl radical (OH), itself formed by the photolysis of ozone, to form peroxide radicals (RO_2). In the presence of NO_x (NO and NO_2) these RO_2 radicals can oxidise NO to NO_2 , which may undergo photodissociation leading to the net formation of tropospheric ozone (Fehsenfeld et al., 1992). Tropospheric ozone can then impact upon human health, forest productivity and crop yields (Royal Society, 2008; Ashmore, 2005). In addition, BVOC species contribute significantly to the formation of secondary organic aerosol (SOA) in the atmosphere. This affects climate both directly and indirectly by the scattering of solar radiation and by acting as cloud condensation nuclei, increasing cloud cover and therefore altering the Earth's albedo (Hallquist et al., 2009).

The Bosco Fontana campaign was carried out as a part of the ÉCLAIRE (Effects of Climate Change on Air Pollution and Response Strategies for European Ecosystems) EC FP7 project to study the surface/atmosphere exchange within a semi-natural forest situated within one of the most polluted regions in Europe, and its interaction with air

ACPD

15, 29213–29264, 2015

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



chemistry. During the Bosco Fontana campaign, VOC fluxes and mixing ratios were measured 4 m above the canopy of a semi-natural forest situated in the Po Valley, northern Italy (45°11'51" N, 10°44'31" E), during June and July 2012. The Po Valley experiences high levels of anthropogenic pollution caused by its proximity to the city of Milan's high levels of industrial and traffic-related emissions of pollutants, intensive agriculture and periods of stagnant air flow caused by the Alps to the north and west and the Apennines to the south (Decesari et al., 2014). A 13 year study of atmospheric pollution in the Po Valley at a site in the town of Modena, approximately 65 km south of the Bosco Fontana nature reserve, recorded very high concentrations of ground level ozone, with average daily maximum concentrations in the summer peaking at ca. 120 $\mu\text{g m}^{-3}$ (Bigi et al., 2011). For comparison European legislation states that the daily 8 h mean should not exceed 120 $\mu\text{g m}^{-3}$.

Here we report the fluxes and mixing ratios of a range of BVOCs recorded from mixed mesophile forest at the Bosco Fontana field site. We compare BVOC flux calculation from above canopy eddy covariance measurements using both a proton transfer reaction-mass spectrometer (PTR-MS) and a proton transfer reaction-time of flight-mass spectrometer (PTR-ToF-MS) with isoprene and monoterpene fluxes obtained by scaling up leaf-level emission data to produce a canopy-scale "bottom-up" modelled flux estimate. We further explore the potential of accounting for the spatial tree species distribution for improving the comparison between top-down and bottom-up approaches.

2 Methods

2.1 Site description

Measurements were taken at a site within the Bosco Fontana natural reserve (45°11'51" N, 10°44'31" E), a 233 ha area of semi-natural woodland situated in the municipality of Marmirolo in the Po Valley. The forest canopy had an average height

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of approximately 28 m and was principally comprised of *Carpinus betulus* (hornbeam) and three oak species *Quercus robur* (pedunculate oak), *Quercus cerris* (turkey oak) and the introduced *Quercus rubra* (northern red oak) (Dalponte et al., 2007). In the centre of the forest there was a cleared area containing a seventeenth century hunting lodge surrounded by hay meadows. The surrounding area was predominantly arable farm land with some pastures to the north and west and a reservoir to the south west. The city of Mantova lies approximately 5 km to the south east, with the small towns of Marmirolo, Soave and Sant'Antonio approximately 2 km north, 1 km west and 3 km east, respectively. A 42 m measurement tower was situated near the centre of the forest to the south west of the central hay meadows. The measurement tower was ca. 760 m from the edge of the forest in the direction of the easterly wind direction that dominated during this measurement period.

2.2 PTR-MS and PTR-ToF-MS setup and measurement procedure

In order to record BVOC fluxes and concentrations, both a high sensitivity PTR-MS (Ionicon Analytik GmbH, Innsbruck, described in detail by Blake et al., 2009; de Gouw and Warneke, 2007; Hansel et al., 1995; Lindinger et al., 1998) and a high resolution PTR-ToF-MS (Ionicon Analytik GmbH, Innsbruck, as described by Graus et al., 2010; Jordan et al., 2009) were used, together with a sonic anemometer (Gill HS, Gill Instruments Ltd, UK). The PTR-MS was equipped with a quadrupole mass analyser, three turbo molecular pumps (Varian) and a heated Silcosteel inlet. The application of PTR-MS to atmospheric measurements has previously been described by Hewitt et al. (2003) and Hayward et al. (2002).

The sonic anemometer was situated 32 m above the ground on the north-west corner of the tower. Both the PTR-MS and the PTR-ToF-MS were housed in an air-conditioned cabin at the base of the tower. The PTR-MS sub-sampled via a 1/8 inch O.D. PTFE tube (I.D. 1 mm) from a 1/2 inch O.D. PTFE common inlet line (I.D. 3/8 inch), heated to avoid condensation, which led from ca. 10 cm below the sonic anemometer to the cabin. Solenoid valves were used to switch between the sample line and zero air which

was generated by passing ambient air through a glass tube packed with platinum catalyst powder heated to 200 °C. The PTR-ToF-MS subsampled via a 3-way valve from the common inlet line; 0.5 L min⁻¹ was pumped through a 1/8 inch (O.D.) and 1/16 inch (O.D.) capillary (together ca. 20 cm long), with 30 mL min⁻¹ entering the instrument and the remaining flow being sent to an exhaust. The common inlet line had a flow rate of ca. 63 L min⁻¹, giving a Reynolds number of ca. 9700 which indicates a turbulent flow. There was no observable influence of the high flow rate on readings from the sonic anemometer, even during periods of relatively low turbulence. Data from both the PTR-MS and the sonic anemometer were logged onto a laptop using a program written in LabVIEW (National Instruments, Austin, Texas, USA).

The PTR-MS was operated continuously throughout the measurement campaign with pauses for optimisation and refill of the water reservoir. PTR-MS settings were controlled so that the reduced electric field strength (E/N , where E is the electric field strength and N the buffer gas density) was held at 122 Td ($1.22 \times 10^{-19} \text{ V m}^{-2}$), with drift tube pressure, temperature and voltage maintained at 2.1 mbar, 45 °C and 550 V respectively. The primary ions and the first water cluster were quantified indirectly from the isotope peaks at m/z 21 ($\text{H}_2^{18}\text{O}^+$) and m/z 39 ($\text{H}_2^{18}\text{O} \cdot \text{H}_2\text{O}^+$), respectively. The inferred count rate of H_3O^+ ions over the course of the campaign varied between 1.33×10^6 and $9.00 \times 10^6 \text{ counts s}^{-1}$. O_2^+ (m/z 32) was kept below 1 % of the primary ion count throughout the campaign in order to limit ionisation of VOCs through charge transfer reactions with O_2^+ and minimise the contribution of the O_2^+ isotope ($^{16}\text{O}^{17}\text{O}^+$) to m/z 33.

During PTR-ToF-MS operation the drift tube temperature was held at 60 °C with 600 V applied across it. The drift tube pressure was 2.3 mbar resulting in an E/N of 130 Td. A more detailed description of the PTR-ToF-MS operation is provided by Schallhart et al. (2015).

The PTR-MS was operated in three modes: zero air, flux and scan alternated in an hourly cycle. The instrument measured zero air for 5 min, followed by 25 min in flux mode, 5 min in scan mode and then a final 25 min in flux mode. While in flux mode, 11

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



protonated masses were monitored sequentially: m/z 21 the hydronium ion isotope, m/z 39 a water cluster isotope and 9 masses relating to VOCs: m/z 33, 45, 59, 61, 69, 71, 73, 81 and 137. The mass spectral peaks at m/z 21 and 39 were analysed with a 0.2 s dwell time (τ). For the nine VOC species $\tau = 0.5$ s was used in order to increase the instrumental sensitivity to these masses. This gave a total scan time of 4.9 s and the acquisition of ca. 306 data points in each 25 min averaging period. The uncertainty caused by disjunct sampling was calculated and found to cause a 0.17 % error in the flux estimation (see Supplement for details).

Identification of the compounds observed at each of these masses is complicated by the fact that PTR-MS only allows the identification of nominal masses, therefore it is impossible to distinguish between isobaric compounds. As such there may be more than one compound contributing to each of the measured masses; Table 1 displays the masses monitored and the compounds likely to be contributing to each mass together with the exact masses observed at each unit mass using the PTR-ToF-MS which has much greater mass resolution than does the quadrupole PTR-MS instrument. It was assumed that the dominant contributions at m/z 33, 45, 59, 61, 69, 71, 73, 81 and 137 were from protonated methanol, acetaldehyde (ethanal), acetone (propanone), acetic acid (ethanoic acid), isoprene (2-methyl-1,3-butadiene), methyl vinyl ketone (MVK, butenone) and methacrolein (MACR, 2-methylprop-2-enal), methyl ethyl ketone (MEK, butanone), a monoterpene mass spectral fragment and monoterpenes respectively. A further contribution to m/z 71, recently identified, are isoprene hydroxy hydroperoxides (ISOPOOH, Rivera-Rios et al., 2014). However, the concentrations of this intermediate are small if NO_x concentrations are high and therefore are likely to be negligible at this site, where NO_x concentrations were large (A. Finco, personal communication, 2015).

2.2.1 PTR-MS calibration

The PTR-MS was calibrated using a gas standard (Ionicon Analytic GmbH, Innsbruck) containing 17 VOCs at a volume mixing ratio of approximately 1×10^{-6} (ca.

1 ppmv). The protonated mass of the VOCs ranged from m/z 31 (formaldehyde, CH_3O^+) to m/z 181 (1,2,4-trichlorobenzene, $\text{C}_6\text{H}_4\text{Cl}_3^+$). Methanol (m/z 33), acetaldehyde (m/z 45), acetone (m/z 59), isoprene (m/z 69), MEK (m/z 73) and the monoterpene α -pinene (m/z 81 and m/z 137) were present in the calibration gas standard, allowing sensitivities to be calculated directly. Due to reduced quadrupole transmission for high masses, monoterpenes were quantified using the fragment ion at m/z 81. For compounds not contained in the gas standard (acetic acid (m/z 61) and MVK and MACR (m/z 71)) empirical sensitivities were calculated. A relative transmission curve was created using the instrumental sensitivities calculated from the masses present in the standard, and from this curve sensitivities for the unknown masses were calculated (Davison et al., 2009; Taipale et al., 2008). Error in calibration using the gas standard was assumed to be below 5 %, whereas relative errors in calibrations using the relative transmission approach are < 30 % (Taipale et al., 2008).

2.2.2 PTR-ToF-MS calibration

Background measurements of the PTR-ToF-MS were made up to three times a day using zero air generated by a custom made catalytic converter. Calibrations were made using a calibration gas (Appel Riemer Environmental Inc., USA) which contained 16 compounds, with masses ranging from 33 to 180 amu. For VOCs not included in the calibration standard, the average instrument sensitivities towards the known C_xH_y , $\text{C}_x\text{H}_y\text{O}_z$ or $\text{C}_x\text{H}_y\text{N}_z$ compound families were used.

2.3 Calculation of volume mixing ratios

Volume mixing ratios were calculated from data generated using the PTR-MS using a program written in LabVIEW (National Instruments, Austin, Texas, USA). Volume mixing ratios (χ_{VOC}) were calculated from the raw PTR-MS data (in counts per second

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(cps)) using a method based on those of Taipale et al. (2008) and Tani et al. (2004).

$$X_{\text{VOC}} = \frac{I(\text{RH}^+)_{\text{norm}}}{S_{\text{norm}}} \quad (1)$$

where S_{norm} is the normalised sensitivity and $I(\text{RH}^+)_{\text{norm}}$ represents the background corrected normalised count rate (ncps) for the protonated compound R which was calculated as shown below.

$$I(\text{RH}^+)_{\text{norm}} = I(\text{RH}^+) \left(\frac{I_{\text{norm}}}{I(\text{H}_3\text{O}^+) + I(\text{H}_3\text{O}^+\text{H}_2\text{O})} \right) \left(\frac{p_{\text{norm}}}{p_{\text{drift}}} \right) - \frac{1}{n} \sum_{i=1}^n I(\text{RH}^+)_{\text{zero},i} \left(\frac{I_{\text{norm}}}{I(\text{H}_3\text{O}^+)_{\text{zero},i} + I(\text{H}_3\text{O}^+\text{H}_2\text{O})_{\text{zero},i}} \right) \left(\frac{p_{\text{norm}}}{p_{\text{drift, zero},i}} \right) \quad (2)$$

where $I(\text{RH}^+)$, $I(\text{H}_3\text{O}^+)$ and $I(\text{H}_3\text{O}^+\text{H}_2\text{O})$ represent the observed count rate for the protonated compound R , H_3O^+ and the $\text{H}_3\text{O}^+\text{H}_2\text{O}$ cluster, respectively. Subscript zero refers to zero air measurements, n is the number of zero air measurement cycles and p_{drift} is the drift tube pressure. The drift tube pressure was normalised to 2 mbar (p_{norm}) and the sum of the primary ion and first water cluster was normalised to a count rate of 10^6 cps (I_{norm}). The compound specific limit of detection (LoD) was calculated using the method described by Karl et al. (2003):

$$\text{LoD} = 2 \times \frac{\sigma_{\text{Background}}}{S_{\text{VOC}}} \quad (3)$$

where S_{VOC} is the instrumental sensitivity to the VOC and $\sigma_{\text{Background}}$ is the mean background normalised count rate.

2.4 Flux calculations from PTR-MS

The 25 min PTR-MS flux files were inspected and incomplete or disrupted files removed. BVOC fluxes were then calculated using a program also written in LabVIEW,

29222

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



based upon the virtual disjunct eddy covariance technique (vDEC) developed by Karl et al. (2002), also termed continuous flow disjunct eddy covariance (Rinne et al., 2008). This method has previously been successfully applied in a number of studies (e.g. Davison et al., 2009; Langford et al., 2009, 2010a, b; Misztal et al., 2011; Rinne et al., 2007).

- 5 This approach allows direct calculation of fluxes of atmospheric constituents, as with standard eddy covariance, yet in this case sampling of scalar concentrations is not continuous. The flux, F_x , for each compound was calculated using a covariance function between the vertical wind velocity, w , and the VOC mixing ratios, χ :

$$F_x(\Delta t) = \frac{1}{N} \sum_{i=1}^N w'(i - \Delta t / \Delta t_w) \chi'(i) \quad (4)$$

- 10 where Δt is the lag time between the PTR-MS concentration measurements and the vertical wind velocity measurements from a sonic anemometer, Δt_w is the sampling interval between wind measurements (0.1 s), N is the number of PTR-MS measurement cycles in each 25 min averaging period (typically 250 in our study) and primes represent the momentary deviations from the mean concentration or vertical wind speed (e.g. $w = w' - \bar{w}$).

- 15 Variations in temperature, pressure and the performance of the sample line pump can cause small deviations in Δt . Therefore these values were calculated using a cross correlation function between w' and χ' . Lag times were calculated individually for each m/z monitored by the PTR-MS by selecting the absolute maximum value of the covariance function within a 30 s time window (MAX method, Taipale et al., 2010). This analysis resulted in a clear isoprene flux but for most masses a high proportion of the data fell below the limit of detection. These data, especially in the case of acetone, showed a significant amount of flux values with the opposite sign, “mirroring” the true flux. These “mirrored” points occur when the measured flux is of comparable magnitude to the total random error of the system (Langford et al., 2015). As the cross-correlation maximum is likely to be an over-estimate when the noise to signal ratio is greater than one, these points were substituted with fluxes calculated using a fixed lag time.

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A histogram of isoprene lag times calculated using the MAX method is displayed in the Supplement showing a clear maximum at 7.5 s. Therefore 7.5 s was chosen as the isoprene fixed lag time and fixed lag times for the other masses were calculated from the isoprene fixed lag time, accounting for the dwell times of the different compounds in the measurement cycle.

2.4.1 Flux quality assessment and potential losses

In order to assess the quality of each 25 min flux file, the resultant fluxes were subjected to three quality checks following a two-dimensional coordinate rotation which was applied to correct for tilting of the sonic anemometer (see Table S1 in the Supplement for summary). Following the criteria of Langford et al. (2010a), data points were labelled if the mean friction velocity (u_*) over the 25 min averaging period was found to be below 0.15 ms^{-1} . Data falling below this threshold predominantly occurred at night when wind velocity reached a minimum. Detection limits for each 25 min flux file were calculated using a method based on that of Wienhold et al. (1994) as applied by Spirig et al. (2005) where the signal of the flux at the true lag is compared to the background noise of the covariance function. The 95th percentile of the covariance function in the lag range 150–180 s was calculated and flux files falling below this value were labelled as having fallen below the LoD. Finally data points underwent a stationarity test as described by Foken and Wichura (1996) and data points found to be generated from periods of non-stationarity were also labelled.

The integral turbulence characteristics were assessed using the FLUXNET criteria described by Foken et al. (2004). The turbulence at the Bosco Fontana field site was well developed with 87 % of the data in the first three categories, defined by Foken et al. (2004) as suitable for fundamental research. Less than 1 % of the data fell into category 9, characterised as data to be excluded under all circumstances.

The flux losses in the virtual disjunct eddy covariance system were assessed. Loss of flux at frequencies higher than the PTR-MS response time and/or dwell time was corrected for using the method described by Horst (1997). Correction factors in the range

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1.01–1.23 were calculated and applied to each 25 min flux file with a mean correction of 8.8 % applied. Rotating the coordinates in order to set the vertical mean vertical wind velocity to 0 for each twenty five minute flux averaging period and block averaging itself act as a high pass flux filter (Moncrieff et al., 2004), leading to the loss of low frequency fluxes. The loss of these low frequency fluxes due to an insufficient averaging period is assessed in the Supplement. Sensible heat flux data were averaged over 50, 75, 100 and 125 min before a coordinate rotation was applied and plotted against the sum of two, three, four and five 25 min coordinate rotated flux files, respectively. The gradient of the fitted line between the two fluxes gives an estimate of the flux lost by the use of twenty five minute averaging periods. As is shown in Fig. S2 of the Supplement, eddies with a time period between 25 and 125 min carry only an additional 2.8 % of the sensible heat flux. Therefore if we assume that the frequency of VOC and sensible heat fluxes are comparable, 1.0–3.6 % of the VOC flux is lost by limiting the averaging period to 25 min. This correction has not been applied to the displayed data as it is so small.

2.5 Flux calculations from PTR-ToF-MS

BVOC fluxes were calculated from PTR-ToF-MS data using the eddy covariance (EC) method similar to that described above for the PTR-MS. The PTR-ToF-MS flux analysis differed in that the cross correlation between w' and χ' was calculated using the method described by Park et al. (2013). Whilst in the PTR-MS measurement, the target compounds are predetermined through the measurement cycle, in the PTR-ToF-MS the entire high resolution mass spectrum can be used to search for compounds that carry a flux. PTR-ToF-MS data were analysed using the TOF Analyzer V2.45 as described by Müller et al. (2013) and ToFTools (Junninen et al., 2010). An automated flux identification routine was then used to calculate the average of the absolute cross covariance functions during a mid-day period. The maximum value was then automatically selected from the averaged spectrum and checked against the manually selected noise level ($10\sigma_{\text{noise}}$) to determine whether a flux was present.

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The fluxes were filtered using the 70 % stationary criteria as presented by Foken and Wichura (1996), as was applied to the PTR-MS data and corrected for loss of high frequency flux Horst (1997). For a more detailed description of the flux calculation from the PTR-ToF-MS see Schallhart et al. (2015).

2.6 Leaf level GC-MS measurements

A portable gas exchange system equipped with a controlled-environment 6 cm² broadleaf cuvette (LI6400, Li-COR, Lincoln, USA) was used to measure net photosynthetic rate (A) and stomatal conductance (g_s) at basal conditions of PAR (1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$), leaf temperature (30 °C) and a CO₂ concentration (400 ppm) from fully expanded leaves. When A reached a steady-state, the outlet tube from the leaf cuvette was replaced with a Teflon tube, and the air stream exiting from the cuvette was used as a sampling port for BVOC (according to the methodology in Loreto et al., 2001) by a silico-steel cartridge packed with 200 mg of tenax (Supelco, PA, USA). Tenax is a very hydrophobic and adsorbent material with high thermal stability generally used for trapping BVOC (Dettmer and Engewald, 2002). The flow rate through the leaf cuvette was maintained at 500 $\mu\text{mol s}^{-1}$, and a subsample of 200 mL min⁻¹ (130 $\mu\text{mol s}^{-1}$) was pumped through the cartridge with an external pump (AP Buck pump VSS-1) for a total volume of 6 L of air. Blank samples of air without a leaf in the cuvette were collected every day before and after the BVOC samplings. Finally the cartridges were sealed and stored at 4 °C until analysis.

The cartridges were analyzed using a Perkin Elmer Clarus 580 gas-chromatograph coupled with a Clarus 560 Mass-Detector and a thermal-desorber Turbo Matrix (Perkin Elmer Inc., Waltham, MA, USA). The gas-chromatograph was equipped with an Elite-5-MS capillary column (30 m length, 250 μm diameter and 0.25 μm film thicknesses). The carrier gas was helium. The column oven temperature was kept at 40 °C for 5 min, then increased with a 5 °C min⁻¹ ramp to 250 °C and maintained at 250 °C for 5 min. BVOC were identified using the NIST library provided with the GC/MS Turbomass software. GC peak retention time was substantiated by analysis of parent ions and main

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



fragments of the spectra. Commercially available reference standards were used to create the calibration curves and to quantify the emissions. To normalize the BVOC results, the quantities of terpenes collected from the empty cuvette (blanks) were subtracted from the plant emission results. The quantification of total BVOC emission was performed using authentic gaseous standards (Rivoira, Milan, Italy) or liquid standards (Sigma Aldrich, Milan, Italy).

2.7 Mapping tree species distribution

Tree species distribution data were obtained from Dalponte et al. (2007) who used a combination of Light Detection and Ranging (LIDAR) and hyperspectral data to develop a high resolution tree species distribution map of the Bosco Fontana natural reserve.

The overall accuracy (kappa coefficient) of this species map is particularly high (0.89), considering the number of classes (23) and the number of training samples (20 % of the data are used in the training set and 80 % in the test set) per class. The LIDAR channels provide relatively sparse information for discriminating between tree species, increasing the overall accuracy of the tree species assignment using the hyperspectral data by only 1 % but the LIDAR data significantly increase the accuracy of understory and underrepresented classes. The kappa coefficient of the main species is also very high (0.88–0.93) showing the effectiveness of this approach for species classification in a very complex forest with 20 different broad-leaves species, some of which, such as *Q. cerris*, *Q. robur* and *Q. rubra*, belong to same genus. For a more detailed discussion of the mapping results and methodology see Dalponte et al. (2007, 2008).

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

[Back](#)

Close

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion



3 Results and discussion

3.1 Meteorological conditions

The measurement campaign at Bosco Fontana ran from 1 June 2012 to 11 July 2012 (41 days) with data recorded using the PTR-MS from the 13 June 2012 to the 11 July 2012. The meteorological conditions recorded at the measurement site during this period are summarised in Figs. 1 and 2, times are reported in central European time (UTC + 1) as used throughout this paper. With the exception of two heavy thunderstorms, the first in the first week of June before measurements began and the second overnight on 6 July, there was no precipitation during the measurement period. The temperature gradually increased from the campaign start until 19 June and then remained more or less constant. During the measurement period ambient temperature varied from a low of 14 °C to a high of 35 °C. Daily photosynthetically active radiation (PAR) peaked within the range 1890–2105 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and the relative humidity during the campaign varied between 29 and 90 %. Winds were generally easterly or north westerly. For most of the campaign wind speeds were below 3.5 ms^{-1} but peaked at 5.6 ms^{-1} on 23 June, with the mean wind speed for the campaign period of 1.6 ms^{-1} .

3.2 BVOC mixing ratios and fluxes

BVOC fluxes were recorded at the Bosco Fontana site using both the PTR-MS and the PTR-ToF-MS. Unless stated, the results displayed here were calculated from measurements made using the PTR-MS. Data analysis was carried out with the aid of the R openair package (Carslaw and Ropkins, 2012; R Core Team, 2012). For a full discussion of all fluxes and concentrations recorded using the PTR-ToF-MS see Schallhart et al. (2015).

The mixing ratios of the eight BVOC species measured in flux mode using the PTR-MS are displayed in Fig. 3 and are summarised in Table 2 (for further details, see Fig. S5 of the Supplement). These mixing ratios were calculated using the high fre-

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



quency flux measurements so the presented mixing ratios are an average over 25 min. The mixing ratio LoDs, calculated as described above (Karl et al., 2003; Langford et al., 2009; Misztal et al., 2011) were in the same range as those calculated on previous campaigns (Langford et al., 2009; Misztal et al., 2011) and, with the exception of isoprene where the mixing ratio dropped towards zero at night, the recorded mixing ratios generally remained above their respective LoD.

Table 3 summarizes the flux data recorded during the Bosco Fontana measurement campaign. Wind speeds decreased at night, leading to a large proportion of the night time data falling below the u_* threshold of 0.15 ms^{-1} . Consequently, average emission fluxes of all eight compounds are reported for the daytime period 10:00–15:00 LT as well as for the whole campaign. Large fluxes of m/z 69 and m/z 81 (assigned to isoprene and monoterpenes respectively) were observed and are shown in Fig. 4. Fluxes of m/z 33, 45, 59, 61, 71 and 73 (assigned to methanol, acetaldehyde, acetone, acetic acid, MVK + MACR and MEK, respectively) were also observed, but these fluxes were weaker, leading to a high percentage of fluxes failing the LoD check. However, as is described by Langford et al. (2015), when these flux data are averaged to show the average diurnal cycle, it is appropriate to use a combined LoD value appropriate for the same period rather than the LoD attached specifically to each 25 min flux file. It is, though, essential that each individual flux period be processed carefully to avoid the introduction of a bias due to the use of the MAX method of time-lag identification. The LoD for the mean ($\overline{\text{LoD}}$) decreases with the square root of the number of samples averaged (N).

$$\overline{\text{LoD}} = \frac{1}{N} \sqrt{\sum_{i=1}^N \text{LoD}^2} \quad (5)$$

Therefore, while the flux time series of methanol, acetaldehyde, acetone, acetic acid, MVK + MACR and MEK are not presented here, the campaign average diurnal fluxes are shown (Fig. 5). As discussed above, 25 min averaged flux files flagged as below the

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



LoD were included in these diurnal averages. Flux files falling below the 0.15 m s^{-1} wind speed threshold were also included to prevent the night time flux being biased high for depositing compounds. For compounds showing emission, night-time fluxes are close to zero anyway and the application has little influence on the results. Data flagged for non-stationarity were excluded. For a more detailed discussion of the fluxes and mixing ratios of each BVOC and comparison made with other temperate and Mediterranean ecosystems, see the Supplement.

The fluxes of isoprene and monoterpenes calculated using both the PTR-MS and the PTR-ToF-MS instruments are displayed in Fig. 4 and summarised in Table 3. The isoprene fluxes calculated using both instruments show very good agreement ($R^2 = 0.91$, slope 1.3 and intercept 0.17). The monoterpene fluxes, calculated using m/z 81 with the PTR-MS and m/z 81.070 with the PTR-ToF-MS show an $R^2 = 0.50$. Three additional mass spectral peaks are observed at m/z 81 in the PTR-ToF-MS: m/z 80.92, 80.99 and 81.03, however statistically significant fluxes from these peaks could not be calculated using the PTR-ToF-MS. Owing to the lower sensitivity of the PTR-MS at m/z 81 and the lower sampling frequency of the disjunct sampling protocol (Rinne and Ammann, 2012), the monoterpene flux calculated using this instrument is significantly noisier than the flux calculated using the PTR-ToF-MS.

PTR-MS and PTR-ToF-MS mass scans were averaged over a ten day period (14–24 June). A comparison of these mass scans over the range m/z 33 to 100 at unit mass resolution is displayed in Fig. 6, with masses reported relative to m/z 59 (acetone). A good agreement between the PTR-MS and PTR-ToF-MS is seen for all masses, except for m/z 33 where the PTR-MS gives a significantly higher signal. As both instruments have comparable sensitivities at this mass (11.6 and ca. $10\text{--}12 \text{ ncps ppbv}^{-1}$ for the PTR-MS and PTR-ToF-MS respectively) this discrepancy must be the result of interference from another ion at this mass. O^{17}O^+ could interfere with the methanol signal at m/z 33 but as a significant peak is not observed at m/z 34 (O^{18}O^+) a large contribution from O^{17}O^+ to m/z 33 is unlikely. This suggests that there is a greater formation of O_2H^+ in the PTR-MS than in the PTR-ToF-MS under these particular op-

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



eration parameters. No major mass spectral peaks are observed in one instrument alone, indicating that there is no artefact formation or unexpected loss of chemical species with either instrument. The mass scans show a much cleaner spectrum than was reported by Misztal et al. (2011) above an oil palm plantation in South-East Asia, suggesting an atmosphere dominated by fewer chemical species at higher concentrations.

3.2.1 BVOC correlations

Scatter plots were used to investigate the relationship between the measured species. Methanol, acetone and MEK (Fig. 7) all showed a bimodal relationship with two linear groupings, one at lower temperature (ca. $< 20^{\circ}\text{C}$) and another at higher temperatures (ca. $> 20^{\circ}\text{C}$). This suggests that either there are two different sources contributing to the mixing ratios (most likely an atmospheric background and a photochemical source at higher temperatures) or that a second compound contributes to the nominal mass at higher temperatures. As few compounds have been reported to contribute to m/z 33 or 59, an additional source at higher temperatures seems more likely.

3.2.2 Short-chain oxygenated BVOCs

A mean methanol mixing ratio of 6.2 ppbv at 4 m above the canopy was recorded over the duration of the campaign, making it the dominant BVOC observed at Bosco Fontana. Mean acetaldehyde, acetone and acetic acid mixing ratios were 3.4, 3.2 and 1.9 ppbv at 4 m above the canopy, respectively. Methanol, acetaldehyde and acetic acid mixing ratios all followed similar diurnal cycles (Fig. 3), with mixing ratios remaining stable through the night before a drop in the morning, probably caused by expansion of the planetary boundary layer after sunrise. Then mixing ratios increased again in the late afternoon as emissions accumulated in a shrinking boundary layer. Acetone mixing ratios remained on average stable throughout the day (Fig. 3).

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The flux of methanol peaked at $0.49 \text{ mg m}^{-2} \text{ h}^{-1}$ with a mean day-time flux of $0.03 \text{ mg m}^{-2} \text{ h}^{-1}$ (Fig. 5). Methanol deposition was observed during the night and mornings followed by a rapid increase in methanol emission in the late morning and peaking in the early afternoon. Bidirectional exchanges of methanol have been reported previously (for example Fares et al., 2012; Karl et al., 2004) with methanol absorption/desorption thought to occur in thin water films within the canopy (Wohlfahrt et al., 2015). The mean morning (06:30–10:30 LT) methanol deposition velocity (V_d) at the measurement height (z_m) was calculated using the relationship (Misztal et al., 2011):

$$V_d(z_m) = -\frac{F}{\chi(z_m)} \quad (6)$$

and was found to be 0.31 cm s^{-1} . The night-time deposition velocity was lower, 0.02 cm s^{-1} , falling at the bottom end of the $0.02\text{--}1.0 \text{ cm s}^{-1}$ range reported by Wohlfahrt et al. (2015) from a review of eight different north hemisphere sites.

Acetic acid deposition was also observed in the morning, but any emission flux in the afternoon remained below the limit of detection, even if aggregated into mean diurnal cycles. The mean diurnal acetaldehyde flux is shown in Fig. 5. The flux increased from below the detection limit in late morning to a peak in the early afternoon before dropping again towards zero at night. The flux peaked at $0.44 \text{ mg m}^{-2} \text{ h}^{-1}$ on 29 June and the campaign mean day-time flux was $0.06 \text{ mg m}^{-2} \text{ h}^{-1}$. As can be seen in Fig. 5, the acetone flux remained below the limit of detection for most of the day with a small positive flux observed in the late afternoon.

3.2.3 MVK + MACR and MEK

MVK and MACR are the main products formed following the first stage of isoprene oxidation in the atmosphere (Atkinson and Arey, 2003), accounting for ca. 80 % of the carbon. MACR can also be directly produced within plants as a by-product in the production of cyanogenic glycosides (Fall, 2003) and experimental observation demonstrated

that emissions of MVK and MACR increase with temperature stress (Jardine et al., 2012). The mid-day (10:00–15:00 LT) mixing ratios of MVK + MACR at 4 m above the canopy showed a positive correlation with those of isoprene ($R^2 = 0.49$), suggesting that the oxidation of isoprene was responsible for the formation of MVK and MACR.

The production of MVK and MACR from isoprene at the Bosco Fontana site has been modelled by Schallhart et al. (2015), who estimated that 4–27 % of the MVK + MACR flux was formed from isoprene oxidation products. MVK and MACR mixing ratios recorded at 4 m above the canopy (Fig. 3) increase in the morning as isoprene concentrations rise, before boundary layer expansion causes them to drop in the middle of the day. The mixing ratios then increase again in the evening as the boundary layer contracts. The flux of MVK + MACR (Fig. 5) peaked in the early afternoon with a mean day-time flux of $0.05 \text{ mg m}^{-2} \text{ h}^{-1}$ comparable to the 0.03 and $0.08 \text{ mg m}^{-2} \text{ h}^{-1}$ observed, respectively, by Kalogridis et al. (2014) and Spirig et al. (2005) over European oak and mixed forests.

MEK may be directly emitted by plants (Fall, 2003) or formed photochemically (Luecken et al., 2012). MEK mixing ratios 4 m above the forest canopy remained stable through the night at ca. 0.6 ppbv before a dropping in the morning, probably caused by expansion of the planetary boundary layer, to ca. 0.3 ppbv and rising again in the evening (Fig. 3). A plot of the mixing ratios of MEK against those of acetone reveals a bimodal distribution suggesting two distinct sinks or sources (Fig. 7), the first occurring at lower temperatures (ca. 12–20 °C) with a MEK to acetone ratio of ca. 0.17 and the second at higher temperatures (ca. 20–34 °C) with a MEK to acetone ratio of ca. 0.06. A relationship between acetone and MEK has been reported by Riemer et al. (1998) who observed an MEK to acetone ratio of 0.07 at temperatures between 20 and 37 °C. This compares well with the observations at Bosco Fontana. This trend was not observed when data were coloured by PAR indicating that the bimodal distribution is not driven by the faster rate of reaction of MEK than of acetone with OH. A low MEK emission flux was observed in the afternoon with a mean day-time flux of $0.02 \text{ mg m}^{-2} \text{ h}^{-1}$.

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2.4 Isoprene and monoterpenes

Isoprene mixing ratios 4 m above the canopy began to rise in the mid-morning from a night-time zero, peaking in the late afternoon at ca. 2 ppbv before falling again to zero in the late evening (Fig. 3). Isoprene fluxes were not observed at night, but increased in the morning to a peak in the mid afternoon before dropping to zero again in the evening (Fig. 5) with a mean day-time flux of $1.9 \text{ mg m}^{-2} \text{ h}^{-1}$.

Isoprene fluxes correlated with leaf temperature (estimated using a method based on that described by Nemitz et al. (2009) and explained in more detail in the Supplement) giving an $R^2 = 0.75$ for an exponential fit, PAR ($R^2 = 0.75$ for an exponential fit) and with sensible heat flux (H) ($R^2 = 0.67$). The relationship between isoprene fluxes and mixing ratios, temperature and PAR is displayed in Fig. 8. An exponential relationship between temperature and both fluxes and mixing ratios was observed for the periods when PAR was greater than zero. Table 4 compares isoprene flux measurements with the fluxes recorded during other field campaigns in the Mediterranean region and the isoprene emission factor under basal conditions. As would be expected the flux of isoprene is shown to be highly dependent on ecosystem type. When fluxes are normalised to standard conditions the fluxes observed on this campaign are lower than those observed over woodland dominated by isoprene emitting oak species due to the lower proportion of isoprene emitting species in the canopy but closer in magnitude to that observed over a mixed pine and oak forest.

The campaign mean monoterpene mixing ratio 4 m above the canopy was 0.2 ppbv. The diurnal profile (Fig. 3) shows a night-time mixing ratio of ca. 0.18 ppbv which increases to ca. 0.21 ppbv in the morning remaining stable through the day and dropping again to ca. 0.18 ppbv at night. The monoterpene flux (Fig. 5) peaked in the early afternoon with a campaign mean mid-day flux of $0.12 \text{ mg m}^{-2} \text{ h}^{-1}$. Monoterpene mixing ratios were not significantly correlated with leaf surface temperature or with PAR ($R^2 = 0.11$ and 0.12 respectively). However, the flux displayed a correlation with both leaf surface temperature and PAR ($R^2 = 0.44$ and 0.39 respectively).

3.3 Calculation of isoprene and monoterpene canopy level emission factors

Although other approaches do exist, isoprene fluxes are widely modelled using the Model of Emissions of Gases and Aerosols from Nature (MEGAN, Guenther et al., 2006). MEGAN calculates isoprene fluxes based on the product of an emission activity factor (γ), a canopy loss and production factor (ρ) and a canopy emission factor (ε). Therefore, plotting isoprene flux against $\gamma \times \rho$ enables the calculation of a canopy-specific isoprene emission factor (Fig. 9), giving a canopy emission factor of $1.68 \text{ mg m}^{-2} \text{ h}^{-1}$ at standard conditions ($1000 \mu\text{mol m}^{-2} \text{ s}^{-1}$ PAR and 303 K) for the campaign period. For the purpose of this work, γ was calculated using the algorithms described by Guenther et al. (2006). Radiative transfer through the canopy was modelled using the canopy model applied by Müller et al. (2008) based on that of Goudriaan and van Laar (1994) and ambient temperature was recorded 4 m above the canopy. The standard light and temperature conditions for MEGAN canopy scale emissions factors are $\sim 1500 \mu\text{mol m}^{-2} \text{ s}^{-1}$ and 303 K (Guenther et al., 2006). In order to allow for direct comparison with the GC-MS data and literature emissions factors the factor which sets the emission activity to unity at standard conditions (C_{CE}) was increased to 1.42 to give standard light and temperature conditions of $1000 \mu\text{mol m}^{-2} \text{ s}^{-1}$ and 303 K, respectively.

The emission factor is lower than those calculated by Kalogridis et al. (2014) and Baghi et al. (2012) from oak (*Quercus pubescens*) dominated forests in southern France (7.4 and $5.4 \text{ mg m}^{-2} \text{ h}^{-1}$, respectively). However, this is to be expected, owing to the high proportion of low or non-isoprene emitting species such as *Carpinus betulus*, *Corylus avellana*, *Sambucus nigra* and *Acer campestre* present in the forest at Bosco Fontana.

Monoterpene emission from plants may take the form of pool or de novo emission. Emission from stored pools is temperature controlled whereas de novo is driven by photosynthesis and is therefore controlled by light as well as temperature (Ghirardo et al., 2010). Emission from stored pools was modelled using the monoterpene–temperature

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



relationship described by Guenther et al. (1995), this model correlated well with the observed monoterpene flux (PTR-ToF-MS) giving R^2 value of 0.55. In order to assess the effect of light on monoterpene emission the residual values from the temperature only model were plotted against PAR. The residuals displayed a correlation with PAR ($R^2 = 0.45$) indicating that light as well as temperature has a significant impact on monoterpene emissions from the forest canopy and therefore that a significant proportion of monoterpene emission takes the form of de novo emission. However, in order to accurately assess the contribution of pool and de novo emissions to the canopy scale monoterpene flux, a species specific leaf level investigation would be required. A monoterpene canopy emission factor calculated using the MEGAN algorithms, which simulate de novo emission, was found to be $0.14 \text{ mg m}^{-2} \text{ h}^{-1}$.

3.4 Speciated bottom-up isoprene and monoterpene flux estimates derived from leaf-level measurements

Tree species distribution data combined with information on leaf-level isoprene and monoterpene emission rates and meteorological data were used to produce a “bottom-up” estimate of the total canopy level flux. Tree species distribution data were obtained from Dalponte et al. (2007), this tree species distribution map reveals an uneven distribution of isoprene emitting species within the forest canopy, with the two main isoprene emitting species (*Q. robur* and *Q. rubra*) concentrated in the south of the forest.

Leaf-level isoprene and monoterpene emissions from the dominant tree species were recorded using GC-MS (Table 5). These species represent 76.6 % of the total vegetation cover. Isoprene emission was dominated by *Q. robur* and *Q. rubra* with *C. avellana* and *C. betulus* the highest monoterpene emitting species. The isoprene emission recorded for both oak species was lower than that previously reported (Karl et al., 2009; Keenan et al., 2009). For species where GC-MS data were not available literature values were used. Leaf-level emission factors for minor species for which no GC-MS measurements were made were taken from Karl et al. (2009) with the exception *Rubus* sp., taken from Owen et al. (2001) and *Acer negundo* and *Morus* sp., taken

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from Benjamin et al. (1996). Emission factors taken from the literature were converted from $\mu\text{g g}_{\text{DW}}^{-1} \text{h}^{-1}$ to $\text{mg m}^{-2} \text{h}^{-1}$ using the mean leaf mass to area ratio, $115 \text{ g}_{\text{DW}} \text{ m}^{-2}$, reported by Niinemets (1999) from a study of ca. 600 species. The leaf-level emissions data were then scaled up to a canopy level using the MEGAN algorithms (Guenther et al., 2006) and incorporating measured PAR and temperature values averaged over 30 min and a single sided leaf area index (LAI, $\text{m}^2 \text{ m}^{-2}$) of 5.5.

The hyperspectral/LIDAR data of Dalponte et al. (2007) was remapped onto a grid centred on the measurement site, with a resolution of 5 m^2 , providing fractional ground cover by each of the 20 tree species within each grid cell. The contribution of each grid cell to each 25 min flux measurement was then calculated at 5 m^2 resolution using a high resolution 2-D footprint model based on Kormann and Meixner (2001) similar to that described by Neftel et al. (2008). Finally, the MEGAN algorithm was applied to all plant species using the 25 min meteorology. The information was combined to provide a bottom-up estimate of the flux that the canopy-scale measurements should have detected, based on the leaf-level data. This footprint and species dependent bottom-up flux estimate showed significantly better agreement with the measured isoprene flux ($R^2 = 0.75$, slope = 0.56) than was observed when the canopy-scale isoprene emission factor calculated above was used ($R^2 = 0.65$, slope = 0.76). This demonstrates the large effect an uneven distribution of isoprene sources can have on the above canopy flux, even within what appears to be a uniform canopy, and the benefit for accounting for spatial species distributions in uniform vegetation canopies.

However, despite capturing the shape of the flux time series, the bottom-up flux underestimated the magnitude of the flux, capturing 56 % of the isoprene flux as measured by vDEC. This could in part be caused by changes in vegetation cover between the tree distribution mapping in 2008 and the flux measurements in 2012. Since 2008, the non-native *Q. rubra* is gradually being removed from the forest. However this does not explain the discrepancy between the vDEC isoprene flux measurements and the bottom-up flux estimate as the reduction in the number of *Q. rubra* trees should have decreased the flux. There are anecdotal reports that *Populus* sp. coverage has in-

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



creased in the understory vegetation but it is unlikely that, despite their high rates of growth, the *Populus* coverage changed significantly in the 4 years between mapping and this campaign. Whilst the hyperspectral/LIDAR tree species data for this site provides a unique opportunity for comparing the canopy-scale measurements with a detailed bottom-up estimate, the hyperspectral/LIDAR data provides information on projected tree species area as seen from above, whilst the flux is regulated by leaf mass and its exposure to radiation. Thus there are uncertainties in the ability of the hyperspectral/LIDAR in detecting understorey vegetation and a single conversion factor was used between projected tree area and leaf mass. However, understorey vegetation is less exposed to sunlight reducing its emission. Indeed, the main reason for the under-estimate of isoprene flux is probably that the leaf level isoprene emission rate recorded from the leaves sampled at ground level (albeit taken at the edge of sun exposed clearings) are not representative of those at the canopy top. Substituting the measured *Q. robur* and *Q. rubra* emission factors with those reported by Karl et al. (2009) caused the bottom-up estimate to give 130 % of the measured flux and improved the correlation between bottom-up estimates and canopy-scale measurements further.

The speciated monoterpene flux also showed good agreement with the above canopy flux ($R^2 = 0.72$) and captured 57 % of the flux. The discrepancy between the magnitude of the speciated monoterpene flux and the above canopy flux was within the range expected to be caused by the loss of monoterpenes within the canopy through oxidation and deposition.

The contribution of different species to the isoprene and monoterpene fluxes over the course of an example day is shown in Fig. 10. As is shown, the isoprene flux was dominated by *Q. robur* but was sensitive to the species composition within the flux footprint. The change in wind direction around 14:00 LT reduced the contribution of *Q. rubra* to the total flux, with the contribution of *Populus × canescens* increasing significantly. The monoterpene flux was predicted to have been dominated by *C. betulus*, the dominant tree species in the canopy at Bosco Fontana. A greater number of tree

species contributed to the monoterpene flux, and emissions were therefore much more uniform across the canopy and less affected by changes in wind direction.

The fit between the above canopy measured isoprene and monoterpene fluxes and the “bottom-up” flux estimate was improved by optimising the leaf-level emission factors, within the constraints displayed in Table 6, using Chi^2 minimisation as implemented by the solver function in Microsoft Excel. Use of the optimised isoprene and monoterpene emission factors gave a good correlation with the measured fluxes with R^2 values of 0.75 and 0.76, respectively for isoprene and monoterpenes. The optimised isoprene and monoterpene emission factors are presented in Table 6 and show a reasonable agreement with literature values (Karl et al., 2009).

4 Conclusions

Direct above-canopy fluxes of methanol, acetaldehyde, acetic acid, isoprene, MVK + MACR, MEK and monoterpenes were calculated using the method of virtual disjunct eddy covariance from mixing ratio data obtained with a PTR-MS above a semi-natural mixed oak and hornbeam forest in northern Italy from 13 June to 11 July 2012. Isoprene was the dominant BVOC emitted with a mean day-time flux of $1.91 \text{ mg m}^{-2} \text{ h}^{-1}$. When normalised to standard conditions (temperature of 30°C , PAR of $1000 \mu\text{mol m}^{-2} \text{ s}^{-1}$) using the MEGAN model (Guenther et al., 2006), a canopy scale emission factor of $1.68 \text{ mg m}^{-2} \text{ h}^{-1}$ was derived. Mixing ratios of VOCs measured at 4 m above the forest canopy were dominated by those of methanol, with a campaign mean mixing ratio of 6.2 ppbv.

The isoprene fluxes obtained using the PTR-MS/vDEC system showed good agreement with those obtained using a direct eddy covariance (with volume mixing ratios measured with a fast response PTR-ToF-MS instrument). Monoterpene fluxes recorded using the PTR-MS were noisier and marginally higher than those recorded using the PTR-ToF-MS due to a lower sensitivity and, probably, the inclusion of isobaric compounds. Comparison of mass scan data generated using the PTR-MS and

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



PTR-ToF-MS showed very good agreement and no significant masses observed in one instrument but not in the other within the mass range m/z 33–100.

Up-scaling leaf-level isoprene and monoterpene emissions to the canopy scale, using a high spatial resolution tree species database and a 2-D footprint model, showed significantly better correlation with the measured above canopy fluxes than was obtained using a canopy scale emission factor. Leaf-level isoprene emissions resulted in an underestimate of the above-canopy isoprene flux and this was assumed to be the result of differences in isoprene emission rates from leaves sampled at ground-level and those at the canopy top.

Overall, the data obtained give confidence in the measurement of biogenic VOC fluxes by the method of virtual disjunct eddy covariance and highlight the importance of using leaf-level emissions data from sun-lit canopy-top leaves when up-scaling leaf-level emissions to produce a “bottom-up” canopy-scale emissions estimate.

**The Supplement related to this article is available online at
doi:10.5194/acpd-15-29213-2015-supplement.**

Acknowledgements. W. J. F. Acton would like to thank Alex Guenther for his advice on the use of MEGAN. This work was funded by the EU FP7 grants ÉCLAIRE (grant 282910) and PEGASOS (grant 265148), as well as by a BBSRC/Ionicon Analytik GmbH Industrial CASE studentship awarded to W. J. F. Acton. We acknowledge access to the measurement site provided by the Italian Corpo forestale dello Stato and provision of the site infrastructure by the Catholic University of Italy at Brescia and in particular by Giacomo Gerosa, Angelo Finco and Riccardo Marzuoli.

**Canopy-scale flux
measurements and
bottom-up emission
estimates**

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Arneeth, A., Monson, R. K., Schurgers, G., Niinemets, Ü., and Palmer, P. I.: Why are estimates of global terrestrial isoprene emissions so similar (and why is this not so for monoterpenes)?, *Atmos. Chem. Phys.*, 8, 4605–4620, doi:10.5194/acp-8-4605-2008, 2008.
- 5 Ashmore, M. R.: Assessing the future global impacts of ozone on vegetation, *Plant Cell Environ.*, 28, 949–964, 2005.
- Atkinson, R. and Arey, J.: Gas-phase tropospheric chemistry of biogenic volatile organic compounds: a review, *Atmos. Environ.*, 37, 197–219, 2003.
- Baghi, R., Durand, P., Jambert, C., Jarnot, C., Delon, C., Serça, D., Striebig, N., Ferlicq, M.,
10 and Keravec, P.: A new disjunct eddy-covariance system for BVOC flux measurements – validation on CO₂ and H₂O fluxes, *Atmos. Meas. Tech.*, 5, 3119–3132, doi:10.5194/amt-5-3119-2012, 2012.
- Benjamin, M. T., Sudol, M., Bloch, L., and Winer, A. M.: Low-emitting urban forests: a taxonomic methodology for assigning isoprene and monoterpene emission rates, *Atmos. Environ.*, 30,
15 1437–1452, 1996.
- Bigi, A., Ghermandi, G., and Harrison, R. M.: Analysis of the air pollution climate at a background site in the Po valley, *J. Environ. Monitor.*, 14, 552–563, 2011.
- Blake, R. S., Monks, P. S., and Ellis, A. M.: Proton-transfer reaction mass spectrometry, *Chem. Rev.*, 109, 861–896, 2009.
- 20 Carslaw, D. C. and Ropkins, K.: Openair – an R package for air quality data analysis, *Environ. Modell. Softw.*, 27–28, 52–61, 2012.
- Dalponte, M., Gianelle, D., and Bruzzone, L.: Use of hyperspectral and LIDAR data for classification of complex forest areas, in: *Canopy Analysis and Dynamics of a Floodplain Forest*, Cierre Edizioni, Verona, Italy, 25–37, 2007.
- 25 Dalponte, M., Bruzzone, L., and Gianelle, D.: Fusion of hyperspectral and LIDAR remote sensing data for classification of complex forest areas, *IEEE T. Geosci. Remote*, 46, 1416–1427, 2008.
- Davison, B., Taipale, R., Langford, B., Misztal, P., Fares, S., Matteucci, G., Loreto, F., Cape, J. N., Rinne, J., and Hewitt, C. N.: Concentrations and fluxes of biogenic volatile
30 organic compounds above a Mediterranean macchia ecosystem in western Italy, *Biogeosciences*, 6, 1655–1670, doi:10.5194/bg-6-1655-2009, 2009.

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Decesari, S., Allan, J., Plass-Duelmer, C., Williams, B. J., Paglione, M., Facchini, M. C., O'Dowd, C., Harrison, R. M., Gietl, J. K., Coe, H., Giulianelli, L., Gobbi, G. P., Lanconelli, C., Carbone, C., Worsnop, D., Lambe, A. T., Ahern, A. T., Moretti, F., Tagliavini, E., Elste, T., Gilge, S., Zhang, Y., and Dall'Osto, M.: Measurements of the aerosol chemical composition and mixing state in the Po Valley using multiple spectroscopic techniques, *Atmos. Chem. Phys.*, 14, 12109–12132, doi:10.5194/acp-14-12109-2014, 2014.
- de Gouw, J. and Warneke, C.: Measurements of volatile organic compounds in the earth's atmosphere using proton transfer-reaction mass spectrometry, *Mass Spectrom. Rev.*, 26, 223–257, 2007.
- de Gouw, J. A., Goldan, P. D., Warneke, C., Kuster, W. C., Roberts, J. M., Marchewka, M., Bertman, S. B., Pszenny, A. A. P., and Keene, W. C.: Validation of proton transfer reaction-mass spectrometry (PTR-MS) measurements of gas-phase organic compounds in the atmosphere during the New England Air Quality Study (NEAQS) in 2002, *J. Geophys. Res.*, 108, D214682, doi:10.1029/2003JD003863, 2003.
- Dettmer, K. and Engewald, W.: Adsorbent materials commonly used in air analysis for adsorptive enrichment and thermal desorption of volatile organic compounds, *Anal. Biochem.*, 373, 490–500, 2002.
- Fall, R.: Abundant oxygenates in the atmosphere: a biochemical perspective, *Chem. Rev.*, 103, 4941–4951, 2003.
- Fares, S., Park, J.-H., Gentner, D. R., Weber, R., Ormeño, E., Karlik, J., and Goldstein, A. H.: Seasonal cycles of biogenic volatile organic compound fluxes and concentrations in a California citrus orchard, *Atmos. Chem. Phys.*, 12, 9865–9880, doi:10.5194/acp-12-9865-2012, 2012.
- Fares, S., Schnitzhofer, R., Jiang, X., Guenther, A., Hansel, A., and Loreto, F.: Observations of diurnal to weekly variations of monoterpene-dominated fluxes of volatile organic compounds from mediterranean forests: implications for regional modeling, *Environ. Sci. Technol.*, 47, 11073–11082, 2013.
- Fehsenfeld, F., Calvert, J., Fall, R., Goldan, P., Guenther, A. B., Hewitt, C. N., Lamb, B., Liu, S., Trainer, M., Westberg, H., and Zimmerman, P.: Emissions of volatile organic compounds from vegetation and the implications for atmospheric chemistry, *Global Biogeochem. Cy.*, 6, 389–430, 1992.
- Foken, T. and Wichura, B.: Tools for quality assessment of surface-based flux measurements, *Agr. Forest Meteorol.*, 78, 83–105, 1996.

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Foken, T., Göckede M., Mauder, M., Mahrt, L., Amiro, B., and Munger, W.: Post-field data quality control, in: Handbook of Micrometeorology: a Guide for Surface Flux Measurement and Analysis, edited by: Lee, W. M. X. and Law, B., Kluwer Academic Publishers, Dordrecht, the Netherlands, 29, 181–203, 2004.

- 5 Ghirardo, A., Koch, K., Taipale, R., Zimmer, I., Schnitzler, J.-P., and Rinne, J.: Determination of de novo and pool emissions of terpenes from four common boreal/alpine trees by $^{13}\text{CO}_2$ labelling and PTR-MS analysis, Plant Cell Environ., 33, 781–792, 2010.

Goudriaan, J. and van Laar, H.: Modelling Potential Crop Growth Processes, Textbook with Exercises, Kluwer Academic Publishers, Dordrecht, the Netherlands, 1994.

- 10 Graus, M., Müller, M., and Hansel, A.: High resolution PTR-TOF: quantification and formula confirmation of VOC in real time, J. Am. Soc. Mass Spectr., 21, 1037–1044, 2010.

Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W. A., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.: A global model for natural volatile organic compound emissions, J. Geophys. Res., 100, 8873–8892, 1995.

- 15 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), Atmos. Chem. Phys., 6, 3181–3210, doi:10.5194/acp-6-3181-2006, 2006.

- 20 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, Geosci. Model Dev., 5, 1471–1492, doi:10.5194/gmd-5-1471-2012, 2012.

- 25 Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, Atmos. Chem. Phys., 9, 5155–5236, doi:10.5194/acp-9-5155-2009, 2009.

- 30 Hansel, A., Jordan, A., Holzinger, R., Prazeller, P., Vogel, W., and Lindinger, W.: Proton transfer reaction mass spectrometry: on-line trace gas analysis at the ppb level, Int. J. Mass Spectrom., 149, 609–619, 1995.

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hayward, S., Hewitt, C. N., Sartin, J. H., and Owen, S. M.: Performance characteristics and applications of a proton transfer reaction-mass spectrometer for measuring volatile organic compounds in ambient air, *Environ. Sci. Technol.*, 36, 1554–1560, 2002.

Hewitt, C. N., Hayward, S., and Tani, A.: The application of proton transfer reaction-mass spectrometry (PTR-MS) to the monitoring and analysis of volatile organic compounds in the atmosphere, *J. Environ. Monitor.*, 5, 1–7, 2003.

Jardine, K. J., Monson, R. K., Abrell, L., Saleska, S. R., Arneth, A., Jardine, A., Ishida, F. Y., Serrano, A. M. Y., Artaxo, P., Karl, T., Fares, S., Goldstein, A., Loreto, F., and Huxman, T.: Within-plant isoprene oxidation confirmed by direct emissions of oxidation products methyl vinyl ketone and methacrolein, *Glob. Change Biol.*, 18, 973–984, 2012.

Jordan, A., Haidacher, S., Hanel, G., Hartungen, E., Märk L., Seehauser, H., Schottkowsky, R., Sulzer, P., and Märk, T. D.: A high resolution and high sensitivity proton-transfer-reaction time-of-flight mass spectrometer (PTR-TOF-MS), *Int. J. Mass Spectrom.*, 286, 122–128, 2009.

Junninen, H., Ehn, M., Petäjä, T., Luosujärvi, L., Kotiaho, T., Kostianen, R., Rohner, U., Gonnin, M., Fuhrer, K., Kulmala, M., and Worsnop, D. R.: A high-resolution mass spectrometer to measure atmospheric ion composition, *Atmos. Meas. Tech.*, 3, 1039–1053, doi:10.5194/amt-3-1039-2010, 2010.

Kalogridis, C., Gros, V., Sarda-Estève, R., Langford, B., Loubet, B., Bonsang, B., Bonnaire, N., Nemitz, E., Genard, A.-C., Boissard, C., Fernandez, C., Ormeño, E., Baisnée, D., Reiter, I., and Lathièrre, J.: Concentrations and fluxes of isoprene and oxygenated VOCs at a French Mediterranean oak forest, *Atmos. Chem. Phys.*, 14, 10085–10102, doi:10.5194/acp-14-10085-2014, 2014.

Karl, M., Guenther, A., Köble, R., Leip, A., and Seufert, G.: A new European plant-specific emission inventory of biogenic volatile organic compounds for use in atmospheric transport models, *Biogeosciences*, 6, 1059–1087, doi:10.5194/bg-6-1059-2009, 2009.

Karl, T., Guenther, A., Spirig, C., Hansel, A., and Fall, R.: Seasonal variation of biogenic VOC emissions above a mixed hardwood forest in northern Michigan. *Geophys. Res. Lett.*, 30, 2186, doi:10.1029/2003GL018432, 2003.

Karl, T., Potosnak, M., Guenther, A., Clark, D., Walker, J., Herrick, J. D., and Geron, C.: Exchange processes of volatile organic compounds above a tropical rain forest: implications for modeling tropospheric chemistry above dense vegetation, *J. Geophys. Res.*, 109, D18306, doi:10.1029/2004JD004738, 2004.

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Karl, T. G., Spirig, C., Rinne, J., Stroud, C., Prevost, P., Greenberg, J., Fall, R., and Guenther, A.: Virtual disjunct eddy covariance measurements of organic compound fluxes from a subalpine forest using proton transfer reaction mass spectrometry, *Atmos. Chem. Phys.*, 2, 279–291, doi:10.5194/acp-2-279-2002, 2002.

- 5 Keenan, T., Niinemets, Ü., Sabate, S., Gracia, C., and Peñuelas, J.: Process based inventory of isoprenoid emissions from European forests: model comparisons, current knowledge and uncertainties, *Atmos. Chem. Phys.*, 9, 4053–4076, doi:10.5194/acp-9-4053-2009, 2009.

Kormann, R. and Meixner, F. X.: An analytical footprint model for non-neutral stratification, *Bound.-Lay. Meteorol.*, 99, 207–224, 2001.

- 10 Langford, B., Davison, B., Nemitz, E., and Hewitt, C. N.: Mixing ratios and eddy covariance flux measurements of volatile organic compounds from an urban canopy (Manchester, UK), *Atmos. Chem. Phys.*, 9, 1971–1987, doi:10.5194/acp-9-1971-2009, 2009.

Langford, B., Misztal, P. K., Nemitz, E., Davison, B., Helfter, C., Pugh, T. A. M., MacKenzie, A. R., Lim, S. F., and Hewitt, C. N.: Fluxes and concentrations of volatile organic compounds from a South-East Asian tropical rainforest, *Atmos. Chem. Phys.*, 10, 8391–8412, doi:10.5194/acp-10-8391-2010, 2010a.

- 15 Langford, B., Nemitz, E., House, E., Phillips, G. J., Famulari, D., Davison, B., Hopkins, J. R., Lewis, A. C., and Hewitt, C. N.: Fluxes and concentrations of volatile organic compounds above central London, UK, *Atmos. Chem. Phys.*, 10, 627–645, doi:10.5194/acp-10-627-2010, 2010b.

Langford, B., Acton, W., Ammann, C., Valach, A., and Nemitz, E.: Eddy-covariance data with low signal-to-noise ratio: time-lag determination, uncertainties and limit of detection, *Atmos. Meas. Tech. Discuss.*, 8, 2913–2955, doi:10.5194/amtd-8-2913-2015, 2015.

- 25 Lindinger, W., Hansel, A., and Jordan, A.: Proton-transfer-reaction mass spectrometry (PTR-MS): on-line monitoring of volatile organic compounds at pptv levels, *Chem. Soc. Rev.*, 27, 347–354, 1998.

Loreto, F., Fischbach, R. J., Schnitzler, J. P., Ciccioli, P., Brancaleoni, E., Calfapietra, C., Seufert, G.: Monoterpene emission and monoterpene synthase activities in the Mediterranean evergreen oak *Quercus ilex* L. grown at elevated CO₂ concentrations, *Glob. Change Biol.*, 7, 709–717, 2001.

- 30 Lueken, D. J., Hutzell, W. T., Strum, M. L., and Pouliot, G. A.: Regional sources of atmospheric formaldehyde and acetaldehyde, and implications for atmospheric modelling, *Atmos. Environ.*, 47, 477–490, 2012.

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Misztal, P. K., Nemitz, E., Langford, B., Di Marco, C. F., Phillips, G. J., Hewitt, C. N., MacKenzie, A. R., Owen, S. M., Fowler, D., Heal, M. R., and Cape, J. N.: Direct ecosystem fluxes of volatile organic compounds from oil palms in South-East Asia, *Atmos. Chem. Phys.*, 11, 8995–9017, doi:10.5194/acp-11-8995-2011, 2011.

5 Monks, P. S., Granier, C., Fuzzi, S., et al.: Atmospheric composition change – global and regional air quality, *Atmos. Environ.*, 43, 5268–5350, doi:10.1016/j.atmosenv.2009.08.021, 2009.

Moncrieff, J., Clement, R., Finnigan, J., and Meyers, T.: Averaging, detrending, and filtering of eddy covariance time series, in: *Handbook of Micrometeorology: a Guide for Surface Flux Measurement and Analysis*, edited by: Lee, W. M. X. and Law, B., Kluwer Academic Publishers, Dordrecht, the Netherlands, 2004.

10 Müller, J.-F., Stavrou, T., Wallens, S., De Smedt, I., Van Roozendaal, M., Potosnak, M. J., Rinne, J., Munger, B., Goldstein, A., and Guenther, A. B.: Global isoprene emissions estimated using MEGAN, ECMWF analyses and a detailed canopy environment model, *Atmos. Chem. Phys.*, 8, 1329–1341, doi:10.5194/acp-8-1329-2008, 2008.

15 Müller, M., Mikoviny, T., Jud, W., D’Anna, B., and Wisthaler, A.: A new software tool for the analysis of high resolution PTR-TOF mass spectra, *Chemometr. Intell. Lab.*, 127, 158–165, 2013.

Neftel, A., Spirig, C., and Ammann, C.: Application and test of a simple tool for operational footprint evaluations, *Environ. Pollut.*, 152, 644–652, 2008.

20 Nemitz, E., Hargreaves, K. J., Neftel, A., Loubet, B., Cellier, P., Dorsey, J. R., Flynn, M., Hensen, A., Weidinger, T., Meszaros, R., Horvath, L., Dämmgen, U., Frühauf, C., Löpmeier, F. J., Gallagher, M. W., and Sutton, M. A.: Intercomparison and assessment of turbulent and physiological exchange parameters of grassland, *Biogeosciences*, 6, 1445–1466, doi:10.5194/bg-6-1445-2009, 2009.

25 Niinemets, Ü: Research review, Components of leaf dry mass per area – thickness and density – alter leaf photosynthetic capacity in reverse directions in woody plants, *New Phytol.*, 144, 35–47, 1999.

Owen, S. M., Boissard, C., and Hewitt, C. N.: Volatile organic compounds (VOCs) emitted from 30 40 Mediterranean plant species: VOC speciation and extrapolation to habitat scale, *Atmos. Environ.*, 35, 5393–5409, 2001.

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Park, J.-H., Goldstein, A. H., Timkovsky, J., Fares, S., Weber, R., Karlik, J., and Holzinger, R.: Active atmosphere-ecosystem exchange of the vast majority of detected volatile organic compounds, *Science*, 341, 643–647, 2013.

R Core Team: R: a Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, available at: <http://www.R-project.org/> (last access: 18 September 2015), 2012.

Riemer, D., Pos, W., Milne, P., Farmer, C., Zika, R., Apel, E., Olszyna, K., Kliendienst, T. Lonneman, W., Bertman, S., Shepson, P., and Starn, T.: Observations of nonmethane hydrocarbons and oxygenated volatile organic compounds at a rural site, *J. Geophys. Res.*, 103, 28111–28128, 1998.

Rinne, J. and Ammann, C.: Disjunct Eddy covariance method, in: *Eddy Covariance: a Practical Guide to Measurement and Data*, edited by: Aubinet, M., Vesala, T., and Papale, D., Springer, New York, USA, 2012.

Rinne, J., Taipale, R., Markkanen, T., Ruuskanen, T. M., Hellén, H., Kajos, M. K., Vesala, T., and Kulmala, M.: Hydrocarbon fluxes above a Scots pine forest canopy: measurements and modeling, *Atmos. Chem. Phys.*, 7, 3361–3372, doi:10.5194/acp-7-3361-2007, 2007.

Rinne, J., Douffet, T., Prigent, Y., and Durand, P.: Field comparison of disjunct and conventional eddy covariance techniques for trace gas flux measurements, *Environ. Pollut.*, 152, 630–635, 2008.

Rivera-Rios, J. C., Nguyen, T. B., Crounse, J. D., Jud, W., St. Clair, J. M., Mikoviny, T., Gilman, J. B., Lerner, B. M., Kaiser, J. B., de Gouw, J., Wisthaler, A., Hansel, A., Wennberg, P. O., Seinfeld, J. H., and Keutsch, F. N.: Conversion of hydroperoxides to carbonyls in field and laboratory instrumentation: observational bias in diagnosing pristine versus anthropogenically controlled atmospheric chemistry, *Geophys. Res. Lett.*, 41, 8645–8651, 2014.

Royal Society: Ground-Level Ozone in the 21st Century: Future Trends, Impacts and Policy Implications: Science Policy report 15/08, The Royal Society, London, UK, 2008.

Schallhart, S., Rantala, P., Nemitz, E., Mogensen, D., Tillmann, R., Mentel, T. F., Rinne, J., and Ruuskanen, T. M.: Characterization of total ecosystem scale biogenic VOC exchange at a Mediterranean oak-hornbeam forest, *Atmos. Chem. Phys. Discuss.*, 15, 27627–27673, doi:10.5194/acpd-15-27627-2015, 2015.

Spirig, C., Neftel, A., Ammann, C., Dommén, J., Grabmer, W., Thielmann, A., Schaub, A., Beauchamp, J., Wisthaler, A., and Hansel, A.: Eddy covariance flux measurements of bio-

genic VOCs during ECHO 2003 using proton transfer reaction mass spectrometry, Atmos. Chem. Phys., 5, 465–481, doi:10.5194/acp-5-465-2005, 2005.

Taipale, R., Ruuskanen, T. M., Rinne, J., Kajos, M. K., Hakola, H., Pohja, T., and Kulmala, M.: Technical Note: Quantitative long-term measurements of VOC concentrations by PTR-MS – measurement, calibration, and volume mixing ratio calculation methods, Atmos. Chem. Phys., 8, 6681–6698, doi:10.5194/acp-8-6681-2008, 2008.

Taipale, R., Ruuskanen, T. M., and Rinne, J.: Lag time determination in DEC measurements with PTR-MS, Atmos. Meas. Tech., 3, 853–862, doi:10.5194/amt-3-853-2010, 2010.

Tani, A., Hayward, S., Hansel, A., and Hewitt, C. N.: Effect of water vapour pressure on monoterpene measurements using proton transfer reaction-mass spectrometry (PTR-MS), Int. J. Mass Spectrom., 239, 161–169, 2004.

Wienhold, F. G., Frahm, H., and Harris, G. W.: Measurements of N₂O fluxes from fertilized grassland using a fast response tunable diode laser spectrometer, J. Geophys. Res., 99, 16557–16567, 1994.

Wohlfahrt, G., Amelynck, C., Ammann, C., Arneth, A., Bamberger, I., Goldstein, A. H., Gu, L., Guenther, A., Hansel, A., Heinesch, B., Holst, T., Hörtnagl, L., Karl, T., Laffineur, Q., Nef, A., McKinney, K., Munger, J. W., Pallardy, S. G., Schade, G. W., Seco, R., and Schoon, N.: An ecosystem-scale perspective of the net land methanol flux: synthesis of micrometeorological flux measurements, Atmos. Chem. Phys. Discuss., 15, 2577–2613, doi:10.5194/acpd-15-2577-2015, 2015.

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Unit masses measured using the PTR-MS during the ÉCLAIRE campaign at Bosco Fontana and the exact masses observed using the PTR-ToF-MS. Where the PTR-MS sensitivity was calculated directly from a compound in the calibration standard this compound is indicated in brackets, at m/z 61 and 71 the sensitivity was calculated from a transmission curve.

Unit mass (PTR-MS)	Exact mass (PTR-ToF-MS)	Contributing compound(s)	Formula	PTR-MS sensitivity (ncps ppbv ⁻¹)
21	21.023	Water isotope	H ₃ ¹⁸ O ⁺	–
33	32.997	Oxygen isotope	O ¹⁷ O ⁺	11.60 (methanol)
	33.033	Methanol	CH ₅ O ⁺	
39	39.033	Water cluster	H ₅ O ¹⁸ O ⁺	–
45	44.997	Protonated carbon dioxide	C ₁ H ₃ O ₂ ⁺	9.90 (acetaldehyde)
	45.033	Acetaldehyde	C ₂ H ₅ O ⁺	
59	59.049	Acetone	C ₃ H ₇ O ⁺	8.82 (acetone)
	59.049	Propanal	C ₃ H ₇ O ⁺	
61	61.028	Acetic acid	C ₂ H ₅ O ₂	8.40 (transmission curve)
69	69.0699	Isoprene	C ₅ H ₉ ⁺	3.80 (isoprene)
	69.0699	2-Methyl-3-buten-2-ol fragment	C ₅ H ₉ ⁺	
	69.0699	Methyl butanal fragment	C ₅ H ₉ ⁺	
71	71.049	Methyl vinyl ketone	C ₄ H ₇ O ⁺	5.29 (transmission curve)
	71.049	Methacrolein	C ₄ H ₇ O ⁺	
	71.085	Unknown	C ₅ H ₁₁ ⁺	
73	73.026	Unknown	C ₃ H ₅ O ₂ ⁺	5.87 (Methyl ethyl ketone)
	73.047	Unknown	Unknown	
	73.065	Methyl ethyl ketone	C ₄ H ₉ O ⁺	
	73.065	Butanal	C ₄ H ₉ O ⁺	
81	80.997	Unknown	C ₄ H ₁ O ₂ ⁺	1.59 (α -pinene fragment)
	81.033	Unknown	C ₅ H ₅ O ⁺	
	81.070	Monoterpene fragment	C ₆ H ₉ ⁺	
	81.070	Hexenal fragment	C ₆ H ₉ ⁺	
137	137.056	Unknown	Unknown	0.16 (α -pinene)
	137.133	Monoterpenes	C ₁₀ H ₁₇ ⁺	

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Table 2. Summary of the bVOC mixing ratios (ppbv) recorded at 4 m above the forest canopy during the Bosco Fontana measurement campaign and limits of detection (LoD, ppbv), based on 25 min averages.

<i>m/z</i>	33	45	59	61	69	71	73	81
Compound	Methanol	Acetaldehyde	Acetone	Acetic acid	Isoprene	MVK + MACR	MEK	Monoterpenes
Max	14.6	3.44	7.31	14.9	4.79	1.95	1.05	0.419
Min	2.13	< LOD	1.18	0.396	< LOD	0.083	0.097	< LOD
Mean	6.16	1.46	3.24	1.92	1.07	0.506	0.454	0.198
Standard deviation	2.52	0.67	0.91	1.09	0.80	0.28	0.21	0.07
Median	5.69	1.30	3.14	1.73	0.934	0.506	0.428	0.199
1st Quartile	4.19	0.964	2.68	1.22	0.409	0.325	0.311	0.140
3rd Quartile	7.53	1.87	3.82	2.31	1.53	1.95	0.568	0.245
LOD	0.436	0.712	0.239	0.141	0.167	0.081	0.048	0.067

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Table 3. Summary of the BVOC fluxes ($\text{mg m}^{-2} \text{h}^{-1}$) recorded during the Bosco Fontana field campaign based on 25 min values. Values in brackets cover the campaign period where data is available from both instruments to enable direct comparison (15 June–6 July 2012 and 15–25 June 2012 for isoprene and monoterpenes, respectively).

<i>m/z</i>	33	45	59	61	69		71	73	81	
Compound	Methanol	Acetaldehyde	Acetone	Acetic acid	Isoprene PTR-MS	Isoprene PTR-ToF-MS	MVK + MACR	MEK	Monoterpenes PTR-MS	Monoterpenes PTR-ToF-MS
Max emission flux	0.492	0.436	0.585	0.328	9.867 (9.867)	9.195 (9.195)	0.641	0.181	0.478 (0.478)	0.609 (0.603)
Max deposition flux	−1.589	−0.335	−0.692	−0.876	−0.238 (−0.238)	−0.305 (−0.305)	−0.457	−0.128	−0.167 (−0.167)	−0.065 (−0.057)
1st Quartile	−0.032	−0.011	−0.029	−0.044	0.005 (0.005)	0.019 (0.019)	−0.012	−0.012	−0.009 (−0.008)	0.005 (0.001)
3rd Quartile	0.070	0.053	0.057	0.033	1.624 (1.796)	2.661 (2.661)	0.054	0.024	0.093 (0.101)	0.159 (0.137)
Mean	0.017	0.024	0.016	−0.007	0.961 (1.003)	1.465 (1.465)	0.025	0.009	0.056 (0.060)	0.098 (0.088)
Standard deviation	0.123	0.067	0.098	0.091	1.369 (1.387)	1.911 (1.911)	0.076	0.039	0.108 (0.111)	0.138 (0.134)
Median	0.010	0.013	0.008	0.000	0.168 (0.199)	0.410 (0.410)	0.011	0.005	0.020 (0.021)	0.036 (0.028)
Mean day-time flux (06:00–18:00 LT)	0.033	0.045	0.030	0.001	1.912 (1.978)	2.917 (2.917)	0.049	0.018	0.117 (0.120)	0.206 (0.207)
Standard deviation	0.161	0.082	0.125	0.096	1.401 (1.383)	1.842 (1.842)	0.095	0.050	0.141 (0.129)	0.141 (0.144)
Median day-time flux (06:00–18:00 LT)	0.038	0.044	0.026	0.001	1.635 (1.790)	2.905 (2.905)	0.041	0.014	0.090 (0.099)	0.192 (0.164)

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Table 4. Non-exhaustive summary of isoprene fluxes recorded in the Mediterranean region and the isoprene emission factor under basal conditions (temperature: 30 °C and PAR: 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$).

Ecosystem	Dominant species	Season	Mean day time isoprene flux ($\text{mg m}^{-2} \text{h}^{-1}$)	Isoprene emission factor under basal conditions ($\text{mg m}^{-2} \text{h}^{-1}$)	Reference
Mixed oak and hornbeam forest	<i>Carpinus betulus</i> <i>Quercus robur</i>	Summer	2.6	1.7	This study
Oak forest	<i>Quercus pubescens</i>	Spring	2.8	7.4	Kalogridis et al. (2014)
Oak forest	<i>Quercus pubescens</i>	Summer	5.4–10.1	5.4	Baghi et al. (2012)
Mixed oak and pine forest	<i>Pinus pinea</i> <i>Quercus ilex</i> <i>Quercus suber</i>	Autumn	ca. 0.13	0.61	Fares et al. (2013)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Table 5. Leaf level isoprene and monoterpene emission ($\text{mg m}^{-2} \text{h}^{-1}$) from single leaves under basal conditions (temperature: 30°C and PAR: $1000 \mu\text{mol m}^{-2} \text{s}^{-1}$). ND signifies not detected.

Tree species	isoprene flux (standard error)	α -pinene flux (standard error)	sabinene flux (standard error)	β -pinene flux (standard error)	limonene flux (standard error)	sum monoterpene flux
<i>Carpinus betulus</i>	2.25×10^{-3} (1.50×10^{-3})	1.07×10^{-2} (6.00×10^{-3})	1.81×10^{-2} (1.36×10^{-2})	5.14×10^{-2} (1.23×10^{-2})	5.83×10^{-1} (2.36×10^{-1})	6.63×10^{-1}
<i>Quercus robur</i>	2.39×10^0 (6.12×10^{-1})	2.81×10^{-2} (1.45×10^{-2})	ND	4.70×10^{-3} (3.08×10^{-3})	2.16×10^{-1} (6.49×10^{-2})	2.49×10^{-1}
<i>Quercus rubra</i>	9.14×10^{-1} (2.02×10^{-1})	ND	ND	7.95×10^{-3} (2.22×10^{-3})	2.34×10^{-2} (7.11×10^{-3})	3.13×10^{-2}
<i>Corylus avellana</i>	4.97×10^{-4} (3.93×10^{-4})	1.30×10^{-2} (8.00×10^{-3})	ND	2.08×10^{-2} (4.80×10^{-3})	7.57×10^{-1} (4.15×10^{-1})	7.90×10^{-1}
<i>Acer campestre</i>	4.40×10^{-4} (3.11×10^{-4})	5.14×10^{-2} (2.95×10^{-2})	ND	2.27×10^{-1} (3.54×10^{-2})	1.07×10^{-1} (1.41×10^{-2})	3.85×10^{-1}
<i>Sambucus nigra</i>	4.09×10^{-3} (3.66×10^{-3})	ND	ND	9.67×10^{-3} (2.69×10^{-3})	2.49×10^{-1} (1.41×10^{-1})	2.59×10^{-1}
<i>Cornus sanguinea</i>	4.00×10^{-1} (4.00×10^{-1})	1.11×10^{-3} (1.11×10^{-3})	ND	1.95×10^{-2} (4.91×10^{-3})	2.28×10^{-1} (1.73×10^{-1})	2.49×10^{-1}

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 6. Species specific isoprene and monoterpene emission factors (for a standard temperature of 30 °C and a PAR value of 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$) derived from optimising the leaf level emission factors to give the best fit with the measured above canopy isoprene and monoterpene fluxes within the constraints displayed.

Species	Isoprene emission factor ($\text{mg m}^{-2} \text{h}^{-1}$)	Isoprene constraint ($\text{mg m}^{-2} \text{h}^{-1}$)	Monoterpene emission factor ($\text{mg m}^{-2} \text{h}^{-1}$)	Monoterpene constraint ($\text{mg m}^{-2} \text{h}^{-1}$)
<i>Acer campestre</i>	0.00	< 1.0	0.15	< 0.50
<i>Acer negundo</i>	0.00	< 1.0	0.33	< 0.64
<i>Alnus glutinosa</i>	0.01	< 1.0	0.22	< 0.50
<i>Carpinus betulus</i>	0.00	< 1.0	0.57	< 0.63
<i>Corylus avellana</i>	0.00	< 1.0	0.23	< 0.50
<i>Fraxinus angustifolia</i>	0.00	< 1.0	0.00	< 0.50
<i>Juglans nigra</i>	0.00	< 1.0	0.12	< 0.50
<i>Juglans regia</i>	0.36	< 1.0	0.15	< 0.50
<i>Morus</i> sp.	0.00	< 1.0	0.19	< 0.50
<i>Platanus hispanica</i>	2.97	< 4.4	0.50	< 0.50
<i>Populus</i> × <i>canescens</i>	10.66	< 16.1	0.29	< 0.50
<i>Populus</i> × <i>hybrida</i>	8.06	< 16.1	0.00	< 0.50
<i>Prunus avium</i>	0.00	< 1.0	0.01	< 0.50
<i>Quercus cerris</i>	0.02	< 1.0	0.07	< 0.50
<i>Quercus robur</i>	7.46	< 16.1	0.19	< 0.50
<i>Quercus rubra</i>	1.38	< 8.1	0.02	< 0.50
<i>Robinia pseudoacacia</i>	1.38	< 2.8	0.01	< 0.50
<i>Rubus</i> sp.	0.00	< 1.0	0.01	< 0.50
<i>Tilia</i> sp.	0.00	< 1.0	0.00	< 0.50
<i>Ulmus minor</i>	0.01	< 1.0	0.01	< 0.50
Grass	0.06	< 1.0	0.06	< 0.15
Not woodland	0.06	< 1.0	0.08	< 0.15
Outside forest	0.06	< 1.0	0.06	< 0.50




Figure 1. Time series of meteorological conditions recorded over the campaign period. From top to bottom: PAR ($\mu\text{mol m}^{-2} \text{s}^{-1}$), air temperature ($^{\circ}\text{C}$), relative humidity (%), wind speed (m s^{-1}) and wind direction ($^{\circ}$).

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.




Figure 2. Wind direction and wind speed observed during the Bosco Fontana field campaign.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Figure 3. Mean 4 m above-canopy diurnal volume mixing ratios of volatile organic compounds measured during the Bosco Fontana field campaign. Error bars represent one standard deviation from the mean and the dashed line denotes limit of detection.

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.




Figure 4. Time series of isoprene (top) and monoterpene (bottom) fluxes ($\text{mg m}^{-2} \text{h}^{-1}$) measured using the method of vDEC. Blue circles, triangles and diamonds represent 25 min averaged flux data from the PTR-MS which respectively passed all tests, fell below the u_* threshold or the LoD. Red circles and lines represent PTR-ToF-MS isoprene and monoterpene fluxes with 30 min averaged flux files failing the stationarity test removed.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Figure 5. Mean diurnal fluxes of volatile organic compounds measured using vDEC. Shaded area represents the limit of detection of the averaged data, and error bars represent one standard deviation between days from the mean.

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.




Figure 6. Comparison of PTR-MS (blue) and PTR-ToF-MS (red) mass scans relative to m/z 59 at unit mass resolution averaged between 14 and 24 June. Compounds recorded in flux mode using the PTR-MS are presented in black with compounds tentatively identified in grey.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 7. Scatter plots displaying the relationship between the volume mixing ratios of methanol, acetone and MEK measured 4 m above the canopy, coloured by temperature.




Figure 8. The exponential relationship between temperature ($^{\circ}\text{C}$) and isoprene fluxes ($\text{mg m}^{-2} \text{h}^{-1}$) and volume mixing ratios (ppbv), coloured according to the magnitude of photosynthetically active radiation ($\mu\text{mol m}^{-1} \text{s}^{-1}$).




Figure 9. Measured isoprene fluxes against the product of γ (emission activity factor, itself the product of the temperature, light and leaf area index activity factors) and ρ (the canopy loss and production factor).

Canopy-scale flux measurements and bottom-up emission estimates

W. J. F. Acton et al.




Figure 10. The contribution of individual tree species to the speciated isoprene and monoterpene flux on the 2 July 2012 with PAR displayed as a yellow line, wind direction as a black line and the flux recorded using the PTR-MS as bold black bars.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

