

## **Emission of monoterpenes from European beech (*Fagus sylvatica* L.) as a function of light and temperature**

T. Dindorf, U. Kuhn, L. Ganzeveld, G. Schebeske, P. Ciccioli, C. Holzke, R. Köble, G. Seufert, J. Kesselmeier

► **To cite this version:**

T. Dindorf, U. Kuhn, L. Ganzeveld, G. Schebeske, P. Ciccioli, et al.. Emission of monoterpenes from European beech (*Fagus sylvatica* L.) as a function of light and temperature. Biogeosciences Discussions, European Geosciences Union, 2005, 2 (1), pp.137-182. <hal-00297729>

**HAL Id: hal-00297729**

**<https://hal.archives-ouvertes.fr/hal-00297729>**

Submitted on 27 Jan 2005

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Biogeosciences Discussions is the access reviewed discussion forum of Biogeosciences

## Emission of monoterpenes from European beech (*Fagus sylvatica* L.) as a function of light and temperature

T. Dindorf<sup>1</sup>, U. Kuhn<sup>1</sup>, L. Ganzeveld<sup>2</sup>, G. Schebeske<sup>1</sup>, P. Ciccioli<sup>3</sup>, C. Holzke<sup>4</sup>, R. Köble<sup>5</sup>, G. Seufert<sup>5</sup>, and J. Kesselmeier<sup>1</sup>

<sup>1</sup>Max Planck Institute for Chemistry, Department of Biogeochemistry, Johann-Joachim Becher Weg 27, D-55128, Mainz, Germany

<sup>2</sup>Max Planck Institute for Chemistry, Department of Airchemistry, Johann-Joachim Becher Weg 27, D-55128, Mainz, Germany

<sup>3</sup>Istituto di Metodologie Chimiche, Area della Ricerca del CNR di Montelibretti, 00019 Monterotondo Scalo, Italy

<sup>4</sup>Research Centre Jülich GmbH, Institute II: Troposphere, 52425 Jülich, Germany

<sup>5</sup>Joint Research Centre Ispra, Institute for Environment and Sustainability, T.P. 051, 21020 Ispra, Italy

Received: 17 December 2004 – Accepted: 25 January 2005 – Published: 27 January 2005

Correspondence to: T. Dindorf (dindorf@mpch-mainz.mpg.de)

© 2005 Author(s). This work is licensed under a Creative Commons License.

### Monoterpene emission from European beech

T. Dindorf et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

EGU

## Abstract

Using a dynamic branch enclosure technique European beech (*Fagus sylvatica* L.) was characterised as a strong emitter of monoterpenes, with sabinene being the predominant compound released. Since monoterpene emission was demonstrated to be a function of light and temperature, application of light and temperature dependent algorithms resulted in reasonable agreement with the measured data. Furthermore, during high temperature periods the depression of net CO<sub>2</sub> exchange during midday (midday depression) was accompanied by a depression of monoterpene emission on one occasion. The species dependent standard emission factor and the light and temperature regulated release of monoterpenes is of crucial importance for European VOC emissions. All measurements were performed within the framework of the ECHO project (Emission and Chemical transformation of biogenic volatile Organic compounds) during two intensive field campaigns in the summers of 2002 and 2003.

## 1. Introduction

The release of biogenic volatile organic compounds (VOCs) represents a substantial input of reactive trace gases into the atmosphere and influences atmospheric chemistry and physics (Andreae and Crutzen, 1997; Atkinson, 2000; Fehsenfeld et al., 1992; Went, 1960). Furthermore, VOC may represent a substantial loss of carbon for the biosphere (Guenther, 2002; Kesselmeier et al., 2002). The exchange (emission and deposition) of volatile organic compounds, involved in the oxidant cycle, aerosol production and carbon budget estimates, plays a crucial role in climate forcing but is poorly understood in view of the high number of different VOC species and their exchange regulations. The emission of isoprenoids, the dominating biogenic VOC fraction consisting mainly of isoprene and monoterpenes, has been investigated intensively during the last decades. However, our knowledge still is full of gaps. Until a decade ago a clear difference between the emission of isoprene and that of monoterpenes was

**BGD**

2, 137–182, 2005

## Monoterpene emission from European beech

T. Dindorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

## Monoterpene emission from European beech

T. Dindorf et al.

Title Page	Introduction	References	Figures
Abstract	Conclusions	Tables	►◄
◄	►	Back	Close
Full Screen / Esc			
Print Version			
Interactive Discussion			

postulated. Isoprene emission was regarded as dependent on light and temperature, whereas monoterpenes were thought to be produced as storage compounds and to be emitted as a function of temperature only. However, within the course of the EU-project “BEMA, Biogenic Emissions in the Mediterranean Area” (for an overview see Seufert et al., 1997), it became obvious that monoterpenes can be released in the same manner as isoprene (see Kesselmeier and Staudt, 1999), an observation which has recently been confirmed for the tropical rainforest (Kuhn et al., 2002b; Rinne et al., 2002). Meanwhile the release of monoterpenes from storage pools as a function of temperature only is discussed rather as a special case (Kesselmeier, 2004). Light dependence of monoterpene emission is in full agreement with the recent knowledge of biosynthesis of isoprenoids and the close relation between photosynthesis and isoprene/monoterpene production within chloroplasts (see Lichtenthaler, 1999). This knowledge has resulted in a substantial number of new questions concerning ecology and evolution (Lerdau and Gray, 2003), regulation and controls of production and emission (Niinemets and Reichstein, 2003; Peñuelas and Llusia, 2001; Wolfertz et al., 2003), as well as what the contribution of different carbon sources to isoprenoid biosynthesis are in order to understand production, accumulation and emission (Schnitzler et al., 2004). Several plant species of high importance for regional or global estimations have not been sufficiently investigated yet but are nevertheless included in budget calculations just by assigning emission rates based on plant family relationship (Karlik and Winer, 2001). Finally, the seasonal development of VOC emission capacity plays a significant role (Kesselmeier et al., 2003; Kuhn et al., 2004) and neglecting environmental effects and plant adaptations may lead to a significant under/overestimation of isoprenoid emission. Within the framework of the German ECHO project (Emission and Chemical transformation of biogenic volatile Organic compounds, AFO, 2000) we performed enclosure measurements during two intensive field campaigns in the summers of 2002 and 2003 in order to investigate the primary emission of volatile organic compounds from naturally growing >100 years old European beech (*Fagus sylvatica* L.). Until now this tree species has been regarded as a non-isoprene, but low emitter of

monoterpenes (see Kesselmeier and Staudt, 1999). As *Fagus sylvatica* L. is one of the dominant deciduous tree species occurring in Europe, the two years of measurements should help to clarify the significance of beech trees for regional and global budgets of VOCs.

## 5 2. Experimental

### 2.1. Site description

The measurement site was located in an urban area near the small city of Jülich, Germany. All experiments were carried out in a deciduous forest stand of about 3.5 km<sup>2</sup> size, partially located on the premises of the Research Centre Jülich. The location is characterised by moderate climatic conditions, with a mean annual precipitation of 685 mm and an average annual temperature of 10°C. Enclosure measurements were performed on top of a 24 m scaffold tower located at 50°54.321' N, 006°25.130' E. The predominant soil type of this forest area is luvic stagnosol that provides a moderate supply of nutrients for the growing plants. The area nearby the tower site was dominated by ~160 year old European beech trees of up to 28 m in height. The leaf area index (LAI) at this tower site showed a maximum density of 4.7 at ground level, quite typical for a European beech stand. A detailed overview of the measurement site is given by Auburn et al. (2005)<sup>1</sup>.

### 2.2. Enclosure measurements and plant material

20 Enclosure measurements for monoterpene emission from European beech (*Fagus sylvatica* L., plant family: *Fagaceae*) were carried out during two intensive field campaigns

<sup>1</sup>Auburn, S., Koppmann, R., Leiti, B., Möllmann-Coers, M., and Schaub, A.: Physical modelling of a complex forest area in a wind tunnel – Comparison with field data, Agricultural and forest Meteorology, submitted, 2005.

**BGD**

2, 137–182, 2005

## Monoterpene emission from European beech

T. Dindorf et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

EGU

in the summers of 2002 and 2003. All experiments were conducted by use of an open, dynamic (flow through) enclosure system, consisting of two identical cuvettes of ~75 l volume each. As proven earlier, the system can be regarded as inert for the relevant volatile organic compounds and allows the investigation of an enclosed branch for several days without visible effect of stress (for a detailed description see Gut et al., 2002; Kesselmeier et al., 1996, 1997, 1998; Kuhn et al., 2002a, b; Schäfer et al., 1992). Enclosure measurements were carried out over a period of 8 days in June 2002 and 16 days in July/August 2003 with the same branch of a ~160 year old beech tree. Leaf samples from the measured branch proved to be “sunlit leaves” by microscopical analysis of the leaf morphology. For measurements, the branch was enclosed in the sample cuvette as shown by Fig. 1, while the other cuvette remained empty as a reference. Leaf area, dry and fresh weight were calculated similar to Kuhn et al. (2002a). An overview of these parameters is given in Table 1. Micrometeorological parameters were recorded by standard sensors. Exchange rates for CO<sub>2</sub> and water vapour were measured by use of an infrared gas analyser (Model Li-7000, Licor, USA). All trace gas exchange rates were calculated by using the difference concentration between the branch enclosing sample cuvette and the empty reference cuvette according to Kuhn et al. (2002a, b). Unless indicated otherwise, leaf gas exchange rates were normalised to leaf dry weight. The stomatal conductance for water vapour was determined according to Pearcy et al. (1989). Uncertainties for gas exchange rates were assessed by conventional Gaussian error propagation. Ambient temperature and rainfall preceding the enclosure measurements were monitored by the meteorological station of the Research Centre Jülich at about 470 m distance from the measurement site on a second scaffolding tower at a height of 20 m.

### 2.3. Measurement of volatile organic compounds

Measurement of volatile organic compounds was performed by the use of solid adsorbents and subsequent analysis of the sampled compounds by GC-FID and GC-MS. Samples from the reference and branch cuvette, as well as from ambient air, were

141

## BGD

2, 137–182, 2005

### Monoterpene emission from European beech

T. Dindorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

collected simultaneously by use of three custom made automatic sampling systems that are described in detail by Kuhn et al. (2005)<sup>2</sup>. For GC-FID measurements samples were collected on Silicosteel cartridges (1/4" OD, 89 mm length, Restek, USA) packed with 130 mg Carbograph 1 (90 m<sup>2</sup> g<sup>-1</sup>) and 130 mg Carbograph 5 (560 m<sup>2</sup> g<sup>-1</sup>) (20–40 mesh each, Lara s.r.l., Italy). Samples were collected for 30 min at flow rates of 150 ml min<sup>-1</sup> resulting in a total sampling volume of 4500 ml. Analysis of the samples by GC-FID measurements was performed in the laboratory at the Max Planck Institute in Mainz. Cartridges were desorbed thermally for 10 min by use of a thermal-desorption system (Model ATD400, Perkin Elmer, Germany) at 260°C, that was connected to a GC-FID (Model AutoSystem XL, Perkin Elmer, Germany). Refocussing prior peak separation was accomplished by a small quartz tube packed with 20 mg Carbograph 1 kept at -30°C that was heated rapidly to 280°C. The separation of peaks was achieved by use of a dimethylpolysiloxane column (model HP-1, 100 m length, 0.25 mm ID, film thickness 0.5 µm, [Agilent Technologies, USA]) at a temperature program ranging from -10 to 40°C (20°C min<sup>-1</sup>), 40 to 145°C (1.5°C min<sup>-1</sup>) and 145 to 220°C (30°C min<sup>-1</sup>). In addition to several volatile organics, 10 different monoterpene compounds were evaluated from these analyses: camphene, Δ3-carene, p-cymene, limonene, myrcene, α-pinene, β-pinene, sabinene, α-terpinene, and γ-terpinene. The detection limit for monoterpene samples was calculated to <10 ppt (corresponding to a detection limit of 12 ng g<sup>-1</sup> h<sup>-1</sup> and 0.9 µg m<sup>-2</sup> h<sup>-1</sup> for *Fagus sylvatica* L.) (for details see Kuhn et al., 2002b). Calibration for this system was accomplished by use of a gaseous standard containing isoprene and several n-alkanes. Unless indicated otherwise, the following paragraphs will report on the total sum of monoterpene compounds measured by GC-FID or GC-MS.

Samples for GC-MS analysis were collected occasionally on glass tubes (6 mm OD, 160 mm length) that were packed sequentially with 118 mg Carbograph 2 (12 m<sup>2</sup> g<sup>-1</sup>),

<sup>2</sup>Kuhn, U., Dindorf, T., Ammann, C., Holzinger, R., Ausma, S., Kennitner, T., Helleis, F., and Kesselmeier, J.: Design and field application of an automated cartridge sampler for VOC concentration and flux measurements, J. Envir. Mon., submitted, 2005

## BGD

2, 137–182, 2005

### Monoterpene emission from European beech

T. Dindorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

60 mg Carbograph 1, and 115 mg Carbograph 5 (20–40 mesh each, Lara s.r.l., Italy). Analysis of the cartridges was carried out in the laboratory of CNR in Rome, Italy. Monoterpene compounds that were evaluated from these analyses were camphene,  $\Delta$ 3-carene, p-cymene, limonene, myrcene,  $\alpha$ -phellandrene,  $\beta$ -phellandrene,  $\alpha$ -pinene,  $\beta$ -pinene, sabinene,  $\alpha$ -terpinene,  $\gamma$ -terpinene, terpinolene,  $\alpha$ -thujene, and tricyclene. A detailed overview of the method used for GC-MS analysis is given by Ciccioni et al. (1992) and Brancaleoni et al. (1999). According to the results of subsequent laboratory tests, it was shown that sabinene partially decomposed to p-cymene,  $\alpha$ -phellandrene,  $\beta$ -phellandrene,  $\alpha$ -terpinene,  $\gamma$ -terpinene, terpinolene, and  $\alpha$ -thujene during the storage time of these cartridges. Thus a correction factor was applied to the relevant compounds.

#### 2.4. Calculation of European VOC emission

To assess the potential implication of monoterpene emissions from *Fagus sylvatica* L. on a European scale, we applied an offline version of the Guenther et al. (1995) VOC emission algorithm, normally applied for global scale studies (Ganzeveld et al., 2002). The algorithm uses the Olson (1992) global ecosystem database, which distinguishes 72 ecosystems at a 0.5 $\times$ 0.5 grid resolution, combined with a 5-year climatology of monthly NDVI (Normalized Differential Vegetation Index) satellite data (Gutman et al., 1995) to infer global surface cover properties. VOC emissions are calculated as a function of ecosystem specific emission factors, surface radiation, temperature, foliar density and its vertical distribution. The latter is required to calculate the within-canopy profiles of photosynthetic active radiation (Weiss and Norman, 1985) and distinguishes four canopy layers based on the sensitivity of the emissions on the vertical resolution (Ganzeveld et al., 2002). Normally, the emission algorithm is applied in the chemistry-climate model ECHAM to calculate the biogenic VOC emissions and their role for atmospheric chemistry online from the model's surface temperature and net radiation. However, for the offline calculations presented, we have applied the temperature and net radiation output fields of a ECHAM T106 (~125 km resolution) simulation for the

## BGD

2, 137–182, 2005

### Monoterpene emission from European beech

T. Dindorf et al.

Title Page	Introduction
Abstract	References
Conclusions	Figures
Tables	
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

EGU

month of July at 6 h time intervals. For a comparison with the global scale algorithm, we have used a high-resolution dataset that describes the European distribution of *Fagus sylvatica* L. at a 1 × 1 km grid resolution (Köble and Seufert, 2001) in combination with the measured monoterpene emission factor.

### 3. Results and discussion

#### 3.1. Diurnal course of plant physiology and monoterpene emission

Figure 2 shows the evolution of micrometeorological and physiological parameters that were measured during the enclosure of *Fagus sylvatica* L. in June 2002 and July/August 2003. Regarding the course of net CO<sub>2</sub> assimilation, transpiration, and stomatal conductance, all parameters exhibited pronounced diurnal characteristics following photosynthetic active radiation (PAR) and leaf temperature. Irradiation and leaf temperatures increased particularly during the course of the measurement period in June 2002 as a result of a short high ambient temperature period. Only a few days were cloudy during both campaigns and saturation of photosynthesis was reached for most of the days. Leaf temperature reached maximum readings of 44°C during the experiments in June 2002 and July/August 2003 and was several times above the temperature optimum of net CO<sub>2</sub> assimilation (see Fig. 3). In close relation to the course of PAR and leaf temperature, monoterpene emission from European beech exhibited pronounced diurnal characteristics during both years. As shown in Fig. 2 monoterpene exchange was measured typically in 1–2 h intervals during three and six days in June 2002 and July/August 2003, respectively. Daytime monoterpene emission for the sum of 10 individual monoterpene compounds, reached maximum exchange rates of up to 33.2 μg g<sup>-1</sup> h<sup>-1</sup> in June 2002 and 9.6 μg g<sup>-1</sup> h<sup>-1</sup> in July/August 2003. Night time monoterpene emission was always close to the detection limit (emission ≤80 ng g<sup>-1</sup> h<sup>-1</sup> for both experiments).

Title Page	Abstract	Introduction
Abstract	Conclusions	References
Conclusions	Tables	Figures
Tables	◀	▶
Figures	◀	▶
◀	Back	Close
▶	Full Screen / Esc	
◀	Print Version	
▶	Interactive Discussion	

### 3.2. Monoterpene emission as a function of light and temperature

According to the diurnal characteristics of monoterpene emission from European beech a correlation of monoterpene exchange to micrometeorological parameters is evident. Night time exchange rates near the detection limit eliminate an exclusive role of leaf temperature as reported for monoterpene emission from coniferous trees. In analogy to the monoterpene emission pattern that was observed for other tree species of the plant family *Fagaceae* (e.g. BEMA-Project, 1997; Bertin and Staudt, 1996; Ciccioni et al., 1997; Loreto et al., 1996; Niinemets et al., 2002a; Owen et al., 2002; Staudt and Seufert, 1995; Staudt and Bertin, 1998), and the laboratory experiments conducted on European beech by Schuh et al. (1997), net CO<sub>2</sub> assimilation and monoterpene emission from *Fagus sylvatica* L. were correlated to PAR and leaf temperature for both years (Fig. 3). While in 2002 the small dataset was limiting, in 2003 both net CO<sub>2</sub> assimilation and monoterpene emission revealed similar saturation effects when correlated to light intensity (saturation trend at light intensities of more than 400  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ). A temperature optimum of net CO<sub>2</sub> assimilation was found between leaf temperatures of 25 to 27°C. Monoterpene emission did not exhibit a temperature optimum and increased exponentially with leaf temperature up to a maximum of 43°C during both campaigns. According to the laboratory experiments conducted by Fischbach et al. (2000, 2002) temperature optima for monoterpene synthesis from *Quercus ilex* L. ranged between 30 to 40°C (in vitro) and enzyme activity was measurable up to 60°C. Staudt and Bertin (1998) reported in vivo optima at 42°C for a variety of monoterpene compounds that were emitted by *Quercus ilex* L. Similar optima (~40°C) were obtained by Niinemets et al. (2002b) with *Quercus ilex* L. and *Quercus coccifera* L. (in vivo), who demonstrated that the shape of in vitro and in vivo temperature dependencies differed. They concluded that monoterpene synthase activity was influenced by the chloroplastic (stromal) pH. As a decrease in photosynthetic activity at temperatures above the optimum of photosynthesis lead to acidification of the stromal pH, a decrease in photosynthesis should favour monoterpene emission since pH

## BGD

2, 137–182, 2005

### Monoterpene emission from European beech

T. Dindorf et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

EGU

**Monoterpene  
emission from  
European beech**

T. Dindorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

|◀

▶|

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

optima of monoterpene synthase were slightly shifted to the acidic range (Bohlmann et al., 1998; Fischbach et al., 2000; Niinemets et al., 2002b). Accordingly, the results obtained in the present study indicate that the temperature optimum for monoterpene synthase from European beech is not below temperatures of 43°C in vivo. Moreover, several authors discussed the relevance of unspecific storage pools for the emission of monoterpenes from *Quercus sp.* (Ciccioni et al., 1997; Delfine et al., 2000; Loreto et al., 1996, 2000; Niinemets et al., 2002b, 2004; Niinemets and Reichstein, 2002). We cannot exclude a relevance of such storage pools for the emission of monoterpenes from *Fagus sylvatica* L., particularly since Schuh et al. (1997) reported significant night time emission of  $\alpha$ -pinene at emission rates of  $24.5 \mu\text{g m}^{-2} \text{h}^{-1}$  ( $\Phi^P$  at 25°C). However, comparable night time emissions (or emissions in the absence of light) should have been detectable in the present study but were not observed. As the results show a clear correlation of monoterpene emission to light and temperature, we applied two algorithms to describe the emission behaviour. One algorithm for isoprene, developed by Guenther et al. (1993, 1995, 1997) and a second one for monoterpene emission described by Schuh et al. (1997) (referred to in the following as G97 and S97, see Appendix A for equations). While both algorithms assume a saturation effect for VOC emission at high light intensity, there are two major differences

1. in correlation to irradiance G97 assumes a hyperbolic increase at low light intensities, while the S97 algorithm assumes an allosteric enzyme regulation resulting in a sigmoidal increase,
2. G97 solely assumes enzymatic processes leading to a temperature optimum of monoterpene emission at 39°C. S97 assumes an additional release of monoterpenes from storage pools.

Scattering in the present dataset (including morning and afternoon exchange rates) does not allow clear conclusions as to whether the increase of monoterpene emission with light intensity was hyperbolic or sigmoidal. However, emission in the absence of light was near detection limit. Thus, the storage pool term of the S97 algorithm was

Title Page	Introduction
Abstract	References
Conclusions	Figures
Tables	
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

neglected as recommended for sabinene emission by Schuh et al. (1997), extracting the term from biosynthesis directly prior to emission ( $\Phi^B$ ) only. Figure 4 shows the linear correlation of monoterpene emission to the product of light and temperature terms from the G97 and S97 models. Outliers from this correlation were often caused by temperature effects (e.g. midday depression (see Sect. 3.5.)). Standard emission factors ( $SEF$ , G97) that were calculated from the linear correlation for the sum of monoterpenes ranged between  $13.1 \mu\text{g g}^{-1} \text{h}^{-1}$  for the experiments performed in June 2002 and  $4.4 \mu\text{g g}^{-1} \text{h}^{-1}$  for the experiments performed in July/August 2003. Application of the S97 model resulted in emission factors ( $\Phi^{B,S}$ ) of  $14.1 \mu\text{g g}^{-1} \text{h}^{-1}$  and  $4.6 \mu\text{g g}^{-1} \text{h}^{-1}$  for the campaigns of 2002 and 2003, respectively. Although standard emission differed by a factor of 3 between both years, differences observed for individual days during one growing season were smaller. Table 2 gives an overview of minimum, maximum, and average standard emission factors that were calculated for single days of the respective year by application of the G97 algorithm. Consistent with the results obtained by the enclosure measurements, ambient monoterpene concentrations (measured at the canopy top) ranged up to 1.8 ppb in June 2002 and up to 1.1 ppb in July/August 2003. This result is indicative of the strong influence of beech trees on atmospheric gases in the vicinity of the tower site.

### 3.3. Artificial shading experiment

A variety of physiological functions is influenced by diurnal rhythms of the plant. Whereas endogenous (circadian) rhythms are characterised mainly by the fact that they obtain periodicity even if environmental factors remain constant, exogenous rhythms are triggered by ambient conditions. Monoterpene emission for European beech was shown to have a diurnal periodicity that correlates to light and temperature (see context above and Schuh et al., 1997). However, since light intensity and temperature exhibited a coupled diurnal periodicity by themselves, a pseudo correlation of monoterpene emission to both factors might be possible. To exclude this and to prove the light dependency of monoterpene emission under a daytime temperature regime, an artificial

shading experiment was conducted in 2003 (see Fig. 5). The artificial darkening of the enclosure started at noon and was completed within 30 min (remaining light intensity  $20\text{--}21\ \mu\text{mol m}^{-2}\ \text{s}^{-1}$ ). Two hours later the artificial plant cover was removed again and irradiation progressed in a conventional daily pattern. When the plant cuvette was coated by a dark cover, cuvette temperature (and as a consequence leaf temperature) decreased after darkening but increased again in the course of the shading period (max. difference in leaf temperature during shading  $13^\circ\text{C}$ ). VOC exchange measured 30 min after complete coverage of the cuvette showed no emission of monoterpene compounds. As soon as the artificial darkening was removed, monoterpene emission progressed with its conventional diurnal characteristics. Assuming the existence of unspecific storage pools for monoterpenes in European beech as reported by Schuh et al. (1997), storage pools must have been emptied after darkening in a time period of 30 min. This is not consistent with the analysis performed by Niinemets and Reichstein (2002) which proved that the decrease in monoterpene emission from storage pools follows a double exponential function. Other authors have also reported a persistence of monoterpene emission from unspecific storage pools of *Quercus sp.*, for several hours to days (Ciccioli et al., 1997; Loreto et al., 1996, 2000; Niinemets et al., 2002a). However, if storage pools would have existed for European beech and were not depleted by the time the cover was removed, a typical burst of monoterpene emission as reported earlier for *Quercus ilex L.* (Loreto et al., 2000) should have been observed for European beech as well (which was not the case). Furthermore, a rapid depletion of potential storage pools would explain the lack of night time emission in the present study, since time resolution of monoterpene measurement was typically 1–2 h. However, the darkening experiment clearly showed the obligatory role of light for monoterpene emission from beech and that emission from storage pools may also be neglected under a high temperature regime.

**Monoterpene  
emission from  
European beech**

T. Dindorf et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

### 3.4. Composition of emitted monoterpene species

Figure 6 shows the diurnal course of individual monoterpene emission measured by GC-MS for one of the measurement days in August 2003. Light and temperature dependency of monoterpene emission was observed for all analysed compounds with exception of tricyclene that scattered at exchange rates below  $1 \text{ ng g}^{-1} \text{ h}^{-1}$ . Sabinene was the predominant monoterpene compound emitted (max.  $8.1 \mu\text{g g}^{-1} \text{ h}^{-1}$ ). Standard emission factors that were calculated for the sum of all monoterpenes for this respective day reached  $7.1 \mu\text{g g}^{-1} \text{ h}^{-1}$  (G97, sum of 15 individual monoterpene compounds) and  $5.9 \mu\text{g g}^{-1} \text{ h}^{-1}$  (G97, sum of 10 individual monoterpene compounds, see GC-FID). Monoterpene emission calculated by the G97 algorithm fitted well to midday and afternoon values but morning and evening monoterpene emission was much lower than calculated. Thus integration of measured and calculated emission for the respective day resulted in an overestimation of +13% by the calculated data. Application of a sigmoidal increase of monoterpene emission with light intensity (as assumed by the S97 algorithm,  $\Phi^B$  only) resulted in a better reproducibility of the observed data in the morning hours. However, both algorithms still overestimated monoterpene emission in the morning and underestimated midday and afternoon emission to a various extent. In order to determine the contribution of different monoterpene species to the sum of monoterpene emission, the fraction of standard emission factors for different monoterpene compounds were calculated as shown in Table 3. Correlation of different monoterpene species to each other resulted in good results for the majority of compounds with the exception of tricyclene. Our results are consistent with most previous studies which report sabinene as being the predominant monoterpene compound released from European beech (see Kahl et al., 1999; Schuh et al., 1997; Tollsten and Müller, 1996). Moreover, reasonable agreement was obtained for the emission of myrcene and limonene. In contrast to these results, the apportionment of other monoterpene compounds was much more inconsistent between the different studies mentioned above. This particularly affects the emission of  $\alpha$ -pinene, that was reported

## BGD

2, 137–182, 2005

### Monoterpene emission from European beech

T. Dindorf et al.

Title Page	Introduction
Abstract	References
Conclusions	Figures
Tables	
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

EGU

to be released to a major (Kahl et al., 1999; Schuh et al., 1997), moderate (König et al., 1995), or minor (this study and Tollsten and Müller, 1996) extent by European beech. Similar contradictory results were obtained for the emission of  $\beta$ -pinene that was reported to be released as a major (König et al., 1995) or minor (this study and Tollsten and Müller, 1996) compound. As discussed by König et al. (1995) temperature effects can be an important factor for the apportionment of monoterpene emission from European beech, resulting in the dominance of different compounds at different enclosure temperatures. Hence, inconsistencies between the different studies mentioned above might not be surprising, since most of them were conducted at different temperatures.

5

### 10 3.5. Midday depression of net CO<sub>2</sub> assimilation and monoterpene emission

The clear light dependence of monoterpene release reflects the close link of emission and production, which is closely coupled to net CO<sub>2</sub> assimilation. A plants primary productivity depends on several factors, with water availability being of essential importance. Increasing leaf temperature and water vapour pressure deficit have a severe effect on the water loss of the leaf epidermis which is an important factor for stomatal aperture (for a detailed overview see Schulze, 1986). According to Backes and Leuschner (2000) *Fagus sylvatica* L. evolved a very sensitive stomatal regulation that allows maintenance of leaf turgor during dry, high temperature conditions. Consequently, on days with high ambient temperature European beech, due to its sensitive stomatal regulation, exhibited a distinct midday depression of net CO<sub>2</sub> assimilation as observed two times in June 2002 and five times in July/August 2003. Figure 7 gives an overview of the daily course of leaf temperature, as well as the correlation of stomatal conductance and photosynthesis to PAR during the measurement period in June 2002. As shown, European beech developed a midday depression of photosynthesis when midday leaf temperatures increased to more than 41°C during the 6<sup>th</sup> and 7<sup>th</sup> day of measurement (Figs. 7f and 7g). As soon as midday leaf temperatures decreased and were below 28°C the next day, midday depression was not observed and plant stomatal conductance and net CO<sub>2</sub> assimilation progressed in a conventional pattern. As

20

150

**BGD**

2, 137–182, 2005

## Monoterpene emission from European beech

T. Dindorf et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

EGU

discussed above, monoterpene emission exhibited light saturation. However, during sunny days the correlation revealed a typical hysteretic structure that was observed to be a function of time (lower monoterpene emission in the morning, see Fig. 8, top graph). The hysteresis was attributed to a time lag phase of monoterpene emission in the early morning. Since the morning increase of monoterpene emission with PAR was less pronounced than the decrease in the early evening, the correlation of monoterpene emission to PAR resulted in a hysteresis. In contrast, during hot sunny days, midday depression of net CO<sub>2</sub> assimilation and stomatal conductance may lead to a hysteretic course of monoterpene emission that follows the time of day in an opposite direction (lower monoterpene emission in the afternoon). Indeed, this effect was observed once, during the measurement period in June 2002 as monoterpene emission was very high and the plant experienced its first high temperature period of the respective growing season (see Fig. 8, bottom graph). As reported by Loreto et al. (1996) and discussed in detail by Niinemets and Reichstein (2003) monoterpenes are emitted through stomata but are not controlled by stomatal aperture. Thus, other physiological processes such as photosynthesis must have limited monoterpene emission during the respective day. As discussed above, a decrease in photosynthesis and consequently in stromal pH, should favour monoterpene emission (Bohlmann et al., 1998; Fischbach et al., 2000; Niinemets et al., 2002b). However, midday depression of monoterpene emission from *Quercus ilex L.* has been observed by other studies as well (Bertin et al., 1997; Kesselmeier et al., 1997; Penuelas and Llusia, 1999). The DOXP Pathway with its substrates glycerine-aldehyde-3-phosphate and pyruvate is known to represent the major source for plasticidic isoprenoids (Lichtenthaler, 1999). As reported by Parry et al. (2002), drought effects may lead to a down regulation of ribulose-1,5-bisphosphate-carboxylase (RUBISCO) by tight binding inhibitors, a mechanism that was discussed recently as a central effect of midday depression of photosynthesis (Griffith and Parry, 2002). Thus, a restriction of RUBISCO might lead to a substrate limitation under high monoterpene production rates. Experiments performed by Kahl et al. (1999) on European beech indicated that 90% of the total sabinene emission was formed de novo from

**Monoterpene emission from European beech**

T. Dindorf et al.

Title Page	Abstract	Conclusions	Tables	1 ◀	▶ 1
Introduction	References	Figures		◀	▶
Back	Close				
Full Screen / Esc					
	Print Version				
	Interactive Discussion				

photosynthetic intermediates. Since sabinene was the predominant compound emitted by *Fagus sylvatica* L. (fraction of 57%) a decrease of photosynthetic activity might also lead to a decrease in the emission of monoterpenes. During the midday depression of monoterpene emission the carbon balance (carbon loss in form of monoterpenes per photosynthetic fixed carbon) exceeded the 1% level (up to 1.7%), while under normal environmental conditions the carbon loss was typically below 0.5%. Since midday depression of monoterpene synthesis was observed only during high monoterpene production rates in 2002, we conclude that monoterpene emission was limited by substrate availability in this special case.

### 10 3.6. Variability of standard emission factors

Standard emission factors that were calculated by application of the G97 algorithm on a dry weight basis differed by a factor of 3 between both years. However, the relatively low standard emission factors calculated for 2003 were confirmed by canopy scale flux measurements performed by Spirig et al. (2004). The importance of variations in developmental stages, seasonality, growth conditions, and habitat for monoterpene emission from *Quercus ilex* L. has been reported recently (e.g. Bertin et al., 1997; Fischbach et al., 2002; Llusia and Peñuelas, 2000; Niinemets et al., 2002a; Peñuelas and Llusia, 1999; Sabillon and Cremades, 2001; Staudt et al., 2002, 2003; Street et al., 1997). Also, European beech is known to develop ecotypes that are adapted to the climatic conditions of the habitat they live in (Peuke et al., 2002). The location selected for our measurements of European beech is characterised by temperate climatic conditions. However, temperatures can display strong variations as observed during the measurement period of 2002 and 2003. Figure 9 gives an overview of ambient temperature and rainfall that were measured prior and during the enclosure of European beech in the summers of 2002 and 2003. Average ambient temperatures measured 30 days preceding the enclosure measurements ranged from moderate temperatures of 16°C in Mai/June 2002 to higher temperatures of 20°C in June/July 2003. As observed by Staudt et al. (2003), the acclimatisation time of standard emission factors to

## BGD

2, 137–182, 2005

### Monoterpene emission from European beech

T. Dindorf et al.

Title Page	Abstract	Introduction
Conclusions	Tables	References
Figures	◀	▶
◀	◀	▶
Back	Close	
Full Screen / Esc		
Print Version		
Interactive Discussion		

EGU

5 previous temperatures was highly variable and ranged between 3 days and 3 weeks for *Quercus ilex* L.. These experiments are not in agreement with our results from European beech, as higher daytime temperatures were observed in the days preceding the measurements in 2003, when exchange rates were much smaller. Thus, other effects might have dominated the observed variability in standard emission between the experiments in 2002 and 2003. One of these effects might have been a long lasting drought period that preceded the measurements in 2003 and resulted in a reduction of average photosynthesis during the latter campaign. Since a reduction of photosynthesis and transpiration have been reported to be indicators of drought to European beech (Peuke et al., 2002; Thomas, 2000), we can not exclude long term effects of drought in 2003. However, transpiration did not decrease in the latter campaign. Bertin and Staudt (1996) demonstrated that long periods of drought restricted monoterpene emissions from *Quercus ilex* L. by two orders of magnitude as the daily net carbon balance approached zero. Likewise, Staudt et al. (2002) reported a reduction of photosynthesis and transpiration, paralleled by a reduction of monoterpene emission of at least 25% due to drought effects. As reported by Backes and Leuschner (2000) and Thomas (2000), physiological properties of European beech are much more sensitive to drought than physiological properties of oak trees. Hence, this sensibility provides an explanation for the stronger reduction of monoterpene emission observed during the present study. Moreover, comparing the intensity of photosynthetic active radiation during clear sky days in 2002 and 2003, light intensity was lessened by some percent during the latter campaign. This effect might have been caused by an increase of biomass that surrounded the investigated branch during the vegetation period. However, light intensity was still high and microscopic analysis of leaf morphology pointed to the typical structure of sunlit leaves. Although the lower light intensity might explain the decrease in LMA values in 2003 and might have contributed to the lower standard emission factors investigated. Furthermore, as measurements started in June during the first campaign but were conducted in July/August in 2003, seasonal effects on monoterpene emission have to be taken into account as well. According to Schuh

**Monoterpene emission from European beech**

T. Dindorf et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

Title Page	Introduction
Abstract	References
Conclusions	Figures
Tables	
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

et al. (1997), monoterpene emission from *Fagus sylvatica* L. decreased by a factor of 16 between spring and autumn. Also König et al. (1995), who investigated beech trees in Austria in late August and early September, reported a decrease of monoterpene emission. For a detailed overview of the seasonal development of monoterpene emission from European beech see Holzke et al. (2005)<sup>3</sup>, who monitored monoterpene emission from *Fagus sylvatica* L. for a time period of 2 years. In contrast to most literature data, the present study revealed *Fagus sylvatica* L. as being a strong monoterpene emitter. For the plant family *Fagaceae*, only a few species that have been investigated so far have exhibited substantial monoterpene emission rates (for review see Kesselmeier and Staudt, 1999). Table 4 gives an overview of experiments that were conducted earlier to examine monoterpene emission from European beech. The majority of these experiments reported that *Fagus sylvatica* L. emitted only low amounts of monoterpenes. Regarding the laboratory experiments performed by Hewitt and Street (1992) and Steinbrecher et al. (1993), monoterpene emission from European beech was below or near the detection limit of the analytical system. Also König et al. (1995) and Tollsten and Müller (1996) who examined European beech trees under field conditions in Austria and Switzerland found only low emission of monoterpenes at 0.2 and 0.3  $\mu\text{g g}^{-1} \text{h}^{-1}$ , respectively. Only laboratory experiments conducted by Schuh et al. (1997) and Kahl et al. (1999) revealed substantial monoterpene emission at 414 and 284  $\mu\text{g m}^{-2} \text{h}^{-1}$  (at 25°C). Monoterpene emissions that were measured during the first year of the present study significantly exceeded these emission rates reported.

### 3.7. Implications for the European budget of monoterpene emission

According to Guenther et al. (1995, 1997), there are two approaches to assign emission factors at an ecosystem scale.

<sup>3</sup>Holzke, C., Dindorf, T., Kuhn, U., Kesselmeier, J., and Koppmann, R.: Terpene emissions from European beech (*Fagus sylvatica* L.): pattern and emission behaviour within two vegetation periods, in preparation, 2005.

1. The first method assigns a landscape type to each location within the model domain. An emission potential, derived by micrometeorological measurement techniques or from general assumptions of species distribution, is associated with each landscape type.

2. The second approach requires an estimate of the composition of plant species for each location in the model domain, as well as a database of specific emission potentials that are derived e.g. by enclosure measurements for each plant species. A landscape average emission potential can then be assigned as the weighted average of all species at each location.

In the global model of Guenther et al. (1995) distinct emission factors have been assigned to various ecosystem types following the first approach described above. Figure 10 shows the mean European monoterpene emission flux (domain 10° W–30° E and 35° N–65° N) for the month of July that was calculated with the default monoterpene emission factors assigned to the Olson ecosystems as described by Guenther et al. (1995, henceforth G95OIs) for a 0.5x0.5 grid resolution. Figure 11 shows the relative increase of monthly mean monoterpene emission in relation to the latter assumption, if the spatial distribution of *Fagus sylvatica* L. is specifically considered (henceforth G95FS). The G95FS flux is calculated from the flux based on the measured monoterpene emission factor ( $13 \mu\text{g g}^{-1} \text{h}^{-1}$ , year 2002) and the flux using the default emission factor of the G95OIs algorithm. In this way weighted average fluxes were calculated, specifically taking into consideration the fraction of *Fagus sylvatica* L. area coverage in every 0.5 x 0.5 grid. The relative difference to the default G95OIs model (see Fig. 11) is calculated according to Eq. (1).

$$100 \times \frac{\text{G95FS} - \text{G95OIs}}{(\text{G95FS} + \text{G95OIs})/2} \quad (1)$$

The spatial distribution of the differences actually reflects the European distribution of *Fagus sylvatica* L., which makes up about 7.1% of the European forest area. Despite

**Monoterpene emission from European beech**

T. Dindorf et al.

Title Page	Abstract	Conclusions	Tables	1 ◀	▶ 1
Introduction	References	Figures		◀	▶
Back	Full Screen / Esc	Print Version	Interactive Discussion		

**Monoterpene  
emission from  
European beech**

T. Dindorf et al.

Title Page	Introduction	References
Abstract	Conclusions	Figures
Tables	◀	▶
◀	◀	▶
Back	Close	
Full Screen / Esc		
	Print Version	
	Interactive Discussion	

the relatively small area covered by European beech there is a significant increase in the European monoterpene emission in July from about 694 to 1017 tC month<sup>-1</sup> if the high standard emission factor and the spatial distribution of *Fagus sylvatica* L. are taken into account. However, in both G95OIs and G95FS monoterpene emission was calculated as a function of temperature only. By also considering the role of light as a controlling parameter of monoterpene emission from *Fagus sylvatica* L. (henceforth FSlight), the total amount of monoterpene release is reduced significantly. In the latter assumption the total emission amounts to 792 tC month<sup>-1</sup>. Nevertheless, as shown by Fig. 12, relative increases > 100% between the FSlight and the G95OIs were estimated on a local scale, when relative changes are calculated in analogy to Eq. (1). Note that the FSlight results reflect a simulation where only the light and temperature attenuation functions for *Fagus sylvatica* L. are applied, whereas the monoterpene emissions for the remaining fraction in each 0.5 x 0.5 grid are calculated considering the role of temperature only. However, if generalized for deciduous tree species, a strong light dependence of monoterpene emission would have a strong impact on predicted magnitude and temporal distribution of monoterpenes. Consistent with previous studies that were based on a similar methodology (e.g. Guenther et al., 1997; Lenz et al., 2001; Solmon et al., 2004) it can be concluded that consideration of a more detailed spatial distribution of a specific land cover type, for which measured emission parameters are available, results in significant changes in the local monoterpene emission flux. However, for the European domain the increase is small considering all uncertainties involved such as biomass estimates and using the surface- versus the actual canopy or leaf temperature. In addition, the results reflect the simulations for the month of July with high radiation intensity and temperature using the high emission flux measured in 2002. Consequently, the results reflect an upper range impact of the observed *Fagus sylvatica* L. emission rate and light dependence for the European domain.

#### 4. Conclusions

The experiments performed in the present study clearly indicate that monoterpene emission from European beech is a function of light and temperature. Even though the temperature optimum was not reached up to leaf temperatures of 43°C, monoterpene release that was calculated by application of light and temperature dependent algorithms (G97 and S97), generated reasonable agreement with the measured data. In reference to the default G95OIs model, the consideration of the substantial emission factor and light intensity as a controlling parameter of monoterpene release, generated a relative increase in the European monoterpene emission up to 14%, yielding monoterpene emissions of 792 tC month<sup>-1</sup>. However, long term drought effects might restrict monoterpene emission and might explain the observed interannual and seasonal variability of monoterpene emission. Furthermore, short term drought effects, like midday depression of photosynthesis, can be crucial for daily variations in VOC emission, particularly when rates of monoterpene productivity are high.

#### Appendix A

Algorithms describing VOC emission as a function of light and temperature.

A.1. Algorithm developed by Guenther et al. (1993, 1995, 1997)

$$\Phi = SEF \cdot \left[ \frac{\alpha \cdot C_L \cdot L}{\sqrt{1 + \alpha^2 \cdot L^2}} \right] \cdot \left[ \frac{\exp\left(\frac{C_{T1} \cdot (T - T_s)}{R \cdot T_s \cdot T}\right)}{C_{T3} + \exp\left(\frac{C_{T2} \cdot (T - T_M)}{R \cdot T_s \cdot T}\right)} \right] \quad (A1)$$

## BGD

2, 137–182, 2005

### Monoterpene emission from European beech

T. Dindorf et al.

Title Page	►
Abstract	►
Conclusions	►
Tables	►
Introduction	►
References	►
Figures	►
Back	Close
Full Screen / Esc	Print Version
Interactive Discussion	

EGU

A.2. Algorithm developed by Schuh et al. (1997)

$$\Phi = \Phi^{P,S} \cdot \exp \left[ \frac{C_{TP}}{R} \cdot \left( \frac{T - T_S}{T \cdot T_S} \right) \right] + \Phi^{B,S} \cdot C_L \cdot \left[ \frac{\alpha \cdot L}{\sqrt{1 + \alpha^2 \cdot L^2}} \right]^2 \cdot \left[ \frac{\exp \left( \frac{C_{T1} \cdot (T - T_S)}{R \cdot T \cdot T_S} \right)}{1 + \exp \left( \frac{C_{T2} \cdot (T - T_M)}{R \cdot T \cdot T_S} \right)} \right] \quad (A2)$$

A.3. Algorithm developed by Schuh et al. (1997), biosynthesis part, denominator modified according to Guenther et al. (1997)

$$\Phi^B = \Phi^{B,S} \cdot C_L \cdot \left[ \frac{\alpha \cdot L}{\sqrt{1 + \alpha^2 \cdot L^2}} \right]^2 \cdot \left[ \frac{\exp \left( \frac{C_{T1} \cdot (T - T_S)}{R \cdot T \cdot T_S} \right)}{C_{T3} + \exp \left( \frac{C_{T2} \cdot (T - T_M)}{R \cdot T \cdot T_S} \right)} \right] \quad (A3)$$

$\alpha$  (empirical coefficient)=0.0027

$C_L$  (empirical coefficient)=1.066

$C_{T1}$  (empirical coefficient [J mol<sup>-1</sup>])=95 000

$C_{T2}$  (empirical coefficient [J mol<sup>-1</sup>])=230 000

$C_{T3}$  (empirical coefficient)=0.961

$C_{TP}$  (empirical coefficient, e.g. 0 for sabinene)

$L$  (photosynthetic active radiation (PAR) [ $\mu\text{mol m}^{-2} \text{s}^{-1}$ ])

$\Phi$  (actual VOC emission [e.g.  $\mu\text{g g}^{-1} \text{h}^{-1}$ ])

$\Phi^{B,S}$  (flow from biosynthesis at standard condition [e.g.  $\mu\text{g g}^{-1} \text{h}^{-1}$ ])

$\Phi^{P,S}$  (flow from pool at standard condition [e.g.  $\mu\text{g g}^{-1} \text{h}^{-1}$ ])

$R$  (universal gas constant [J K<sup>-1</sup> mol<sup>-1</sup>])=8.314

$SEF$  (standard emission factor [e.g.  $\mu\text{g g}^{-1} \text{h}^{-1}$ ])

$T$  (leaf temperature [°K])

$T_M$  (empirical coefficient [°K])=314

$T_S$  (leaf temperature at standard condition [°K])=303

158

**BGD**

2, 137–182, 2005

## Monoterpene emission from European beech

T. Dindorf et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

EGU

*Acknowledgements.* We would like to thank the staff of the Research Centre Jülich for providing excellent facilities. Especially we appreciate the help of A. Schaub, M. Komenda, and R. Koppmann for the coordination and preparation of both field experiments. We would like to thank A. Knapps for the supply of meteorological data recorded prior our measurement campaigns. F. Steindel, N. Knothe, G. Chaparro, C. Tritsch, and M. Scheibe were of great support in setting up and maintaining the equipment on the measurement site. The valuable comments on manuscript preparation by T. W. Andreae were highly appreciated. Finally we would like to thank C. Ammann for his advice as well as E. Brancaleoni and M. Frattoni for their support in analysing the GC-MS samples. This work was supported by the Max Planck Society and the German Atmospheric Research Program 2000, project ECHO (Emission and Chemical transformation of biogenic volatile Organic compounds, FZK:07ATF47).

## References

- Andreae, M. O. and Crutzen, P. J.: Atmospheric aerosols: Biogeochemical sources and role in atmospheric chemistry, *Science*, 276, 1052–1058, 1997.
- Atkinson, R.: Atmospheric chemistry of VOCs and NO<sub>x</sub>, *Atmos. Environ.*, 34, 2063–2101, 2000.
- Backes, K. and Leuschner, C.: Leaf water relations of competitive *Fagus sylvatica* and *Quercus petraea* trees during 4 years differing in soil drought, *Canadian Journal of Forest Research- Revue Canadienne De Recherche Forestiere*, 30, 335–346, 2000.
- BEMA-Project: Special Issue, *Atmos. Environ.*, 31 (SI), 1997.
- Bertin, N. and Staudt, M.: Effect of water stress on monoterpene emissions from young potted holm oak (*Quercus ilex* L.) trees, *Oecologia*, 107, 456–462, 1996.
- Bertin, N., Staudt, M., Hansen, U., Seufert, G., Ciccioli, P., Foster, P., Fugit, J. L., and Torres, L.: Diurnal and seasonal course of monoterpene emissions from *Quercus ilex* (L.) under natural conditions – Applications of light and temperature algorithms, *Atmos. Environ.*, 31, 135–144, 1997.
- Bohmann, J., Meyer-Gauen, G., and Croteau, R.: Plant terpenoid syntheses: Molecular biology and phylogenetic analysis, *Proceedings of the National Academy of Sciences of the United States of America*, 95, 4126–4133, 1998.
- Brancaleoni, E., Scovaventi, M., Frattoni, M., Mabilia, R., and Ciccioli, P.: Novel family of multi-layer cartridges filled with a new carbon adsorbent for the quantitative determination

## BGD

2, 137–182, 2005

### Monoterpene emission from European beech

T. Dindorf et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

EGU

of volatile organic compounds in the atmosphere, *Journal of Chromatography A*, 845, 317–328, 1999.

Ciccioli, P., Cecinato, A., Brancaleoni, E., Frattoni, M., and Liberti, A.: Use of Carbon Adsorption Traps Combined with High-Resolution Gas-Chromatography-Mass-Spectrometry for the Analysis of Polar and Nonpolar C-4-C-14 Hydrocarbons Involved in Photochemical Smog Formation, *Hrc-Journal of High Resolution Chromatography*, 15, 75–84, 1992.

Ciccioli, P., Fabozzi, C., Brancaleoni, E., Cecinato, A., Frattoni, M., Loreto, F., Kesselmeier, J., Schäfer, L., Bode, K., Torres, L., and Fugit, J. L.: Use of the isoprene algorithm for predicting the monoterpene emission from the Mediterranean holm oak *Quercus ilex* L.: Performance and limits of this approach, *J. Geophys. Res.-A*, 102, 23 319–23 328, 1997.

Delfine, S., Csiky, O., Seufert, G., and Loreto, F.: Fumigation with exogenous monoterpenes of a non-isoprenoid-emitting oak (*Quercus suber*): monoterpene acquisition, translocation, and effect on the photosynthetic properties at high temperatures, *New Phytologist*, 146, 27–36, 2000.

Fehsenfeld, F., Calvert, J., Fall, R., Goldan, P., Guenther, A., Hewitt, C. N., Lamb, B., Liu, S., Trainer, M., Westberg, H., and Zimmermann, P.: Emissions of volatile organic compounds from vegetation and the implications for atmospheric chemistry, *Global Biogeochem. Cycles*, 6, 389–430, 1992.

Fischbach, R. J., Zimmer, I., Steinbrecher, R., Pfichner, A., and Schnitzler, J. P.: Monoterpene synthase activities in leaves of *Picea abies* (L.) Karst. and *Quercus ilex* L., *Phytochemistry*, 54, 257–265, 2000.

Fischbach, R. J., Staudt, M., Zimmer, I., Rambal, S., and Schnitzler, J. P.: Seasonal pattern of monoterpene synthase activities in leaves of the evergreen tree *Quercus ilex*, *Physiologia Plantarum*, 114, 354–360, 2002.

Ganzeveld, L. N., Lelieveld, J., Dentener, F. J., Krol, M. C., Bouwman, A. J., and Roelofs, G. J.: Global soil-biogenic NO<sub>x</sub> emissions and the role of canopy processes, *J. Geophys. Res.-A*, 107, doi:10.1029/2001JD001289, 2002.

Griffith, H. and Parry, M. A. J.: Preface: Plant responses to water stress, *Annals of Botany*, 89, 801–802, 2002.

Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W.A., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.: A Global-Model of Natural Volatile Organic-Compound Emissions, *J. Geophys. Res.-A*, 100, 8873–8892, 1995.

**BGD**

2, 137–182, 2005

## Monoterpene emission from European beech

T. Dindorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

- Guenther, A.: Seasonal and spatial variations in natural volatile organic compound emissions, *Ecological Applications*, 7, 34–45, 1997.
- Guenther, A.: The contribution of reactive carbon emissions from vegetation to the carbon balance of terrestrial ecosystems, *Chemosphere*, 49, 837–844, 2002.
- Guenther, A. B., Zimmerman, P. R., Harley, P. C., Monson, R. K., and Fall, R.: Isoprene and Monoterpene Emission Rate Variability – Model Evaluations and Sensitivity Analyses, *J. Geophys. Res.-A*, 98, 12609–12617, 1993.
- Gut, A., Scheibe, M., Rottenberger, S., Rummel, U., Welling, M., Ammann, C., Kirkman, G. A., Kuhn, U., Meixner, F.X., Kesselmeier, J., Lehmann, B. E., Schmidt, W., Müller, E., and Piedade, M.T.F.: Exchange fluxes of NO<sub>2</sub> and O<sub>3</sub> at soil and leaf surfaces in an Amazonian rain forest, *J. Geophys. Res.-A*, 107, doi:10.1029/2001JD000654, 2002.
- Gutman, G., Tarpley, D., Ignatov, A., and Olson, S.: The Enhanced Noaa Global Land Dataset from the Advanced Very High-Resolution Radiometer, *Bulletin of the American Meteorological Society*, 76, 1141–1156, 1995.
- Hewitt, C. N. and Street, R. A.: A Qualitative Assessment of the Emission of Nonmethane Hydrocarbon Compounds from the Biosphere to the Atmosphere in the Uk – Present Knowledge and Uncertainties, *Atmospheric Environment Part a-General Topics*, 26, 3069–3077, 1992.
- Kahl, J., Hoffmann, T., and Klockow, D.: Differentiation between de novo synthesized and constitutively released terpenoids from *Fagus sylvatica*, *Phytochemistry*, 51, 383–388, 1999.
- Karlik, J. F. and Winer, A. M.: Measured isoprene emission rates of plants in California landscapes: comparison to estimates from taxonomic relationships, *Atmos. Environ.*, 35, 1123–1131, 2001.
- Kesselmeier, J., Schäfer, L., Ciccioli, P., Brancaleoni, E., Cecinato, A., Frattoni, M., Foster, P., Jacob, V., Denis, J., Fugit, J. L., Dutaur, L., and Torres, L.: Emission of monoterpenes and isoprene from a Mediterranean oak species *Quercus ilex* L. measured within the BEMA (Biogenic Emissions in the Mediterranean Area) project, *Atmos. Environ.*, 30, 1841–1850, 1996.
- Kesselmeier, J., Bode, K., Hofmann, U., Müller, H., Schäfer, L., Wolf, A., Ciccioli, P., Brancaleoni, E., Cecinato, A., Frattoni, M., Foster, P., Ferrari, C., Jacob, V., Fugit, J.L., Dutaur, L., Simon, V., and Torres, L.: Emission of short chained organic acids, aldehydes and monoterpenes from *Quercus ilex* L. and *Pinus pinea* L. in relation to physiological activities, carbon budget and emission algorithms, *Atmos. Environ.*, 31, 119–133, 1997.

**Monoterpene emission from European beech**

T. Dindorf et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
	Print Version
	Interactive Discussion

Kesselmeier, J. and Staudt, M.: Biogenic volatile organic compounds (VOC): An overview on emission, physiology and ecology, *J. Atmos. Chem.*, **33**, 23–88, 1999.

Kesselmeier, J., Ciccioli, P., Kuhn, U., Stefani, P., Biesenthal, T., Rottenberger, S., Wolf, A., Vitullo, M., Valentini, R., Nobre, A., Kabat, P., and Andreae, M. O.: Volatile organic compound emissions in relation to plant carbon fixation and the terrestrial carbon budget, *Global Biogeochem. Cycles*, **16**, doi:10.1029/2001GB001813, 2002.

Kesselmeier, J., Rottenberger, S., Kuhn, U., and Ciccioli, P.: Ecosystem seasonality affects biogenic emission of volatile organic compounds (VOC): Flooding and dry/wet season transition in tropical regions as an elicitor for emission changes at regular intervals?, in: *Integrated Land Ecosystem-Atmosphere Processes Study (ILEAPS)*, International Open Science Conference, edited by: Arneth, A., Korhonen, H., Kulmala, M., Raivonen, M., Ruuskanen, T., and Suni, T., Finnish Association for Aerosol Research, Helsinki, Finland, 187–192, 2003.

Kesselmeier, J.: Interactive comment on “Seasonal cycles of isoprene concentrations in the Amazonian rainforest” by Trostdorf, C. R. et al., *Atmos. Chem. Phys. Discuss.*, **4**, S528–S530, 2004.

Kleindienst, T. E., Corse, E. W., Blanchard, F. T., and Lonneman, W. A.: Evaluation of the performance of DNPH-coated silica gel and C-18 cartridges in the measurement of formaldehyde in the presence and absence of ozone, *Environ. Sci. Technol.*, **32**, 124–130, 1998.

Köble, R. and Seufert, G.: Novel maps for forest tree species in Europe, in: *A changing atmosphere: 8th European symposium on the physico-chemical behaviour of atmospheric pollutants*, Torino, 2001.

König, G., Brunda, M., Puxbaum, H., Hewitt, C. N., Duckham, S. C., and Rudolph, J.: Relative Contribution of Oxygenated Hydrocarbons to the Total Biogenic Voc Emissions of Selected Mid-European Agricultural and Natural Plant-Species, *Atmos. Environ.*, **29**, 861–874, 1995.

Kuhn, U., Rottenberger, S., Biesenthal, T., Ammann, C., Wolf, A., Schebeske, G., Oliva, S., T., Tavares, T. M., and Kesselmeier, J.: Exchange of short-chain monocarboxylic acids by vegetation at a remote tropical forest site in Amazonia, *J. Geophys. Res.-A*, **107**, doi:10.1029/2000JD000303, 2002a.

Kuhn, U., Rottenberger, S., Biesenthal, T., Wolf, A., Schebeske, G., Ciccioli, P., Brancaleoni, E., Frattoni, M., Tavares, T. M., and Kesselmeier, J.: Isoprene and monoterpene emissions of Amazonian tree species during the wet season: Direct and indirect investigations on controlling environmental functions, *J. Geophys. Res.-A*, **107**, doi:10.1029/2001JD000978, 2002b.

Kuhn, U., Rottenberger, S., Biesenthal, T., Wolf, A., Schebeske, G., Ciccioli, P., Brancaleoni,

**BGD**

2, 137–182, 2005

**Monoterpene emission from European beech**

T. Dindorf et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

EGU

**Monoterpene  
emission from  
European beech**

T. Dindorf et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

EGU

- E., Frattoni, M., Tavares, T. M., and Kesselmeier, J.: Seasonal differences in isoprene and light-dependent monoterpene emission by Amazonian tree species, *Global Change Biology*, 10, 663–682, 2004.
- 5 Lenz, R., Köble, R., and Seufert, G.: Species-based mapping of biogenic emissions in Europe case study Italy, in: *A changing atmosphere: 8th European symposium on the physico-chemical behaviour of atmospheric pollutants*, Torino, 2001.
- Lerdau, M. and Gray, D.: Ecology and evolution of light-dependent and light-independent phyto-  
togenic volatile organic carbon, *New Phytologist*, 157, 199–211, 2003.
- 10 Lichtenthaler, H. K.: The 1-deoxy-D-xylulose-5-phosphate pathway of isoprenoid biosynthesis in plants, *Annual Review of Plant Physiology and Plant Molecular Biology*, 50, 47–65, 1999.
- Llusia, J. and Peñuelas, J.: Seasonal patterns of terpene content and emission from seven Mediterranean woody species in field conditions, *American Journal of Botany*, 87, 133–140, 2000.
- Loreto, F., Ciccioli, P., Cecinato, A., Brancaleoni, E., Frattoni, M., and Tricoli, D.: Influence of environmental factors and air composition on the emission of alpha-pinene from Quercus  
ilex leaves, *Plant Physiology*, 110, 267–275, 1996.
- 15 Loreto, F., Ciccioli, P., Brancaleoni, E., Frattoni, M., and Delfino, S.: Incomplete C-13 labeling of alpha-pinene content of *Quercus ilex* leaves and appearance of unlabelled C in alpha-pinene emission in the dark, *Plant Cell and Environment*, 23, 229–234, 2000.
- 20 Niinemets, U., Hauff, K., Bertin, N., Tenhunen, J. D., Steinbrecher, R., and Seufert, G.: Monoterpene emissions in relation to foliar photosynthetic and structural variables in Mediterranean evergreen Quercus species, *New Phytologist*, 153, 243–256, 2002a.
- Niinemetts, U., Seufert, G., Steinbrecher, R., and Tenhunen, J. D.: A model coupling foliar  
monoterpene emissions to leaf photosynthetic characteristics in Mediterranean evergreen  
Quercus species, *New Phytologist*, 153, 257–275, 2002b.
- 25 Niinemets, U. and Reichstein, M.: A model analysis of the effects of nonspecific monoterpeneoid storage in leaf tissues on emission kinetics and composition in Mediterranean sclerophyllous Quercus species, *Global Biogeochem. Cycles*, 16, doi:10.1029/2002GB001927, 2002.
- Niinemetts, U. and Reichstein, M.: Controls on the emission of plant volatiles through stomata: Differential sensitivity of emission rates to stomatal closure explained, *J. Geophys. Res.-A*,  
108, doi:10.1029/2002JD002626, 2003.
- 30 Niinemets, U., Loreto, F., and Reichstein, M.: Physiological and physicochemical controls on foliar volatile organic compound emissions, *Trends in Plant Science*, 9, 180–186, 2004.

Olson, J.: World ecosystems (WE1.4): Digital raster data on a 10 min geographic 1080'2160 grid square, in: Global Ecosystem Database, Version 1.0: Disc A Ed. NOAA National Geophysical Data Center, Boulder, 1992.

Owen, S. M., Harley, P., Guenther, A., and Hewitt, C. N.: Light dependency of VOC emissions from selected Mediterranean plant species, *Atmos. Environ.*, 36, 3147–3159, 2002.

Parry, M. A. J., Andralojc, P. J., Khan, S., Lea, P. J., and Keys, A. J.: Rubisco activity: Effects of drought stress, *Annals of Botany*, 89, 833–839, 2002.

Pearcy, R. W., Schulze, E. D., and Zimmermann, R.: Measurement of transpiration and leaf conductance, in: *Plant Physiological Ecology*, Chapman and Hall, New York, 141–142, 1989.

Penuelas, J. and Llusia, J.: Short-term responses of terpene emission rates to experimental changes of PFD in *Pinus halepensis* and *Quercus ilex* in summer field conditions, *Environmental and Experimental Botany*, 42, 61–68, 1999.

Peñuelas, J. and Llusia, J.: Seasonal emission of monoterpenes by the Mediterranean tree *Quercus ilex* in field conditions: Relations with photosynthetic rates, temperature and volatility, *Physiologia Plantarum*, 105, 641–647, 1999.

Peñuelas, J. and Llusia, J.: The complexity of factors driving volatile organic compound emissions by plants, *Biologia Plantarum*, 44, 481–487, 2001.

Peuke, A. D., Schraml, C., Hartung, W., and Rennenberg, H.: Identification of drought-sensitive beech ecotypes by physiological parameters, *New Phytologist*, 154, 373–387, 2002.

Rinne, H. J. I., Guenther, A. B., Greenberg, J. P., and Harley, P. C.: Isoprene and monoterpene fluxes measured above Amazonian rainforest and their dependence on light and temperature, *Atmos. Environ.*, 36, 2421–2426, 2002.

Sabillon, D. and Cremades, L. V.: Diurnal and seasonal variation of monoterpene emission rates for two typical Mediterranean species (*Pinus pinea* and *Quercus ilex*) from field measurements-relationship with temperature and PAR, *Atmos. Environ.*, 35, 4419–4431, 2001.

Schäfer, L., Kesselmeier, J., and Helas, G.: Formic and Acetic acid emission from conifers measured with a “cuvette” technique, in: *CeC Air Pollution Research 39: Field Measurements and Interpretation of Species Related to Photooxidants and Acid Deposition*, edited by: Beilke, S., Angeletti, G., and Slanina, J., Brussels, 319–323, 1992.

Schnitzler, J. P., Graus, M., Kreuzwieser, J., Heizmann, U., Rennenberg, H., Wisthaler, A., and Hansel, A.: Contribution of different carbon sources to isoprene biosynthesis in poplar leaves, *Plant Physiology*, 135, 152–160, 2004.

**Monoterpene emission from European beech**

T. Dindorf et al.

Title Page	Abstract	Introduction
	Conclusions	References
	Tables	Figures
◀	▶	▶▶
◀◀	▶▶▶	▶▶▶▶
Back	Close	
Full Screen / Esc		
Print Version		
Interactive Discussion		

- Schuh, G., Heiden, A. C., Hoffmann, T., Kahl, J., Rockel, P., Rudolph, J., and Wildt, J.: Emissions of volatile organic compounds from sunflower and beech: Dependence on temperature and light intensity, *J. Atmos. Chem.*, 27, 291–318, 1997.
- Schulze, E. D.: Carbon-Dioxide and Water-Vapor Exchange in Response to Drought in the Atmosphere and in the Soil, *Annual Review of Plant Physiology and Plant Molecular Biology*, 37, 247–274, 1986.
- Seufert, G., Bartzis, J., Bomboi, T., Ciccioli, P., Cieslik, S., Dlugi, R., Foster, P., Hewitt, C.N., Kesselmeier, J., Kotzias, D., Lenz, R., Manes, F., Pastor, R. P., Steinbrecher, R., Torres, L., Valentini, R., and Versino, B.: An overview of the Castelporziano experiments, *Atmos. Environ.*, 31, 5–17, 1997.
- Solmon, F., Sarrat, C., Serça, D., Tulet, P., and Rosset, R.: Isoprene and monoterpenes biogenic emissions in France: modeling and impact during a regional pollution episode, *Atmos. Environ.*, 38, 3853–3865, 2004.
- Spirig, C., Neffel, A., Ammann, C., Dommen, J., Grabmer, W., Thielmann, A., Schaub, A., Beauchamp, J., Wisthaler, A., and Hansel, A.: Flux measurements of biogenic VOCs during ECHO 2003, *Atmos. Chem. Phys. Discuss.*, 4, 6603–6643, 2004, [SRef-ID: 1680-7375/acpd/2004-4-6603](https://doi.org/10.5194/acpd/2004-4-6603).
- Staudt, M. and Seufert, G.: Light-Dependent Emission of Monoterpenes by Holm Oak (*Quercus-ilex* L.), *Naturwissenschaften*, 82, 89–92, 1995.
- Staudt, M. and Bertin, N.: Light and temperature dependence of the emission of cyclic and acyclic monoterpenes from holm oak (*Quercus ilex* L.) leaves, *Plant Cell and Environment*, 21, 385–395, 1998.
- Staudt, M., Rambal, S., Joffre, R., and Kesselmeier, J.: Impact of drought on seasonal monoterpene emissions from *Quercus ilex* in southern France, *J. Geophys. Res.-A*, 107, doi:10.1029/2001JD002043, 2002.
- Staudt, M., Joffre, R., and Rambal, S.: How growth conditions affect the capacity of *Quercus ilex* leaves to emit monoterpenes, *New Phytologist*, 158, 61–73, 2003.
- Steinbrecher, R., Schürmann, W., Schreiner, A. M., and Ziegler, H.: Terpenoid emissions from common oak (*Quercus robur* L.) and Norway spruce (*Picea abies* L. Karst), in: *Proceedings of Joint Workshop of CEC/BIATEX of EUROTRAC*, 251–257, 1993
- Street, R. A., Owen, S., Duckham, S. C., Boissard, C., and Hewitt, C. N.: Effect of habitat and age on variations in volatile organic compound (VOC) emissions from *Quercus ilex* and *Pinus pinea*, *Atmos. Environ.*, 31, 89–100, 1997.

## Monoterpene emission from European beech

T. Dindorf et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

- Thomas, F. M.: Growth and water relations of four deciduous tree species (*Fagus sylvatica* L., *Quercus petraea* [MATT.] LIEBL., *Q-pubesces* WILLD., *Sorbus aria* [L.] CR.) occurring at Central-European tree-line sites on shallow calcareous soils: physiological reactions of seedlings to severe drought, *Flora*, 195, 104–115, 2000.
- <sup>5</sup> Tollsten, L. and Muller, P. M.: Volatile organic compounds emitted from beech leaves, *Phytochemistry*, 43, 759–762, 1996.
- Tollsten, L. and Müller, P. M.: Volatile organic compounds emitted from beech leaves, *Phytochemistry*, 43, 759–762, 1996.
- Weiss, A. and Norman, J. M.: Partitioning Solar-Radiation into Direct and Diffuse, Visible and near-Infrared Components, *Agricultural and Forest Meteorology*, 34, 205–213, 1985.
- <sup>10</sup> Went, F. W.: Blue Hazes in the Atmosphere, *Nature*, 187, 641–643, 1960.
- Wolfertz, M., Sharkey, T. D., Boland, W., Kuhnemann, F., Yeh, S., and Weise, S. E.: Biochemical regulation of isoprene emission, *Plant Cell and Environment*, 26, 1357–1364, 2003.

**BGD**

2, 137–182, 2005

---

**Monoterpene  
emission from  
European beech**

T. Dindorf et al.

---

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀

▶

◀

▶

Back Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

**Monoterpene  
emission from  
European beech**

T. Dindorf et al.

**Table 1.** Reference values for branch enclosures. Abbreviations: (a) from originally enclosed leaves, (b) calculated from reference leaves taken from the same branch.

Measurement period	Leaf area (m <sup>2</sup> )	Fresh weight (g)	Dry weight (g)	Water content (%)	Leaf dry mass per area (g m <sup>-2</sup> )
June 2002	0.17 (a)	32.62 (b)	18.35 (b)	44 (b)	108
July/August 2003	0.14 (a)	21.33 (a)	10.73 (a)	50 (a)	77

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Print Version

Interactive Discussion

**Monoterpene  
emission from  
European beech**

T. Dindorf et al.

**Table 2.** Standard emission factor (G97, monoterpenes, sum of 10 individual compounds) for European Beech as measured during the growing season in the years of 2002 and 2003. Standard temperature 30°C, standard light intensity (PAR) 1000  $\mu\text{mol m}^{-2} \text{s}^{-1}$ .

Measurement period	Standard emission factor ( $\mu\text{g g}^{-1} \text{h}^{-1}$ ) ( $\mu\text{mol m}^{-2} \text{s}^{-1}$ )		
	Maximum	Minimum	Average
June 2002	13.5 (3.0)	9.0 (2.0)	13.1 (2.9)
July/August 2003	5.6 (0.9)	3.1 (0.5)	4.4 (0.7)

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Print Version

Interactive Discussion

**Monoterpene emission from European beech**

T. Dindorf et al.

**Table 3.** Standard emission factors (G97) for 15 monoterpene species measured by GC-MS during one single day in August 2003. Standard temperature 30°C, standard light intensity (PAR) 1000  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . Abbreviations: (a) partially decomposed, (b) partial decomposition product.

Monoterpene species	Monoterpene emission, SEF ( $\text{ng g}^{-1} \text{h}^{-1}$ )	Monoterpene emission, SEF (%)
sabinene (a)	4036	56.83
$\alpha$ -thujene (b)	584	8.22
$\gamma$ -terpinene (b)	541	7.62
$\alpha$ -terpinene (b)	483	6.80
p-cymene (b)	365	5.14
$\beta$ -phellandrene (b)	336	4.73
myrcene	166	2.34
limonene	154	2.17
terpinolene (b)	142	2.00
$\alpha$ -phellandrene (b)	113	1.59
$\alpha$ -pinene	99	1.39
$\beta$ -pinene	73	1.03
camphene	6	0.08
$\Delta$ 3-carene	3	0.04
tricyclene	1	0.01

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

**Monoterpene emission from European beech**

T. Dindorf et al.

**Table 4.** Monoterpene emission from European beech as reported by several authors. Abbreviations: *a* (age of trees), *bdl* (below detection limit), *e* (enclosure), *f* (field experiment), *l* (laboratory experiment), *ldm* (leaf disc method), max. (maximum), — (not specified), PAR (photosynthetic active radiation), *r* (radiation), *t* (temperature).

Reference	Monoterpene emission		Comment
	( $\mu\text{g g}^{-1} \text{h}^{-1}$ )	( $\mu\text{g m}^{-2} \text{h}^{-1}$ )	
Hewitt and Street (1992)	<i>bdl</i>	<i>bdl</i>	<i>ldm</i> , <i>a</i> (young/adult) <i>t</i> (—), PAR (max. 900 $\mu\text{mol m}^{-2} \text{s}^{-1}$ )
Steinbrecher et al. (1993)	—	~0.49	<i>e</i> , <i>l</i> , <i>a</i> (2 years) <i>t</i> (2–14°C), PAR (max. 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ )
König et al. (1995)	0.19	—	<i>e</i> , <i>f</i> , <i>a</i> (adult), <i>t</i> (20°C), PAR (—)
Tollsten and Muller (1996)	0.25	—	<i>e</i> , <i>f</i> , <i>a</i> (adult), <i>t</i> (—), PAR (—)
Schuh et al. (1997)	—	414	<i>e</i> , <i>l</i> , <i>a</i> (—), $\phi^{P+B}$ at <i>t</i> (25°C), PAR (various)
Kahl et al. (1999)	—	284	<i>e</i> , <i>l</i> , <i>a</i> (6 years), <i>t</i> (25°C), <i>r</i> (max. 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ )
This study	4–13	334–1415	<i>e</i> , <i>f</i> , <i>a</i> (~160 years), <i>t</i> (30°C), PAR (max. 1736 $\mu\text{mol m}^{-2} \text{s}^{-1}$ )

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Print Version

Interactive Discussion

**BGD**

2, 137–182, 2005

---

**Monoterpene  
emission from  
European beech**

T. Dindorf et al.

---

Title Page	Introduction
Abstract	References
Conclusions	Figures
Tables	
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

EGU




**Fig. 1.** Enclosure of European Beech (*Fagus sylvatica* L.) during the experiments performed in June 2002.

**Monoterpene emission from European beech**

T. Dindorf et al.


Title Page	Abstract	Conclusions	Tables	◀	▶	Back	Full Screen / Esc	Print Version	Interactive Discussion
	Introduction	References	Figures	◀	▶	Close			



**Fig. 2.** Diurnal course of micrometeorological and physiological parameters during both field experiments in June 2002 (left panel) and July/August 2003 (right panel). Top graph: Development of leaf temperatures (red line) and monoterpene emission. Monoterpene emission that was calculated by application of the G97 algorithm is indicated by the grey line, monoterpene emission that was measured by GC-FID analysis is indicated by red circles. Middle graph: Diurnal course of net CO<sub>2</sub> assimilation (green line) and transpiration (blue line). Bottom graph: Photosynthetic active radiation (PAR) (black line).

**Monoterpene emission from European beech**

T. Dindorf et al.




**Fig. 3.** Light saturation and temperature dependence of net CO<sub>2</sub> assimilation and monoterpene emission during both measurement campaigns. Figures A and C indicate measurements performed in the year 2002 (left panel). Figures B and D indicate measurements performed in the year 2003 (right panel). Net CO<sub>2</sub> assimilation is indicated by green diamonds. Monoterpene emission was normalised with the respective light and temperature function of the G97 algorithm and is indicated by red circles. Monoterpene emission calculated by the respective term of the G97 algorithm is indicated by the black line.

Title Page	Abstract	Conclusions	Tables	◀	▶	Full Screen / Esc
Introduction	References	Figures	◀	▶	Close	Print Version
						Interactive Discussion

Monoterpene emission from European beech

T. Dindorf et al.




**Fig. 4.** Monoterpene emission as a function of the G97 and S97 algorithm for the year 2002 (top graph) and 2003 (bottom graph). Red circles indicate monoterpene emission correlated to the G97 function. Black diamonds indicate monoterpene emission correlated to the S97 ( $\Phi^B$ ) function. Red solid lines represent the linear fit to the G97 function for each dataset. Black solid lines represent the linear fit to the S97 function. The corresponding equation of the linear fit is indicated in the graph.

Title Page	Introduction	References	Figures
Abstract	Conclusions	Tables	◀   ▶
◀	▶	Back	Close
Full Screen / Esc			
Print Version			
Interactive Discussion			

**Monoterpene emission from European beech**

T. Dindorf et al.


Title Page	Abstract	Conclusions	Tables	◀	▶	Back	Full Screen / Esc
Introduction	References	Figures	◀	▶	Close	Print Version	Interactive Discussion



**Fig. 5.** Course of photosynthetic active radiation (black solid line) and leaf temperature (red solid line) during the artificial shading of the branch enclosure in July 2003. Monoterpene emission measured prior, during and after the artificial darkening is indicated by red circles.

**Monoterpene emission from European beech**

T. Dindorf et al.




**Fig. 6.** Course of photosynthetic active radiation (PAR), leaf temperature, and monoterpene emission measured during one single day in August 2003. Top graph: Diurnal course of PAR (black squares plus dashed line) and leaf temperature (red circles plus dotted line). Data show the appropriate 30 min average during that VOC cartridges were collected. Bottom graph: Diurnal course and composition of monoterpene emission. Data show monoterpene emission measured by GC-MS analysis (stacked bars, for caption see graph), the respective emission calculated by the G97 algorithm (grey stars plus solid line), and the S97 algorithm (emission from biosynthesis only, red diamonds plus dotted line).

Title Page	Abstract	Conclusions	Tables	◀	▶	Full Screen / Esc
Introduction	References	Figures	◀	▶	Close	Print Version
				◀	▶	Interactive Discussion

**Monoterpene emission from European beech**

T. Dindorf et al.

Title Page	Abstract	Conclusions	Tables	◀	▶	Back	Full Screen / Esc
Introduction	References	Figures	◀	▶	Close	Print Version	Interactive Discussion




**Fig. 7.** Midday depression of plant physiology in June 2002. Figures A to H show the correlation of net CO<sub>2</sub> assimilation (green diamonds) and stomatal conductance (grey stars) to photosynthetic active radiation (PAR) for each single day of the measurement period (12–19 June 2002). The development of the respective daily leaf temperature is indicated by the small graphs as a function of time.



**Monoterpene emission from European beech**

T. Dindorf et al.

Title Page	Abstract	Introduction
	Conclusions	References
	Tables	Figures
◀	▶	▶▶
◀	▶	
Back	Close	
Full Screen / Esc		
Print Version		
Interactive Discussion		




**Fig. 9.** Ambient temperature and rainfall prior and during the enclosure measurements of European beech in the years of 2002 (top graph) and 2003 (bottom graph). Grey areas indicate the respective measurement period of each year. Solid lines represent the daily average temperature. Dashed lines show the daily minima and maxima temperature. Grey bars indicate the respective amount of rainfall. Standard emission factors (SEF) for monoterpene exchange rates were calculated for single measurement days and are indicated by grey stars.

**Monoterpene  
emission from  
European beech**

T. Dindorf et al.

Title Page	Abstract	Introduction	Conclusions	Tables	Figures
1.1	0.9	0.8	0.7	0.6	0.5
1.1	1.1	1.3	1.4	1.5	1.6
1.7	1.8	1.9	2.0	2.1	2.2
2.3	2.4	2.5	2.6	2.7	2.8
2.9	3.0	3.1	3.2	3.3	3.4
3.5	3.6	3.7	3.8	3.9	4.0
4.1	4.2	4.3	4.4	4.5	4.6
4.7	4.8	4.9	5.0	5.1	5.2
5.3	5.4	5.5	5.6	5.7	5.8
5.9	6.0	6.1	6.2	6.3	6.4
6.5	6.6	6.7	6.8	6.9	7.0
7.1	7.2	7.3	7.4	7.5	7.6
7.7	7.8	7.9	8.0	8.1	8.2
8.3	8.4	8.5	8.6	8.7	8.8
8.9	9.0	9.1	9.2	9.3	9.4
9.5	9.6	9.7	9.8	9.9	10.0
10.1	10.2	10.3	10.4	10.5	10.6
10.7	10.8	10.9	11.0	11.1	11.2
11.3	11.4	11.5	11.6	11.7	11.8
11.9	12.0	12.1	12.2	12.3	12.4
12.5	12.6	12.7	12.8	12.9	13.0
13.1	13.2	13.3	13.4	13.5	13.6
13.7	13.8	13.9	14.0	14.1	14.2
14.3	14.4	14.5	14.6	14.7	14.8
14.9	15.0	15.1	15.2	15.3	15.4
15.5	15.6	15.7	15.8	15.9	16.0
16.1	16.2	16.3	16.4	16.5	16.6
16.7	16.8	16.9	17.0	17.1	17.2
17.3	17.4	17.5	17.6	17.7	17.8
17.9	18.0	18.1	18.2	18.3	18.4
18.5	18.6	18.7	18.8	18.9	19.0
19.1	19.2	19.3	19.4	19.5	19.6
19.7	19.8	19.9	20.0	20.1	20.2
20.3	20.4	20.5	20.6	20.7	20.8
20.9	21.0	21.1	21.2	21.3	21.4
21.5	21.6	21.7	21.8	21.9	22.0
22.1	22.2	22.3	22.4	22.5	22.6
22.7	22.8	22.9	23.0	23.1	23.2
23.3	23.4	23.5	23.6	23.7	23.8
23.9	24.0	24.1	24.2	24.3	24.4
24.5	24.6	24.7	24.8	24.9	25.0
25.1	25.2	25.3	25.4	25.5	25.6
25.7	25.8	25.9	26.0	26.1	26.2
26.3	26.4	26.5	26.6	26.7	26.8
26.9	27.0	27.1	27.2	27.3	27.4
27.5	27.6	27.7	27.8	27.9	28.0
28.1	28.2	28.3	28.4	28.5	28.6
28.7	28.8	28.9	29.0	29.1	29.2
29.3	29.4	29.5	29.6	29.7	29.8
29.9	30.0	30.1	30.2	30.3	30.4
30.5	30.6	30.7	30.8	30.9	31.0
31.1	31.2	31.3	31.4	31.5	31.6
31.7	31.8	31.9	32.0	32.1	32.2
32.3	32.4	32.5	32.6	32.7	32.8
32.9	33.0	33.1	33.2	33.3	33.4
33.5	33.6	33.7	33.8	33.9	34.0
34.1	34.2	34.3	34.4	34.5	34.6
34.7	34.8	34.9	35.0	35.1	35.2
35.3	35.4	35.5	35.6	35.7	35.8
35.9	36.0	36.1	36.2	36.3	36.4
36.5	36.6	36.7	36.8	36.9	37.0
37.1	37.2	37.3	37.4	37.5	37.6
37.7	37.8	37.9	38.0	38.1	38.2
38.3	38.4	38.5	38.6	38.7	38.8
38.9	39.0	39.1	39.2	39.3	39.4
39.5	39.6	39.7	39.8	39.9	40.0
40.1	40.2	40.3	40.4	40.5	40.6
40.7	40.8	40.9	41.0	41.1	41.2
41.3	41.4	41.5	41.6	41.7	41.8
41.9	42.0	42.1	42.2	42.3	42.4
42.5	42.6	42.7	42.8	42.9	43.0
43.1	43.2	43.3	43.4	43.5	43.6
43.7	43.8	43.9	44.0	44.1	44.2
44.3	44.4	44.5	44.6	44.7	44.8
44.9	45.0	45.1	45.2	45.3	45.4
45.5	45.6	45.7	45.8	45.9	46.0
46.1	46.2	46.3	46.4	46.5	46.6
46.7	46.8	46.9	47.0	47.1	47.2
47.3	47.4	47.5	47.6	47.7	47.8
47.9	48.0	48.1	48.2	48.3	48.4
48.5	48.6	48.7	48.8	48.9	49.0
49.1	49.2	49.3	49.4	49.5	49.6
49.7	49.8	49.9	50.0	50.1	50.2
50.3	50.4	50.5	50.6	50.7	50.8
50.9	51.0	51.1	51.2	51.3	51.4
51.5	51.6	51.7	51.8	51.9	52.0
52.1	52.2	52.3	52.4	52.5	52.6
52.7	52.8	52.9	53.0	53.1	53.2
53.3	53.4	53.5	53.6	53.7	53.8
53.9	54.0	54.1	54.2	54.3	54.4
54.5	54.6	54.7	54.8	54.9	55.0
55.1	55.2	55.3	55.4	55.5	55.6
55.7	55.8	55.9	56.0	56.1	56.2
56.3	56.4	56.5	56.6	56.7	56.8
56.9	57.0	57.1	57.2	57.3	57.4
57.5	57.6	57.7	57.8	57.9	58.0
58.1	58.2	58.3	58.4	58.5	58.6
58.7	58.8	58.9	59.0	59.1	59.2
59.3	59.4	59.5	59.6	59.7	59.8
59.9	60.0	60.1	60.2	60.3	60.4
60.5	60.6	60.7	60.8	60.9	61.0
61.1	61.2	61.3	61.4	61.5	61.6
61.7	61.8	61.9	62.0	62.1	62.2
62.3	62.4	62.5	62.6	62.7	62.8
62.9	63.0	63.1	63.2	63.3	63.4
63.5	63.6	63.7	63.8	63.9	64.0
64.1	64.2	64.3	64.4	64.5	64.6
64.7	64.8	64.9	65.0	65.1	65.2
65.3	65.4	65.5	65.6	65.7	65.8
65.9	66.0	66.1	66.2	66.3	66.4
66.5	66.6	66.7	66.8	66.9	67.0
67.1	67.2	67.3	67.4	67.5	67.6
67.7	67.8	67.9	68.0	68.1	68.2
68.3	68.4	68.5	68.6	68.7	68.8
68.9	69.0	69.1	69.2	69.3	69.4
69.5	69.6	69.7	69.8	69.9	70.0
70.1	70.2	70.3	70.4	70.5	70.6
70.7	70.8	70.9	71.0	71.1	71.2
71.3	71.4	71.5	71.6	71.7	71.8
71.9	72.0	72.1	72.2	72.3	72.4
72.5	72.6	72.7	72.8	72.9	73.0
73.1	73.2	73.3	73.4	73.5	73.6
73.7	73.8	73.9	74.0	74.1	74.2
74.3	74.4	74.5	74.6	74.7	74.8
74.9	75.0	75.1	75.2	75.3	75.4
75.5	75.6	75.7	75.8	75.9	76.0
76.1	76.2	76.3	76.4	76.5	76.6
76.7	76.8	76.9	77.0	77.1	77.2
77.3	77.4	77.5	77.6	77.7	77.8
77.9	78.0	78.1	78.2	78.3	78.4
78.5	78.6	78.7	78.8	78.9	79.0
79.1	79.2	79.3	79.4	79.5	79.6
79.7	79.8	79.9	80.0	80.1	80.2
80.3	80.4	80.5	80.6	80.7	80.8
80.9	81.0	81.1	81.2	81.3	81.4
81.5	81.6	81.7	81.8	81.9	82.0
82.1	82.2	82.3	82.4	82.5	82.6
82.7	82.8	82.9	83.0	83.1	83.2
83.3	83.4	83.5	83.6	83.7	83.8
83.9	84.0	84.1	84.2	84.3	84.4
84.5	84.6	84.7	84.8	84.9	85.0
85.1	85.2	85.3	85.4	85.5	85.6
85.7	85.8	85.9	86.0	86.1	86.2
86.3	86.4	86.5	86.6	86.7	86.8
86.9	87.0	87.1	87.2	87.3	87.4
87.5	87.6	87.7	87.8	87.9	88.0
88.1	88.2	88.3	88.4	88.5	88.6
88.7	88.8	88.9	89.0	89.1	89.2
89.3	89.4	89.5	89.6	89.7	89.8
89.9	90.0	90.1	90.2	90.3	90.4
90.5	90.6	90.7	90.8	90.9	91.0
91.1	91.2	91.3	91.4	91.5	91.6
91.7	91.8	91.9	92.0	92.1	92.2
92.3	92.4	92.5	92.6	92.7	92.8
92.9	93.0	93.1	93.2	93.3	93.4
93.5	93.6	93.7	93.8	93.9	94.0
94.1	94.2	94.3	94.4	94.5	94.6
94.7	94.8	94.9	95.0	95.1	95.2
95.3	95.4	95.5	95.6	95.7	95.8
95.9	96.0	96.1	96.2	96.3	96.4
96.5	96.6	96.7	96.8	96.9	97.0
97.1	97.2	97.3	97.4	97.5	97.6
97.7	97.8	97.9	98.0	98.1	98.2
98.3	98.4	98.5	98.6	98.7	98.8
98.9	99.0	99.1	99.2	99.3	99.4
99.5	99.6	99.7	99.8	99.9	100.0




**Fig. 10.** Mean European monoterpene emission fluxes for the month of July in  $\mu\text{g C g}^{-1} \text{h}^{-1}$ . Monoterpene emissions were calculated by the default monoterpene emission factors, considering temperature dependence only (G95OIs).



**Monoterpene  
emission from  
European beech**

T. Dindorf et al.

Title Page	Abstract	Conclusions	Tables	1◀	▶1	Full Screen / Esc
Introduction	References	Figures	▶	▶	Close	Print Version
			Back	◀		Interactive Discussion



**Fig. 12.** Relative increase of mean monoterpene emission in % for the month of July if the high standard emission factor and the spatial distribution of European beech are considered as a function of light and temperature.