

Multiscale approach to explore the relationships between connectivity and function in whole brain simulations

S. Diaz¹, C. Nowke², A. Peyser¹, B. Weyers², B. Hentschel², A. Morrison^{1,3}, T. W. Kuhlen²

1. Simulation Laboratory Neuroscience – Bernstein Facility for Simulation and Database Technology.

Institute for Advanced Simulation, Jülich Forschungszentrum, Jülich, Germany


2. IT Center, RWTH Aachen University, Aachen, Germany

3. Institute of Cognitive Neuroscience, Faculty of Psychology, Ruhr-University Bochum, Bochum, Germany

Motivation

- We want to better understand the relationship between connectivity and function in the brain at different scales.
- To achieve this, in this work we use point-neuron network simulations to complement connectivity information for whole brain simulations based on a dynamic neuron mass model.
- The results of subsequent simulations will be compared with experimental data in order to find correlations between function and structure.

Multiscale single region models



Multiscale approach

- We simulate a whole brain parcellated into 68 regions based on [1].
- Each region is modeled as a dynamic neuron mass, and in parallel, as small 200 point-neuron populations in NEST [2].
- Structural plasticity in NEST is then used to calculate the inner inhibitory connectivity required to match experimentally observed firing rates inside each region.
- The point-neuron network self-generates the inner inhibitory connectivity using simple homeostatic rules.
- With the resulting connectivity data from the NEST simulations and experimentally obtained DTI inter-region connectivity, simulations of the whole brain producing results comparable to experimental fMRI data are performed.


Acknowledgments

The Excellence Initiative of the German federal and state governments, JARA-HPC, the CRCNS grant and the EU Seventh Framework Programme (FP7/2007-2013) grant 604202 (HBP).

Interactive steering and visualization

- We use an interactive tool based on the NETT framework developed for visualization and steering of the structural plasticity algorithm to bring all regions to their ideal firing activity.
- The user can see the evolution of the firing rate and the creation/ deletion of connections, modify parameters and visualize the impact of the modifications while the simulation progresses.
- This steering tool can be used interactively on supercomputers for larger number of regions or neurons per region.

Multiple regions view



Changes in Inhibitory Connections guided by the structural plasticity algorithm

Evolution of the average firing rate in each region




Change multiple parameters and store status

Discussion

- Fitting and parameter space exploration:
 - is around 10 times faster than iterative fitting algorithms
 - is more robust since it progressively achieves global stability
 - aids a better understanding of the parameter space
- Our multiscale approach supports a new way to explore the impact of connectivity in function at different scales.

How does the underlying *structural* connectivity affect our ability to predict experimental *functional* data with our simulations?



References

[1] Deco, Gustavo, et al. "How local excitation-inhibition ratio impacts the whole brain dynamics." The Journal of Neuroscience 34.23 (2014): 7886-7898.

[2] Bos, Hannah et al.. (2015). NEST 2.10.0. Zenodo. 10.5281/zenodo.44222. [3] Deco, Gustavo, et al. "Resting-state functional connectivity emerges from

structurally and dynamically shaped slow linear fluctuations." The Journal of Neuroscience 33.27 (2013): 11239-11252.