

Genetic architecture of plant stress resistance: multi-trait genome-wide association mapping

Manus P. M. Thoen^{1,2,3*}, Nelson H. Davila Olivas^{1*}, Karen J. Kloth^{1,2,3*}, Silvia Coolen^{4*}, Ping-Ping Huang^{5*}, Mark G. M. Aarts⁵, Johanna A. Bac-Molenaar^{2,5}, Jaap Bakker⁶, Harro J. Bouwmeester², Colette Broekgaarden^{4,7}, Johan Bucher⁷, Jacqueline Busscher-Lange^{1,2,3}, Xi Cheng², Emilie F. Fradin², Maarten A. Jongsma³, Magdalena M. Julkowska^{8,9}, Joost J. B. Keurentjes⁵, Wilco Ligterink², Corné M. J. Pieterse⁴, Carolien Ruyter-Spira², Geert Smant⁶, Christa Testerink^{8,9}, Björn Usadel¹⁰, Joop J. A. van Loon¹, Johan A. van Pelt⁴, Casper C. van Schaik⁶, Saskia C. M. van Wees⁴, Richard G. F. Visser⁷, Roeland Voorrips⁷, Ben Vosman⁷, Dick Vreugdenhil², Sonja Warmerdam⁶, Gerrie L. Wiegiers^{1,3}, Joost van Heerwaarden¹¹, Willem Kruijer¹¹, Fred A. van Eeuwijk¹¹ and Marcel Dicke¹

¹Laboratory of Entomology, Wageningen University and Research, PO Box 16, 6700 AA Wageningen, the Netherlands; ²Laboratory of Plant Physiology, Wageningen University and Research, PO Box 16, 6700 AA Wageningen, the Netherlands; ³Business Unit Bioscience, Wageningen Plant Research, Wageningen University and Research, PO Box 16, 6700 AA Wageningen, the Netherlands; ⁴Plant–Microbe Interactions, Department of Biology, Utrecht University, PO Box 800.56, 3508 TB Utrecht, the Netherlands; ⁵Laboratory of Genetics, Wageningen University and Research, PO Box 16, 6700 AA Wageningen, the Netherlands; ⁶Laboratory of Nematology, Wageningen University and Research, PO Box 8213, 6700 ES Wageningen, the Netherlands; ⁷Wageningen University and Research Plant Breeding, Wageningen University and Research, PO Box 386, 6700 AJ Wageningen, the Netherlands; ⁸Section of Plant Physiology, Swammerdam Institute for Life Sciences, University of Amsterdam, PO Box 94215, 1090 GE Amsterdam, the Netherlands; ⁹Section of Plant Cell Biology, Swammerdam Institute for Life Sciences, University of Amsterdam, PO Box 94215, 1090 GE Amsterdam, the Netherlands; ¹⁰Institute for Biology I, RWTH Aachen University, Worringer Weg 3, 52074 Aachen, Germany; ¹¹Biometris, Wageningen University and Research, PO Box 16, 6700 AA Wageningen, the Netherlands

Summary

Author for correspondence:

Marcel Dicke

Tel: +31317484311

Email: marcel.dicke@wur.nl

Received: 26 June 2016

Accepted: 17 August 2016

New Phytologist (2017) 213: 1346–1362

doi: 10.1111/nph.14220

Key words: abiotic stress, biotic stress, genetic architecture, genome-wide association mapping, multiple stresses.

- Plants are exposed to combinations of various biotic and abiotic stresses, but stress responses are usually investigated for single stresses only.
- Here, we investigated the genetic architecture underlying plant responses to 11 single stresses and several of their combinations by phenotyping 350 *Arabidopsis thaliana* accessions. A set of 214 000 single nucleotide polymorphisms (SNPs) was screened for marker-trait associations in genome-wide association (GWA) analyses using tailored multi-trait mixed models.
- Stress responses that share phytohormonal signaling pathways also share genetic architecture underlying these responses. After removing the effects of general robustness, for the 30 most significant SNPs, average quantitative trait locus (QTL) effect sizes were larger for dual stresses than for single stresses.
- Plants appear to deploy broad-spectrum defensive mechanisms influencing multiple traits in response to combined stresses. Association analyses identified QTLs with contrasting and with similar responses to biotic vs abiotic stresses, and below-ground vs above-ground stresses. Our approach allowed for an unprecedented comprehensive genetic analysis of how plants deal with a wide spectrum of stress conditions.

Introduction

In nature, plants face variable environments that impose a wide range of biotic and abiotic stresses. These include, for example, below-ground and above-ground stresses, stresses imposed by unicellular and multicellular organisms, and short- and long-lasting stresses. Under natural conditions, these stresses do not occur in isolation but are commonly present simultaneously

(Rizhsky *et al.*, 2004; Bergelson & Roux, 2010; Mittler & Blumwald, 2010; Vile *et al.*, 2012; Prasad & Sonnewald, 2013; Rasmussen *et al.*, 2013; Kissoudis *et al.*, 2014; Rivero *et al.*, 2014; Sewelam *et al.*, 2014; Suzuki *et al.*, 2014). Thus, plants are under strong selection to adapt to local conditions and have evolved sophisticated mechanisms to withstand multiple adverse environmental conditions (Howe & Jander, 2008; Bergelson & Roux, 2010; Pieterse *et al.*, 2012; Stam *et al.*, 2014; Brachi *et al.*, 2015; Julkowska & Testerink, 2015; Kerwin *et al.*, 2015). Yet, investigating this in a targeted experimental way is a major challenge

*These authors contributed equally to this work.

owing to the complexity of multiple stress exposure. To gain an insight into the adaptation of plants to the wide variety of stress-inducing conditions they face, genetic variation and mechanisms underlying stress resistance should be studied (Alonso-Blanco *et al.*, 2009; Brachi *et al.*, 2015; Kerwin *et al.*, 2015). The responses of plants to stresses have traditionally been investigated for individual stresses (Howe & Jander, 2008), but the research focus is currently shifting towards plant responses to combinations of stresses (Holopainen & Gershenzon, 2010; Pierik & Testerink, 2014; Stam *et al.*, 2014; Suzuki *et al.*, 2014; Kissoudis *et al.*, 2015). The emerging picture is that responses to stress combinations cannot be predicted reliably from the responses to individual stresses (De Vos *et al.*, 2006; Makumburage *et al.*, 2013). For instance, the majority of transcriptional responses of *Arabidopsis* to combinations of two abiotic stresses could not be predicted from responses to the individual stresses (Rasmussen *et al.*, 2013). Moreover, phenotype expression in response to two biotic stresses could not be predicted on the basis of existing information regarding interactions between underlying signaling pathways (De Vos *et al.*, 2006). Phytohormones are major players in a signaling network, mediating responses to both biotic and abiotic stresses (Pieterse *et al.*, 2009). For instance, chewing insect herbivores particularly elicit the jasmonic acid (JA), abscisic acid (ABA) and ethylene (ET) signaling pathways; phloem-sucking insects and biotrophic microbial pathogens particularly elicit the salicylic acid (SA) pathway; and drought elicits the ABA pathway (Pieterse *et al.*, 2009). The phytohormonal responses exhibit extensive crosstalk, resulting in specific changes in plant phenotype in response to individual stresses (De Vos *et al.*, 2005; Pieterse *et al.*, 2012).

In plant breeding, resistance and tolerance to multiple stresses are a common selection target (Braun *et al.*, 1996). A well-known strategy to achieve resistance and tolerance is by evaluation of candidate varieties in multi-environment trials, that is, field trials at multiple locations during several years (van Eeuwijk *et al.*, 2010; Malosetti *et al.*, 2013). In such trials, multiple stresses can occur, but their occurrence and the intensity with which they occur are not guaranteed and, therefore, plant breeders developed the concept of managed stress trials in which specific and well-defined stress conditions are imposed for a single stress or a small number of stresses (Cooper & Hammer, 1996; Cooper *et al.*, 2014). Recently, the urge to manage environmental factors even more precisely has led to the development of phenotyping platforms, where, again, mainly single stresses are investigated (Fiorani & Schurr, 2013; Granier & Vile, 2014; Kloth *et al.*, 2015).

Most studies, outside plant breeding, that have examined plant responses to multiple stresses included only one or a few genotypes (Holopainen & Gershenzon, 2010; Rasmussen *et al.*, 2013; Pierik & Testerink, 2014; Stam *et al.*, 2014; Suzuki *et al.*, 2014; Kissoudis *et al.*, 2015). To obtain a further understanding of the genetic architecture of complex traits such as plant adaptation to a diversity of stresses, extensive study of the natural genetic variation within a species is instrumental. Genome-wide association (GWA) analysis is an important tool for this, requiring a large number of well-genotyped plant accessions. Yet, although the interest in natural variation and GWA mapping is rapidly

increasing (Wijnen & Keurentjes, 2014; Ogura & Busch, 2015), a large-scale evaluation of natural genetic variation in resistance of plants to the full diversity of stresses to which they are exposed, including pathogens, herbivores and abiotic stresses and their interactions, has not been done to date. To elucidate the genetic architecture of plant stress resistance, an integrated approach is needed that models the genetics of responses to a range of single and combined stresses, including the interaction between those responses. Here, we have adopted a comprehensive and integrated approach to investigate the genetics underlying plant responses to 15 carefully standardized single stresses or stress combinations (Table 1), making use of a global population of 350 *Arabidopsis* accessions that have been genotyped for 214 000 single nucleotide polymorphisms (SNPs) (Baxter *et al.*, 2010; Li *et al.*, 2010). The standardization of these 15 stress conditions is an important element of the study, because it allows for phenotyping of well-defined stress responses. We developed a tailored multi-trait GWA analysis that allowed the identification of candidate genes associated with plant responses to multiple stresses that were validated by gene expression and mutant analyses.

Materials and Methods

Arabidopsis thaliana population

In this study we included 350 *Arabidopsis thaliana* (L.) Heynh. accessions from the HapMap population (<http://bergelson.uchicago.edu/wp-content/uploads/2015/04/Justins-360-lines.xls>). The HapMap population has been genotyped for 250 000 bi-allelic SNPs (Baxter *et al.*, 2010; Platt *et al.*, 2010; Chao *et al.*, 2012) and after quality control and imputation this SNP set was reduced to a set of 214 051 SNPs.

Definition of the target traits

For every experiment, the target traits were derived from the individual plant data using the following strategy. First, when residuals deviated from normality, a logarithmic, arcsine or square root transformation was applied to the original observations. Second, genotypic (accession) means for each treatment were calculated using a mixed model to account for design effects. Different mixed models were used in the experiments, reflecting the different designs. In all cases, accession effects were modeled as fixed, and the accession means were the best linear unbiased estimator (BLUE) of these effects. Third, for traits measured in treatment and control conditions, differences or residuals (when regressing treatment on control values) were defined, in order to obtain a measure of stress tolerance that was corrected for the expression of the same trait under control conditions. Finally, within each experiment, the traits were replaced by the first principal component if the latter explained more than half of the variation in all traits in this experiment; in all other cases, the original traits were retained. An overview of final traits and their corresponding sections in the Supporting Information Methods can be found in Table 1. In case of replacement by the first principal component, original traits and the variance explained by the first principal

Table 1 Phenotypes assessed

	Stress	Trait name	Section of Supporting Information	Trait phenotype	Treatment
Abiotic stresses	Salt	Salt_1	Methods S1	Main root length, number of lateral roots and straightness	75 mM NaCl
		Salt_2	Methods S1	Main root length	125 mM NaCl
		Salt_3	Methods S1	Number of lateral roots	125 mM NaCl
		Salt_4	Methods S1	Main root angle	125 mM NaCl
		Salt_5	Methods S2	Biomass	25 mM NaCl
	Drought	Drought_1	Methods S2	Biomass	Drought
		Drought_2	Methods S3	Biomass	Drought
	Osmotic	Osmotic	Methods S2	Biomass	PEG8000
	Heat	Heat	Methods S2	Number of siliques	35°C
Biotic stresses	Parasitic plant	Parasitic plant	Methods S4	Attachments	<i>Phelipanche ramosa</i>
	Nematode	Nematode	Methods S5	Offspring, egg mass	<i>Meloidogyne incognita</i>
	Whitefly	Whitefly_1	Methods S6	Survival, whiteflies	<i>Aleyrodes proletella</i>
		Whitefly_2	Methods S6	Reproduction, eggs	<i>A. proletella</i>
	Aphid	Aphid_1	Methods S7	Behavior T1, probing	<i>Myzus persicae</i>
		Aphid_2	Methods S7	Behavior T2, probing	<i>M. persicae</i>
		Aphid_3	Methods S7	Offspring, aphids	<i>M. persicae</i>
	Thrips	Thrips_1	Methods S8	Feeding damage	<i>Frankliniella occidentalis</i>
		Thrips_2	Methods S8	Behavior T1	<i>F. occidentalis</i>
		Thrips_3	Methods S8	Behavior T2	<i>F. occidentalis</i>
	Caterpillar	Caterpillar_1	Methods S9	Leaf area consumed	<i>Pieris rapae</i>
		Caterpillar_2	Methods S3	Biomass	<i>P. rapae</i>
		Caterpillar_3	Methods S3	Number of damaged leaves and feeding sites	<i>P. rapae</i>
	Fungus	Fungus	Methods S10	Number of spreading lesions	<i>Botrytis cinerea</i>
	Double stress	Fungus and caterpillar_1	Methods S3	Biomass	<i>B. cinerea</i> and <i>P. rapae</i>
		Fungus and caterpillar_2	Methods S3	Number of damaged leaves and feeding sites	<i>B. cinerea</i> and <i>P. rapae</i>
		Caterpillar and fungus	Methods S10	Number of spreading lesions	<i>P. rapae</i> and <i>B. cinerea</i>
Abiotic and biotic stress	Double stress	Drought and fungus	Methods S10	Number of spreading lesions	Drought and <i>B. cinerea</i>
		Drought and caterpillar	Methods S3	Number of damaged leaves and feeding sites	Drought and <i>P. rapae</i>
		Caterpillar and osmotic_1	Methods S9	Projected leaf area	<i>P. rapae</i> and PEG8000
		Caterpillar and osmotic_2	Methods S9	Biomass	<i>P. rapae</i> and PEG8000

The dataset contains three plant stress categories applied to *Arabidopsis thaliana*: abiotic stress, biotic stress and combinations of both abiotic and biotic stress. Phenotype assessments that were performed under similar environmental conditions have similar background shading (light and dark gray). ‘Phenotype’ refers to different phenotypic assessments (in some cases the first principal component of a group of phenotypes). ‘Treatment’ refers to the sort of stress that was applied. Additional information on traits can be found in Supporting Information Methods S1–S10. Yellow, abiotic stress; green, biotic stress; blue, combinations of biotic and abiotic stress.

component are listed (Supporting Information Methods Tables M1–M5 in Methods S1–S3, S7 & S9). In total, phenotypic data for 73 individual traits were obtained by 10 different research groups. All calculations were performed in R, unless stated otherwise. Mixed-model analysis was performed with the R package ASREML (Butler *et al.*, 2009). In all equations, the term *E* denotes residual error. All other terms represent fixed effects unless stated otherwise. A colon (:) is used to define interactions between terms.

Statistics

Genetic correlation networks and heritability Pairwise marker-based genetic correlations between traits, genomic correlations, were estimated using a multi-trait mixed model (MTMM) (Korte

et al., 2012). Residuals were assumed to be uncorrelated for traits that were measured on different plants. For some pairs of traits, the likelihood was monotone, which can also occur in single-trait mixed models (Kruijer *et al.*, 2015). In this case, the genetic correlation was estimated by the (Pearson) correlation between the univariate G-BLUPs (De los Campos *et al.*, 2013) estimated for these traits. A network between predefined groups of traits was constructed by connecting groups whose average genetic correlation across pairs of traits was > 0.2.

Narrow sense heritability (Table S1) was estimated using the mixed model $Y_i = \mu + A_i + E_i$ where Y_i represents the phenotypic means of accessions ($i = 1, \dots, 350$), and A_i and E_i are random genetic and residual effects. The vector of additive genetic effects follows a multivariate normal distribution with covariance $\sigma_A^2 K$, K being a marker-based relatedness matrix. The residual errors

are independent, with variance σ_E^2 . We obtained restricted maximum likelihood (REML) estimates of σ_A^2 and σ_E^2 , and estimated heritability as $h^2 = \sigma_A^2 / (\sigma_A^2 + \sigma_E^2)$. This is an estimate of narrow-sense heritability, as the model for the genetic effects only captures additive effects, and σ_E^2 is the sum of environmental and nonadditive genetic effects (see e.g. Kruijer *et al.*, 2015).

Multi-trait mixed models Following Zhou & Stephens (2014), we assume the MTMM, $Y = XB + G + E$, with Y being the genotypes by traits ($n \times p$) matrix of phenotypic observations. The terms XB , G and E stand for, respectively, the fixed effects (including trait-specific intercepts and SNP effects) and the random genetic and environmental effects. G follows a zero mean matrix-variate normal distribution with row-covariance (marker-based kinship) matrix K and column (trait) covariance matrix V_g . V_g is a $p \times p$ matrix modeling the genetic correlations between traits. This is equivalent to $g = \text{vec}(G)$ (the vector containing the columns of G) being multivariate normal with a covariance matrix defined by the Kronecker product $V_g \otimes K$ (Zhou & Stephens, 2014). Similarly, $\text{vec}(E)$ follows a zero mean normal distribution with covariance $V_e \otimes I_n$, where V_e accounts for the nongenetic correlations between traits.

Factor-analytic models As V_g and V_e contain a total of $p(p+1)$ parameters, the MTMM becomes difficult to fit for > 10 traits (Zhou & Stephens, 2014). For V_g we therefore assumed a factor analytic model, which is well known in the context of quantitative trait locus (QTL) mapping for experimental populations with limited numbers of markers (Boer *et al.*, 2007), but has not been used in the context of multivariate GWA studies (GWAS). As almost all traits were derived from measurements on different plants, a diagonal model $V_e = \text{diag}(\sigma_{e,1}^2, \dots, \sigma_{e,p}^2)$ was chosen for the environmental covariances. For V_g , a second-order factor analytic structure was chosen $V_g = \sigma_g^2(\lambda\lambda^t + \text{diag}(\tau_1^2, \dots, \tau_p^2))$, where σ_g^2 represents a scale parameter, the magnitude of genetic effects, the $p \times 2$ matrix λ contains the trait-specific scores belonging to the factor analytic part of the model that provides a rank two variance-covariance structure between traits, and $\text{diag}(\tau_1^2, \dots, \tau_p^2)$ provides trait-specific residual genetic variances (Piepho, 1997; Meyer, 2009). The model was fitted with the R package ASReml (Butler *et al.*, 2009).

Compressed kinship Factor analytic models have been successfully applied to experimental populations with a simple genetic relatedness structure (Boer *et al.*, 2007; Malosetti *et al.*, 2008; Alimi *et al.*, 2013), but currently available software could not perform REML estimation for the HapMap population. The kinship matrix was therefore replaced with a compressed kinship matrix (Bradbury *et al.*, 2007; Zhang *et al.*, 2010), modeling the genetic relatedness between a number of internally homogeneous groups. Assuming there are m such groups, containing $n_1, \dots, n_m$ accessions each, the original kinship matrix K is replaced by ZK_CZ^t , where K_C is the kinship matrix for the groups, and Z is the $n \times m$ incidence matrix assigning each of the n accessions to one of the m groups. The groups were created by a procedure that

restricted the marker data to be linear combinations of environmental covariates representing the conditions at the place of origin of the accessions, as explained later.

Compressed kinship was calculated as the average kinship within genetic groups. Genotypes were assigned to k genetic groups by performing Ward clustering based on the squared Euclidean distance along the first $k-1$ principal components calculated from a matrix of standardized SNP scores, followed by cutting the resulting dendrogram into k distinct clusters (van Heerwaarden *et al.*, 2012, 2013; Odong *et al.*, 2013).

The use of a compressed kinship matrix requires a choice of the degree of compression, as determined by the number of genetic groups over which the individual kinship is averaged. This choice needs to balance the gain in computational efficiency with model fit (Zhang *et al.*, 2010) and the ability of the compressed matrix to capture the correlation between genetic dissimilarity and phenotypic differences, which is ultimately the reason for including a kinship matrix in the association model. There are currently no standard methods to determine the optimum degree of compression, at least not when used in a multi-trait setting. We determined the appropriate degree of compression for each association model based on the model likelihood, convergence and correspondence between kinship and phenotypic and geographical similarity. The latter was quantified as the Frobenius norm of the difference between the complement of the compressed kinship matrix, expanded to a block matrix of full rank, and the Euclidean distance matrix of phenotypic traits or geographic coordinates. We considered a range of four to 100 groups. Correspondence with phenotypic and geographical dissimilarity increased steeply from four to c. 35 groups, after which correspondence with geographic distance increased more slowly and the correspondence with phenotypic distance showing a local decrease until 58 groups. Model likelihood was relatively stable above four groups, but convergence was erratic depending on the modeled contrasts. For each model the number of groups was therefore chosen to be the minimum number of groups needed to achieve a degree of correspondence approximating that found at 35 groups, under condition of model convergence.

Multi-trait GWAS Traits (columns of Y) were standardized. Along the genome, MTMMs of the type $Y = XB + G + E$ were fitted with initially for each marker trait-specific QTL effects $\beta_1, \dots, \beta_p$ (contained in B). To identify general QTLs with trait-specific effects, for individual markers, the null hypothesis $\beta_1 = \beta_2 = \dots \beta_p = 0$ was tested by a Wald test against the alternative hypothesis that at least one of the trait-specific effects was nonzero (Zhou & Stephens, 2014). To identify consistent QTLs, the null hypothesis $\beta_1 = \beta_2 = \dots \beta_p = \beta \neq 0$ was tested. To identify potentially adaptive QTLs, contrasts defined on the trait-specific QTL effects were tested. For example, suppose the first p_1 of the full set of p traits represents responses measured under abiotic stresses, while the second p_2 traits represent responses under biotic stresses. A contrast can now be defined to test the hypothesis of whether the QTL effect for abiotic stresses differs from that for biotic stresses: $\beta_1 = \beta_2 = \dots \beta_{p_1} = \alpha_{\text{abiotic}}$; $\beta_{p_1+1} = \beta_{p_1+2} = \dots \beta_p = \alpha_{\text{biotic}}$ and $H_0: \alpha_{\text{abiotic}} = \alpha_{\text{biotic}}$ vs $H_A:$

$\alpha_{\text{abiotic}} \neq \alpha_{\text{biotic}}$. For the Wald test for the hypothesis $\beta_1 = \dots = \beta_p$ we first fit the MTMM $Y = XB + G + E$ with XB only containing trait-specific means $\mu_1, \dots, \mu_p$, and then we test hypotheses on the marker effects. The contrast is defined through a partitioning of the traits in two groups (e.g. resistance against biotic or abiotic stress). Using the R package ASREML (Butler *et al.*, 2009) we perform Wald tests for the following hypotheses:

- 1 $H_0: \beta = 0$, in the constrained model $\beta_1 = \dots = \beta_p = \beta$.
- 2 $H_0: \alpha_1 = \alpha_2$, in the constrained model where α_1 is the effect on all traits in the first group, and α_2 is the effect on traits in the second group.

Simulations to compare power for full MTMM, contrast MTMM and univariate analysis We further compared the different Wald tests using simulations, described in more detail in Methods S12. Specifically, we compared the performance of the general MTMM (i.e. testing the hypothesis $\beta_1 = \beta_2 = \dots = \beta_p = 0$) with the MTMM used for the contrasts (i.e. $H_0: \alpha_{\text{group1}} = \alpha_{\text{group2}}$, where, within two predefined groups of traits, all SNP effects equal α_{group1} and α_{group2} , respectively). We simulated phenotypic data for given genotypic data, either assuming the SNP effects were positive (but not equal) within one group of traits and negative for the other (scenario A), or choosing the sign of each SNP effect randomly (scenario B). The simulation results as presented in Fig. S11 (see later) clearly indicate that the Wald test for the contrast has superior power under scenario A, while the general MTMM performs best under scenario B. In both cases, univariate analysis of the trait with the highest heritability is outperformed by at least one of the MTMM analyses. As a consequence, univariate GWAS and GWAS with the general and contrast MTMM give different rankings of SNPs.

Selecting candidate genes

A significance threshold of $P < 0.0001$ was chosen after implementation of genomic control (see below in the section 'Correction for genomic inflation'). For MTMM this resulted in 43 SNPs meeting this criterion. Such a threshold of 0.0001 is not uncommon in studies involving single-trait GWAS (e.g. El-Soda *et al.*, 2015; van Rooijen *et al.*, 2015; Kooke *et al.*, 2016). Given the total number of SNPs analyzed (i.e. 199 589 SNPs having a minor allele frequency > 0.05) and under the null hypothesis of no QTLs and independence of the markers, we arrive at a naive estimate for the expected number of false positives of $c. 20$, which is considerably smaller than the 43 SNPs with $P < 0.0001$ recorded in the full MTMM, suggesting that about half of the significant SNPs must be true positives. Furthermore, following the procedure described by Benjamini & Hochberg (1995), we estimated the false discovery rate to be 0.45, a number very comparable to our naive estimate earlier. SNPs within a 20 kb region were considered to be part of one linkage disequilibrium (LD) block. This resulted in 30 genomic regions. For presentation purposes, each LD block was represented in figures and heat maps by the SNP with the strongest (absolute) effect, on average, across all traits. For the GWA contrast analyses, the same procedure was followed to define LD blocks and representative SNPs.

Correcting for genomic inflation

The Wald test is known to suffer from some inflation (Zhou & Stephens, 2014), which we correct for using genomic control (Devlin & Roeder, 1999; Devlin *et al.*, 2001), which divides the observed test statistics $T_1, \dots, T_p$ by the genomic inflation factor. For both the unconstrained MTMM and the MTMM for contrasts described earlier, we observed inflation for small as well as large P -values (i.e. also more P -values close to 1 than expected). Consequently, the usual genomic control procedures based on the observed vs expected median of test statistics gave overly optimistic inflation factors. We therefore applied an alternative genomic control procedure, in which we regress the observed $-\log_{10}(P)$ values on the expected ones, and correct the observed $-\log_{10}(P)$ values for the slope. The genomic inflation factor was 1.24 for the full MTMM, with similar values for the other MTMM analyses (between 1.07 and 1.38). For the full MTMM without correction for population structure (i.e. taking the kinship to be the identity matrix), the inflation factor was 2.36.

Results

The phenotypic response of a population of 350 *Arabidopsis* accessions to an extensive set of stress-inducing conditions was quantified *relative to* the respective control treatments. Correcting for the respective control means that in the residual signal for a trait, effects of earliness, flowering time, general robustness, vigor, and so on, have been removed already. Therefore, the traits as analyzed represent a kind of stress *per se* response from which all kinds of disturbances have already been eliminated. Thirty traits, including, for example, root length, number of damaged leaves or number of pathogen-inflicted spreading lesions (Table 1), were quantified when the plants were exposed to 15 different stresses, that is, four abiotic stresses (drought, salt stress, osmotic stress and heat), seven biotic stresses (parasitic plant, phloem-feeding aphid, phloem-feeding whitefly, cell-content feeding thrips, leaf-chewing caterpillar, root-feeding nematode and necrotrophic fungus) and four stress combinations (fungus and caterpillar, drought and fungus, drought and caterpillar, caterpillar and osmotic stress). For detailed information on the carefully standardized stress treatments, the trait definitions and phenotyping, see Supporting Information Methods S1–S10.

Heritability of responses to biotic and abiotic stresses

The phenotypic analysis resulted in a wide range of marker-based, narrow-sense heritability (Kruijer *et al.*, 2015) estimates with 15 traits of low ($h^2 < 0.2$), 10 of moderate heritability ($0.2 < h^2 < 0.5$) and five of high ($h^2 > 0.5$) heritability (Fig. S1). The number of abiotic stress traits per heritability category was similar, while the number of traits related to biotic and combined stresses decreased with increasing heritability class. The most heritable traits were responses to feeding damage by thrips (Thrips_1; $h^2 = 0.8$), nematodes ($h^2 = 0.7$) and responses to salt (Salt_1 and Salt_3; resp. $h^2 = 0.6$ and $h^2 = 0.7$) and heat (Heat; $h^2 = 0.6$) (Table S1). The traits related to combined stresses have

predominantly low heritabilities; however, it should be emphasized that the combined stresses particularly relate to combinations involving fungal and caterpillar stress.

Genetic commonality underlying responses to different stresses


To analyze the phenotypic variation between *Arabidopsis thaliana* accessions as a function of molecular marker variation, we used various mixed-model approaches (see the Materials and Methods section). We estimated marker-based genetic correlations, that is, correlations based on the genome-wide commonality of SNP effects underlying pairs of traits (see the Materials and Methods section), to investigate the magnitude of genetic commonality underlying resistance mechanisms in response to a range of biotic and abiotic stresses. For brevity, we will refer to these marker-based genetic correlations as genetic correlations. Such genetic correlations can be interpreted as upper bounds to the joint determination of pairs of traits by genetic factors. Genetic correlation analysis revealed a strong connection between the responses to parasitic plants and to aphids ($r=0.8$), which were both negatively associated with other stress responses (Fig. 1). Parasitic plants and aphids have in common that they target phloem and xylem tissue (Tjallingii & Hogen Esch, 1993; Dorr & Kollmann, 1995), and induce the SA phytohormonal pathway (De Vos *et al.*, 2005; Runyon *et al.*, 2008). By contrast, the biotic stress responses that were negatively associated with the responses to parasitic plants and aphids, that is, responses to necrotrophic fungi, caterpillars, and thrips, represent JA-inducing stresses (De Vos *et al.*, 2005; Pieterse *et al.*, 2009, 2012). Because the SA and JA pathways predominantly interact through negative crosstalk (Pieterse *et al.*, 2009), the two main clusters resulting from the genetic correlation analysis represent different phytohormonal signaling response mechanisms. We also observed a strong

genetic correlation between plant responses to osmotic stress and root-feeding nematodes. This supports the notion that root-knot nematodes trigger a differentiation of root cells to multinucleate giant cells with severely altered water potential and osmotic pressure (Baldacci-Cresp *et al.*, 2015). While the correlations between traits at the phenotypic level were generally rather low, the genetic correlation analysis revealed a common genetic basis underlying the responses to sets of single and combined stresses (Fig. S2).

Candidate genes underlying responses to stresses

To identify individual candidate genes that contributed most to the pattern of genetic correlations, we fitted multi-trait QTL mixed models (MTMMs) to the total set of 30 traits, using a 214 000 SNP set that is commonly used for GWAS in *Arabidopsis* (Kim *et al.*, 2007; Atwell *et al.*, 2010; Li *et al.*, 2010; Horton *et al.*, 2012; Bac-Molenaar *et al.*, 2015). Our multi-trait GWA approach closely follows the modeling framework developed by Zhou & Stephens (2014) and generalizes the use of MTMMs as described previously (Boer *et al.*, 2007; Maloressi *et al.*, 2008; Alimi *et al.*, 2013) for classical biparental offspring populations to association panels. This GWA analysis identified 30 chromosome regions with multiple, significant SNP–trait associations. From each of those regions, the significant SNP with the strongest effect was chosen to represent the locus (Fig. 2; Table S2). Clustering of stresses by estimated SNP-effect profiles (Fig. 2) indicates that multiple SNPs were associated with response to more than one stress. Stress combinations induced large QTL allele substitution effects in the MTMM mapping (Fig. 2; Table S2), indicating that combinations of stresses trigger broad-spectrum defensive mechanisms. A total of 125 genes were in LD with the 30 most significant SNPs from the GWA analysis. Twenty of these genes were stress-related according to gene

Fig. 1 Mean genetic correlations between responses of *Arabidopsis thaliana* to abiotic (red) and biotic (dark blue) plant stresses. The thickness of lines represents the strength of mean genome-wide correlations, annotated with r -values (orange, positive; blue, negative correlation). The more shared genetic associations between stresses, the higher the absolute genetic correlation. Correlations are negative when alleles have opposite effects, that is, resulting in increased resistance to one stress, but decreased resistance to the other stress. Values in balloons represent mean within-group correlation (not shown for groups consisting of a single trait). Mean between-group correlations are not shown if they are below an absolute value of $r=0.2$. Two clusters can be distinguished: parasitic plants and aphids; and the other stresses, except whiteflies.






Fig. 2 Multi-trait mixed-model (MTMM) genome-wide association (GWA) mapping with 30 different stress responses of *Arabidopsis thaliana*. The top panel shows the 214 000 single nucleotide polymorphisms (SNPs) with their corresponding $-\log_{10}(P)$ values for the five chromosomes. The lower panel depicts the trait-specific effect sizes of the rare alleles for significant SNPs ($P < 0.0001$) as estimated by the full MTMM. When several SNPs were located within the 20 kb linkage disequilibrium half-windows around the most significant SNP in a region, the effects for the SNP with the strongest absolute average effects are shown (red-flagged in the Manhattan plot). SNPs are named by chromosome number and position on the chromosome. Negative effect sizes (blue) correspond to reduced plant resistance as a result of the rare allele, and positive effect sizes (yellow) correspond to increased resistance as a result of the rare allele. Stress responses were clustered hierarchically according to their effect, using Ward's minimum variance method. The key shows the frequency distribution for the effect sizes of the SNPs.

ontology annotation data (Table S3). Of these 20 genes, six have been functionally characterized by at least one study (Table 2a). For these six genes, we explored expression data to evaluate the biological relevance of these genes in stress-responsive mechanisms of *Arabidopsis* (Fig. S3). Of special interest were SNPs chr5.7493620, chr5.22041081 and chr4.6805259, which were in LD with *WRKY38* (encoding a WRKY transcription factor involved in SA-dependent disease resistance) (Kim *et al.*, 2008), *AtCNGC4* (involved in pathogen resistance) (Chin *et al.*, 2013) and *RMG1* (coding for disease resistance protein) (Yu *et al.*, 2013), respectively.

Phytohormonal signaling underlying contrasts in stress responses

The MTMM framework allowed constraints to be imposed on the values of the estimated QTL effects, thereby providing a powerful testing framework for QTLs that have a common effect for the stresses belonging to one particular group of stresses, as contrasted with the effect for another group of stresses (see the Materials and Methods section 'Multi-trait GWAS'). We investigated whether polymorphisms for genes involved in SA and JA biosynthesis or genes responsive to signals from these pathways

were the cause of the negative genetic correlations between the groups of traits sharing one or the other phytohormonal signaling pathway. To this end, we performed multi-trait GWA mapping to test the contrast between parasitic plant and aphid response vs the most negatively correlated traits, that is, fungus, caterpillar, thrips and drought response (Fig. 1). Fifteen SNPs were significantly associated with contrasting effects between the two trait clusters (Fig. S4). Seven of these SNPs were in LD with one or more genes known to be involved in JA-, SA- or resistance-related signal transduction (Table S4). Among these genes are *LOX5*, whose product is involved in facilitating aphid feeding (Nalam *et al.*, 2012a,b), *MYB107* encoding a transcription factor responsive to SA (Stracke *et al.*, 2001; Chen *et al.*, 2006), the JA-inducible genes *TPS02* and *TPS03* encoding terpene synthases (Huang *et al.*, 2010), and *MES16*, encoding a methyl jasmonate esterase (Christ *et al.*, 2012). Using TAIR10 annotations, we found that in total there are 371 genes that have an annotation related to JA and SA signaling (JA-SA genes). Our GWA analysis identified significant SNPs inside or in a 20 kb neighborhood of five of those. In the remainder of the genome (i.e. non JA-SA), we identified 162 genes close to or with significant SNPs. So, in candidate regions for JA-SA, we had a ratio of $5/371 = 1.35\%$ significant genes, while in noncandidate regions, we found 162/

Table 2 Candidate *Arabidopsis thaliana* genes resulting from (a) multi-trait mixed-model (MTMM) analysis of all 30 stress responses as presented in Fig. 2; and (b) contrast-specific analysis with MTMM for contrasting effects of biotic and abiotic stresses as presented in Fig. 3

Marker*	Gene in LD	Gene name	Gene description [†]	Responsiveness	References
(a)					
chr2.11659416	AT2G27250	CLV3	One of the three <i>CLAVATA</i> genes controlling the size of the shoot apical meristem (SAM) in <i>Arabidopsis</i>	Unknown	Clark <i>et al.</i> (1996); Fletcher <i>et al.</i> (1999); Shinohara & Matsubayashi (2010)
chr3.19804402	AT3G53420	PIP2	A member of the plasma membrane intrinsic protein subfamily PIP2	Heat, salt and heat, heat and silwet	Martiniere <i>et al.</i> (2012); Peret <i>et al.</i> (2012); Rasmussen <i>et al.</i> (2013); Sanchez-Romera <i>et al.</i> (2014)
chr4.6805259	AT4G11170	RMG1	Encodes RMG1 (Resistance Methylated Gene 1), an NB-LRR disease resistance protein with a Toll/interleukin-1 receptor (TIR) domain at its N terminus	Flagellin	Yu <i>et al.</i> (2013)
chr5.7493620	AT5G22570	WRKY38	Member of WRKY Transcription Factor; Group III	SA, <i>Pseudomonas</i>	Mare <i>et al.</i> (2004); Kim <i>et al.</i> (2008)
chr5.22041081	AT5G54250	CNGC4	Member of cyclic nucleotide gated channel family, a downstream component of the signaling pathways leading to hypersensitive response (HR) resistance. Mutant plants exhibit gene-for-gene disease resistance against avirulent <i>Pseudomonas syringae</i> despite the near-complete absence of the HR. Salicylic acid accumulation in <i>dnd2</i> mutants is completely <i>PAD4</i> -independent	Cold, flagellin	Jurkowski <i>et al.</i> (2004); Keisa <i>et al.</i> (2011); Chin <i>et al.</i> (2013); Rasmussen <i>et al.</i> (2013)
chr5.23302987	AT5G57560	TCH4	Encodes a cell wall modifying enzyme, rapidly up-regulated in response to environmental stimuli	Heat, heat and silwet, heat and salt, heat and high light, high light, high light and cold, high light and salt	Braam & Davis (1990); Xu <i>et al.</i> (1996); Purugganan <i>et al.</i> (1997); Iliev <i>et al.</i> (2002); Rasmussen <i>et al.</i> (2013)
(b)					
chr1.30381439	AT1G80820	CCR2	<i>CINNAMOYL COA REDUCTASE</i> . Encodes a cinnamoyl CoA reductase isoform. Involved in lignin biosynthesis	Cold and flagellin and silwet	Luderitz & Grisebach (1981); Lauvergeat <i>et al.</i> (2001); Zhou <i>et al.</i> (2010); Rasmussen <i>et al.</i> (2013)
chr1.30381439	AT1G80840	WRKY40	Pathogen-induced transcription factor. Binds W-box sequences <i>in vitro</i> . Forms protein complexes with itself and with WRKY60. Coexpression with <i>WRKY18</i> or <i>WRKY60</i> made plants more susceptible to both <i>P. syringae</i> and <i>Botrytis</i>	Cold and flagellin and silwet	Chen <i>et al.</i> (2010a); Pandey <i>et al.</i> (2010); Liu <i>et al.</i> (2012); Rasmussen <i>et al.</i> (2013)
chr1.6038270	AT1G17610	CHS1	<i>CHILLING SENSITIVE 1</i> , mutant accumulates steryl-esters at low temperature	Cold and high light	Rasmussen <i>et al.</i> (2013); Wang <i>et al.</i> (2013); Zbierzak <i>et al.</i> (2013)
chr5.171177	AT5G17640	ASG1	<i>ABIOTIC STRESS GENE 1</i> ; expression of this gene is induced by ABA and salt stress	ABA, salt	Coste <i>et al.</i> (2008); Batelli <i>et al.</i> (2012)
chr5.23247572	AT5G57380	VIN3	Encodes a plant homeodomain protein <i>VERNALIZATION INSENSITIVE 3 (VIN3)</i> . <i>In planta</i> VIN3 and VRN2, <i>VERNALIZATION 2</i> , are part of a large protein complex that can include the polycomb group (PcG) proteins <i>FERTILIZATION INDEPENDENT ENDOSPERM (FIE)</i> , <i>CURLY LEAF (CLF)</i> , and <i>SWINGER (SWN or EZA1)</i> . The complex has a role in establishing <i>FLC (FLOWERING LOCUS C)</i> repression during vernalization	Cold	Sung <i>et al.</i> (2007); Bond <i>et al.</i> (2009); Finnegan <i>et al.</i> (2011)

Table 2 (Continued)

Marker*	Gene in LD	Gene name	Gene description†	Responsiveness	References
chr5.23293119	AT5G57560	TCH 4	Encodes a cell wall-modifying enzyme	Heat, heat and silwet, heat and salt, heat and high light, high light, high light and cold, high light and salt	Braam & Davis (1990); Xu <i>et al.</i> (1996); Purugganan <i>et al.</i> (1997); Iliev <i>et al.</i> (2002); Rasmussen <i>et al.</i> (2013)
chr5.23293870	AT5G57490	VDAC4	Encodes a voltage-dependent anion channel (VDAC: AT3G01280/VDAC1)	<i>Pseudomonas</i>	Lee <i>et al.</i> (2009); Tateda <i>et al.</i> (2011)
chr5.23366252	AT5G57685	GDU3	Encodes a member of the GDU (glutamine dumper) family proteins involved in amino acid export: At4g31730 (GDU1)	Unknown	Chen <i>et al.</i> (2010b)

NB-LRR, nucleotide binding site–leucine-rich repeat.

*Markers derived from MTMM analysis (see Fig. 2).

†Based on information on <http://www.arabidopsis.org/tools/bulk/go/index.jsp>.

27863 = 0.58%. This is an enrichment of 2.33 times, significant at $\alpha = 0.05$ (Fisher's exact probability test, mid- P value < 0.046; Rivals *et al.*, 2007). Following Atwell *et al.* (2010), an upper bound for the false discovery rate is then $1/2.33 = 0.43$.

In addition to screening for SNPs with contrasting effects, we screened for SNPs with a similar effect across the earlier-mentioned trait clusters (Fig. S5) and found candidate genes involved in oxidative stress and plant responses to salinity and pathogens (Table S5).

QTLs underlying contrasts in responses to biotic and abiotic stresses


We expected a negative correlation between the responses to abiotic and biotic stresses as a result of antagonistic interactions between ABA and the SA and JA-ET pathways (Anderson *et al.*, 2004; Fujita *et al.*, 2006; De Torres Zabala *et al.*, 2009; Kissoudis *et al.*, 2015). Testing for this contrast within the GWA analysis using our MTMM approach significantly identified 43 SNPs with a QTL effect that changed sign between biotic and abiotic conditions. For presentation purposes, traits were grouped by a cluster analysis across SNPs, while SNPs were grouped by clustering across traits. Fig. 3 shows the SNPs with the strongest overall effects, identified in 18 LD intervals. The minor alleles of nine of these SNPs displayed a positive effect on biotic stress response traits and a negative effect on abiotic response traits. The remaining nine SNPs displayed the opposite effect (Fig. 3). Several candidate genes were identified in LD with the SNPs that are specific for plant responses to either abiotic or biotic stresses (Table 2b), such as *TCH4* (encoding a cell wall-modifying enzyme), *AtCCR2* (involvement in lignin biosynthesis) and *ASG1* (a gene induced by ABA and salt stress). Transcription data (Fig. S6) support the notion that these genes play a contrasting role in responses to abiotic and biotic stresses and reveal an antagonistic responsiveness between ABA and JA treatment (*TCH4*) or a specific responsiveness to either ABA (*AtCCR2*, *ASG1*, *ATVDAC4*) or JA (*ATWRKY40*). This is in line with the hypothesis that there are antagonistic effects between abiotic stress

responses, predominantly involving the ABA pathway, and wound and biotic stress responses involving the JA-ET or SA pathways (Kissoudis *et al.*, 2015). Previous studies have, however, also revealed an overlap in abiotic and biotic plant responses, such as similar transcriptomic perturbations after salinity and pathogen stress (Ma *et al.*, 2006). A screen for QTLs with similar effects on resistance to biotic and abiotic stress (Fig. S7) identified three genes annotated to be responsive to stress stimuli (Table S6). Transcriptional data show that these genes respond differentially to different (a)biotic stresses and phytohormones (Fig. S8). *ARGAH2*, encoding an arginase enzyme with a role in the metabolism of polyamines and nitric oxide, is involved in both SA- and JA-mediated resistance to both biotrophic and necrotrophic pathogens, and is also responsive to abiotic stimuli such as temperature, salt and light intensity (Fig. S8) (Jubault *et al.*, 2008; Grivot *et al.*, 2012; Rasmussen *et al.*, 2013). *PKS1* is known to be involved in adaptation in plant growth in response to light (Fankhauser *et al.*, 1999; Molas & Kiss, 2008), but also seems to be responsive to *Botrytis* (Fig. S8). These genes are promising candidates for consistent effects across biotic and abiotic stresses.

QTLs underlying contrasts in responses to below- and above-ground stresses

We expected a negative correlation between responses to below- and above-ground stresses. A strong QTL signal was found on chromosome 1 for this contrasting response (Fig. S9). The associated marker (chr1.13729757) had 12 genes in LD with it, of which 11 are annotated as pseudogenes. Transcriptional data on abiotic stresses for the only protein coding gene (*AT1G36510*) show an up-regulation in above-ground tissues, yet a down-regulation in the root tissues (Winter *et al.*, 2007). Marker chr5.16012837 showed the strongest signal for similar effects on responses to below- and above-ground stresses (Fig. S10) for which the *pathogenesis-related thaumatin superfamily protein* (*AT5G40020*) is the most promising candidate gene.

Fig. 3 Genetic associations specific for contrasting responses of *Arabidopsis thaliana* to abiotic and biotic stresses. Genetic associations (in red) were estimated with a contrast-specific genome-wide association analysis using a multi-trait mixed model (MTMM). For exploratory purposes, significant single nucleotide polymorphisms (SNPs) ($P \leq 10^{-4}$) for the biotic–abiotic contrast were clustered on their trait-specific effect sizes as estimated in the full MTMM, that is, without imposing a contrast restriction on the SNP effects. If there was another SNP in LD that had a higher effect size, this SNP was used as a representative of the LD block. Negative effects (blue) were cases where the rare allele was associated with a detrimental effect on the plants, while positive effects (yellow) were cases where the rare allele was associated with increased resistance to the stress. The rare alleles of the top nine SNPs are associated with enhanced resistance to abiotic stresses and reduced resistance to biotic stresses; the bottom nine SNPs show the inverse. Stresses were clustered on the basis of SNP effects using Ward's minimum variance method. The key shows the frequency distribution of SNPs across effect sizes.



Validation of identified QTLs

To obtain experimental support for the most interesting QTLs resulting from the MTMM, we tested homozygous T-DNA insertion lines for candidate genes *RMG1* and *WRKY38* (both resulting from the MTMM analysis), and *TCH4* (from MTMM analysis on biotic vs abiotic contrast) for several of the stresses addressed in this study. Two independent *rmg1* T-DNA insertion lines showed a phenotype that was different from the wild-type (Col-0) for some of the stress conditions (Fig. 4; Methods S11), being more resistant to caterpillar feeding and osmotic stress (Fig. 4). *RMG1* (AT4G11170) encodes a nucleotide binding site–leucine-rich repeat disease resistance protein, which acts as a pattern-recognition receptor that recognizes evolutionarily conserved pathogen-derived signatures, and transcription is induced by the bacterial peptide flg22 (Yu *et al.*, 2013). The rare allele of the corresponding marker chr4.6805259 is associated with enhanced resistance to salt stress and the combined stresses ‘caterpillar and drought’ and ‘caterpillar and fungus’ and with enhanced susceptibility to drought stress. Gene expression data show that *RMG1* is up-regulated by several abiotic and biotic stresses (Fig. 4). In addition, gene ontology enrichment analysis of the coexpression network of *RMG1* shows an overrepresentation of genes involved in immune responses and maintenance of ion homeostasis. The latter is based upon coexpression with five

genes encoding glutamate receptors (*GLR1.2*, *GLR1.3*, *GLR2.5*, *GLR2.8*, and *GLR2.9*), putatively involved in ion-influx-mediated long-distance signaling of wound, pathogen and salt stress (Ma *et al.*, 2006; Mousavi *et al.*, 2013; Choi *et al.*, 2014; Kissoudis *et al.*, 2015). T-DNA insertion lines for *TCH4* and *WRKY38* did not show a phenotype different from the wild-type (Col-0) for any of the tested stress conditions. Whether this is dependent on the genetic background used remains to be investigated.

Summarizing, our multi-trait GWA methodology facilitated a detailed analysis of the genetic architecture of resistance in *Arabidopsis* to a wide diversity of biotic and abiotic stresses. Application of this methodology revealed novel candidate genes associated with multiple stress responses, where specific contrasts were identified with some genes positively associated with the resistance to one set of stresses while being negatively associated with another set of stresses. In plant breeding (Brady *et al.*, 2005; Ballesteros *et al.*, 2015), such genes are classified as adaptive. Alternatively, other genes were identified with consistent effects across a wide spectrum of stress conditions. Such genes are labeled as constitutive in the plant breeding literature (Brady *et al.*, 2005; Ballesteros *et al.*, 2015). Both adaptive and constitutive QTLs are important factors to contribute to improved stress resistance and tolerance in commercial crop species (Brady *et al.*, 2005; Ballesteros *et al.*, 2015).




Fig. 4 Phenotypes of *RMG1* T-DNA mutant screenings for *Arabidopsis thaliana*. Phenotypes are given for two T-DNA lines in the *RMG1* gene and for Col-0 as control. (a) Number of thrips feeding spots on a detached leaf at 6 d postinfestation ($n = 24$). (b) Leaf area consumed by *Pieris rapae* caterpillars ($n = 6$). (c) Number of nematode egg masses ($n = 23$). (d) Number of *Myzus persicae* aphid offspring ($n = 10$ –17). (e) Percentage survival of adult whiteflies (*Aleyrodes proletella*) ($n = 10$). (f) Plant FW after osmotic treatment in comparison with control (% relative to control) ($n = 4$). (g) Plant DW after 75 mM salt treatment in comparison with control (ratio) ($n = 7$ –10; mean $\pm$ SE; *, $P < 0.05$; **, $P < 0.01$ (difference in comparison with Col-0)). (h, i) Relative expression fold-change for *RMG1* compared with untreated control plants in above-ground (h) and below-ground (i) tissue. Expression data from *Arabidopsis* eFP browser (<http://bbc.botany.utoronto.ca>).

Discussion

We developed a novel mixed-model approach to multi-trait GWA mapping with a special feature for testing contrasts between groups of stresses to identify the genetic architecture underlying a total of 30 stress response traits in *Arabidopsis*. The strength of our statistical approach was that our multi-trait mixed model accounted simultaneously for dependencies between genotypes and between traits, providing a natural and appropriate correction for multiple testing, while maximizing power for the detection of QTLs for the stress contrast under study. As we addressed a large number of stresses, our phenotyping experiments were distributed across a series of laboratories and were not performed simultaneously. To mitigate as much as possible the occurrence of QTLs induced purely by experiment-specific differences in plant management and environmental control, our phenotypic responses were defined in terms of control-corrected responses. This type of correction will emphasize QTLs for resistance and tolerance *per se* and will decrease detection power for QTLs related to development and viability.

The extensive phenotyping executed in this study was done under carefully controlled conditions in climate chambers. Ideally, phenotyping should be done in nature because that is where genetic variation is exposed to natural selection (Bergelson & Roux, 2010; Brachi *et al.*, 2010, 2013). Here, we have phenotyped the plant population to 15 different stresses under laboratory conditions and our data show an interesting pattern based on genetic correlations that matches with phytohormonal signalling underlying stress responses (Fig. 1). This indicates that the genetic architecture recorded here is biologically relevant. Drought and salt stress responses share signal transduction mechanisms (Zhu, 2002) which are represented by the genetic correlations recorded (Fig. 1). Insect damage is commonly associated with drought or osmotic stress and this is also clear from overlap in underlying phytohormonal signalling (Pieterse *et al.*, 2012). Fig. 1 shows that drought stress and osmotic stress correlate with insect stresses. Extending studies of genetic variation and the genetic architecture underlying responses to multiple stresses to natural conditions will be an important next step (Bergelson & Roux, 2010).

Through the approach developed here, candidate genes for stress responses were identified that are involved in contrasting responses when comparing biotic and abiotic stresses, above- and below-ground stresses, and attack by phloem feeders vs other biotic stresses. Among these genes many are involved in phytohormone-mediated processes, supporting the notion that the phytohormonal regulatory network plays an important role in plant stress responses (Pieterse *et al.*, 2012). The MTMM approach further showed that certain SNPs were associated with multiple stress responses and that transcriptional patterns of genes to which the SNPs were linked, as well as the phenotype expressed upon knocking out one of these genes, matched the observed stress responses of the plants. The *RMG1* gene that was identified through this procedure has relevant effects on plant phenotype in the context of responses to individual stresses. *RMG1* is a bacterium-inducible resistance gene whose activity is

modulated by the plant through RNA-directed DNA methylation (RdDM) (Yu *et al.*, 2013). *RMG1* expression activates the SA pathway (Yu *et al.*, 2013). Thus, the increased resistance against caterpillars in *rmg1* mutants may be the result of elimination of SA-mediated interference with JA-induced resistance to caterpillars (Pieterse *et al.*, 2012). *RMG1* appears to be inducible by several stresses and deserves further in-depth analysis for its role in plant response to multiple stresses. Our data show that for the 30 most significant SNPs resulting from the MTMM analysis, the average absolute effect size for double stresses is higher, on average, than that for single stresses ($P < 0.007$, Table S2). This suggests that resistance mechanisms involved in countering dual stresses are of a more general nature, in contrast to the rather specific resistance mechanisms involved in single stress responses. However, the combined stresses included in this study particularly involve fungal and caterpillar stresses. Future studies including other combined stresses are needed to further investigate the suggested pattern.

The MTMM framework that we used for GWA mapping provides unbiased estimates for QTL allele substitution effects together with correct standard errors for these effects. Within the same framework we developed unique facilities to test hypotheses on QTL $\times$ stress interactions in multi-trait models, which are not available in competing meta-analysis approaches (Zhu *et al.*, 2015). The variance-covariance structure that we used for the polygenic term protects against inflated type I error, that is, too many false-positive SNP–trait associations, as a consequence of population structure and kinship on the genotypic side and genetic correlations between traits on the trait side. The inclusion of trait correlations will, for most QTLs, improve the power of detection in comparison to single-trait GWA mapping (Korte *et al.*, 2012; Zhou & Stephens, 2014; see ‘Multi-trait GWAS’ in the Materials and Methods section). For a comparison of the MTMM analysis with single-trait analyses, see ‘Simulations to compare power for full MTMM, contrast MTMM and univariate analysis’ in the Materials and Methods section, Methods S12 and Figs S11 and S12. Our choice for the variance-covariance structure of the polygenic term as a Kronecker product of a compressed kinship on the genotypes with an approximated unstructured variance-covariance model on the environments is sometimes used in plant breeding for genomic prediction models (Burgueno *et al.*, 2012). However, implementation of such models in GWA mapping and especially on the scale that we present here, with 30 traits, is unprecedented and is practically far from straightforward. It required substantial work on preparatory phenotypic analyses as well as fine-tuning of the genotypic and trait variance-covariance structures to achieve convergence of the mixed models.

The MTMM analyses identified candidate genes associated with contrasting responses to biotic and abiotic stresses. Stress combinations appeared to have a strong influence on the MTMM outcome, indicative for significant interactions between different stresses when occurring simultaneously, and underlining the importance of studying the resistance of plants to combinations of stress. Transcriptional data and phenotyping of mutants provide initial support for the role of several of the candidate

genes identified. Studies of plant responses to a diverse set of biotic stresses show that the transcriptional pattern is stress-specific and that phytohormonal signaling pathways can explain up to 70% of the induced gene regulation (De Vos *et al.*, 2005). Taking the outcome of the MTMM analyses to investigate the involvement of identified candidate genes in the resistance of plants to several stresses, not only in *Arabidopsis* but also in related crop species, such as, for example, *Brassica* species, will be valuable in the breeding by design of future crops to protect them against combinations of stresses, including biotic and abiotic stresses. This will be of great value for next-generation crops.

Acknowledgements

This research is supported by the Perspective Programme 'Learning from Nature' of the Dutch Technology Foundation STW, which is part of the Netherlands Organisation for Scientific Research (NWO) and is partly funded by the Ministry of Economic Affairs.

Author contributions

M.P.M.T., N.H.D.O., K.J.K., S.C., P-P.H., J.A.B-M., C.B., J. Bucher, J.B-L., X.C., E.F.F., M.M.J., W.L., J.A.v.P., S.W. and G.L.W. phenotyped the plants; M.P.M.T., N.H.D.O., K.J.K., S.C., P-P.H., M.G.M.A., J.A.B-M., J. Bakker, H.J.B., J. Bucher, C.B., X.C., E.F.F., M.A.J., M.M.J., J.J.B.K., W.L., C.M.J.P., C.R-S., G.S., C.T., J.J.A.v.L., J.A.v.P., C.C.v.S., S.C.M.v.W., R.G.F.V., R.V., B.V., D.V., S.W., G.L.W. and M.D. designed the phenotyping experiments and made initial analyses; W.K., J.v.H. and F.A.v.E. developed the multi-trait mixed model; W.K., J.v.H., M.P.M.T., N.H.D.O., K.J.K., S.C., P-P.H., B.U., F.A.v.E. and M.D. analyzed the total dataset. M.D. and F.A.v.E. coordinated the study as a whole. M.P.M.T., N.H.D.O., K.J.K., S.C., P-P.H., W.K., J.v.H., F.A.v.E. and M.D. wrote the manuscript with input from J.J.B.K., C.J.M.P. and M.G.M.A. and all authors proofread the final version of the manuscript.

References

- Alimi NA, Bink MCAM, Dieleman JA, Magán JJ, Wubs AM, Palloix A, Van Eeuwijk FA. 2013. Multi-trait and multi-environment QTL analyses of yield and a set of physiological traits in pepper. *Theoretical and Applied Genetics* 126: 2597–2625.
- Alonso-Blanco C, Aarts MGM, Bentsink L, Keurentjes JJB, Reymond M, Vreugdenhil D, Koornneef M. 2009. What has natural variation taught us about plant development, physiology, and adaptation? *Plant Cell* 21: 1877–1896.
- Anderson JP, Badruzsaufari E, Schenk PM, Manners JM, Desmond OJ, Ehlert C, Maclean DJ, Ebert PR, Kazan K. 2004. Antagonistic interaction between abscisic acid and jasmonate-ethylene signaling pathways modulates defense gene expression and disease resistance in *Arabidopsis*. *Plant Cell* 16: 3460–3479.
- Atwell S, Huang YS, Vilhjalmsón BJ, Willems G, Horton M, Li Y, Meng D, Platt A, Tarone AM, Hu TT *et al.* 2010. Genome-wide association study of 107 phenotypes in *Arabidopsis thaliana* inbred lines. *Nature* 465: 627–631.
- Bac-Molenaar JA, Fradin EF, Becker FFM, Rienstra JA, van der Schoot J, Vreugdenhil D, Keurentjes JJB. 2015. Genome-wide association mapping of fertility reduction upon heat stress reveals developmental stage-specific QTLs in *Arabidopsis thaliana*. *Plant Cell* 27: 1857–1874.
- Baldacci-Cresp F, Maucourt M, Deborde C, Pierre O, Moing A, Brouquisse R, Favery B, Frendo P. 2015. Maturation of nematode-induced galls in *Medicago truncatula* is related to water status and primary metabolism modifications. *Plant Science* 232: 77–85.
- Ballesteros DC, Mason RE, Addison CK, Acuna MA, Arguello MN, Subramanian N, Miller RG, Sater H, Gbur EE, Miller D *et al.* 2015. Tolerance of wheat to vegetative stage soil waterlogging is conditioned by both constitutive and adaptive QTL. *Euphytica* 201: 329–343.
- Batelli G, Massarelli I, Van Oosten M, Nurchio R, Vannini C, Raimondi G, Leone A, Zhu JK, Maggio A, Grillo S. 2012. *Asg1* is a stress-inducible gene which increases stomatal resistance in salt stressed potato. *Journal of Plant Physiology* 169: 1849–1857.
- Baxter I, Brazelton JN, Yu DN, Huang YS, Lahner B, Yakubova E, Li Y, Bergelson J, Borevitz JO, Nordborg M *et al.* 2010. A coastal cline in sodium accumulation in *Arabidopsis thaliana* is driven by natural variation of the sodium transporter AtHKT1;1. *PLoS Genetics* 6: e1001193.
- Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)* 57: 289–300.
- Bergelson J, Roux F. 2010. Towards identifying genes underlying ecologically relevant traits in *Arabidopsis thaliana*. *Nature Reviews Genetics* 11: 867–879.
- Boer MP, Wright D, Feng L, Podlich DW, Luo L, Cooper M, Van Eeuwijk FA. 2007. A mixed-model quantitative trait loci (QTL) analysis for multiple-environment trial data using environmental covariables for QTL-by-environment interactions, with an example in maize. *Genetics* 177: 1801–1813.
- Bond DM, Wilson IW, Dennis ES, Pogson BJ, Jean Finnegan E. 2009. VERNALIZATION INSENSITIVE 3 (VIN3) is required for the response of *Arabidopsis thaliana* seedlings exposed to low oxygen conditions. *Plant Journal* 59: 576–587.
- Braam J, Davis RW. 1990. Rain-induced, wind-induced, and touch-induced expression of calmodulin and calmodulin-related genes in *Arabidopsis*. *Cell* 60: 357–364.
- Brachi B, Faure N, Bergelson J, Cuguen J, Roux F. 2013. Genome-wide association mapping of flowering time in *Arabidopsis thaliana* in nature: genetics for underlying components and reaction norms across two successive years. *Acta Botanica Gallica* 160: 205–218.
- Brachi B, Faure N, Horton M, Flahauw E, Vazquez A, Nordborg M, Bergelson J, Cuguen J, Roux F. 2010. Linkage and association mapping of *Arabidopsis thaliana* flowering time in nature. *PLoS Genetics* 6: 17.
- Brachi B, Meyer CG, Villoutreix R, Platt A, Morton TC, Roux F, Bergelson J. 2015. Coselected genes determine adaptive variation in herbivore resistance throughout the native range of *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* 112: 4032–4037.
- Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES. 2007. Tassel: software for association mapping of complex traits in diverse samples. *Bioinformatics* 23: 2633–2635.
- Brady KU, Kruckeberg AR, Bradshaw HD. 2005. Evolutionary ecology of plant adaptation to serpentine soils. *Annual Review of Ecology Evolution and Systematics* 36: 243–266.
- Braun H-J, Rajaram S, van Ginkel M. 1996. CIMMYT's approach to breeding for wide adaptation. *Euphytica* 92: 175–183.
- Burgueno J, de los Campos G, Weigel K, Crossa J. 2012. Genomic prediction of breeding values when modeling genotype x environment interaction using pedigree and dense molecular markers. *Crop Science* 52: 707–719.
- Butler DG, Cullis BR, Gilmour AR, Gogel BJ. 2009. *ASReml-R reference manual*. Brisbane, Qld, Australia: Queensland Department of Primary Industries and Fisheries.
- Chao DY, Silva A, Baxter I, Huang YS, Nordborg M, Danku J, Lahner B, Yakubova E, Salt DE. 2012. Genome-wide association studies identify heavy metal ATPase3 as the primary determinant of natural variation in leaf cadmium in *Arabidopsis thaliana*. *PLoS Genetics* 8: e1002923.
- Chen H, Lai Z, Shi J, Xiao Y, Chen Z, Xu X. 2010a. Roles of *Arabidopsis* WRKY18, WRKY40 and WRKY60 transcription factors in plant responses to abscisic acid and abiotic stress. *BMC Plant Biology* 10: 281.

- Chen H, Zhang Z, Teng K, Lai J, Zhang Y, Huang Y, Li Y, Liang L, Wang Y, Chu C *et al.* 2010b. Up-regulation of LSB1/GDU3 affects geminivirus infection by activating the salicylic acid pathway. *Plant Journal* 62: 12–23.
- Chen YH, Yang XY, He K, Liu MH, Li JG, Gao ZF, Lin ZQ, Zhang YF, Wang XX, Qiu XM *et al.* 2006. The MYB transcription factor superfamily of arabidopsis: expression analysis and phylogenetic comparison with the rice MYB family. *Plant Molecular Biology* 60: 107–124.
- Chin K, DeFalco TA, Moeder W, Yoshioka K. 2013. The Arabidopsis cyclic nucleotide-gated ion channels AtCNGC2 and AtCNGC4 work in the same signaling pathway to regulate pathogen defense and floral transition. *Plant Physiology* 163: 611–624.
- Choi WG, Toyota M, Kim SH, Hilleary R, Gilroy S. 2014. Salt stress-induced Ca^{2+} waves are associated with rapid, long-distance root-to-shoot signaling in plants. *Proceedings of the National Academy of Sciences, USA* 111: 6497–6502.
- Christ B, Schelbert S, Aubry S, Sussenbacher I, Muller T, Krautler B, Hortensteiner S. 2012. MES16, a member of the methyl esterase protein family, specifically demethylates fluorescent chlorophyll catabolites during chlorophyll breakdown in Arabidopsis. *Plant Physiology* 158: 628–641.
- Clark SE, Jacobsen SE, Levin JZ, Meyerowitz EM. 1996. The CLAVATA and SHOOT MERISTEMLESS loci competitively regulate meristem activity in Arabidopsis. *Development* 122: 1567–1575.
- Cooper M, Hammer GL. 1996. *Plant adaptation and crop improvement*. International Rice Research Institute. Wallingford, UK: CAB International.
- Cooper M, Messina CD, Podlich D, Totir LR, Baumgarten A, Hausmann NJ, Wright D, Graham G. 2014. Predicting the future of plant breeding: complementing empirical evaluation with genetic prediction. *Crop & Pasture Science* 65: 311–336.
- Coste AT, Ramsdale M, Ischer F, Sanglard D. 2008. Divergent functions of three *Candida albicans* zinc-cluster transcription factors (CTA4, ASG1 and CTF1) complementing pleiotropic drug resistance in *Saccharomyces cerevisiae*. *Micobiology* 154: 1491–1501.
- De los Campos G, Hickey JM, Pong-Wong R, Daetwyler HD, Calus MPL. 2013. Whole-genome regression and prediction methods applied to plant and animal breeding. *Genetics* 193: 327–345.
- De Torres Zabala M, Bennett MH, Truman WH, Grant MR. 2009. Antagonism between salicylic and abscisic acid reflects early host–pathogen conflict and moulds plant defence responses. *Plant Journal* 59: 375–386.
- De Vos M, Van Oosten VR, Van Poecke RMP, Van Pelt JA, Pozo MJ, Mueller MJ, Buchala AJ, Metraux JP, Van Loon LC, Dicke M *et al.* 2005. Signal signature and transcriptome changes of *Arabidopsis* during pathogen and insect attack. *Molecular Plant–Microbe Interactions* 18: 923–937.
- De Vos M, Van Zaanen W, Koornneef A, Korzelius JP, Dicke M, Van Loon LC, Pieterse CMJ. 2006. Herbivore-induced resistance against microbial pathogens in Arabidopsis. *Plant Physiology* 142: 352–363.
- Devlin B, Roeder K. 1999. Genomic control for association studies. *Biometrics* 55: 997–1004.
- Devlin B, Roeder K, Wasserman L. 2001. Genomic control, a new approach to genetic-based association studies. *Theoretical Population Biology* 60: 155–166.
- Dorr I, Kollmann R. 1995. Symplasmic sieve element continuity between *Orobanche* and its host. *Botanica Acta* 108: 47–55.
- van Eeuwijk FA, Bink MC, Chenu K, Chapman SC. 2010. Detection and use of QTL for complex traits in multiple environments. *Current Opinion in Plant Biology* 13: 193–205.
- El-Soda M, Kruijer W, Malosetti M, Koornneef M, Aarts MGM. 2015. Quantitative trait loci and candidate genes underlying genotype by environment interaction in the response of *Arabidopsis thaliana* to drought. *Plant, Cell & Environment* 38: 585–599.
- Fankhauser C, Yeh KC, Lagarias JC, Zhang H, Elich TD, Chory J. 1999. PKS1, a substrate phosphorylated by phytochrome that modulates light signaling in Arabidopsis. *Science* 284: 1539–1541.
- Finnegan EJ, Bond DM, Buzas DM, Goodrich J, Helliwell CA, Tamada Y, Yun JY, Amasino RM, Dennis ES. 2011. Polycomb proteins regulate the quantitative induction of VERNALIZATION INSENSITIVE 3 in response to low temperatures. *Plant Journal* 65: 382–391.
- Fiorani F, Schurr U. 2013. Future scenarios for plant phenotyping. *Annual Review of Plant Biology* 64: 267–291.
- Fletcher JC, Brand U, Running MP, Simon R, Meyerowitz EM. 1999. Signaling of cell fate decisions by CLAVATA3 in Arabidopsis shoot meristems. *Science* 283: 1911–1914.
- Fujita M, Fujita Y, Noutoshi Y, Takahashi F, Narusaka Y, Yamaguchi-Shinozaki K, Shinozaki K. 2006. Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signaling networks. *Current Opinion in Plant Biology* 9: 436–442.
- Granier C, Vile D. 2014. Phenotyping and beyond: modelling the relationships between traits. *Current Opinion in Plant Biology* 18: 96–102.
- Gravot A, Deleu C, Wagner G, Lariagon C, Lugan R, Todd C, Wendehenne D, Delourme R, Bouchereau A, Manzanera-Dauleux MJ. 2012. Arginase induction represses gall development during clubroot infection in Arabidopsis. *Plant and Cell Physiology* 53: 901–911.
- van Heerwaarden J, Hufford MB, Ross-Ibarra J. 2012. Historical genomics of North American maize. *Proceedings of the National Academy of Sciences, USA* 109: 12420–12425.
- van Heerwaarden J, Odong TL, van Eeuwijk FA. 2013. Maximizing genetic differentiation in core collections by PCA-based clustering of molecular marker data. *Theoretical and Applied Genetics* 126: 763–772.
- Holopainen JK, Gershenzon J. 2010. Multiple stress factors and the emission of plant VOCs. *Trends in Plant Science* 15: 176–184.
- Horton MW, Hancock AM, Huang YS, Toomajian C, Atwell S, Auton A, Mulyati NW, Platt A, Sperone FG, Vilhjalmsdottir BJ *et al.* 2012. Genome-wide patterns of genetic variation in worldwide *Arabidopsis thaliana* accessions from the RegMap panel. *Nature Genetics* 44: 212–216.
- Howe GA, Jander G. 2008. Plant immunity to insect herbivores. *Annual Review of Plant Biology* 59: 41–66.
- Huang MS, Abel C, Sohrabi R, Petri J, Haupt I, Cosimano J, Gershenzon J, Tholl D. 2010. Variation of herbivore-induced volatile terpenes among Arabidopsis ecotypes depends on allelic differences and subcellular targeting of two terpene synthases, TPS2 and TPS3. *Plant Physiology* 153: 1293–1310.
- Iliev EA, Xu W, Polisensky DH, Oh MH, Torisky RS, Clouse SD, Braam J. 2002. Transcriptional and posttranscriptional regulation of Arabidopsis *TCH4* expression by diverse stimuli. Roles of *cis* regions and brassinosteroids. *Plant Physiology* 130: 770–783.
- Jubault M, Hamon C, Gravot A, Lariagon C, Delourme R, Bouchereau A, Manzanera-Dauleux MJ. 2008. Differential regulation of root arginine catabolism and polyamine metabolism in clubroot-susceptible and partially resistant Arabidopsis genotypes. *Plant Physiology* 146: 2008–2019.
- Julkowska MM, Testerink C. 2015. Tuning plant signaling and growth to survive salt. *Trends in Plant Science* 20: 586–594.
- Jurkowski GI, Smith RK Jr, Yu IC, Ham JH, Sharma SB, Klessig DF, Fengler KA, Bent AF. 2004. Arabidopsis DND2, a second cyclic nucleotide-gated ion channel gene for which mutation causes the “defense, no death” phenotype. *Molecular Plant–Microbe Interactions* 17: 511–520.
- Keisa A, Kanberga-Silina K, Nakurte I, Kunga L, Rostoks N. 2011. Differential disease resistance response in the barley necrotic mutant nec1. *BMC Plant Biology* 11: 66.
- Kerwin R, Feusier J, Corwin J, Rubin M, Lin C, Muok A, Larson B, Li BH, Joseph B, Francisco M *et al.* 2015. Natural genetic variation in Arabidopsis thaliana defense metabolism genes modulates field fitness. *eLife* 4: e05604.
- Kim KC, Lai ZB, Fan BF, Chen ZX. 2008. Arabidopsis WRKY38 and WRKY62 transcription factors interact with histone deacetylase 19 in basal defense. *Plant Cell* 20: 2357–2371.
- Kim S, Plagnol V, Hu TT, Toomajian C, Clark RM, Ossowski S, Ecker JR, Weigel D, Nordborg M. 2007. Recombination and linkage disequilibrium in *Arabidopsis thaliana*. *Nature Genetics* 39: 1151–1155.
- Kissoudis C, Chowdhury R, van Heusden S, van de Wiel C, Finkers R, Visser RF, Bai Y, van der Linden G. 2015. Combined biotic and abiotic stress resistance in tomato. *Euphytica* 202: 317–332.
- Kissoudis C, van de Wiel C, Visser RGF, van der Linden G. 2014. Enhancing crop resilience to combined abiotic and biotic stress through the dissection of physiological and molecular crosstalk. *Frontiers in Plant Science* 5: 207.
- Kloth KJ, ten Broeke CJM, Thoen MPM, den Brink MHV, Wiegers GL, Krips OE, Noldus L, Dicke M, Jongsma MA. 2015. High-throughput phenotyping of plant resistance to aphids by automated video tracking. *Plant Methods* 11: 4.

- Kooke R, Kruijer W, Bours R, Becker F, Kuhn A, van de Geest H, Buntjer J, Doeswijk T, Guerra J, Bouwmeester H *et al.* 2016. Genome-wide association mapping and genomic prediction elucidate the genetic architecture of morphological traits in *Arabidopsis*. *Plant Physiology* 170: 2187–2203.
- Korte A, Vilhjalmsón BJ, Segura V, Platt A, Long Q, Nordborg M. 2012. A mixed-model approach for genome-wide association studies of correlated traits in structured populations. *Nature Genetics* 44: 1066–1071.
- Kruijer W, Boer MP, Malosetti M, Flood PJ, Engel B, Kooke R, Keurentjes JJB, van Eeuwijk FA. 2015. Marker-based estimation of heritability in immortal populations. *Genetics* 199: 379–393.
- Lauvergeat V, Lacomme C, Lacombe E, Lasserre E, Roby D, Grima-Pettenati J. 2001. Two cinnamoyl-CoA reductase (CCR) genes from *Arabidopsis thaliana* are differentially expressed during development and in response to infection with pathogenic bacteria. *Phytochemistry* 57: 1187–1195.
- Lee SM, Hoang MH, Han HJ, Kim HS, Lee K, Kim KE, Kim DH, Lee SY, Chung WS. 2009. Pathogen inducible voltage-dependent anion channel (AtVDAC) isoforms are localized to mitochondria membrane in *Arabidopsis*. *Molecules and Cells* 27: 321–327.
- Li Y, Huang Y, Bergelson J, Nordborg M, Borevitz JO. 2010. Association mapping of local climate-sensitive quantitative trait loci in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* 107: 21199–21204.
- Liu ZQ, Yan L, Wu Z, Mei C, Lu K, Yu YT, Liang S, Zhang XF, Wang XF, Zhang DP. 2012. Cooperation of three WRKY-domain transcription factors WRKY18, WRKY40, and WRKY60 in repressing two ABA-responsive genes ABI4 and ABI5 in *Arabidopsis*. *Journal of Experimental Botany* 63: 6371–6392.
- Luderitz T, Grisebach H. 1981. Enzymic synthesis of lignin precursors. Comparison of cinnamoyl-CoA reductase and cinnamyl alcohol:NADP⁺ dehydrogenase from spruce (*Picea abies* L.) and soybean (*Glycine max* L.). *European Journal of Biochemistry* 119: 115–124.
- Ma SS, Gong QQ, Bohnert HJ. 2006. Dissecting salt stress pathways. *Journal of Experimental Botany* 57: 1097–1107.
- Makumburage GB, Richbourg HL, LaTorre KD, Capps A, Chen CX, Stapleton AE. 2013. Genotype to phenotype maps: multiple input abiotic signals combine to produce growth effects via attenuating signaling interactions in maize. *G3 – Genes Genomes. Genetics* 3: 2195–2204.
- Malosetti M, Ribaut JM, van Eeuwijk FA. 2013. The statistical analysis of multi-environment data: modeling genotype-by-environment interaction and its genetic basis. *Frontiers in Physiology* 4: 17.
- Malosetti M, Ribaut JM, Vargas M, Crossa J, Van Eeuwijk FA. 2008. A multi-trait multi-environment QTL mixed model with an application to drought and nitrogen stress trials in maize (*Zea mays* L.). *Euphytica* 161: 241–257.
- Mare C, Mazzucotelli E, Crosatti C, Francia E, Stanca AM, Cattivelli L. 2004. Hv-WRKY38: a new transcription factor involved in cold- and drought-response in barley. *Plant Molecular Biology* 55: 399–416.
- Martiniere A, Li X, Runions J, Lin J, Maurel C, Luu DT. 2012. Salt stress triggers enhanced cycling of *Arabidopsis* root plasma-membrane aquaporins. *Plant Signaling & Behavior* 7: 529–532.
- Meyer K. 2009. Factor-analytic models for genotype × environment type problems and structured covariance matrices. *Genetics Selection Evolution* 41: 21.
- Mittler R, Blumwald E. 2010. Genetic engineering for modern agriculture: challenges and perspectives. *Annual Review of Plant Biology* 61: 443–462.
- Molas ML, Kiss JZ. 2008. PKS1 plays a role in red-light-based positive phototropism in roots. *Plant, Cell & Environment* 31: 842–849.
- Mousavi SAR, Chauvin A, Pascaud F, Kellenberger S, Farmer EE. 2013. GLUTAMATE RECEPTOR-LIKE genes mediate leaf-to-leaf wound signalling. *Nature* 500: 422–426.
- Nalam VJ, Keeretaweep J, Shah J. 2012b. The green peach aphid, *Myzus persicae*, acquires a LIPOXYGENASE5-derived oxylipin from *Arabidopsis thaliana*, which promotes colonization of the host plant. *Plant Signaling & Behavior* 8: e22735.
- Nalam VJ, Keeretaweep J, Sarowar S, Shah J. 2012a. Root-derived oxylipins promote green peach aphid performance on *Arabidopsis* foliage. *Plant Cell* 24: 1643–1653.
- Odong TL, van Heerwaarden J, van Hintum TJJ, van Eeuwijk FA, Jansen J. 2013. Improving hierarchical clustering of genotypic data via principal component analysis. *Crop Science* 53: 1546–1554.
- Ogura T, Busch W. 2015. From phenotypes to causal sequences: using genome wide association studies to dissect the sequence basis for variation of plant development. *Current Opinion in Plant Biology* 23: 98–108.
- Pandey SP, Roccaro M, Schon M, Logemann E, Somssich IE. 2010. Transcriptional reprogramming regulated by WRKY18 and WRKY40 facilitates powdery mildew infection of *Arabidopsis*. *Plant Journal* 64: 912–923.
- Peret B, Li G, Zhao J, Band LR, Voss U, Postaire O, Luu DT, Da Ines O, Casimiro I, Lucas M *et al.* 2012. Auxin regulates aquaporin function to facilitate lateral root emergence. *Nature Cell Biology* 14: 991–998.
- Piepho HP. 1997. Analyzing genotype-environment data by mixed models with multiplicative effects. *Biometrics* 53: 761–766.
- Pierik R, Testerink C. 2014. The art of being flexible: how to escape from shade, salt, and drought. *Plant Physiology* 166: 5–22.
- Pieterse CMJ, Leon-Reyes A, Van der Ent S, Van Wees SCM. 2009. Networking by small-molecule hormones in plant immunity. *Nature Chemical Biology* 5: 308–316.
- Pieterse CMJ, Van der Does D, Zamioudis C, Leon-Reyes A, Van Wees SCM. 2012. Hormonal modulation of plant immunity. *Annual Review of Cell and Developmental Biology* 28: 489–521.
- Platt A, Horton M, Huang YS, Li Y, Anastasio AE, Mulyati NW, Agren J, Bossdorf O, Byers D, Donohue K *et al.* 2010. The scale of population structure in *Arabidopsis thaliana*. *Plos Genetics* 6: e1000843.
- Prasch CM, Sonnewald U. 2013. Simultaneous application of heat, drought, and virus to *Arabidopsis* plants reveals significant shifts in signaling networks. *Plant Physiology* 162: 1849–1866.
- Purugganan MM, Braam J, Fry SC. 1997. The *Arabidopsis* TCH4 xyloglucan endotransglycosylase – substrate specificity, pH optimum, and cold tolerance. *Plant Physiology* 115: 181–190.
- Rasmussen S, Barah P, Suarez-Rodriguez MC, Bressendorff S, Friis P, Costantino P, Bones AM, Nielsen HB, Mundy J. 2013. Transcriptome responses to combinations of stresses in *Arabidopsis*. *Plant Physiology* 161: 1783–1794.
- Rivals I, Personnaz L, Taing L, Potier MC. 2007. Enrichment or depletion of a GO category within a class of genes: which test? *Bioinformatics* 23: 401–407.
- Rivero RM, Mestre TC, Mittler R, Rubio F, Garcia-Sanchez F, Martinez V. 2014. The combined effect of salinity and heat reveals a specific physiological, biochemical and molecular response in tomato plants. *Plant, Cell & Environment* 37: 1059–1073.
- Rizhsky L, Liang H, Shuman J, Shulaev V, Davletova S, Mittler R. 2004. When defense pathways collide. The response of *Arabidopsis* to a combination of drought and heat stress. *Plant Physiology* 134: 1683–1696.
- van Rooijen R, Aarts MGM, Harbinson J. 2015. Natural genetic variation for acclimation of photosynthetic light use efficiency to growth irradiance in *Arabidopsis*. *Plant Physiology* 167: 1412–1429.
- Runyon JB, Mescher MC, De Moraes CM. 2008. Parasitism by *Cuscuta pentagona* attenuates host plant defenses against insect herbivores. *Plant Physiology* 146: 987–995.
- Sanchez-Romera B, Ruiz-Lozano JM, Li G, Luu DT, Martinez-Ballesta Mdel C, Carvajal M, Zamarreno AM, Garcia-Mina JM, Maurel C, Aroca R. 2014. Enhancement of root hydraulic conductivity by methyl jasmonate and the role of calcium and abscisic acid in this process. *Plant, Cell & Environment* 37: 995–1008.
- Sewelam N, Oshima Y, Mitsuda N, Ohme-Takagi M. 2014. A step towards understanding plant responses to multiple environmental stresses: a genome-wide study. *Plant, Cell & Environment* 37: 2024–2035.
- Shinohara H, Matsubayashi Y. 2010. Arabinosylated glycopeptide hormones: new insights into CLAVATA3 structure. *Current Opinion in Plant Biology* 13: 515–519.
- Stam JM, Kroes A, Li YH, Gols R, van Loon JJA, Poelman EH, Dicke M. 2014. Plant interactions with multiple insect herbivores: from community to genes. *Annual Review of Plant Biology* 65: 689–713.
- Stracke R, Werber M, Weisshaar B. 2001. The R2R3-MYB gene family in *Arabidopsis thaliana*. *Current Opinion in Plant Biology* 4: 447–456.
- Sung S, Schmitz RJ, Amasino R. 2007. The role of VIN3-LIKE genes in environmentally induced epigenetic regulation of flowering. *Plant Signaling & Behavior* 2: 127–128.
- Suzuki N, Rivero RM, Shulaev V, Blumwald E, Mittler R. 2014. Abiotic and biotic stress combinations. *New Phytologist* 203: 32–43.

- Tateda C, Watanabe K, Kusano T, Takahashi Y. 2011. Molecular and genetic characterization of the gene family encoding the voltage-dependent anion channel in *Arabidopsis*. *Journal of Experimental Botany* 62: 4773–4785.
- Tjallingii WF, Hogen Esch T. 1993. Fine structure of aphid stylet routes in plant tissues in correlation with EPG signals. *Physiological Entomology* 18: 317–328.
- Vile D, Pervent M, Belluau M, Vasseur F, Bresson J, Muller B, Granier C, Simonneau T. 2012. *Arabidopsis* growth under prolonged high temperature and water deficit: independent or interactive effects? *Plant, Cell & Environment* 35: 702–718.
- Wang Y, Zhang Y, Wang Z, Zhang X, Yang S. 2013. A missense mutation in CHS1, a TIR-NB protein, induces chilling sensitivity in *Arabidopsis*. *Plant Journal* 75: 553–565.
- Wijnen CL, Keurentjes JJB. 2014. Genetic resources for quantitative trait analysis: novelty and efficiency in design from an *Arabidopsis* perspective. *Current Opinion in Plant Biology* 18: 103–109.
- Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, Provart NJ. 2007. An “electronic fluorescent pictograph” browser for exploring and analyzing large-scale biological data sets. *PLoS ONE* 2: 12.
- Xu W, Campbell P, Vargheese AK, Braam J. 1996. The *Arabidopsis* XET-related gene family: environmental and hormonal regulation of expression. *Plant Journal* 9: 879–889.
- Yu A, Lepere G, Jay F, Wang JY, Bapaume L, Wang Y, Abraham AL, Penterman J, Fischer RL, Voinnet O *et al.* 2013. Dynamics and biological relevance of DNA demethylation in *Arabidopsis* antibacterial defense. *Proceedings of the National Academy of Sciences, USA* 110: 2389–2394.
- Zbiezka AM, Porfirova S, Griebel T, Melzer M, Parker JE, Dormann P. 2013. A TIR-NBS protein encoded by *Arabidopsis Chilling Sensitive 1 (CHS1)* limits chloroplast damage and cell death at low temperature. *Plant Journal* 75: 539–552.
- Zhang Z, Ersoz E, Lai CQ, Todhunter RT, Tiwari HK, Gore MA, Bradbury PJ, Yu J, Arnett DK, Ordovas JM *et al.* 2010. Mixed linear model approach adapted for genome-wide association studies. *Nature Genetics* 42: 355–360.
- Zhou R, Jackson L, Shadle G, Nakashima J, Temple S, Chen F, Dixon RA. 2010. Distinct cinnamoyl CoA reductases involved in parallel routes to lignin in *Medicago truncatula*. *Proceedings of the National Academy of Sciences, USA* 107: 17803–17808.
- Zhou X, Stephens M. 2014. Efficient multivariate linear mixed model algorithms for genome-wide association studies. *Nature Methods* 11: 407–409.
- Zhu JK. 2002. Salt and drought stress signal transduction in plants. *Annual Review of Plant Biology* 53: 247–273.
- Zhu XF, Feng T, Tayo BO, Liang JJ, Young JH, Franceschini N, Smith JA, Yanek LR, Sun YV, Edwards TL *et al.* 2015. Meta-analysis of correlated traits via summary statistics from GWASs with an application in hypertension. *American Journal of Human Genetics* 96: 21–36.

Supporting Information

Additional Supporting Information may be found online in the Supporting Information tab for this article:

Fig. S1 Narrow-sense heritability for *Arabidopsis thaliana* resistance to abiotic and biotic stresses.

Fig. S2 Genetic and phenotypic correlation matrix.

Fig. S3 Expression data of six candidate genes (resulting from MTMM) in plants exposed to biotic or abiotic stress factors, relative to control conditions.

Fig. S4 Genetic associations specific for plant responses to the main clusters of the genetic correlation network (see Fig. 1): parasitic plant and aphid vs fungus, caterpillar, thrips and drought.

Fig. S5 Genetic associations common for plant response to the main clusters of the genetic correlation network: parasitic plant and aphid, on the one hand, vs fungus, caterpillar, thrips and drought on the other.

Fig. S6 Expression data of six candidate genes (resulting from MTMM analysis) in plants exposed to biotic or abiotic stress factors, relative to control conditions.

Fig. S7 Genetic associations common for plant responses to abiotic and biotic stresses.

Fig. S8 Expression data of three candidate genes (resulting from MTMM) in plants exposed to biotic or abiotic stress factors, relative to control conditions.

Fig. S9 Genetic associations specific for plant responses to either below- or above-ground stress.

Fig. S10 Genetic associations common for plant responses to below- and above-ground stresses.

Fig. S11 Power of MTMM in simulations.

Fig. S12 Comparison of SNPs identified by MTMM and univariate GWAS.

Table S1 Data overview on phenotyping the 350 *Arabidopsis thaliana* accessions of the HapMap collection

Table S2 Summed effect sizes of 30 most significant SNPs in MTMM per trait

Table S3 125 candidate genes derived from the Multitrait Mixed Model analysis

Table S4 Genes in linkage with SNPs with $-\log_{10}(P)$ score > 4 (20 kb half-window size) in the contrast-specific GWA mapping of parasitic plants and aphids, on the one hand, vs fungus, caterpillar, thrips and drought on the other

Table S5 Candidate genes in linkage with SNPs with $-\log_{10}(P)$ score > 4 (20 kb half-window size) that have common effects on plant response to parasitic plants and aphids, on the one hand, vs fungus, caterpillar, thrips and drought on the other

Table S6 Candidate genes in linkage with SNPs with $-\log_{10}(P)$ score > 4 (20 kb half-window size) that have common effects on biotic and abiotic stress responses

Methods S1 Salt.

Methods S2 Abiotic.

Methods S3 Caterpillar – combinatory stress.

Methods S4 Parasitic plants.

Methods S5 Nematodes.

Methods S6 Whiteflies.

Methods S7 Aphids.

Methods S8 Thrips.

Methods S9 Drought – combinatory stress.

Methods S10 Fungus – combinatory stress.

Methods S11 Screening of T-DNA lines.

Methods S12 Simulations to compare power for full MTMM, contrast MTMM and univariate analysis.

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.



About *New Phytologist*

- *New Phytologist* is an electronic (online-only) journal owned by the New Phytologist Trust, a **not-for-profit organization** dedicated to the promotion of plant science, facilitating projects from symposia to free access for our Tansley reviews.
- Regular papers, Letters, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as ready' via *Early View* – our average time to decision is <28 days. There are **no page or colour charges** and a PDF version will be provided for each article.
- The journal is available online at Wiley Online Library. Visit **www.newphytologist.com** to search the articles and register for table of contents email alerts.
- If you have any questions, do get in touch with Central Office (np-centraloffice@lancaster.ac.uk) or, if it is more convenient, our USA Office (np-usaoffice@lancaster.ac.uk)
- For submission instructions, subscription and all the latest information visit **www.newphytologist.com**