


Change of Fractal Dimension during the early stages of Lysozyme Crystallization

R. J. Heigl¹, J. Stellbrink², A. Radulescu¹, R. Schweins³, T. E. Schrader¹, D. Richter²

¹Jülich Centre for Neutron Science (JCNS) at Heinz Maier-Leibnitz Zentrum (MLZ), Forschungszentrum Jülich GmbH, Lichtenbergstr.1, 85748 Garching, Germany, ²Jülich Centre for Neutron Science JCNS and Institute for Complex Systems ICS, Forschungszentrum Jülich GmbH, 52525 Jülich, Germany, ³Heinz Maier-Leibnitz-Zentrum (MLZ), Lichtenbergstr. 1, 85748 Garching, ³ Institut Laue – Langevin (ILL), DS / LSS, 71 avenue des Martyrs, 38000 Grenoble, France

Motivation / chosen model system

The BioDiff instrument at MLZ in Garching:




Necessary crystal size:
At least 0.5 mm³

- Deeper understanding of the underlying crystallization mechanism is required

For a deeper understanding of the enzymatic action of some proteins, hydrogen atoms play a key role. Be it due to their involvement in a critical hydrogen bonding network or as part of a proton transport pathway. Often x-ray crystallography fails to reveal the atom positions of these important hydrogen atoms. Here, neutron protein crystallography as for example performed with the instrument BioDiff provides a solution, because neutrons scatter from nuclei and hydrogen has a similar scattering cross section as carbon or oxygen. The big drawback of this method is the need for a comparatively large protein crystal in order to obtain a decent signal. The aim of this study was to develop a microscopic picture of protein crystal growth. Especially the early phase of crystal growth decides whether many crystal seeds lead to the growth of many but small crystals or few but large crystals. The latter being disireable for neutron protein crystallography.

Chosen crystallization conditions


- Lysozyme 60 mg/ml in D₂O, pH adjusted with 1M NaAc 0,02 µm filtered
- NaCl 6wt% in D₂O Puffer 10mM NaAc HAC 0,02 µm filtered
- **1:1 mixture:**
Lysozyme 30 mg/ml + NaCl 3 wt% in D₂O buffer @ pH 4.35



Methods



SANS





DLS





SLS

Since the crystal growth strongly depends on parameters like Temperature, pH, protein and salt concentration the reproducibility of the crystallization speed between two crystallization runs is sometimes limited. This makes it attractive to obtain as much information from one run as possible and use this information to align the results from the SLS and SANS curves.




Results

Results from the dynamic light scattering measurements





Combination of static light scattering with Small angle neutron scattering curves





The new results from this study were an observation of the change in fractal deminsion in the early phase of the crystallisation process. This can be deduced from fitting the static light scattering data (see figures on the right). This has not been observed previously since structural resolution was missing in earlier studies. A model based approach for interpreting the time dependence of the growth rate of the hydrodynamic radii deduced from the DLS data can only give an average fractal dimension (see figure on the left).

Fitting of the static light scattering curves Reveals a change in fractal dimension in the Early phase of crystal growth




Reproducibility of the measurements



The speed of the crystallisation process varies slightly due to small variations in temperature or salt/protein concentration. This can be corrected by a time scaling factor based on the DLS results.

Resulting model



Y. Georgalis, P. Umbach, J. Raptis and Wolfram Saenger, Acta Cryst. 53 (1997) 703-712

Based on the results shown above one can improve the models for the growth of lysozyme crystals. Referring to the model proposed by Georgalis et al. (shown on the left in black) we see more a single track with a co-existence of the fractals and the growing crystals in the final equilibrium state (see figure on the right). In the early phase the rising fractal dimension indicates a filling up of the spaces between the branches of the earliest fractals of dimension 1.0. This later results then in a complete space filling and a subsequent crystal growth or in a slow fractal growth presumably when the space filling is not in accordance with the crystal structure.

