

MINI-REVIEW ARTICLE



## Copper(II) and the pathological H50Q $\alpha$ -synuclein mutant: Environment meets genetics

Anna Villar-Piqué<sup>a</sup>, Giulia Rossetti<sup>b,c,d</sup>, Salvador Ventura<sup>e</sup>, Paolo Carloni<sup>b</sup>, Claudio O. Fernández<sup>f,g</sup>, and Tiago Fleming Outeiro<sup>a,h</sup>

<sup>a</sup>Department of Neurodegeneration and Restorative Research, University Medical Centre Göttingen, Göttingen, Germany; <sup>b</sup>Computational Biomedicine, Institute for Advanced Simulation IAS-5 and Institute of Neuroscience and Medicine INM-9, Jülich, Germany; <sup>c</sup>Department of Oncology, Hematology, and Stem Cell Transplantation, Medical School, Rheinisch-Westfälische Technische Hochschule Aachen University, Aachen, Germany; <sup>d</sup>Simulation Laboratory Biology - Jülich Supercomputing Centre (JSC), Jülich, Germany; <sup>e</sup>Institut de Biotecnologia i Biomedicina and Departament de Bioquímica i Biologia Molecular, Universitat Autònoma de Barcelona, Barcelona, Spain; <sup>f</sup>Max Planck Laboratory for Structural Biology, Chemistry, and Molecular Biophysics of Rosario, Universidad Nacional de Rosario, Ocampo y Esmeralda, Santa Fe, Argentina; <sup>g</sup>Instituto de Investigaciones para el Descubrimiento de Fármacos de Rosario UNR-Consejo Nacional de Investigaciones Científicas y Técnicas de Argentina, Universidad Nacional de Rosario, Ocampo y Esmeralda, Rosario, Argentina; <sup>h</sup>Max Planck Institute for Experimental Medicine, Göttingen, Germany

### ABSTRACT

Copper is one of the metals described to bind the Parkinson disease-related protein  $\alpha$ -synuclein (aSyn), and to promote its aggregation. Although histidine at position 50 in the aSyn sequence is one of the most studied copper-anchoring sites, its precise role in copper binding and aSyn aggregation is still unclear. Previous studies suggested that this residue does not significantly affect copper-mediated aSyn aggregation. However, our findings showed that the aggregation of the pathological H50Q aSyn mutant is enhanced by copper hints otherwise. Despite the inexistence of a model for aSyn H50Q-copper complexation, we discuss possible mechanisms by which this metal contributes to the misfolding and self-assembly of this particular aSyn mutant. Considering the genetic association of the H50Q mutation with familial forms of Parkinson disease, and the fact that copper homeostasis is deregulated in this disorder, understanding the interplay between both factors will shed light into the molecular and cellular mechanisms triggering the development and spreading of the aSyn pathology.

### ARTICLE HISTORY

Received 7 November 2016  
Revised 27 November 2016  
Accepted 1 December 2016

### KEYWORDS

$\alpha$ -synuclein; amyloid; copper; H50Q mutation; Parkinson disease; protein aggregation

The abnormal accumulation of  $\alpha$ -synuclein (aSyn) is a characteristic of various neurodegenerative diseases, such as Parkinson disease (PD), that are collectively known as synucleinopathies.<sup>1</sup> However, the pathological consequences of this process are still not clear, neither the molecular mechanisms promoting and/or causing aSyn aggregation. Factors that promote aSyn misfolding and self-assembly include point mutations, found in genetic forms of synucleinopathies, and the presence of environmental stimuli, such as pro-aggregation metals.<sup>2,3</sup> We recently exploited the unique combination of the H50Q pathological aSyn mutation and the presence of copper(II) ( $\text{Cu}^{2+}$ ), a metal whose homeostasis is deregulated in many neurodegenerative conditions, and found that the neuronal damage induced by the aSyn species used in our study is not correlated with the accumulation

of intracellular aSyn inclusions, supporting the dissociation between toxicity and the accumulation of large inclusion bodies.<sup>4</sup>


Since the recent discovery of the H50Q mutation in familial forms of PD, various studies reported that it enhances the aggregation of aSyn in vitro.<sup>5-7</sup> Additionally,  $\text{Cu}^{2+}$  and other metals are known to accelerate aSyn fibrillation.<sup>8</sup>  $\text{Cu}^{2+}$  interacts with the negatively-charged C-terminal region of the protein, but binds the N-terminal region with higher affinity.<sup>9,10</sup> Two main possible complexation modes in the N-terminus are suggested: one consider the first few residues and His50 (together with neighboring residues) as 2 independent binding sites;<sup>10</sup> the other considers the formation of a long-range bridge in the polypeptide chain through the binding of a single metal ion, in which His 50 would act as a coordination switch (Fig. 1).<sup>9</sup>  $\text{Cu}^{2+}$

**CONTACT** Prof. Dr. Tiago Fleming Outeiro  [touteir@gwdg.de](mailto:touteir@gwdg.de)  University Medical Centre of Göttingen, Waldweg 33, 37073 Göttingen, Germany.

Mini-Review to: Villar-Piqué A, Lopes da Fonseca T, Sant'Anna R, Szegő ÉM, Fonseca-Ornelas L, Pinho R, Carija A, Gerhardt E, Masaracchia C, Abad Gonzalez E, Rossetti G, Carloni P, Fernández CO, Foguel D, Milosevic I, Zweckstetter M, Ventura S, Outeiro TF. Environmental and genetic factors support the dissociation between  $\alpha$ -synuclein aggregation and toxicity. *Proc Natl Acad Sci U S A*. 113(42):E6506-E6515.; <http://dx.doi.org/10.1073/pnas.1606791113>

© 2017 Anna Villar-Piqué, Giulia Rossetti, Salvador Ventura, Paolo Carloni, Claudio O. Fernández, and Tiago Fleming Outeiro. Published with license by Taylor & Francis.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.



**Figure 1.** The H50 residue is key for anchoring  $\text{Cu}^{2+}$  binding to aSyn. Quantum Mechanics / Molecular Mechanics (QM/MM) simulations<sup>21</sup> (21) were performed to test the stability of all the possible coordination geometries of  $\text{Cu}^{2+}$  at H50 site, fitting available experimental evidences.<sup>14,17,20</sup> Models A and B results to be the most plausible. The  $\text{Cu}^{2+}$  ion binds to H50 side chain in both of them. The 3 additional ligands are the H50 amide group, a water molecule, and either V48 carbonyl O (model A) or V49 deprotonated amide (model B). All of the coordination bonds exhibited relatively small fluctuations around their average values, except for the  $\text{Cu}^{2+}$ -V48 carbonyl and  $\text{Cu}^{2+}$ -V49 amide. This suggests an interconversion between the 2 forms. Residues and water molecules are shown in ball-and-stick representation: red, blue, gray and white for oxygen, nitrogen, carbon and hydrogen, respectively.  $\text{Cu}^{2+}$  is shown as a golden ball.

coordination provokes compaction in aSyn.<sup>11,12</sup> These conformational changes, even if subtle, may reduce the free energy barrier and trigger aggregation. However, these hypotheses are challenged by the recent finding that the aggregation-inducing activity of  $\text{Cu}^{2+}$  is stronger in the H50Q mutant than in the wild-type (WT).<sup>4</sup> This conclusion is derived not only from cell-free aggregation experiments but also from a cellular model where a panel of pathological mutations was expressed and exposed to different metals and only the combination of  $\text{Cu}^{2+}$  and H50Q resulted in the formation of inclusion bodies.<sup>4</sup> This uncovers a plethora of questions regarding the binding mode of  $\text{Cu}^{2+}$  in this particular mutant of aSyn, and why such event causes the rapid formation of aSyn aggregates (Table 1).

Existing data indicate that the binding profile of aSyn- $\text{Cu}^{2+}$  is not significantly disturbed in the H50Q mutant, despite the absence of the His50 site, suggesting the absence of coordination of  $\text{Cu}^{2+}$  with Gln50.<sup>5</sup> However, the faster aggregation of H50Q in the presence of  $\text{Cu}^{2+}$ , when compared with WT aSyn, indicates that position 50 likely plays a role in modulating the effect of  $\text{Cu}^{2+}$ . One possibility is that His50 lies on a “strategical” position in the sequence, in such a way that the binding of  $\text{Cu}^{2+}$  to this residue may induce a local turn-like structure, thereby favoring  $\beta$ -sheet nucleation.<sup>13</sup> The lack of this process in the H50Q variant, due to the lack of  $\text{Cu}^{2+}$  binding at position 50, would trigger other misfolding mechanisms, resulting in different aggregation characteristics. Another possibility is that, indeed, the differences between  $\text{Cu}^{2+}$ -promoted aggregation of WT and H50Q

do not depend on conformational changes in the beginning of the aggregation process, but rather on the subsequent elongation stages. However, previous studies reported that, at least with WT aSyn,  $\text{Cu}^{2+}$  mainly influences the lag phase of aggregation, promoting accelerated nucleation.<sup>14</sup> Similarly, although the  $\text{Cu}^{2+}$  effect on WT aSyn seems restricted to intramolecular phenomena<sup>11</sup> promotion of intermolecular aberrant interactions could occur in the H50Q. Additionally, it is also possible that the increased flexibility of the C-terminal region in the H50Q mutant<sup>15</sup> could change the  $\text{Cu}^{2+}$ -binding features in this domain and, thereby affect the overall aggregation process. Although the absence of the imidazole ring of

**Table 1.** Effects of  $\text{Cu}^{2+}$  on H50Q vs. WT aSyn.

Metal binding profile	No major alterations detected besides the loss of the His50 binding site (measured by electrospray ionization–ion mobility spectrometry–mass spectrometry, <sup>18</sup> electron paramagnetic resonance <sup>19</sup> and nuclear magnetic resonance <sup>5</sup> )
In vitro aggregation	Faster aggregation rate <sup>4,18</sup>
Aggregate morphology	Fibrillar morphology is replaced by amorphous aggregates <sup>4</sup>
Seeding activity	Mature aggregates display very low homo-seeding capacity <sup>4</sup>
Amyloid structure	Reduced thioflavin-T binding and $\beta$ -sheet structure <sup>4</sup>
Cellular aggregation	Stronger inclusion formation capacity of H50Q aSyn in the presence of $\text{Cu}^{2+}$ <sup>4</sup>
Exposure of neurons to exogenous aSyn aggregates	H50Q aggregates formed with $\text{Cu}^{2+}$ are stronger inducers of intracellular aSyn aggregation, but display reduced neurotoxicity <sup>4</sup>

His50 may disturb  $\text{Cu}^{2+}$ -triggered aSyn aggregation, Gln50 may also promote a toxic gain-of-function role. The aggregation of different aSyn point-mutants at position 50 revealed that, while H50Q was the more aggregation-prone among the mutants tested, a positive charge at this site (WT or H50R) probably protects from fibrilization.<sup>15</sup> A similar experiment performed in the presence of  $\text{Cu}^{2+}$  would contribute to defining the effect of Gln50 in  $\text{Cu}^{2+}$  binding. Even if the Gln50 is not directly involved in metal coordination, it could play a role in stabilizing protein-metal complexation.<sup>16</sup>

Electron microscopy confirmed that fibrils formed by the H50Q mutant resemble those formed by WT aSyn.<sup>4-6</sup>  $\text{Cu}^{2+}$  also does not alter the morphology of fibers formed by WT aSyn.<sup>14</sup> However, the combination of the H50Q mutation and  $\text{Cu}^{2+}$  results in the formation of amorphous aggregates.<sup>4</sup> Although this could be the mere consequence of the very rapid self-assembly rate, it will be worthwhile exploring this issue further to understand when and where those morphological changes arise.

Finally, certain physiologic circumstances may also cause the loss of  $\text{Cu}^{2+}$ -binding, such as N-terminal acetylation of aSyn or a reduction in the micro-environmental pH.<sup>17,18</sup> The latter would result in protonation of the His50 imidazole ring, impairing metal binding. In this case, the H50Q substitution could create a similar scenario, forcing  $\text{Cu}^{2+}$  to coordinate only to the N-terminal cluster. The lack of a potential intramolecular bridge supported by a  $\text{Cu}^{2+}$  ion might alter the entire conformation, probably inducing more open  $\text{Cu}^{2+}$ -aSyn complexation modes, affecting the aggregation process and eventually the toxicity of the resultant species.

In the future, when further addressing the role of  $\text{Cu}^{2+}$  as an enhancer of the H50Q aggregation, physiologic micro-environments and posttranslational modifications will have to be considered. A molecular understanding of the interplay between genetic factors, such as aSyn mutations and environmental factors, such as the presence of  $\text{Cu}^{2+}$ , might greatly contribute for our understanding of the pathological events causing PD.

## Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

## Funding

AV-P is supported by the Georg-August-Universität Göttingen under the Dorothea Schlözer Program. PC and GR were supported in this project by the Deutsche Forschungsgemeinschaft (DFG) (CA 973/14-1), as well as, by 270,000 core

hours on the JUROPA supercomputer in the Jülich Supercomputing Center through the NIC program (NIC: 5720 and 6695). TFO is supported by the DFG Center for Nanoscale Microscopy and Molecular Physiology of the Brain. S.V. has been granted an ICREA ACADEMIA award. TFO and SV are supported by a grant from Fundación La Marato de TV3 (Ref. 20144330).

## References

- [1] McCann H, Stevens CH, Cartwright H, Halliday GM.  $\alpha$ -Synucleinopathy phenotypes. *Park Relat Disord* 2014; 20:S62-7; [http://dx.doi.org/10.1016/S1353-8020\(13\)70017-8](http://dx.doi.org/10.1016/S1353-8020(13)70017-8)
- [2] Santner A, Uversky VN. Metalloproteomics and metal toxicology of alpha-synuclein. *Metallomics* 2010; 2:378-92; PMID:21072383; <http://dx.doi.org/10.1039/b926659c>
- [3] Petrucci S, Ginevrino M, Valente EM. Phenotypic spectrum of alpha-synuclein mutations: New insights from patients and cellular models. *Parkinsonism Relat Disord* 2016; 22:S16-20; PMID:26341711; <http://dx.doi.org/10.1016/j.parkreldis.2015.08.015>
- [4] Villar-Piqué A, Lopes da Fonseca T, Sant'Anna R, Szegő ÉM, Fonseca-Ornelas L, Pinho R, Carija A, Gerhardt E, Masaracchia C, Abad Gonzalez E, et al. Environmental and genetic factors support the dissociation between  $\alpha$ -synuclein aggregation and toxicity. *Proc Natl Acad Sci USA* 2016; In press. <http://www.pnas.org/content/early/2016/10/04/1606791113.abstract>
- [5] Khalaf O, Fauvet B, Oueslati A, Dikiy I, Mahul-Mellier AL, Ruggeri FS, Mbefo MK, Vercruysse F, Dietler G, Lee SJ, et al. The H50Q mutation enhances alpha-synuclein aggregation, secretion, and toxicity. *J Biol Chem* 2014; 289:21856-76; PMID:24936070; <http://dx.doi.org/10.1074/jbc.M114.553297>
- [6] Rutherford NJ, Moore BD, Golde TE, Giasson BI. Divergent effects of the H50Q and G51D SNCA mutations on the aggregation of  $\alpha$ -synuclein. *J Neurochem* 2014; 131:859-67; PMID:24984882; <http://dx.doi.org/10.1111/jnc.12806>
- [7] Ghosh D, Mondal M, Mohite GM, Singh PK, Ranjan P, Anoop A, Ghosh S, Jha NN, Kumar A, Maji SK. The parkinson's disease-associated H50Q mutation accelerates  $\alpha$ -synuclein aggregation in vitro. *Biochemistry* 2013; 52:6925-7; PMID:24047453; <http://dx.doi.org/10.1021/bi400999d>
- [8] Uversky VN, Li J, Fink AL. Metal-triggered structural transformations, aggregation, and fibrillation of human alpha-synuclein: A possible molecular link between parkinson's disease and heavy metal exposure. *J Biol Chem* 2001; 276:44284-96; PMID:11553618; <http://dx.doi.org/10.1074/jbc.M105343200>
- [9] Dudzik CG, Walter ED, Millhauser GL. Coordination features and affinity of the  $\text{Cu}(2+)$  site in the alpha-synuclein protein of Parkinson's disease. *Biochemistry* 2011; 50:1771-7; PMID:21319811; <http://dx.doi.org/10.1021/bi101912q>
- [10] Sung YH, Rospigliosi C, Eliezer D. NMR mapping of copper binding sites in alpha-synuclein. *Biochim Biophys Acta* 2006; 1764:5-12; PMID:16338184; <http://dx.doi.org/10.1016/j.bbapap.2005.11.003>

- [11] Ahmad A, Burns CS, Fink AL, Uversky VN. Peculiarities of copper binding to alpha-synuclein. *J Biomol Struct Dyn* 2012; 29:825-42; PMID:22208282; <http://dx.doi.org/10.1080/073911012010525023>
- [12] Binolfi A, Rodriguez EE, Valensin D, D'Amelio N, Ippoliti E, Obal G, Duran R, Magistrato A, Pritsch O, Zweckstetter M, et al. Bioinorganic chemistry of Parkinson's disease: Structural determinants for the copper-mediated amyloid formation of alpha-synuclein. *Inorg Chem* 2010; 49:10668-79; PMID:20964419; <http://dx.doi.org/10.1021/ic1016752>
- [13] Rose F, Hodak M, Bernholc J. Mechanism of copper(II)-induced misfolding of Parkinson's disease protein. *Sci Rep* 2011; 1:11; PMID:22355530; <http://dx.doi.org/10.1038/srep00011>
- [14] Rasia RM, Bertoncini CW, Marsh D, Hoyer W, Cherny D, Zweckstetter M, Griesinger C, Jovin TM, Fernández CO. Structural characterization of copper(II) binding to alpha-synuclein: Insights into the bioinorganic chemistry of Parkinson's disease. *Proc Natl Acad Sci U S A* 2005; 102:4294-9; PMID:15767574; <http://dx.doi.org/10.1073/pnas.0407881102>
- [15] Chi YC, Armstrong GS, Jones DNM, Eisenmesser EZ, Liu CW. Residue histidine 50 plays a key role in protecting  $\alpha$ -Synuclein from aggregation at physiological pH. *J Biol Chem* 2014; 289:15474-81; PMID:24742669; <http://dx.doi.org/10.1074/jbc.M113.544049>
- [16] Chiera NM, Rowinska-Zyrek M, Wieczorek R, Guerrini R, Witkowska D, Remelli M, Kozłowski H. Unexpected impact of the number of glutamine residues on metal complex stability. *Metallomics* 2013; 5:214-21; PMID:23370132; <http://dx.doi.org/10.1039/c3mt20166j>
- [17] De Ricco R, Valensin D, Dell'Acqua S, Casella L, Dorlet P, Faller P, Hureau C. Remote His50 acts as a coordination switch in the high-affinity N-terminal centered copper(II) Site of  $\alpha$ -Synuclein. *Inorg Chem* 2015; 54:4744-51; PMID:25926427; <http://dx.doi.org/10.1021/acs.inorgchem.5b00120>
- [18] Mason RJ, Paskins AR, Dalton CF, Smith DP. Copper binding and subsequent aggregation of alpha-synuclein are modulated by N-terminal acetylation and ablated by the H50Q missense mutation. *Biochemistry* 2016; 55:4737-41; PMID:27517125; <http://dx.doi.org/10.1021/acs.biochem.6b00708>
- [19] Proukakis C, Dudzik CG, Brier T, MacKay DS, Cooper JM, Millhauser GL, Houlden H, Schapira AH. A novel alpha-synuclein missense mutation in Parkinson disease. *Neurology* 2013; 80:1062-4; PMID:23427326; <http://dx.doi.org/10.1212/WNL.0b013e31828727ba>
- [20] Valensin D, Camponeschi F, Luczkowski M, Baratto MC, Remelli M, Valensin G, Kozłowski H. The role of His-50 of alpha-synuclein in binding Cu(II): pH dependence, speciation, thermodynamics and structure. *Metallomics* 2011; 3(3):292-302; PMID:21212878; <http://dx.doi.org/10.1039/c0mt00068j>
- [21] Laio A, VandeVondele J, Rothlisberger U. A Hamiltonian electrostatic coupling scheme for hybrid Car-Parrinello molecular dynamics simulations. *J Chem Phys* 2002; 116(16):6941-6947