

SCIENTIFIC REPORTS

OPEN

Hair Cortisol in Twins: Heritability and Genetic Overlap with Psychological Variables and Stress-System Genes

Liz Rietschel^{1,2}, Fabian Streit³, Gu Zhu⁴, Kerrie McAloney⁴, Josef Frank³, Baptiste Couvy-Duchesne^{4,5}, Stephanie H. Witt³, Tina M. Binz⁶, CORtisolNETwork (CORNET) Consortium*, Major Depressive Disorder Working Group of the Psychiatric Genomics Consortium (PGC)*, John McGrath^{5,7,8}, Ian B. Hickie⁹, Narelle K. Hansell⁵, Margaret J. Wright^{5,10}, Nathan A. Gillespie^{4,11}, Andreas J. Forstner^{12,13,14,15}, Thomas G Schulze^{3,16,17,18,19}, Stefan Wüst^{3,20}, Markus M. Nöthen^{12,13}, Markus R. Baumgartner⁶, Brian R. Walker²¹, Andrew A. Crawford^{21,22}, Lucía Colodro-Conde⁴, Sarah E. Medland⁴, Nicholas G. Martin⁴ & Marcella Rietschel³

Hair cortisol concentration (HCC) is a promising measure of long-term hypothalamus-pituitary-adrenal (HPA) axis activity. Previous research has suggested an association between HCC and psychological variables, and initial studies of inter-individual variance in HCC have implicated genetic factors. However, whether HCC and psychological variables share genetic risk factors remains unclear. The aims of the present twin study were to: (i) assess the heritability of HCC; (ii) estimate the phenotypic and genetic correlation between HPA axis activity and the psychological variables perceived stress, depressive symptoms, and neuroticism; using formal genetic twin models and molecular genetic methods, i.e. polygenic risk scores (PRS). HCC was measured in 671 adolescents and young adults. These included 115 monozygotic and 183 dizygotic twin-pairs. For 432 subjects PRS scores for plasma cortisol, major depression, and neuroticism were calculated using data from large genome wide association studies. The twin model revealed a heritability for HCC of 72%. No significant phenotypic or genetic correlation was found between HCC and the three psychological variables of interest. PRS did not explain variance in HCC. The present data suggest that HCC is highly heritable. However, the data do not support a strong biological link between HCC and any of the investigated psychological variables.

¹University Hospital of Child and Adolescent Psychiatry and Psychotherapy, Research Department, University of Bern, Bern, Switzerland. ²SRH University Heidelberg, Academy for Psychotherapy, Heidelberg, Germany. ³Department of Genetic Epidemiology in Psychiatry, Central Institute of Mental Health, Medical Faculty Mannheim, University of Heidelberg, Mannheim, Germany. ⁴Genetics & Computational Biology Department, QIMR Berghofer Medical Research, Brisbane, Australia. ⁵Queensland Brain Institute, University of Queensland, Brisbane, Australia. ⁶Zurich Institute of Forensic Medicine, Centre for Forensic Hair Analysis, University of Zurich, Zurich, Switzerland. ⁷Queensland Centre for Mental Health Research, The Park Centre for Mental Health, Wacol, Australia. ⁸National Centre for Register-Based Research, Aarhus University, Aarhus, Denmark. ⁹Brain and Mind Centre, University of Sydney, Sydney, Australia. ¹⁰Centre for Advanced Imaging, University of Queensland, Brisbane, Australia. ¹¹Virginia Institute for Psychiatric and Behavioral Genetics, Virginia Commonwealth University, Richmond, VA, USA. ¹²Institute of Human Genetics, University of Bonn, Bonn, Germany. ¹³Life & Brain Center, Department of Genomics, University of Bonn, Bonn, Germany. ¹⁴Department of Psychiatry (UPK), University of Basel, Basel, Switzerland. ¹⁵Human Genomics Research Group, Department of Biomedicine, University of Basel, Basel, Switzerland. ¹⁶Department of Psychiatry and Behavioral Sciences, Johns Hopkins University, Baltimore, USA. ¹⁷Institute of Psychiatric Phenomics and Genomics (IPPG), Medical Center of the University of Munich, Campus Innenstadt, Munich, DE, Germany. ¹⁸Human Genetics Branch, NIMH Division of Intramural Research Programs, Bethesda, USA. ¹⁹Department of Psychiatry and Psychotherapy, University Medical Center Göttingen, Goettingen, DE, Germany. ²⁰Institute of Experimental Psychology, University of Regensburg, Regensburg, Germany. ²¹British Heart Foundation Centre for Cardiovascular Science, Queen's Medical Research Institute, University of Edinburgh, Edinburgh, UK. ²²Medical Research Council Integrative Epidemiology Unit, School of Social and Community Medicine, University of Bristol, Bristol, UK. Liz

Received: 18 May 2017

Accepted: 30 August 2017

Published online: 10 November 2017

Rietschel and Fabian Streit contributed equally to this work. Nicholas G. Martin and Marcella Rietschel jointly supervised this work. *A comprehensive list of consortium members appears at the end of the paper. Correspondence and requests for materials should be addressed to L.R. (email: Liz.Rietschel@srh.de)

Research has generated robust evidence that chronic stress is a risk factor for mental disorders^{1,2}. Understanding the mechanisms through which stress impacts mental health is therefore an important aim of epidemiological research. A core element of the biological stress response is the hypothalamus-pituitary-adrenal (HPA) axis. HPA axis activity is typically measured according to its end product, the steroid hormone cortisol. While prompt HPA axis activation in response to acute stressors is adaptive, long-term dysregulation of basal HPA axis activity and reactivity can have deleterious effects on physiology^{2,3}. Alterations in HPA axis regulation are observed in subjects suffering from psychiatric disorders, and have been suggested as not only a consequence but also as a premorbid vulnerability factor^{4,5}. Research has shown that variation in HPA axis regulation is influenced by both environmental and genetic factors⁶. Furthermore, authors have suggested that the effects of genetic and environmental risk factors for psychiatric disorders might be mediated in part via dysregulation of HPA axis activity^{2,7}. Investigating if HPA axis regulation shares underlying genetic factors with psychological or psychiatric phenotypes can inform about a true biological link between these, potentially indicating a causal involvement of the HPA axis in the vulnerability for psychiatric disorders.

Cortisol is usually measured in saliva, urine, or blood⁸. However, single cortisol measures are strongly influenced by factors such as circadian rhythm, physical activity, and nutrition. Thus the assessment of long-term alterations in HPA axis regulation requires the meticulous assessment of cortisol at multiple time points. In recent years, the assessment of hair cortisol concentration (HCC) has been established as a marker of long-term cumulative HPA axis activity^{9,10}. HCC is usually measured in a 3 centimeter (cm) hair segment cut as close as possible to the scalp. Since hair grows, on average, one cm per month, HCC in a 3 cm sample is considered to reflect cortisol secretion during the preceding 3 month period¹¹. Hair cortisol is assumed to reflect free cortisol, which is the biologically active share of cortisol not bound to corticosteroid binding globulin (CBG)^{12,13}. Studies investigating the relationship of HCC with cortisol levels in other tissues show the highest correlations (up to $r = 0.61$) with cumulative or average cortisol measures acquired over several days or weeks^{14–17}. While these studies which assessed multiple measurement-points have been mainly carried out using saliva and urine, studies which used serum and plasma blood samples, have been mainly based on single assessments, and the observed correlations were lower or non-significant (e.g. refs^{18–20}). Studies in pregnant women^{14,21} and in subjects with Cushing's syndrome^{20,22} -conditions characterized by pronounced alterations in circulating cortisol- have revealed altered HCC levels, and therefore indicate that HCC is a marker of long-term changes in circulating cortisol. HCC therefore represents an efficient method for the retrospective assessment of long-term cortisol secretion, and thus long-term HPA axis activity.

While HCC would thus appear to be an ideal biomarker for stress-related phenotypes, to date, studies of the association between HCC and measures of stress and mental health have generated inconsistent results (for review see refs^{9,10,23}). A recent meta-analysis based on aggregated data from a total of 124 samples, and comprising 10,289 subjects, showed no consistent associations with mood disorders or self-reported perceived stress and depressiveness, but found that stress-exposed groups as a whole exhibit 22% increased HCC²⁴. While stressful environmental factors play a major role in HPA axis activation, twin studies have indicated that genetic factors have a substantial impact on the secretion of cortisol, especially morning cortisol. Predominantly moderate heritability estimates have been reported in adults (as reviewed in refs^{25–27}) and adolescents^{25,27,28}. The observed heritability of HPA axis regulation suggests a contribution of genetically determined biological mechanisms. To date, the only large genome wide association study (GWAS) systematically investigating the genes underlying HPA axis activity used total plasma cortisol levels in the morning: Bolton *et al.* (2014) identified associated genetic variants in a region which contains the genes encoding CBG and $\alpha 1$ -antitrypsin²⁹. If a strong heritability for HCC could be confirmed, HCC would represent a promising target for genetic studies into long-term HPA axis activity and its relationship to mental health.

The first study on heritability of HCC was conducted in a colony of female vervet monkeys, and heritability estimates of ~30% in high and low stress environments were reported³⁰. In humans, Tucker-Drob *et al.* (2017) demonstrated heritability of HCC for the first time by investigating an ethnically and socioeconomically diverse sample of 1070 children and adolescents including 533 twin pairs³¹. The authors showed that 65% of the total variability of HCC was explained by additive genetic effects, and that genetic influences on HCC decreased with age. In subjects with low socioeconomic status, a non-significant trend was observed towards increased genetic influences and reduced shared environmental influences on HCC. The authors hypothesized that genetic influences may be stronger under high stress conditions.

As mentioned above, a genetic overlap between HPA axis activity and psychological or psychiatric phenotypes would suggest an involvement of the HPA axis in the vulnerability to psychiatric disorder. However, previous studies investigating a possible genetic overlap between cortisol secretion and psychological variables have been limited in both number and size: A study in 29 monozygotic female twin pairs suggested around 40–45% of the total variance in morning and evening saliva cortisol levels is shared by monozygotic twins³². Furthermore, increased ($p = 0.06$) mean cortisol levels were observed in those twins with a history of major depression (MDD). Notably, intermediate cortisol levels were observed in twins without history of MDD from pairs discordant for history of MDD, however these observations were not significant. A study conducted in 125 female twin pairs demonstrated a heritability of 55% for neuroticism and a heritability of up to 69% for morning cortisol secretion in saliva³³. However, the authors found no phenotypic or genotypic association between cortisol levels and neuroticism.

In a previous twin study, the present authors demonstrated that continuous measures of perceived stress, depressive symptoms, and neuroticism were heritable and showed strong phenotypic and genetic correlations in healthy adolescent and young adult twins³⁴. In line with this, recent large genome-wide studies have demonstrated that genetic variants make a substantial contribution to the development of neuroticism^{35,36}, depressive symptoms³⁶, and MDD³⁷. These studies have also revealed that these risk variants show partial overlap. In a subsequent pilot study, the present authors measured HCC and the three psychological variables in a sample of 109 children and young adults, which included eight monozygotic- and 21 dizygotic twin pairs³⁸. Due to the small sample size, no reliable heritability estimates for HCC could be generated. However, the findings suggested that HCC and the assessed psychological variables may share a common genetic basis.

The present larger study aimed (i) to estimate the heritability of HCC and (ii) to investigate the question of whether HPA axis activity shows a genetic overlap with the continuous psychological measures perceived stress, depressive symptoms, and neuroticism, using formal genetic (twin models), and molecular genetic methods i.e. polygenic risk score (PRS) analyses. PRS provide a quantitative measure of genetic risk or vulnerability for a given trait. PRS estimation uses GWAS results to predict genetic risk for each individual in an independent genotyped sample. Investigations can then be performed to determine whether this risk is associated with potentially related phenotypes. However, while significant associations can be found using this approach, the explained variance is limited. For the purposes of the present study, PRS were calculated based on results from GWAS of HPA axis activity²⁹; MDD³⁷; and neuroticism³⁵. Since the GWAS of HPA axis activity was based on plasma cortisol levels, the association between HCC and this PRS was explored i.e. the association between HCC and the genetic variants influencing concentration of cortisol in another tissue.

Materials and Methods

Subjects. Our samples consisted of adolescent and young adult twins from the Brisbane area recruited mainly for studies of the genetics of melanoma risk factors and cognition; we neither selected nor excluded participants for any particular phenotype, nor did we systematically obtain data on medical history and treatment. Hair samples were collected from 674 adolescents/young adults. After exclusion of four hair samples (see *hair sampling and HCC analysis*), the final cohort comprised 671 subjects (age mean = 14.5 ± 2.4 years; range = 10.1–31.1 years; 419 females). The cohort included 116 monozygotic (MZ) and 187 dizygotic (DZ) twin pairs (3 families had two sets of DZ twins), and 14 sets of trizygotic triplets. For the purposes of the present analyses, each set of triplets was considered to be a DZ twin pair with one additional singleton. The cohort included a total of 65 singletons, who were derived from the divided triplets (n = 14) and the siblings of the MZ and DZ twin pairs (n = 51) (for details see Supp. Table 1).

All subjects had participated in at least one phase of the Brisbane Longitudinal Twin Study^{39,40}. This ongoing, longitudinal study of adolescent/young adult twins and siblings from the general population of the Brisbane area (Australia), is conducted in several phases and investigates somatic- and mental health and related phenotypes (for details see Supp. Text and Supp. Table 2). For 146 subjects (including 29 MZ and 42 DZ twin pairs), hair samples were collected at two time-points (73 subjects at 12 and 14 years; 73 subjects at 14 and 16 years; for details see Supp. Table 1b). Stability of HCC and the psychological variables was assessed in terms of correlations between the two time points (for details see Supp. Table 1b). For 432 subjects (age mean = 15.5 ± 2.4 years; range = 10–31 years; 268 females), genome-wide genotype data were available (for details see Supp. Table 1c). The study was approved by the Human Research Ethics Committee of the Queensland Institute of Medical Research (QIMR) and conducted in accordance with the Declaration of Helsinki. Written informed consent was obtained from all subjects, and from legal guardians in the case of minors, prior to inclusion and sample collection.

Psychological variables. For participants less than 16 years old, perceived stress was measured using the 30-item 'Daily Life and Stressors Scale' (DLSS⁴¹) and neuroticism was measured using the respective 20 items of the 'Junior Eysenck Personality Questionnaire' (JEPQ⁴²) as the DLSS and JEPQ are validated for children and adolescents. For participants aged 16 years or older, perceived stress was measured using the 10-item 'Perceived Stress Scale' (PSS⁴⁴). Additionally, 168 subjects between 16 and 19 years completed both the PSS and the DLSS, and this overlap was used to harmonize the two scales using item response theory (IRT; see below). For participants aged 16 years or older, neuroticism was measured using the respective 12 items of the 'NEO-Five Factor Inventory revised version' (NEO-FFI-R⁴⁴). At all ages, depressive symptoms were assessed using the 34-item 'Somatic and Psychological Health Report' (SPHERE⁴⁵). For all subjects, measures of stress (PSS or DLSS), neuroticism, and depressive symptoms were obtained at the time of hair sampling. For some subjects, data were unavailable for perceived stress (n = 1); neuroticism (n = 51); and depressive symptoms (n = 55), since subjects did not fill out the respective questionnaires. In the subgroup that underwent assessment at two time-points, one subject had missing data for perceived stress and depressive symptoms at the second time point.

Hair sampling and HCC analysis. Using fine scissors, a 3 cm hair swatch of approximately 3 millimeters in diameter was cut as close as possible to the skin from the posterior vertex of the scalp. Hair cortisol was analyzed by TMB and MRB at the Institute of Forensic Medicine, Centre for Forensic Hair Analysis, University of Zurich. Cortisol concentration was measured using liquid chromatography-tandem mass spectrometry (LC-MS/MS), as described by Binz *et al.*⁴⁶ (for details see Supp. Text). Prior to HCC measurement, hair samples were randomly assigned to batches (irrespective of time-point or twin-pair) in order to minimize the effect of batch differences on twin correlations. To assess technical error, 106 hair samples were assessed in duplicate and 27 in triplicate. The hair samples were assayed between April and July 2016 in a total of 35 batches (comprising 27 samples respectively). Samples with extreme high or low values (n = 26; 13 ≤ 0.2 and 13 ≥ 26.3) were re-assayed in order to confirm the extreme values.

Statistical Analysis. *Treatment of psychological variables and HCC.* To harmonize neuroticism scores, the neuroticism sum-scores of the NEO-FFI-R and the JEPQ scores were separately z-transformed and then combined, as described in previous studies^{47,48}. To harmonize data from the two stress rating scales DLSS and PSS, Item Response Theory (IRT) analysis was performed (for raw values and details see Supp. Text and Supp. Table 3). IRT models have the advantage that the difficulty and discriminability of each item is taken into account by modeling a normally distributed liability based on the responses to the individual questionnaire items. It is thus superior to a simple sum score that assumes all items have the same discriminating ability with respect to the underlying liability being measured and is thus particularly useful if widely different scales are being combined—as here for perceived stress. We made use of the overlapping cases who had completed both scales in order to put both measures on the same liability scale. IRT analysis was also applied to the 34 items from the SPHERE in order to produce a single liability measure for depressive symptoms.

To account for skewness, raw HCC values (mean = 6.29 pg/mg; SD = 28.92; range = 0–560) were log transformed. Here, the lowest measured value (0.1) was added to each value prior to \log_{10} transformation. Five outliers $>3SD$ ($HCC > 64.70$ pg/mg) were winsorized to $3SD$ on the \log_{10} scale. Analyses were also conducted to test the effect of experimental variables reported in previous studies of HCC. The analyses included: (i) batch number ($n = 35$); (ii) study phase ($n = 6$); (iii) storage time, defined as time between date of collection and date of assay (range = 0.50–4.02 years) and divided into $n = 5$ groups according to increasing storage time; and (iv) sun exposure, operationalized according to month of assessment ($n = 12$) and self- and maternal ratings of sun exposure. Since all aforementioned variables changed the fit of the model significantly, they were regressed out from the HCC measurement using a linear model, including the dummy coded variables as fixed effects. Subsequent analyses were carried out with the residuals of this model. The HCC value from the first time point was used in the following analyses; the second time point was only used for assessing stability of HCC over a two-year period in the subset of longitudinally assessed subjects ($n = 146$).

Twin correlations, heritability of HCC, and shared covariance with psychological variables. To estimate twin correlations and the heritability of HCC and the three psychological variables of interest, the twin and sibling data were used to generate structural equation models. Model parameters were estimated using the full maximum likelihood method implemented in Mx⁴⁹. This makes use of all data points - including those of unpaired twins and singletons - in order to improve estimation of sample means and variances. This approach allows partitioning of the variation into: additive genetic influences (A); shared environmental influences (C); and unique environmental influences (E). Using likelihood ratio chi-square tests, sub-models with only two factors (AE and CE models) were compared with the three-factor models (ACE) in order to estimate the sources of variance and select the most parsimonious variance structure of the traits.

To investigate the influence of genetic and environmental factors on HCC and the psychological variables of interest, as well as genetic and environmental correlations between these variables, multivariate analysis was performed. This involved use of a simultaneous Cholesky decomposition. A Cholesky decomposition is a good initial multivariate method to use in the absence of a clear model of the factor structure relating a set of correlated variables. Furthermore, taking advantage of having both MZ and DZ twins one can fit a 3-part Cholesky model including decompositions of (co)variance due to A (additive genetics), C (shared environment) and E (unique environment), since one cannot assume that factor loadings between these three sources of variation will be proportional for each of the examined variables. In fact, there are known cases of opposite-signed loadings of factors A and C (e.g., for items of Eysenck's Psychoticism scale⁵⁰). The source of variation, C, captures cultural environment shared by twins within a family regardless of zygosity and could include factors such as sharing the same school, neighborhood, and exposure to infectious diseases. C is distinguished from unique environment, E, which is specific to individuals and may include factors such as accidents; however, it is important note that E also includes measurement error, which is often the major contributor to this source of variance. An ACE Cholesky decomposition model was compared with an AE and a CE Cholesky decomposition. Perceived stress was used as the first, depressive symptoms as the second, neuroticism as the third, and HCC as the last latent factor in order to estimate (i) the genetic variance of HCC that is shared with genes affecting the psychological variables and (ii) the independent genetic variance for HCC after removing the effects of the genes with the primary influence on the psychological variables (for further details see Supp. Text).

The fit of each model was assessed according to the differences in log likelihood between the sub and the full models. The most parsimonious model was chosen for the purposes of data interpretation. Sex and puberty which - generally starts earlier in girls- influence the cortisol secretion⁵¹. As puberty status was not assessed in the study, sex, age, age², sex x age, and sex x age² were included as covariates in all models, to allow for different age effects in boys and girls and the fact that these may be curvilinear. Body mass index ($r = 0.06$, $p = 0.11$), socio-economic index ($r = -0.07$, $p = 0.13$), and hair dyeing ($r = 0.01$, $p = 0.81$) showed no significant associations with HCC in the present sample and were thus not included as covariates. Since previous results suggest that heritability of HCC is age-dependent³¹, a separate analysis was performed in the younger and the older half of the sample, as defined by a median split to test the heritability of HCC in the different age groups. Further details of the twin design and analytical methods, including assumption testing and multivariate modeling, are provided elsewhere⁵².

Genotyping, quality control, and imputation. Genotyping was performed using the Illumina Human610-Quad and Core+Exome SNP chips. Quality control included inspection of pedigree, sex, Mendelian errors, and ancestry, as well as filtering for genotyping quality (GenCall < 0.7); SNP and individual call rates (< 0.95); Hardy-Weinberg equilibrium failure ($P < 10^{-6}$); and minor allele frequency (< 0.01). Subjects were imputed to the Haplotype Reference Consortium (HRC.1.1)⁵³ on the Michigan Imputation Server (<https://imputationserver.sph.umich.edu/i>). Imputation was carried out in two separate waves for the Illumina Human610-Quad and the

	Total (n = 671)	Male (n = 252)	Female (n = 419)
Age (years)	14.49 (2.43)	14.42 (2.59)	14.53 (2.32)
Perceived stress ^a	0.01 (0.58)	-0.02 (0.61)	0.03 (0.57)
Depressive symptoms ^b	0.01 (0.92)	-0.09 (0.94)	0.07 (0.90)
Neuroticism ^c	0.01 (1.01)	-0.12 (1.02)	0.08 (0.99)
Hair cortisol (pg/mg)	5.31 (14.23)	5.68 (18.42)	5.09 (10.98)

Table 1. Means (SD) for age, psychological variables and HCC. Abbreviations: SD = standard deviation. ^aIRT-scores deriving from perceived stress questionnaires (DLS and PSS); ^bIRT-scores of the SPHERE z -standardized values of the neuroticism questionnaires.

Core+Exome SNP chips. To account for population stratification (i.e. allele frequency differences between subjects due to systematic ancestry differences) in the PRS analysis, genetic principal components (PC) reflecting the respective ancestry were calculated using EigenSoft 6.0.1 (<http://www.hsph.harvard.edu/alkes-price/software/>). Further details on genotyping, quality control, and imputation are provided in the Supp. Text.

Polygenic risk score analysis. PLINK 1.90 (version 3, May 2016, <https://www.cog-genomics.org/plink2/>) was used to compute PRS in accordance with the procedure described by Wray *et al.*⁵³. For PRS estimation, the results of large GWAS (discovery sample) are used to calculate an aggregated genetic risk score for each individual in an independent genotyped sample (target sample). The PRS represents the sum of the risk alleles, as weighted by their respective estimated effect sizes. PRS provide a quantitative measure of the genetic risk or vulnerability for a given trait: The higher the score, the higher the predisposed genetic risk of the individual for the trait in question. In the present study, PRS were calculated using summary statistics from recent GWAS or GWAS meta-analyses of: (i) plasma cortisol (CORTisol NETwork (CORNET) Consortium²⁹, comprising 12,597 subjects); (ii) MDD (PGC-MDD²⁷ minus QIMR samples, comprising a total of 49,524 cases and 110,074 controls, for details see Supp. Table 4); and (iii) neuroticism (UK Biobank³⁵, comprising 91,370 subjects). The SNP-sets used to compute the PRS were selected using eight different p -value thresholds (5e-8, 1e-5, 0.001, 0.01, 0.05, 0.1, 0.5, 1.0) in the respective discovery sample.

To take family structure into account, associations of PRS with HCC and the three psychological variables of interest were tested using linear mixed regression models in GCTA (Genome-wide Complex Trait Analysis v. 1.26)⁵⁵. Here, the following were used as covariates: sex; age; age²; sex \times age; sex \times age²; the first five genetic principal components (PC); and the imputation wave. One-sided p -values are reported, according to the hypothesis of a positive association of the PRS with HCC and each of the psychological variables. Further details of the PRS analysis are provided in the Supp. Text.

Data availability. The datasets generated during and/or analyzed during the current study are not publicly available due to privacy regulations but are available from the authors on reasonable request.

Results

Clinical characteristics of the cohort. HCC was measured in 671 subjects. HCC, neuroticism and depressive symptom scores were higher in females than in males. Furthermore, perceived stress, depressive symptoms and neuroticism showed an association with age, and perceived stress and neuroticism showed an interaction of sex and age. Details can be found in Supp. Table 5. Distributions for age and all variables are shown in Table 1.

Influence of experimental covariates on HCC. Quality control of HCC assessed in duplicate and triplicate revealed a high correlation of the log transformed HCC values for samples re-assayed once ($n = 106$; $r = 0.89$) and samples assayed three times ($n = 27$; $r = 0.98$ with the 1st and $r = 0.98$ with the 2nd assay).

Analysis of the influence of experimental covariates resulted in a final model with a total of 54 deviations for experimental effects (34 for batch number, 4 for storage time, 11 for month, and 5 for the respective study phase). All of these experimental covariates were highly significant. Notably, the analysis demonstrated: (i) a decrease in HCC with increasing storage time; and (ii) maximum values in March (end of the Australian summer) and minimum values in September (end of the Australian winter). Dropping any one of the covariates from the full model (which includes all four) increased variance by 5.9% for batch effects, 1.8% for storage effects, 3.8% for month effects, and 6.4% for study effects (details see Supp. Table 6).

Heritability. For HCC, a DZ correlation of $r = 0.42$, and a MZ correlation of $r = 0.66$ were observed. This corresponded to a heritability estimate of $h^2 = 0.72$ in the multivariate model including HCC and the psychological variables. Slightly lower DZ and MZ twin correlations were observed for the psychological variables. Here, the heritability estimates were $h^2 = 0.54$ for perceived stress; $h^2 = 0.55$ for depressive symptoms; and $h^2 = 0.56$ for neuroticism (see Table 2).

Stability over time. As shown in Table 2, stability over the two-year period ranged between $r = 0.51$ and $r = 0.61$ for the psychological variables, while HCC stability was $r = 0.32$.

Correlation of HCC with psychological variables. High phenotypic correlations were found between the three psychological measures, ranging from $r = 0.59$ to $r = 0.64$ (Supp. Table 7). In contrast, negligible

	rMZ (115 pairs)	rDZ (183 pairs)	Heritability ^a	Two year Stability ^b (n = 146)
Perceived stress	0.51 (0.37–0.61)	0.29 (0.17–0.40)	0.54 (0.42–0.64)	0.61 (0.50–0.69)
Depressive symptoms	0.55 (0.43–0.65)	0.27 (0.13–0.40)	0.55 (0.42–0.65)	0.51 (0.37–0.62)
Neuroticism	0.53 (0.42–0.62)	0.31 (0.17–0.44)	0.56 (0.45–0.67)	0.58 (0.46–0.67)
Hair cortisol	0.66 (0.56–0.74)	0.42 (0.31–0.51)	0.72 (0.63–0.79)	0.32 (0.16–0.45)

Table 2. Monozygotic (MZ) and dizygotic (DZ) twin correlations (95% CI), heritability and stability over two years, for the psychological variables and HCC. Abbreviations: CI = confidence interval, rDZ = correlation between dizygotic twins, rMZ = correlation between monozygotic twins. All calculations were corrected for sex, age, age², sex x age, sex x age². ^aHeritabilities were estimated using the multivariate model (see Fig. 1). ^bTwo year stability was calculated as the correlation between time point 1 and time point 2.




Figure 1. Cholesky Decomposition for the AE Model. Latent factor loadings are standardized to unit variance and must be squared to obtain standardized variance components. A1–A4 additive genetic factors, E1–E4 unique environmental factors. Abbreviations: STR = perceived Stress, DEP = depressive symptoms, NEU = neuroticism, HCC = hair cortisol concentration.

correlations were observed between the three psychological measures and HCC, with $r = 0.04$ for perceived stress; $r = 0.07$ for depressive symptoms; and $r = 0.08$ for neuroticism. None of these correlations differed significantly from zero.

As counterbalancing genetic and environmental correlations of opposite sign resulting in a small or zero phenotypic correlation have been observed for psychological measures (e.g. psychoticism)⁴⁹, these associations were further explored with a multivariate Cholesky decomposition (Fig. 1). The C matrix, which accounts for shared environmental influences, could be dropped from the model without worsening fit ($\Delta\chi^2 = 13.66$, $p = 0.19$). The AE model revealed low and non-significant genetic correlations (r_A) between HCC and the three psychological variables ($r_A = 0.14$ for perceived stress; $r_A = 0.12$ for depressive symptoms; $r_A = 0.19$ for neuroticism) (Table 3).

Separate AE Cholesky analysis was then performed for the younger (mean age = 12.37 (SD = 0.54) (Supp. Table 8), and the older (mean age = 15.70 (SD = 2.32)) (Supp. Table 9) halves of the cohort (divided by median; Mdn = 14.01 years). Comparable heritability estimates for HCC were found in the younger ($h^2 = 0.74$) and older ($h^2 = 0.69$) halves of the cohort. Heritabilities for the psychological variables were also broadly consistent between the younger and older halves of the sample (see Supp. Tables 8 and 9).

Association of polygenic risk for plasma cortisol, MDD, and neuroticism with HCC and psychological variables.

The PRS for plasma cortisol showed no significant association with the HCC at any of the chosen thresholds (Fig. 2D). Furthermore, the PRS for plasma cortisol did not predict any of the three psychological variables (Fig. 2A–C). The PRS for MDD and the PRS for neuroticism showed positive associations with the psychological variables. For several p-value thresholds, in particular thresholds > 0.01 , these associations reached nominal significance (see Fig. 2A–C). No significant association was found between HCC and the PRS for MDD or the PRS for neuroticism (Fig. 2D). Details of the PRS regression analyses are provided in Supp. Tables 10–21.

Discussion

To our knowledge, the present study is the first to assess the heritability of HCC together with its phenotypic and genetic association with perceived stress, depressive symptoms, and neuroticism.

The analyses generated a heritability estimate for HCC of ~70%, with no significant contribution being found for shared environment. These estimates are nearly identical to those reported by Tucker-Drob and colleagues³¹

		(I) Standardized Parameters %				(II) Correlations			
		1	2	3	4	1	2	3	4
Genetic (A)	1-Perceived stress	53.89				1			
	2-Depressive symptoms	32.95	22.10			0.77	1		
	3-Neuroticism	33.78	2.79	19.81		0.77	0.74	1	
	4-Hair cortisol	1.33	0.03	1.20	69.37	0.14	0.12	0.19	1
Unshared Environment (E)	1-Perceived stress	46.12				1			
	2-Depressive symptoms	6.65	38.30			0.35	1		
	3-Neuroticism	12.15	5.00	26.48		0.53	0.52	1	
	4-Hair cortisol	0.68	0.07	0.21	27.02	0.16	0.11	0.19	1

Table 3. Results of the Cholesky Decomposition for the AE Model, with (I) Standardized Parameters and (II) Genetic and environmental correlations between HCC and psychological measures.




Figure 2. Association of polygenic risk scores (PRS) for plasma cortisol, neuroticism, and major depression (MDD) with: (A) perceived stress; (B) depressive symptoms; (C) neuroticism; and (D) hair cortisol concentration. Negative R^2 indicates a negative direction of the association of PRS with the respective phenotype; p one-sided: ** $p < 0.01$; * $p < 0.05$; # $p < 0.1$.

and at the upper end of those reported for measures of cortisol in other tissues (e.g. saliva and urine). This is not surprising, given that HCC is an integrated, rather than a point measure of HPA axis activity. It can now be taken as a fact that in adolescents from the general population, a substantial proportion of HCC variance is attributable to genetic factors. At first glance, this may appear surprising for a parameter that is considered a potential biomarker for stress. However, this finding does not preclude environmentally induced changes. Furthermore, the stability of HCC, as measured in a subgroup over a two-year period, is relatively low ($r = 0.32$). These stability measures are comparable to those found in students aged 17–21 years assessed at three time points over one year ($r = 0.25–0.39$)⁵⁶ and to those found in children aged 1, 3, 5, and 8 years, in whom HCC was assessed over periods of two and three years ($r = 0.30–0.44$)⁵⁷. In samples with a higher mean age (>30 years), in which intervals of assessment ranged between one month and one year, higher stability measures ($r = 0.68–0.86$) have been reported^{17,58}. These findings are in line with the hypothesis of greater changes of HPA axis functioning during childhood and adolescence which are periods marked by dramatic physical, cognitive, social and emotional changes⁵⁶. Unfortunately, no conclusions can be drawn from the present analyses concerning the heritability of the intra-individual change in HCC over time, since the respective subsample was small and lacked sufficient statistical power for this analysis. However, it is of interest to note, that the heritability estimate of HCC did not differ between the younger and older half of the sample.

Regarding the phenotypic and genetic correlations of HCC with psychological variables, the present analyses demonstrated that shared genetic factors underlie the association between the psychological variables perceived stress, depressive symptoms, and neuroticism. The heritability of neuroticism and depression^{59–61}, as well as their genetic overlap^{62,63}, are well established, and the present data are consistent with those of previous reports.

However, contrary to the results of our previous pilot study³⁸, no significant phenotypic or genetic overlap was found between HCC and any of the three psychological variables. Although no study to date has investigated the association between HCC and neuroticism, the present results are consistent with a previous investigation of 125 twin pairs, which showed heritability for neuroticism and morning cortisol secretion in saliva but no phenotypic or genetic overlap between these two variables³³.

The present findings must be viewed with caution, as they are derived from a cohort of relatively healthy adolescents from the general population. It is possible that the association between HCC and psychological variables may only become apparent for more extreme psychological phenotypes. While studies of subjects with a history of chronic or traumatic psychological disturbance, i.e., conditions which are known to alter HPA axis functioning (e.g., shift work, earthquake, or civil war), have repeatedly reported HCC alterations^{64–66}, previous reports on the association between HCC and psychological variables in healthy subjects are inconsistent (e.g. refs^{9,67–72}). Recently, this was reflected in a meta-analytic study, which found no evidence for a positive association of HCC with subjective stress and depression²⁴. However, the meta-analysis confirmed that HCC is increased in conditions of chronic ongoing stress, and therefore represents a marker for the assessment of long-term alterations in cortisol levels in response to environmental influences²⁴.

In the present sample, we observed values within the normal range of the applied psychological scales (see Supp. Table 3). If significant alterations in HCC only occur in response to major (and chronic) stressors, genetic and phenotypic correlations between HCC and psychological variables may only become observable in subjects who display more pronounced phenotypes, such as full-blown psychiatric disorders, or who are subjected to extreme levels of stress. Therefore, future genetic studies should investigate groups that are extreme in terms of perceived stress (e.g. after traumatic events), or twin pairs in which at least one twin has severe mental health problems.

Additionally, such studies should aim to further dissect the factors underlying HPA axis deregulation by including -besides basal measures such as HCC- dynamic measures of HPA axis (re-)activity, such as circadian rhythm, and reactivity to psychological, physical and pharmacological challenges, which are commonly assessed using multiple saliva or blood samples within a defined sampling scheme. Those measures have been shown to be altered in psychiatric disorders, partially independent of or even contrary to alterations in basal cortisol levels (e.g. as measured in saliva or blood)^{73,74}. Another aspect future studies should address is the interplay of HCC with other hormones known to affect HPA axis activity, which can also be measured in hair, such as cortisone or gonadal steroids.

The molecular genetic approach i.e. using polygenic risk scores to investigate the association between HCC and psychological measures generated several interesting results. First, the observation of nominally significant positive associations of the PRS for MDD and the PRS for neuroticism with the psychological variables is consistent with reported formal genetic correlations. This supports the validity of the approach, even in a sample as small as that used in the present analyses. However, the relatively low degree of explained variance, and the fact that the associations only achieved nominal significance at some of the selected thresholds, highlights that our approach lacks sufficient power. Second, the observation that neither the PRS for MDD nor the PRS for neuroticism predicted HCC, and that the PRS for plasma cortisol did not predict perceived stress, depressive symptoms, or neuroticism, parallels the genetic results from twin models, which suggests a lack of genetic overlap between HCC and the psychological variables. Interestingly, no overlap was found between the PRS for plasma cortisol and HCC. For the interpretation of this result it is important to note, that hair cortisol is assumed to reflect free (unbound) cortisol, while the GWAS investigated total (bound and unbound) plasma cortisol concentrations^{12,13}. The main signal of the GWAS was observed in the region coding for CBG, the main regulator of the ratio of free and total cortisol²⁹ and might thus differentially affect measures of free and total cortisol. Additionally, HCC represents a measure of accumulated long-term cortisol secretion, while the GWAS was based on a one-time measure of morning plasma cortisol. As described, the correlations between HCC and one time measures of cortisol in saliva and blood are inconsistent. However, in view of the limited power, the results of the PRS analyses must be interpreted with caution, and large future studies of easier-to-recruit, unrelated subjects might generate insights into the associations. Additionally, even though each of the GWAS considered in the present analyses involved cohorts in excess of 10,000 subjects and identified genome-wide genetic variants, larger studies are required. Research has demonstrated that for complex phenotypes, GWAS involving several 100,000 subjects are needed to identify the majority of the common polygenic variation^{54,75}. The power of PRS calculated using future GWAS will be increased due to a superior signal to noise ratio in these larger datasets.

Limitations

The present findings should be interpreted with caution, since the study had several limitations. First, the sample size was relatively small in terms of the establishment of twin models, particularly in the case of the exploratory analyses investigating subsamples divided by age, and the detection of small correlations between HCC measures and psychological phenotypes. Second, generalizability of the results to the general population is limited, since a young and relatively healthy cohort was investigated using self-rating questionnaires. Phenotypic and genetic correlations with psychological variables may only become evident in cohorts with more pronounced or specified environmental impacts (e.g. chronic severe stress) and more extreme phenotypes (e.g. psychiatric disorders, biologically relevant endophenotypes). Under the challenge of more adverse environments, stronger variance might occur in those phenotypes, partially driven by distinct genetic factors. Third, we did control for age and sex and interactions in our analyses. However, there is evidence that the influence of puberty processes on the associations between HPA axis activity with stress and depressive symptoms is best accounted for by assessing pubertal status and timing (e.g. refs^{76–78}). Future studies of HCC in adolescents should consider including those measures. Fourth, the self-rating questionnaires for perceived stress and depression do address shorter time frames (last week to last few weeks) than the time frame reflected in the 3 cm segments of hair analyzed for hair cortisol (~3

months). However, we observed a high heritability and stability (over two years) of the psychological measures in our sample. This indicates that these measures largely assess more stable components of the underlying psychological constructs and not merely short term fluctuations. Here, however, we must acknowledge the difficulty for researchers in this area of finding state effects convincingly independent of trait disposition. Fifth, the PRS scores for neuroticism, MDD, and plasma cortisol were derived from adult cohorts, and statistical power was small due to the limited size of the learning and present cohorts. Replication studies in much larger cohorts are required before further conclusions can be drawn, in particular as regards the genetic overlap between plasma cortisol and HCC. Sixth, although the present analyses demonstrate that HCC can be assessed with a high degree of reliability, the quality control analysis demonstrated that various factors influenced HCC, including batch number, season, storage time, and study phase, and had to be statistically controlled for. Fifth, these corrections may have contributed to an overcorrection, and thus to the overlooking of true findings. Seventh, we did not systematically obtain data on medical history or treatment, including oral contraceptives. However, as the major part of our sample is less than 16 years of age, it is unlikely that oral contraceptives and other medications represent a major confounder. Furthermore, unlike cortisol assessed in other tissues⁶, there is no strong evidence that oral contraceptives have a substantial effect on HCC (e.g. refs^{23,24,79}). Eighth, hair dyeing did not affect HCC in the present sample. However, further variables such as frequency of hair washing were not assessed.

Conclusions

Our study demonstrates a high heritability of HCC, but no evidence for a genetic overlap with depressive symptoms, perceived stress and neuroticism. HCC is a reliable measure of long-term HPA axis activity and genetic effects play a major role in inter-individual variability. This knowledge will inform future research, and is of particular relevance in terms of the interpretation of data from cross-sectional studies. If a genetic or phenotypic correlation does exist between HCC and perceived stress, depressive symptoms, or neuroticism, the present analyses have demonstrated that this is difficult to identify in a relatively small sample of young adults from the general population. Further studies are warranted to investigate whether this is also the case in samples with more extreme psychological phenotypes. Particularly in studies for which blood or saliva sampling over several days and multiple time-points is difficult to implement, HCC represents a promising alternative measure for the assessment of long-term HPA axis activation. The high heritability, easy accessibility and cost-effectiveness of HCC render it a promising target for future large scale GWAS of the biological pathways that underlie long-term cortisol secretion and its links to stress-related phenotypes.

References

- Mah, L., Szabuniewicz, C. & Fiocco, A. J. Can anxiety damage the brain? *Current opinion in psychiatry* **29**, 56–63 (2016).
- de Kloet, E. R., Joels, M. & Holsboer, F. Stress and the brain: from adaptation to disease. *Nat Rev Neurosci* **6**, 463–475, <https://doi.org/10.1038/nrn1683> (2005).
- Chrousos, G. P. Stress and disorders of the stress system. *Nat Rev Endocrinol* **5**, 374–381, <https://doi.org/10.1038/nrendo.2009.106> (2009).
- Lupien, S. J., McEwen, B. S., Gunnar, M. R. & Heim, C. Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nat Rev Neurosci* **10**, 434–445, <https://doi.org/10.1038/nrn2639> (2009).
- Studte-Schmiedgen, S., Kirschbaum, C., Alexander, N. & Stalder, T. An integrative model linking traumatization, cortisol dysregulation and posttraumatic stress disorder: Insight from recent hair cortisol findings. *Neuroscience & Biobehavioral Reviews* **69**, 124–135, <https://doi.org/10.1016/j.neubiorev.2016.07.015> (2016).
- Kudielka, B. M., Hellhammer, D. H. & Wüst, S. Why do we respond so differently? Reviewing determinants of human salivary cortisol responses to challenge. *Psychoneuroendocrinology* **34**, 2–18, <https://doi.org/10.1016/j.psyneuen.2008.10.004> (2009).
- Holtzman, C. W. *et al.* Stress and neurodevelopmental processes in the emergence of psychosis. *Neuroscience* **249**, 172–191, <https://doi.org/10.1016/j.neuroscience.2012.12.017> (2013).
- Levine, A., Zagoory-Sharon, O., Feldman, R., Lewis, J. G. & Weller, A. Measuring cortisol in human psychobiological studies. *Physiol Behav* **90**, 43–53, <https://doi.org/10.1016/j.physbeh.2006.08.025> (2007).
- Staufenbiel, S. M., Penninx, B. W., Spijker, A. T., Elzinga, B. M. & van Rossum, E. F. Hair cortisol, stress exposure, and mental health in humans: a systematic review. *Psychoneuroendocrinology* **38**, 1220–1235, <https://doi.org/10.1016/j.psyneuen.2012.11.015> (2013).
- Meyer, J. S. & Novak, M. A. Minireview: hair cortisol: a novel biomarker of hypothalamic-pituitary-adrenocortical activity. *Endocrinology* **153**, 4120–4127 (2012).
- Gow, R., Thomson, S., Rieder, M., Van Uum, S. & Koren, G. An assessment of cortisol analysis in hair and its clinical applications. *Forensic Sci Int* **196**, 32–37, <https://doi.org/10.1016/j.forsciint.2009.12.040> (2010).
- Dettenborn, L., Tietze, A., Kirschbaum, C. & Stalder, T. The assessment of cortisol in human hair: associations with sociodemographic variables and potential confounders. *Stress* **15**, 578–588, <https://doi.org/10.3109/10253890.2012.654479> (2012).
- Stalder, T. & Kirschbaum, C. Analysis of cortisol in hair—state of the art and future directions. *Brain Behav Immun* **26**, 1019–1029, <https://doi.org/10.1016/j.bbi.2012.02.002> (2012).
- D'Anna-Hernandez, K. L., Ross, R. G., Natvig, C. L. & Laudenslager, M. L. Hair cortisol levels as a retrospective marker of hypothalamic-pituitary axis activity throughout pregnancy: comparison to salivary cortisol. *Physiol Behav* **104**, 348–353, <https://doi.org/10.1016/j.physbeh.2011.02.041> (2011).
- Xie, Q. *et al.* Correlation of cortisol in 1-cm hair segment with salivary cortisol in human: hair cortisol as an endogenous biomarker. *Clin Chem Lab Med* **49**, 2013–2019, <https://doi.org/10.1515/cclm.2011.706> (2011).
- van Ockenburg, S. L. *et al.* The relationship between 63 days of 24-h urinary free cortisol and hair cortisol levels in 10 healthy individuals. *Psychoneuroendocrinology* **73**, 142–147, <https://doi.org/10.1016/j.psyneuen.2016.07.220> (2016).
- Short, S. J. *et al.* Correspondence between hair cortisol concentrations and 30-day integrated daily salivary and weekly urinary cortisol measures. *Psychoneuroendocrinology* **71**, 12–18, <https://doi.org/10.1016/j.psyneuen.2016.05.007> (2016).
- Vanaelst, B. *et al.* Intercorrelations between serum, salivary, and hair cortisol and child-reported estimates of stress in elementary school girls. *Psychophysiology* **49**, 1072–1081, <https://doi.org/10.1111/j.1469-8986.2012.01396.x> (2012).
- Sauve, B., Koren, G., Walsh, G., Tokmakejian, S. & Van Uum, S. H. Measurement of cortisol in human hair as a biomarker of systemic exposure. *Clin Invest Med* **30**, E183–191 (2007).
- Hodes, A. *et al.* Hair cortisol in the evaluation of Cushing syndrome. *Endocrine* **56**, 164–174, <https://doi.org/10.1007/s12020-017-1231-7> (2017).

21. Kirschbaum, C., Tietze, A., Skoluda, N. & Dettenborn, L. Hair as a retrospective calendar of cortisol production—Increased cortisol incorporation into hair in the third trimester of pregnancy. *Psychoneuroendocrinology* **34**, 32–37, <https://doi.org/10.1016/j.psyneuen.2008.08.024> (2009).
22. Manenshijn, L. *et al.* A novel tool in the diagnosis and follow-up of (cyclic) Cushing's syndrome: measurement of long-term cortisol in scalp hair. *J Clin Endocrinol Metab* **97**, E1836–1843, <https://doi.org/10.1210/jc.2012-1852> (2012).
23. Wosu, A. C., Valdimarsdóttir, U., Shields, A. E., Williams, D. R. & Williams, M. A. Correlates of cortisol in human hair: implications for epidemiologic studies on health effects of chronic stress. *Annals of epidemiology* **23**, 797–811. e792 (2013).
24. Stalder, T. *et al.* Stress-related and basic determinants of hair cortisol in humans: A meta-analysis. *Psychoneuroendocrinology* **77**, 261–274, <https://doi.org/10.1016/j.psyneuen.2016.12.017> (2017).
25. Bartels, M., V den Berg, M., Sluyter, F., Boomsma, D. I. & de Geus, E. J. Heritability of cortisol levels: review and simultaneous analysis of twin studies. *Psychoneuroendocrinology* **28**, 121–137 (2003).
26. Van Hulle, C. A., Shirtcliff, E. A., Lemery-Chalfant, K. & Goldsmith, H. H. Genetic and environmental influences on individual differences in cortisol level and circadian rhythm in middle childhood. *Hormones and behavior* **62**, 36–42 (2012).
27. Ouellet-Morin, I. *et al.* Evidence of a unique and common genetic etiology between the CAR and the remaining part of the diurnal cycle: A study of 14 year-old twins. *Psychoneuroendocrinology* **66**, 91–100, <https://doi.org/10.1016/j.psyneuen.2015.12.022> (2016).
28. Steptoe, A., van Jaarsveld, C. H., Semmler, C., Plomin, R. & Wardle, J. Heritability of daytime cortisol levels and cortisol reactivity in children. *Psychoneuroendocrinology* **34**, 273–280, <https://doi.org/10.1016/j.psyneuen.2008.09.006> (2009).
29. Bolton, J. L. *et al.* Genome wide association identifies common variants at the SERPINA6/SERPINA1 locus influencing plasma cortisol and corticosteroid binding globulin. *PLoS Genet* **10**, e1004474, <https://doi.org/10.1371/journal.pgen.1004474> (2014).
30. Fairbanks, L. A. *et al.* Heritability and genetic correlation of hair cortisol in vervet monkeys in low and higher stress environments. *Psychoneuroendocrinology* **36**, 1201–1208, <https://doi.org/10.1016/j.psyneuen.2011.02.013> (2011).
31. Tucker-Drob, E. M. *et al.* Genetic influences on hormonal markers of chronic hypothalamic-pituitary-adrenal function in human hair. *Psychol Med*, 1–13, doi:<https://doi.org/10.1017/s0033291716003068> (2017).
32. Young, E. A., Aggen, S. H., Prescott, C. A. & Kendler, K. S. Similarity in saliva cortisol measures in monozygotic twins and the influence of past major depression. *Biol Psychiatry* **48**, 70–74 (2000).
33. Riese, H., Rijdsdijk, F. V., Rosmalen, J. G., Snieder, H. & Ormel, J. Neuroticism and morning cortisol secretion: both heritable, but no shared genetic influences. *J Pers* **77**, 1561–1575, <https://doi.org/10.1111/j.1467-6494.2009.00592.x> (2009).
34. Rietschel, L. *et al.* Perceived Stress has Genetic Influences Distinct from Neuroticism and Depression. *Behav Genet*, <https://doi.org/10.1007/s10519-013-9636-4> (2013).
35. Smith, D. J. *et al.* Genome-wide analysis of over 106 000 individuals identifies 9 neuroticism-associated loci. *Mol Psychiatry* **21**, 749–757, <https://doi.org/10.1038/mp.2016.49> (2016).
36. Okbay, A. *et al.* Genetic variants associated with subjective well-being, depressive symptoms and neuroticism identified through genome-wide analyses. *Nat Genet* **48**, 624–633, <https://doi.org/10.1038/ng.3552> (2016).
37. Consortium, M. D. D. W. G. o. t. P. G. Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depressive disorder. (submitted).
38. Rietschel, L. *et al.* Hair Cortisol and Its Association With Psychological Risk Factors for Psychiatric Disorders: A Pilot Study in Adolescent Twins. *Twin Research and Human Genetics* **19**, 438–446 (2016).
39. Gillespie, N. A., *et al.* The Brisbane Longitudinal Twin Study: pathways to Cannabis Use, Abuse, and Dependence project—current status, preliminary results, and future directions. *Twin Research and Human Genetics*, **16**(1), 21–33 ISO 690 (2013).
40. Wright, M. J. & Martin, N. G. Brisbane adolescent twin study: outline of study methods and research projects. *Australian Journal of Psychology* **56**, 65–78 (2004).
41. Kearney, C. A., Drabman, R. S. & Beasley, J. F. The trials of childhood: The development, reliability, and validity of the Daily Life Stressors Scale. *Journal of Child and Family Studies* **2**, 371–388 (1993).
42. Eysenck, S. Junior Eysenck Personality Inventory: EdITS/Educational and Industrial Testing Service. *PO Box 7234*, 92167 (1972).
43. Cohen, S., Kamarck, T. & Mermelstein, R. A global measure of perceived stress. *Journal of health and social behavior*, 385–396 (1983).
44. McCrae, R. R. & Costa, P. T. A contemplated revision of the NEO Five-Factor Inventory. *Personality and individual differences* **36**, 587–596 (2004).
45. Hickie, I. B. *et al.* Development of a simple screening tool for common mental disorders in general practice. *The Medical journal of Australia* **175**, S10–17 (2001).
46. Binz, T. M., Braun, U., Baumgartner, M. R. & Kraemer, T. Development of an LC-MS/MS method for the determination of endogenous cortisol in hair using (13)C3-labeled cortisol as surrogate analyte. *J Chromatogr B Analyt Technol Biomed Life Sci* **1033–1034**, 65–72, <https://doi.org/10.1016/j.jchromb.2016.07.041> (2016).
47. Rietschel, L. *et al.* Perceived stress has genetic influences distinct from neuroticism and depression. *Behavior genetics* **44**, 639–645 (2014).
48. Hansell, N. K. *et al.* Genetic co-morbidity between neuroticism, anxiety/depression and somatic distress in a population sample of adolescent and young adult twins. *Psychol Med* **42**, 1249–1260, <https://doi.org/10.1017/s0033291711002431> (2012).
49. Neale, M., Boker, S., Xie, G. & Maes, H. Mx: Statistical modeling. Richmond, VA: Department of Psychiatry. Virginia Institute for Psychiatric and Behavior Genetics, Virginia Commonwealth University (2003).
50. Heath, A. C. & Martin, N. G. Psychoticism as a dimension of personality: a multivariate genetic test of Eysenck and Eysenck's psychoticism construct. *J Pers Soc Psychol* **58**, 111–121 (1990).
51. van der Voorn, B., Hollanders, J. J., Ket, J. C. F., Rotteveel, J. & Finken, M. J. J. Gender-specific differences in hypothalamus-pituitary-adrenal axis activity during childhood: a systematic review and meta-analysis. *Biology of Sex Differences* **8**, 3, <https://doi.org/10.1186/s13293-016-0123-5> (2017).
52. Neale, M. & Cardon, L. *Methodology for genetic studies of twins and families*. Vol. 67 (Springer Science & Business Media, 1992).
53. McCarthy, S. *et al.* A reference panel of 64,976 haplotypes for genotype imputation. *bioRxiv*, 035170 (2016).
54. Wray, N. R. *et al.* Research review: Polygenic methods and their application to psychiatric traits. *J Child Psychol Psychiatry* **55**, 1068–1087, <https://doi.org/10.1111/jcpp.12295> (2014).
55. Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. Genome-wide complex trait analysis (GCTA): methods, data analyses, and interpretations. *Methods Mol Biol* **1019**, 215–236, https://doi.org/10.1007/978-1-62703-447-0_9 (2013).
56. Zhang, Q., Chen, Z., Chen, S., Xu, Y. & Deng, H. Intraindividual stability of cortisol and cortisone and the ratio of cortisol to cortisone in saliva, urine and hair. *Steroids* **118**, 61–67, <https://doi.org/10.1016/j.steroids.2016.12.008> (2017).
57. Karlen, J., Frostell, A., Theodorsson, E., Faresjo, T. & Ludvigsson, J. Maternal influence on child HPA axis: a prospective study of cortisol levels in hair. *Pediatrics* **132**, e1333–1340, <https://doi.org/10.1542/peds.2013-1178> (2013).
58. Stalder, T. *et al.* Intraindividual stability of hair cortisol concentrations. *Psychoneuroendocrinology* **37**, 602–610, <https://doi.org/10.1016/j.psyneuen.2011.08.007> (2012).
59. McGuffin, P., Katz, R., Watkins, S. & Rutherford, J. A hospital-based twin register of the heritability of DSM-IV unipolar depression. *Arch Gen Psychiatry* **53**, 129–136 (1996).
60. Jang, K. L., Livesley, W. J. & Vernon, P. A. Heritability of the big five personality dimensions and their facets: a twin study. *J Pers* **64**, 577–591 (1996).

61. Sullivan, P. F., Neale, M. C. & Kendler, K. S. Genetic epidemiology of major depression: review and meta-analysis. *Am J Psychiatry* **157**, 1552–1562 (2000).
62. Hettema, J. M., Neale, M. C., Myers, J. M., Prescott, C. A. & Kendler, K. S. A population-based twin study of the relationship between neuroticism and internalizing disorders. *Am J Psychiatry* **163**, 857–864, <https://doi.org/10.1176/ajp.2006.163.5.857> (2006).
63. Khan, A. A., Jacobson, K. C., Gardner, C. O., Prescott, C. A. & Kendler, K. S. Personality and comorbidity of common psychiatric disorders. *Br J Psychiatry* **186**, 190–196, <https://doi.org/10.1192/bjp.186.3.190> (2005).
64. Etwel, F., Russell, E., Rieder, M. J., Van Uum, S. H. & Koren, G. Hair cortisol as a biomarker of stress in the 2011 Libyan war. *Clin Invest Med* **37**, E403–408 (2014).
65. Gao, W. *et al.* Temporal features of elevated hair cortisol among earthquake survivors. *Psychophysiology* **51**, 319–326, <https://doi.org/10.1111/psyp.12179> (2014).
66. Manenschi, L., van Kruysbergen, R. G., de Jong, F. H., Koper, J. W. & van Rossum, E. F. Shift work at young age is associated with elevated long-term cortisol levels and body mass index. *J Clin Endocrinol Metab* **96**, E1862–1865, <https://doi.org/10.1210/jc.2011-1551> (2011).
67. Heinze, K., Lin, A., Reniers, R. L. & Wood, S. J. Longer-term increased cortisol levels in young people with mental health problems. *Psychiatry research* **236**, 98–104 (2016).
68. Streit, F. *et al.* Perceived stress and hair cortisol: Differences in bipolar disorder and schizophrenia. *Psychoneuroendocrinology* **69**, 26–34 (2016).
69. Rippe, R. C. *et al.* Splitting hair for cortisol? Associations of socio-economic status, ethnicity, hair color, gender and other child characteristics with hair cortisol and cortisone. *Psychoneuroendocrinology* **66**, 56–64, <https://doi.org/10.1016/j.psychneuen.2015.12.016> (2016).
70. Simmons, J. G. *et al.* The lifetime experience of traumatic events is associated with hair cortisol concentrations in community-based children. *Psychoneuroendocrinology* **63**, 276–281 (2016).
71. Vliegthart, J. *et al.* Socioeconomic status in children is associated with hair cortisol levels as a biological measure of chronic stress. *Psychoneuroendocrinology* **65**, 9–14 (2016).
72. Vives, A. H. *et al.* Cortisol levels in fingernails and neurocognitive performance in euthymic bipolar I patients and healthy controls. *Psychoneuroendocrinology* **61**, 70, <https://doi.org/10.1016/j.psychneuen.2015.07.583> (2015).
73. Pochigaeva, K. *et al.* Hair cortisol as a marker of hypothalamic-pituitary-adrenal Axis activity in female patients with major depressive disorder. *Metab Brain Dis* **32**, 577–583, <https://doi.org/10.1007/s11011-017-9952-0> (2017).
74. Kuehl, L. K. *et al.* Hair cortisol and cortisol awakening response are associated with criteria of the metabolic syndrome in opposite directions. *Psychoneuroendocrinology* **51**, 365–370, <https://doi.org/10.1016/j.psychneuen.2014.09.012> (2015).
75. Dudbridge, F. Power and predictive accuracy of polygenic risk scores. *PLoS Genet* **9**, e1003348 (2013).
76. Sun, Y. *et al.* Longitudinal pattern of early maturation on morning cortisol and depressive symptoms: Sex-specific effects. *Psychoneuroendocrinology* **71**, 58–63, <https://doi.org/10.1016/j.psychneuen.2016.04.026> (2016).
77. Trepanier, L. *et al.* Early menarche predicts increased depressive symptoms and cortisol levels in Quebec girls ages 11 to 13. *Dev Psychopathol* **25**, 1017–1027, <https://doi.org/10.1017/s0954579413000345> (2013).
78. King, L. S. *et al.* The impact of the severity of early life stress on diurnal cortisol: The role of puberty. *Psychoneuroendocrinology* **77**, 68–74, <https://doi.org/10.1016/j.psychneuen.2016.11.024> (2017).
79. Staufenbiel, S. M., Penninx, B. W., de Rijke, Y. B., van den Akker, E. L. & van Rossum, E. F. Determinants of hair cortisol and hair cortisone concentrations in adults. *Psychoneuroendocrinology* **60**, 182–194, <https://doi.org/10.1016/j.psychneuen.2015.06.011> (2015).

Acknowledgements

The authors thank the participants for their cooperation and samples. We are grateful to Natalie Garden and Reshika Chand for data collection and to Kerrie McAloney for study coordination. We thank Alisha Hall and Christine Schmal for their critical reading of, and suggestions for, the manuscript. GWAS results for calculating neuroticism polygenic risk scores were provided by the UK Biobank. BCD is supported by a University of Queensland International scholarship. LCC is supported by a QIMR Berghofer fellowship. The CORTisol NETwork Consortium was funded by the Chief Scientist Office of the Scottish Government (grant CZB-4-733) and the British Heart Foundation (grant RG11/4/28734). Andrew Crawford is funded by the Wellcome Trust (Senior Investigator Award to BR Walker; 107049/Z/15/Z). Funding for the collection of twin hair samples was provided by Australian NHMRC (National Health and Medical Research Council (AU)) grants to NGM (APP1049911) and to MJW (APP1009064). Hair cortisol assays were funded by a grant to MR from the German Federal Ministry of Education and Research (BMBF), through the Integrated Network IntegraMent (Integrated Understanding of Causes and Mechanisms in Mental Disorders, grant BMBF01ZX1314G, BMBF01ZX1614G), under the auspices of the e:Med Programme.

Author Contributions

Contributed to conception and design: L. Rietschel, F. Streit, S. Wüst, M.M. Nöthen, A.J. Forstner, T.G. Schulze, S.H. Witt, N.A. Gillespie, J. McGrath, K. McAloney, I.B. Hickie, N.K. Hansell, M.J. Wright, N.G. Martin, B.R. Walker, A.A. Crawford, M. Rietschel. Collected data: N.A. Gillespie, J. McGrath, K. McAloney, I.B. Hickie, N.K. Hansell, M.J. Wright, N.G. Martin. Analyzed the data: L. Rietschel, F. Streit, G. Zhu, L. Colorado Conde, B. Couvy-Duchesne, S.E. Medland, T.M. Binz, M.R. Baumgartner, J. Frank, N.G. Martin, M. Rietschel. Wrote the manuscript: L. Rietschel, F. Streit, L. Colodro-Conde, N.G. Martin, M. Rietschel. Provided GWAS results for the calculation of the PRS: CORTisolNETwork (CORNET) Consortium# and Major Depressive Disorder Working Group of the PGC. All authors critically revised the manuscript and have approved the final article.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-017-11852-3>.

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017

Consortia

CORTisolNETwork (CORNET) Consortium

Jennifer L. Bolton²¹, Caroline Hayward²³, Nese Direk^{24,25}, Anna Anderson²¹, Jennifer Huffman²³, James F. Wilson²⁶, Harry Campbell²⁶, Igor Rudan²⁶, Alan Wright²³, Nicholas Hastie²³, Sarah H. Wild²⁶, Fleur P. Velders²⁴, Albert Hofman²⁴, Andre G. Uitterlinden^{24,27}, Jari Lahti²⁸, Katri Räikkönen²⁸, Eero Kajantie²⁹, Elisabeth Widen³⁰, Aarno Palotie^{30,31}, Johan G. Eriksson^{29,32,33,34,35}, Marika Kaakinen³⁶, Marjo-Riitta Järvelin^{36,37,38,39}, Nicholas J. Timpson⁴⁰, George Davey Smith⁴⁰, Susan M. Ring⁴¹, David M. Evans⁴⁰, Beate St Pourcain⁴¹, Toshiko Tanaka⁴², Yuri Milaneschi^{42,43}, Stefania Bandinelli⁴⁴, Luigi Ferrucci⁴², Pim van der Harst^{45,46,47}, Judith GM Rosmalen⁴⁸, Stephen JL Bakker⁴⁹, Niek Verweij⁴⁵, Robin PF Dullaart⁴⁹, Anubha Mahajan⁵⁰, Cecilia M. Lindgren⁵⁰, Andrew Morris⁵⁰, Lars Lind⁵¹, Erik Ingelsson⁵¹, Laura N. Anderson⁵², Craig E. Pennell⁵³, Stephen J. Lye⁵², Stephen G. Matthews⁵⁴, Joel Eriksson⁵⁵, Dan Mellstrom⁵⁵, Claes Ohlsson⁵⁵, Jackie F. Price²⁶, Mark WJ Strachan²¹, Rebecca M. Reynolds²¹ & Henning Tiemeier^{24,56,57}

Major Depressive Disorder Working Group of the Psychiatric Genomics Consortium (PGC)

Stephan Ripke^{58,59,60}, Manuel Mattheisen^{61,62,63}, Abdel Abdellaoui⁶⁴, Mark J. Adams⁶⁵, Esben Agerbo^{66,8,67}, Tracy M. Air⁶⁸, Till FM Andlauer^{69,70}, Silviu-Alin Bacanu⁷¹, Marie Bækvad-Hansen^{67,72}, Aartjan TF Beekman⁴³, David A. Bennett⁷³, Klaus Berger⁷⁴, Tim B. Bigdeli^{71,11}, Jonas Bybjerg-Grauholm^{67,72}, Enda M. Byrne⁵, Na Cai⁷⁵, Enrique Castela⁷⁶, Toni-Kim Clarke⁶⁵, Jonathan RI Coleman⁷⁷, Converge Consortium⁷⁸, Nick Craddock⁷⁹, Udo Dannlowski^{80,81}, Gareth Davies⁸², Gail Davies⁸³, Eco. J. C. de Geus^{64,84}, Philip De Jager⁸⁵, Ian J. Deary⁸³, Franziska Degenhardt^{12,13}, Erin C. Dunn^{86,87,88}, Erik A. Ehli⁸², Thalia C. Eley⁷⁷, Valentina Escott-Price⁸⁹, Tõnu Esko^{58,90,91,92}, Hilary K. Finucane^{93,94}, Michael Gill⁹⁵, Scott D. Gordon⁹⁶, Jakob Grove^{61,62,67,97}, Lynsey S. Hall^{65,98}, Thomas F. Hansen^{99,100}, Christine Søholm Hansen^{67,72}, Thomas F. Hansen¹⁰¹, Andrew C. Heath¹⁰², Anjali K. Henders⁵, Stefan Herms^{12,13,15}, Per Hoffmann^{12,13,15}, Georg Homuth¹⁰³, Carsten Horn¹⁰⁴, Jouke- Jan Hottenga⁶⁴, David Hougaard^{67,72}, Hailiang Huang^{59,86,105}, Marcus Ising¹⁰⁶, Rick Jansen⁴³, Eric Jorgenson¹⁰⁷, Stefan Kloiber^{108,109}, James A Knowles¹¹⁰, Warren W. Kretschmar⁵⁰, Jesper Krogh¹¹¹, Zoltán Kutalik^{112,113}, Maren Lang³, Glyn Lewis¹¹⁴, Yihan Li⁵⁰, Donald J. MacIntyre^{115,116}, Pamela AF Madden¹⁰², Jonathan Marchini¹¹⁷, Hamdi Mbarek^{118,64}, Peter McGuffin⁷⁷, Divya Mehta¹¹⁹, Andres Metspalu^{92,120}, Christel M. Middeldorp⁶⁴, Evelin Mihailov^{92,121}, Lili Milani⁹², Grant W. Montgomery¹²², Sara Mostafavi^{123,124}, Niamh Mullins⁷⁷, Matthias Nauck^{125,126}, Bernard Ng¹²⁴, Merete Nordentoft^{67,127}, Dale R. Nyholt¹²⁸, Michael C. O'Donovan¹²⁹, Paul F. O'Reilly⁷⁷, Hogni Oskarsson¹³⁰, Michael J. Owen¹²⁹, Sara A. Paciga¹³¹, Carsten Bøcker Pedersen^{66,67,8}, Marianne Giørtz Pedersen^{66,67,8}, Nancy L. Pedersen¹³², Michele L. Pergadia¹³³, Roseann E. Peterson^{11,71}, Erik Pettersson¹³⁴, Wouter J. Peyrot⁴³, David J. Porteous¹³⁵, Danielle Posthuma^{136,137}, James B. Potash¹³⁸, Jorge A. Quiroz¹³⁹, John P. Rice¹⁰², Brien P. Riley⁷¹, Margarita Rivera^{77,140}, Douglas M. Ruderfer¹⁴¹, Saira Saeed Mirza²⁴, Robert Schoevers¹⁴², Ling Shen¹⁰⁷, Jianxin Shi¹⁴³, Engilbert Sigurdsson¹⁴⁴, Grant CB Sinnamoni¹⁴⁵, Johannes H. Smit⁴³, Daniel J. Smith¹⁴⁶, Jordan W. Smoller⁸⁶, Hreinn Stephansson¹⁴⁷, Stacy Steinberg¹⁴⁷, Jana Strohmaier³, Katherine E. Tansey¹⁴⁸, Alexander Teumer¹⁴⁹, Wesley Thompson^{100,135,150,151,152}, Pippa A. Thomson¹³⁵, Thorgeir E. Thorgeirsson¹⁴⁷, Jens Treutlein³, Maciej Trzaskowski¹⁵³, Daniel Umbricht¹⁵⁴, Sandra van der Auwera¹⁵⁵, Gerard van Grootheest⁴³, Albert M. van Hemert¹⁵⁶, Alexander Viktorin¹³², Henry Völzke¹⁴⁹, Yunpeng Wang^{67,100,151}, Bradley T. Webb¹⁵⁷, Myrna M. Weissman^{158,159}, Jürgen Wellmann⁷⁴, Gonneke Willemsen⁶⁴, Hualin S. Xi¹⁶⁰, Bernhard T. Baune⁶⁸, Douglas H. R. Blackwood⁶⁵, Dorret I. Boomsma⁶⁴, Anders D. Børglum^{61,62,67}, Henriette N. Buttenschøn^{62,67,161}, Sven Cichon^{12,162,163,164}, Enrico Domenici¹⁶⁵, Jonathan Flint^{50,166}, Hans J. Grabe¹⁵⁵, Steven P. Hamilton¹⁶⁷, Kenneth S. Kendler⁷¹, Qingqin S. Li¹⁶⁸, Susanne Lucae¹⁰⁶, Patrik K. Magnusson¹³², Andrew M. McIntosh^{65,83}, Ole Mors^{67,169}, Preben Bo Mortensen^{62,8,67}, Bertram Müller-Myhsok^{69,70,170}, Brenda WJH Penninx⁴³, Roy H. Perlis^{87,171}, Martin Preisig⁷⁶, Catherine Schaefer¹⁰⁷, Jordan W. Smoller^{87,88}, Kari Stephansson¹⁴⁷, Henning Tiemeier^{24,56,57}, Rudolf Uher¹⁷², Thomas Werge^{100,150,173}, Ashley R. Winslow^{174,175}, Gerome Breen^{77,176}, Douglas F. Levinson¹⁷⁷, Cathryn M. Lewis^{77,178}, Naomi R. Wray^{5,153} & Patrick F. Sullivan^{134,179,180}

²³MRC Human Genetics Unit, Institute for Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, EH4 2XU, UK. ²⁴Department of Epidemiology, Erasmus Medical Centre, Rotterdam, Netherlands. ²⁵Psychiatry, Dokuz Eylul University School Of Medicine, Izmir, TR, Turkey. ²⁶Centre for Population Health Sciences, Institute for Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, EH8 9AG, UK. ²⁷Internal Medicine, Erasmus MC, Rotterdam, NL, Netherlands. ²⁸Institute of Behavioural Sciences, University of Helsinki, Helsinki, Finland. ²⁹National Institute for Health and Welfare, Helsinki, Finland. ³⁰Institute for Molecular Medicine Finland (FIMM), University of Helsinki, Helsinki, Finland. ³¹Department of Medical Genetics, University of Helsinki and University Central Hospital, Helsinki, Finland. ³²Department of General Practice and Primary Health Care, University of Helsinki, Helsinki, Finland. ³³Helsinki University Central Hospital, Unit of General Practice, Helsinki, Finland. ³⁴Folkhalsan Research Centre, Helsinki, Finland. ³⁵Vasa Central Hospital, Vasa, Finland. ³⁶Institute of Health Sciences and Biocenter Oulu, University of Oulu, Oulu, Finland. ³⁷Department of Children and Yond People and Families, National Institute for Health andelfare, Oulu, Finland. ³⁸Department of Epidemiology and Biostatistics, MRC-HPA Centre for Environment and Health, Imperial College London, London, UK. ³⁹Unit of Primary Care, Oulu University Hospital, Oulu, Finland. ⁴⁰MRC Centre for Causal Analyses in Translational Epidemiology, School of Social and Community Medicine, University of Bristol, Bristol, UK. ⁴¹School of Social and Community Medicine, University of Bristol, Bristol, UK. ⁴²Longitudinal Studies Section, Clinical Research Branch, National Institute on Aging, Baltimore, MD, USA. ⁴³Department of Psychiatry, VU University Medical Center/GGZ inGeest, Amsterdam, Netherlands. ⁴⁴Geriatric Unit, ASF, Florence, Italy. ⁴⁵University of Groningen, University Medical Center Groningen, Department of Cardiology, Groningen, Netherlands. ⁴⁶University of Groningen, University Medical Center Groningen, Department of Genetics, Groningen, Netherlands. ⁴⁷Durrer Center for Cardiogenetic Research, ICIN-Netherlands Heart Institute, Utrecht, Netherlands. ⁴⁸University of Groningen, University Medical Center Groningen, Interdisciplinary Center for Psychiatric Epidemiology, Groningen, Netherlands. ⁴⁹University of Groningen, University Medical Center Groningen, Department of Internal Medicine, Groningen, Netherlands. ⁵⁰Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, UK. ⁵¹Department of Medical Sciences, Uppsala University, Uppsala, Sweden. ⁵²Samuel Lunenfeld Research Institute, Mount Sinai Hospital, Toronto, Ontario, Canada. ⁵³School of Women's and Infant's Health, The University of Western Australia, Crawley, Australia. ⁵⁴Department of Physiology, University of Toronto, Toronto, Ontario, Canada. ⁵⁵Center for Bone and Arthritis Research, Institute of Medicin, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden. ⁵⁶Child and Adolescent Psychiatry, Erasmus MC, Rotterdam, Netherlands. ⁵⁷Psychiatry, Erasmus MC, Rotterdam, Netherlands. ⁵⁸Medical and Population Genetics, Broad Institute, Cambridge, USA. ⁵⁹Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, USA. ⁶⁰Department of Psychiatry and Psychotherapy, Universitätsmedizin Berlin Campus Charité Mitte, Berlin, Germany. ⁶¹Department of Biomedicine, Aarhus University, Aarhus, Denmark. ⁶²iSEQ, Centre for Integrative Sequencing, Aarhus University, Aarhus, Denmark. ⁶³iSPYCH, The Lundbeck Foundation Initiative for Integrative Psychiatric Research, Aarhus, Denmark. ⁶⁴Dept of Biological Psychology, VU University Amsterdam, Amsterdam, Netherlands. ⁶⁵Division of Psychiatry, University of Edinburgh, Edinburgh, UK. ⁶⁶Centre for Integrated Register-based Research, Aarhus University, Aarhus, Denmark. ⁶⁷iPSYCH, The Lundbeck Foundation Initiative for Integrative Psychiatric Research, Aarhus, Denmark. ⁶⁸Discipline of Psychiatry, University of Adelaide, Adelaide, Australia. ⁶⁹Department of Translational Research in Psychiatry, Max Planck Institute of Psychiatry, Munich, Germany. ⁷⁰Munich Cluster for Systems Neurology (SyNergy), Munich, Germany. ⁷¹Department of Psychiatry, Virginia Commonwealth University, Richmond, USA. ⁷²Center for Neonatal Screening, Department for Congenital Disorders, Statens Serum Institut, Copenhagen, Denmark. ⁷³Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, USA. ⁷⁴Institute of Epidemiology and Social Medicine, University of Muenster, Muenster, UK. ⁷⁵Human Genetics, Wellcome Trust Sanger Institute, Cambridge, UK. ⁷⁶Department of Psychiatry, University Hospital of Lausanne, Prilly, Switzerland. ⁷⁷MRC Social Genetic and Developmental Psychiatry Centre, King's College London, London, UK. ⁷⁸University of Oxford, Oxford, UK. ⁷⁹Psychological Medicine, Cardiff University, Cardiff, UK. ⁸⁰Department of Psychiatry, University of Marburg, Marburg, Germany. ⁸¹Department of Psychiatry, University of Münster, Münster, Germany. ⁸²Avera Institute for Human Genetics, Sioux Falls, USA. ⁸³Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh, UK. ⁸⁴EMGO+ Institute, VU University Medical Center, Amsterdam, Netherlands. ⁸⁵Neurology, Brigham and Women's Hospital, Boston, USA. ⁸⁶Stanley Center for Psychiatric Research, Broad Institute, Cambridge, USA. ⁸⁷Department of Psychiatry, Massachusetts General Hospital, Boston, USA. ⁸⁸Psychiatric and Neurodevelopmental Genetics Unit (PNGU), Massachusetts General Hospital, Boston, USA. ⁸⁹Neuroscience and Mental Health Research Institute, Cardiff University, Cardiff, UK. ⁹⁰Division of Endocrinology, Children's Hospital Boston, Boston, USA. ⁹¹Department of Genetics, Harvard Medical School, Boston, USA. ⁹²Estonian Genome Center, University of Tartu, Tartu, Estonia. ⁹³Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, USA. ⁹⁴Department of Mathematics, Massachusetts Institute of Technology, Cambridge, USA. ⁹⁵Department of Psychiatry, Trinity College Dublin, Dublin, Ireland. ⁹⁶Genetics and Computational Biology, QIMR Berghofer Medical Research Institute, Brisbane, Australia. ⁹⁷Bioinformatics Research Centre (BiRC), Aarhus University, Aarhus, Denmark. ⁹⁸Institute of Genetic Medicine, Newcastle University, Newcastle upon Tyne, UK. ⁹⁹Danish Headache Centre, Department of Neurology, Rigshospitalet Glostrup, Glostrup, Denmark. ¹⁰⁰Institute of Biological Psychiatry, Mental Health Center Sct. Hans, Mental Health Services Capital Region of Denmark, Roskilde, Denmark. ¹⁰¹iPSYCH, The Lundbeck Foundation Initiative for Psychiatric Research, Copenhagen, Denmark. ¹⁰²Department of Psychiatry, Washington University in Saint Louis School of Medicine, Saint Louis, USA. ¹⁰³Interfaculty Institute for Genetics and Functional Genomics, Department of Functional Genomics, University Medicine and Ernst Moritz Arndt University Greifswald, Greifswald, Germany. ¹⁰⁴Roche Pharmaceutical Research and Early Development, Pharmaceutical Sciences, Roche Innovation Center Basel, F. Hoffmann-La Roche Ltd, Basel, Switzerland. ¹⁰⁵Department of Medicine, Harvard Medical School, Boston, USA. ¹⁰⁶Max Planck Institute of Psychiatry, Munich, Germany. ¹⁰⁷Division of Research, Kaiser Permanente Northern California, Oakland, USA. ¹⁰⁸Centre for Addiction and Mental Health, Toronto, Canada. ¹⁰⁹Department of Psychiatry, University of Toronto, Toronto, Canada. ¹¹⁰Psychiatry & The Behavioral Sciences, University of Southern California, Los Angeles, USA. ¹¹¹Department of Endocrinology at Herlev

University Hospital, University of Copenhagen, Copenhagen, Denmark. ¹¹²Swiss Institute of Bioinformatics, Lausanne, Switzerland. ¹¹³Institute of Social and Preventive Medicine (IUMSP), Lausanne University Hospital, Lausanne, Switzerland. ¹¹⁴Division of Psychiatry, University College London, London, UK. ¹¹⁵Mental Health NHS 24, Glasgow, UK. ¹¹⁶Division of Psychiatry, Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK. ¹¹⁷Statistics, University of Oxford, Oxford, UK. ¹¹⁸EMGO+ Institute for Health and Care Research, Amsterdam, Netherlands. ¹¹⁹School of Psychology and Counseling, Queensland University of Technology, Brisbane, Australia. ¹²⁰Institute of Molecular and Cell Biology, University of Tartu, Tartu, Estonia. ¹²¹Estonian Biocentre, Tartu, Estonia. ¹²²Institute for Molecular Biology, University of Queensland, Brisbane, Australia. ¹²³Medical Genetics, University of British Columbia, Vancouver, Canada. ¹²⁴Statistics, University of British Columbia, Vancouver, Canada. ¹²⁵DZHK (German Centre for Cardiovascular Research), Partner Site Greifswald, University Medicine, Matthias Nauck, Greifswald, Germany. ¹²⁶Institute of Clinical Chemistry and Laboratory Medicine, University Medicine Greifswald, Greifswald, Germany. ¹²⁷Mental Health Centre Copenhagen, Copenhagen University Hospital, Copenhagen, Denmark. ¹²⁸Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane, Australia. ¹²⁹MRC Centre for Neuropsychiatric Genetics and Genomics, Cardiff University, Cardiff, UK. ¹³⁰Humus, Reykjavik, Iceland. ¹³¹Human Genetics and Computational Biomedicine, Pfizer Global Research and Development, Groton, USA. ¹³²Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden. ¹³³Charles E. Schmidt College of Medicine, Florida Atlantic University, Boca Raton, USA. ¹³⁴Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden. ¹³⁵Medical Genetics Section, CGEM, IGMM, University of Edinburgh, Edinburgh, UK. ¹³⁶Complex Trait Genetics, VU University Amsterdam, Amsterdam, Netherlands. ¹³⁷Clinical Genetics, VU University Medical Center, Amsterdam, Netherlands. ¹³⁸Psychiatry, University of Iowa, Iowa City, USA. ¹³⁹Solid GT, Boston, USA. ¹⁴⁰Department of Biochemistry and Molecular Biology II, Institute of Neurosciences, Center for Biomedical Research, University of Granada, Granada, Spain. ¹⁴¹Psychiatry, Icahn School of Medicine at Mount Sinai, New York, USA. ¹⁴²Department of Psychiatry, University of Groningen, University Medical Center Groningen, Groningen, Netherlands. ¹⁴³Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, USA. ¹⁴⁴Faculty of Medicine, Department of Psychiatry, School of Health Sciences, University of Iceland, Reykjavik, Iceland. ¹⁴⁵School of Medicine and Dentistry, James Cook University, Townsville, Australia. ¹⁴⁶Institute of Health and Wellbeing, University of Glasgow, Glasgow, UK. ¹⁴⁷deCODE Genetics/Amgen, Reykjavik, Iceland. ¹⁴⁸College of Biomedical and Life Sciences, Cardiff University, Cardiff, UK. ¹⁴⁹Institute for Community Medicine, University Medicine Greifswald, Greifswald, Germany. ¹⁵⁰PSYCH, The Lundbeck Foundation Initiative for Integrative Psychiatric Research, Copenhagen, Denmark. ¹⁵¹KG Jebsen Centre for Psychosis Research, Norway Division of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway. ¹⁵²Department of Psychiatry, University of California, San Diego, San Diego, USA. ¹⁵³Institute for Molecular Bioscience, The University of Queensland, Brisbane, Australia. ¹⁵⁴Roche Pharmaceutical Research and Early Development, Neuroscience, Ophthalmology and Rare Diseases Discovery & Translational Medicine Area, Roche Innovation Center Basel, F. Hoffmann-La Roche Ltd, Basel, Switzerland. ¹⁵⁵Department of Psychiatry and Psychotherapy, University Medicine Greifswald, Greifswald, Germany. ¹⁵⁶Department of Psychiatry, Leiden University Medical Center, Leiden, Netherlands. ¹⁵⁷Virginia Institute of Psychiatric & Behavioral Genetics, Virginia Commonwealth University, Richmond, USA. ¹⁵⁸Psychiatry, Columbia University College of Physicians and Surgeons, New York, USA. ¹⁵⁹Division of Epidemiology, New York State Psychiatric Institute, New York, USA. ¹⁶⁰Computational Sciences Center of Emphasis, Pfizer Global Research and Development, Cambridge, USA. ¹⁶¹Department of Clinical Medicine, Translational Neuropsychiatry Unit, Aarhus University, Aarhus, Denmark. ¹⁶²Institute of Neuroscience and Medicine (INM-1), Research Center Juelich, Juelich, Germany. ¹⁶³Department of Biomedicine, University of Basel, Basel, CH, Switzerland. ¹⁶⁴Division of Medical Genetics, University of Basel, Basel, CH, Switzerland. ¹⁶⁵Centre for Integrative Biology, Università degli Studi di Trento, Trento, Italy. ¹⁶⁶Psychiatry, University of California Los Angeles, Los Angeles, USA. ¹⁶⁷Psychiatry, Kaiser Permanente Northern California, San Francisco, USA. ¹⁶⁸Neuroscience Therapeutic Area, Janssen Research and Development, LLC, Titusville, USA. ¹⁶⁹Psychosis Research Unit, Aarhus University Hospital, Risskov, Aarhus, Denmark. ¹⁷⁰Institute of Translational Medicine, University of Liverpool, Liverpool, UK. ¹⁷¹Psychiatry, Harvard Medical School, Boston, USA. ¹⁷²Psychiatry, Dalhousie University, Halifax, Canada. ¹⁷³Institute of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark. ¹⁷⁴Human Genetics and Computational Biomedicine, Pfizer Global Research and Development, Cambridge, USA. ¹⁷⁵Orphan Disease Center, Perelman School of Medicine, University of Pennsylvania, Philadelphia, USA. ¹⁷⁶NIHR BRC for Mental Health, King's College London, London, UK. ¹⁷⁷Psychiatry & Behavioral Sciences, Stanford University, Stanford, USA. ¹⁷⁸Department of Medical & Molecular Genetics, King's College London, London, UK. ¹⁷⁹Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, USA. ¹⁸⁰Department of Psychiatry, University of North Carolina at Chapel Hill, Chapel Hill, USA.