

Whole genome expression analysis in human Jurkat cells after exposure to I-123-iododeoxyuridine, γ -rays and α -particles

Marcus Unverricht-Yeboah¹, Sonja Boldt², Ulrich Giesen³, Ekkehard Pomplun¹, Olaf Wolkenhauer², Ralf Kriehuber¹

¹*Department of Safety and Radiation Protection, Forschungszentrum Jülich, D-52425 Jülich, Germany*

²*Department of Computer Science, Systems Biology and Bioinformatics Group, University of Rostock, D-18051 Rostock, Germany*

³*Physikalisch-Technische Bundesanstalt (PTB), Bundesallee 100, D-38116 Braunschweig, Germany*

Introduction: In order to develop a gene expression profile-based method for biodosimetry purposes we used the human p53-deficient T-lymphoma Jurkat cell line to study whether gene signatures exist allowing the discrimination of radiation quality.

Methods: Equi-effect doses, i.e. radiation doses and exposure conditions causing the same biological effect level, were determined with regard to micronucleus formation, γ -H2AX foci intensity and apoptosis induction for the radiation qualities of γ -rays (Cs-137) and α -particles (Am-241) as well as for the Auger electron emitter I-123. Prior to the DNA-microarray based gene expression experiments, Jurkat cells were either irradiated with 0.8 and 5 Gy γ -rays, respectively with 0.1 and 0.5 Gy α -particles or were exposed to 4 - 200 kBq I-123-iododeoxyuridine (I-123-UdR) per 10E6 cells. I-123-UdR was incorporated into the DNA for 20 h. After quantification of cellular uptake and calculation of accumulated decays the absorbed radiation dose was assessed based on the 3-D geometry of the cells. RNA-isolation was performed 6 h and 24 h post-exposure. Whole human genome DNA-microarrays (Agilent) were processed and expression profiles were analyzed. Genes showing significant expression changes after irradiation were identified by one-way ANOVA and Tukey-HSD post-hoc testing. The biological functions of significantly regulated genes were further investigated.

Results: Preliminary results of the gene expression analysis indicate that the expression of more and different genes is significantly altered after exposure to I-123-UdR and α -particles when compared to γ -irradiation. The functional analysis of significantly changed genes reveals that apoptosis relevant genes are enriched after exposure to I-123-UdR and α -particles in comparison to γ -irradiation.

Conclusions: I-123-UdR and α -particles induce pronounced alterations in gene expression when compared to γ -rays. Changes in the gene expression of p53-dependent apoptosis-related genes were observed suggesting p53-independent back-up pathways for apoptosis signalling in Jurkat cells.

Funded by Bundesministerium für Bildung und Forschung (BMBF), Project No.: 02NUK005A and 02NUK005D