

Microaggregate stability and storage of organic carbon is affected by clay content in arable Luvisols

Lars Krause^{a*}, Andrei Rodionov^b, Steffen A. Schweizer^c, Nina Siebers^a, Eva Lehndorff^b, Erwin Klumpp^a, Wulf Amelung^{a,b}

^a Forschungszentrum Jülich GmbH, Agrosphere (IBG-3) Institute of Bio- and Geosciences, Wilhelm Johnen Straße, 52425 Jülich, Germany

^b University of Bonn, Institute of Crop Science and Resource Conservation - Soil Science and Soil Ecology, Nussallee 13, D-53115 Bonn, Germany

^c Technical University of Munich, Department Ecology and Ecosystem Management, Chair of Soil Science, Emil-Ramann-Straße 2, D-85354 Freising-Weihenstephan, Germany

* Corresponding author, e-mail: l.krause@fz-juelich.de

Keywords: free, occluded, particle size distribution

Abstract

As soil microaggregates (<250 µm, SMA) usually withstand long-term tillage, we hypothesized that i) elevated clay contents in arable soil support aggregation already at microaggregate level, leading to ii) increasing organic carbon (SOC) enrichment in smaller SMA size fractions. To test these hypotheses we sampled the topsoil (5-20 cm) of Luvisols with a long history of agricultural management at the Scheyern experiment station (Germany) in quintuplicates from each of five subsites with different clay contents (19-34%). The field-fresh topsoil was fractionated into macroaggregates (8000-250 µm), large SMA (250-20 µm), and small SMA (<20 µm) and the mass distribution was recorded. In addition, the water stable

macroaggregates ($>250\text{ }\mu\text{m}$) were dispersed ultrasonically to study occluded SMA and single building units. Finally, we analyzed the size distribution of the small SMA by laser diffraction analysis. The total mass distribution of free and occluded SMA grouped soils into those with small (19, 22, and 24%) and large (32 and 34%) clay contents. The finer textured soils exhibited larger portions of occluded SMA, with a gamma size distribution of small SMA peaking at $6\text{ }\mu\text{m}$. Yet the occluded small SMA in the finer textured soils showed an additional enrichment of colloids $<1\text{ }\mu\text{m}$. The SOC was indeed enriched in finer fractions, but more in the small SMA of the coarse textured sites than in the finer textured ones, whereas the opposite was true for the large SMA. We conclude, therefore, that elevated contents of clay-sized particles promote SMA formation and stabilization, therewith shifting SOC enrichment from small to larger SMA.

1 Introduction

Soil microaggregates (SMA) having a size $<250\text{ }\mu\text{m}$ are one of the key structures that influence soil properties such as the storage and stabilization of carbon (C), diversity of microbial habitats, diffusion of nutrients and gases, cycling of trace elements, or water retention as recently reviewed by Totsche et al. (2017). Hence, quantifying SMA and their properties has become an essential task in order to predict dynamic changes in soil ecosystems due to management and global change (Six et al., 2004).

Despite the wealth of information on SMA and their role for SOC protection or formation of organo-mineral-associations as building units (Totsche et al., 2017), we still lack information to systematize microaggregate formation across gradients in soil properties. Up to 40% of organic matter may be stored within SMA (Carter, 1995),

and the turnover of organic matter is associated with SMA stability (Goebel et al., 2009; Mueller et al., 2012). The stability of microaggregates increases with decreasing diameter of SMA (Golchin et al., 1994; Six et al., 2002; Tisdall, 1996). As a result, SMA are divided into two size fractions, large (250-20 μm) and small (<20 μm) SMA (Oades and Waters, 1991; Tisdall and Oades, 1982; Totsche et al., 2017). Small SMA are believed to be mainly stabilized by short-range van-der-Waals forces and electrostatic binding as reviewed in Totsche et al. (2017). They are stable against a range of forces, and cannot be further disintegrated into smaller building units (Kaiser and Berhe, 2014; Kaiser et al., 2012). This is different for larger aggregate units (Kaiser and Berhe, 2014), which contain small SMA as subunits, held together by roots, hyphae, or extracellular polymeric substances as temporary binding agents (Kleber et al., 2007; Six et al., 2000).

It is well established that the amount of clay-sized particles affects the stability and hierarchy of such larger aggregate units (Denef and Six, 2005; Goldberg, 1989; Wagner et al., 2007). Furthermore extensive studies were made about the influence of clay content on the composition and distribution of free and occluded organic matter. These studies showed that clay has an positive effect on the storage of organic carbon in occluded particulate organic matter, which has been linked with increased aggregation in the soils with larger clay content (Kölbl and Kögel-Knabner, 2004). In arable soils, stable organic carbon is mainly associated with fine particle size fractions (Kiem et al., 2002). In turn, organic carbon contributes to the stabilization of microaggregates as well (Kölbl et al., 2005). In soils without tillage and soil management these effects are stronger resulting in an increased soil aggregation and stability as well as improved storage of organic carbon (Ayoubi et al., 2012; Karchegani et al., 2012), which can breakdown during land use change (Schweizer et al., 2017). Furthermore, the stabilization of SOC can also be governed by the

surface area of oxides influencing the interaction of organic compounds with minerals (Kiem and Kögel-Knabner, 2002). However, detailed studies on the role of clay for the properties of the small SMA ($< 20 \mu\text{m}$) as major building units are largely lacking.

We hypothesize that for a given arable land-use system i) elevated clay contents support aggregation not only at macro- but already at microaggregate level, and that ii) stabilization of the smaller size fractions through increased amounts of SOC shifts the overall particle-size distribution to larger equivalent mean weight diameter than at subsites with lower clay contents. In order to test these hypotheses we collected topsoil samples with increasing clay contents from a heterogeneous arable field at Scheyern experiment station, Germany, and investigated the size and mass distribution of small SMA as major building units of their larger counterparts.

2 Materials and Methods

2.1 Soil sampling and sample characteristics

We selected five adjacent sample sites on the same arable plot (Table A.1; Supplementary Materials) at the research station Scheyern ($48^{\circ} 29' 36'' \text{ N}$, $11^{\circ} 26' 15'' \text{ E}$) situated in the rural area of the hill slopes north of Munich (Germany) for sampling. Plot selection was facilitated by using gamma spectrometry for sensing textural variations within a heterogeneous field site (Heggemann et al., 2017). The study site was characterized by a mean annual temperature of 7.4°C and a mean annual precipitation of 803 mm (Schröder et al., 2002). The soils were sandy to loamy textured Luvisols, developed on sandy molasse sediments with silty to clayey molasse deposits. All samples were free of carbonates and dominated by illitic clay mineralogy, which did not show systematic differences across the texture gradient (K.

U. Totsche, personal communication, 2018). Other clay minerals detected were Fe-chlorite, kaolinite and smectite, the latter however only in samples with 12 and 18.5% clay, the other samples contained vermiculite instead (K. U Totsche, personal communication).

Overall 25 samples were used for analysis from each of five sampling sites with five replicates each, taken in a radial sampling scheme (1 m around each subsite; Wilding (1985)). Samples were taken after removal of the upper Ap layer prone to crust formation (first 4-5 cm) with a large core cutter (diameter: 16 cm, length: 15 cm, volume: $\sim 3000 \text{ cm}^3$). Clay contents were determined based on texture analysis using H_2O_2 oxidation. About 10 g of each sample was treated with an excess of H_2O_2 (30%) until no further oxidation could be observed. The oxidized samples were put into an oven at 60 °C with additional 500 mL H_2O for 12 hours to remove excess H_2O_2 . Then, the samples were sieved $<63 \mu\text{m}$, freeze-dried and 3 g were mixed with 80 mL 0.0125 M $\text{Na}_4\text{P}_2\text{O}_7$. The clay content was determined by X-ray attenuation in a Sedigraph III Plus (Micromeritics, Georgia, USA). The clay contents were 19, 22, 24, 32, and 34%, respectively. Additionally 100 cm^3 cores were sampled for field moisture and bulk density determination. The fresh soil samples were gently sieved $<8 \text{ mm}$ and stored at 4 °C. The Fe, Al, and Mn phases (Fe_D , Al_D , Mn_D) were extracted using dithionite-citrate-bicarbonate (DCB) according to Mehra and Jackson (1958), serving as a measure for the abundance of pedogenic oxides. The extracts were analyzed with inductively coupled plasma optical emission spectroscopy (ICP–OES) Vista-Pro (Varian, California, USA). The total C contents in each aggregate fraction were determined using elemental analysis (Vario MICRO cube, Elementar Analysesysteme GmbH, Langenselbold, Germany), according to ISO 10694 (1995).

2.2 Wet fractionation and density fractionation

The fractionation was conducted on field-moist soil in triplicates, using a combination of wet sieving and pressure filtration (Fig. 1). In order to avoid re-aggregation, the samples were kept submerged in deionized water during the whole process of fractionation until they were shock frozen in liquid nitrogen and lyophilized. In short, 50 g of 8 mm field fresh sieved samples were placed on a glass fiber filter being on the top of a sieve tower and pre-wetted for five minutes. The sample was then sieved in a sieve-tower using 6 sieves having a mesh size between 2800 to 20 μm (Fig. 1) under constant shaking (30 U min^{-1}) for 10 minutes. Collected fractions were categorized into free and occluded SMA. Free SMA (250-53, 53-20, and 20-0.22 μm) were collected without further treatment. In order to isolate occluded SMA, remnants of wet sieving being $>250 \mu\text{m}$ were merged, suspended in H_2O and subjected to an ultrasonic treatment using 60 J mL^{-1} (Branson Ultrasonics corp., Connecticut, USA) as recommended by Amelung and Zech (1999). After ultrasonic treatment the suspension was wet sieved using the sieve tower as described above starting with the 2800 μm sieve and obtained size fractions being >250 , 250-53, 53-20, and $<20 \mu\text{m}$ using the sieving tower described by Kösters et al. (2013).

Wet sieving resulted in approximately 50 L suspensions each containing $<20 \mu\text{m}$ fractions of free and occluded SMA. The collection of small SMA was performed through pressure filtration. Homogenized aliquots were pressure filtered (Pressure filter Holder Type 16274, Sartorius, Göttingen, Germany) with 2-3 bar N_2 through a 0.22 μm PVDF filter (Membrane filter PVDF hydrophilic, 1201322, Berrytec GmbH, Oberbayern, Germany). After filtration the remnants were removed from the membrane filter with H_2O , the resulting suspension was shock frosted in liquid nitrogen and freeze-dried. The collected $<0.22 \mu\text{m}$ fraction was considered as

nanoparticle fraction. The process of fractionation resulted in an average total mass recovery of 97%.

Since the collected large SMA and macroaggregates still contained primary mineral particles, a density fractionation with sodium metatungstate was performed with pooled field replicates according to Virto et al. (2008). With this method, primary sand and silt particles with a density of $>2.5 \text{ g cm}^{-3}$ were removed from the affected size fractions. Particulate organic matter (POM) with a density of $1.6\text{-}1.8 \text{ g cm}^{-3}$ was not removed in an additional step since the mass contribution of the POM fractions was limited to a minor proportion (Kölbl and Kögel-Knabner, 2004).

2.3 Particle size distribution

The size distribution of small SMA ($<20 \text{ }\mu\text{m}$) was measured with a laser particle-size analyzer (LA-950, Horiba, Kyōto, Japan). Before each measurement the sample was suspended in water and ultrasonicated (13 J mL^{-1}) using the internal ultrasonic finger of the instrument. During the measurement the SMA were kept in suspension through stirring. The scattered light data was transformed into a volume based size distribution by a software algorithm using the mie-theory provided by the manufacturers' software.

2.4 Statistical tests

Calculation of Pearsons correlation coefficients between clay content and measured elements was performed with Excel (Microsoft Excel 2010, Microsoft corporation, Washington, USA). Statistical analysis was performed with Origin (OriginPro 2017, Originlab, Massachusetts, USA). The Shapiro Wilk test was used for testing the

normal distribution. With the one-way ANOVA the statistical significance was tested at a significance level of $\alpha=0.05$ and in a post-hoc test the means were compared with Tukey's method. The tests were applied for the SOC and mass percentage data, each pooled into soils with small (19-24%) and large (33-34%) clay content. Overall four levels of significance were defined and indicated by asterisks (not significant, $p>0.05$, no symbol; significant, $p\leq 0.05$, *; very significant, $p\leq 0.01$, **; highly significant, $p\leq 0.001$, ***; extremely significant, $p\leq 0.0001$, ****).

3 Results

Generally, the subsampling of the sites was successful in yielding samples with different texture: as clay content increased from 19-34%, mass proportions of sand decreased respectively (Fig. 2). When NaOCl was used instead of H_2O_2 , the respective clay contents were slightly lower, but changed in similar direction (Table A.2; Supplementary Materials). Intriguingly, the Fe_D content did not show significant differences along this transect, being in average 8.3 g kg^{-1} in soils with 19-24% clay, and 8.5 g kg^{-1} in soils with larger clay content (32-34%), respectively (Table A.2; Supplementary Materials). In turn, correlation coefficients of 0.83 and 0.87 indicate that the concentrations of Al_D and Mn_D increased with clay content (Table A.2; Supplementary Materials). With density fractionation the mass of primary mineral particles with a density of $>2.5 \text{ g cm}^{-3}$ was evaluated. For the 8000-250 μm fractions, the average mass percentage of primary mineral particles reached 94% and 96% soil mass, being slightly larger in soils with larger clay content compared to soils with smaller clay content (Table A.2; supplementary materials). For all 250-20 μm fractions, i.e., the large SMA fractions, the mass percentage of primary mineral particles averaged 80% without noticeable differences between subsites of different clay contents. Intriguingly, the amount of primary particles in "free" fractions was

lower (75%) than in the occluded fractions (84%; data not shown). The small SMA fraction was not corrected for primary particle contents.

3.1 SMA mass distribution

Unlike the textural gradient, the evaluation of isolated fractions indicated that the mass distribution of soil aggregates was not linearly related to the amount of clay-sized particles, but rather grouped into soils having small and large clay content, respectively (Fig. 3a). Samples with large clay content showed large amounts of occluded small SMA, while samples with small clay contents showed less occluded small SMA and higher content of free SMA (Fig. 3b). The main differences between samples with small and large clay content were thus the amount of occluded small SMA, the proportion of which was significantly larger in soils with $\geq 32\%$ clay (Fig. 4). The mass-proportion of free small SMA was also affected by clay content and significantly larger in soils with smaller clay contents.

3.2 Size distribution of the small SMA

The volume weighted size distribution of small SMA followed mainly the gamma distribution (Fig. A.1; Supplementary Materials). More than 90% of the measured fraction was $\leq 10 \mu\text{m}$ (Fig. 5). The most abundant size class of small SMA was found in the range of 5 to 7 μm (Fig. 5a). Intriguingly, this size distribution was similar for free and occluded small SMA, which can be clearly seen by comparison of the cumulative percentages (Fig. 5b). In addition, however, soils with larger clay content ($\geq 32\%$) showed an additional abundance of particles $\leq 1 \mu\text{m}$ within the occluded small SMA (Fig. 5c, d), pointing to a potential role of colloids in SMA occlusion. To better

understand the underlying mechanisms, we assessed the concentrations of SOC inside the different SMA fractions.

3.3 SOC content of different SMA fractions

The SOC contents did not strongly correlate with the gradient of clay content but showed some differences between small (19-24%) and large (32-34%) clay content. Apparently, the relationship between SOC-enrichment in clay and clay-content was not linear (see also Christensen (1992)). Compared to the SMA fraction with small clay content the total SOC concentration of SMA from soils with larger clay content had increased significantly in the free and decreased in occluded SMA (Fig. 6a). There were no significant differences in SOC contents between the free and occluded SMA from soils with small clay content. However, the relationship of the SOC content of the SMA to the clay content was not a linear, i.e., the data grouped similarly as did the mass distribution of SMA, we keep the comparison of SOC within SMA of sites with low ($\leq 24\%$ clay) and large clay content ($\geq 32\%$), respectively.

Overall in the soil sequence, larger SOC concentrations were found in the small SMA fraction than in the large SMA. It contained larger amounts of SOC in occluded SMA than in the free ones, and larger amounts of SOC in the coarser than in the finer textured soils (Fig. 6b). In contrast to the observations from small SMA, the large SMA fractions had significantly larger SOC concentrations in the free than in the occluded SMA, with now also larger SOC contents in the samples with larger clay contents than in the coarser textured ones (Fig. 6c). Hence, the distribution of SOC into occluded and free aggregates as well as among the differently textured soils just behaved opposite in the small SMA relative to the large SMA.

4 Discussion

The applied fractionation scheme enabled us deeper insights into the dynamic aggregation processes at different scales ranging from macroaggregates to small SMA. The disruption energy of 60 J mL^{-1} has been considered as suitable for evaluating the SMA size distribution and associated organic matter (Kölbl et al., 2005) preserving largely their natural state (Amelung and Zech, 1999). Differentiation between occluded and free aggregates supported our first hypotheses that elevated clay contents support aggregation not only at macro- but already at microaggregate level. Specifically, elevated clay contents promoted the formation of occluded SMA (Fig. 4), likely by higher aggregate stability through interactions with more abundant colloids $< 1 \mu\text{m}$ (Fig. 5). In this regard, our data are in support of the concept of aggregate hierarchy (Oades and Waters, 1991), in that macroaggregates are built from microaggregates which can be formed from smaller building units. Here we show a novel size distribution measurement of the smaller building units (0.1 to $1 \mu\text{m}$), which was significant higher at sites with elevated clay content. The occluded and free small SMA reveals a gamma like size distribution (Fig. A.1; Supplementary Materials) peaking at $4\text{-}6 \mu\text{m}$ (Fig. 5). The gamma distribution model showed the best fit in comparison to other models which is in agreement with the work of Yang et al. (2012).

The similar size distribution of free and occluded particles in soils with low clay content (Fig. 5a) indicated that there is no preferential accumulation of specific sized free small SMA during aggregation. The evaluation of the total masses of soil aggregate fractions showed that the macroaggregate/SMA ratios became narrower when moving to subsites with larger clay content. This agrees with Yu et al. (2015), who found similar changes in macroaggregate/SMA ratios as a function of silt and clay content. In soils with large clay content the abundance of occluded small SMA

increased, which indicates larger amounts of water stable aggregates and thus an essential role of the clay-sized fraction not only for the formation but also for the stability of SMA.

It has been suggested that a minimum clay content of 15% is required for the abiotic formation of aggregates (Horn, 1990; Oades, 1993). The percentage of the analyzed low clay content soils (19-24%) was above this minimum value. An additional factor contributing to more occluded SMA in soils with large clay content could be the significantly larger concentrations of Al- and Mn-bearing phases (Appendix Table A.2); at least for aluminum oxides it is known that they promote the stabilization of clay minerals and organic matter on mineral surfaces (Goldberg, 1989; Kögel-Knabner et al., 2008).

Another essential factor promoting SMA formation is organic matter. Our results showed that especially small SMA were rich in SOC (Fig 6b, c), which confirms previous studies showing that organic carbon mainly accumulates in the smallest fractions with the largest specific surface area (Kahle et al., 2002; Kennedy et al., 2002). In addition, the SOC concentrations were higher in occluded small SMA than in the free ones (Fig. 6b), confirming previous findings that only clayey soils supplied with organic matter were able to aggregate and that there is a strong correlation between SOC accrual and aggregate stability (Al-Kaisi et al., 2014; Imhoff et al., 2002; Wagner et al., 2007). In turn, more stable aggregates also protect SOC better from degradation as aggregates that disintegrate. Through occlusion, for instance, SOC is better protected against microbial decomposition in the <20 μm fraction than in larger SMA (Hassink, 1997). Physical stabilization affects the turnover of labile molecules like carbohydrates and proteins (Krull et al., 2003) that can act as gluing agents for aggregates. Previous studies showed that this SOC protection works in

addition to chemical and biochemical stabilization (Six et al., 2002) due to the preservation of polysaccharides and phenolic C in the clay fraction (De Nobili et al., 2008; Schöning et al., 2005). The microbial cycling of SOC nearby mineral surfaces fosters the accrual of carbohydrate and other microbial residues (Amelung et al., 1997; Zhang et al., 1999), which might act as glue within microaggregate cores (Ladd et al., 1993; Lynch and Bragg, 1985; Rodionov et al., 2001). As a result, soil organic matter composition may vary with aggregate size (Golchin et al., 1994; Yu et al., 2015). In this regard, a higher SOC concentration in occluded small SMA relative to the free SMA (Fig. 6b) might be explained by an enrichment of microbial-altered SOC (Bingham and Cotrufo, 2016; Cotrufo et al., 2013; Rodionov et al., 2001), whereas fungal hyphae and fine root debris might be part of free SOC, enriched within the free large SMA fraction (Fig. 6c). According to Amelung et al. (2002), SOC accumulation in microaggregates may well happen at the outer surface, i.e., it does not necessarily have to result in an additional occlusion process. Nevertheless, this different direction of SOC enrichment hardly altered the size distribution of SMA with increasing clay content (Fig. 5).

5 Conclusion

Results indicated that the abundance of clay-sized particles as building unit is a superior requirement for aggregate formation and stability. Elevated clay contents promote the formation and stability of SMA with different size, resulting in larger concentrations of occluded SMA than of free ones as clay contents rose. Notably, SOC enrichment followed the opposite direction, i.e., the additional clay mostly trapped SOC in free, large SMAs. We attribute this to an interaction loop: the preferred formation of occluded aggregates with elevated clay contents leads to

321 lower amounts of “free” particles, which therewith concentrate the remaining SOC at
322 their surface.

323 **6 Acknowledgements**

324 We acknowledge A. Lindecke for performing the C/N analysis. Furthermore we want
325 to thank Dr. Daniel Emil Mack and Andrea Hilgers for the support of the particle size
326 analysis, and I. Kögel-Knaber for helpful discussions. This work is associated to the
327 MAD Soil project, which was funded by the DFG (Deutsche
328 Forschungsgemeinschaft, Research Unit 2171

7 References

- Al-Kaisi, M.M., Douelle, A., Kwaw-Mensah, D., 2014. Soil microaggregate and macroaggregate decay over time and soil carbon change as influenced by different tillage systems. *Journal of Soil and Water Conservation* 69, 574-580.
- Amelung, W., Kaiser, K., Kammerer, G., Sauer, G., 2002. Organic carbon at soil particle surfaces—evidence from x-ray photoelectron spectroscopy and surface abrasion. *Soil Science Society of America Journal* 66, 1526-1530.
- Amelung, W., Zech, W., 1999. Minimisation of organic matter disruption during particle-size fractionation of grassland epipedons. *Geoderma* 92, 73-85.
- Amelung, W., Zech, W., Flach, K.W., 1997. Climatic Effects on Soil Organic Matter Composition in the Great Plains. *Soil Science Society of America Journal* 61, 115-123.
- Ayoubi, S., Mokhtari Karchegani, P., Mosaddeghi, M.R., Honarjoo, N., 2012. Soil aggregation and organic carbon as affected by topography and land use change in western Iran. *Soil and Tillage Research* 121, 18-26.
- Bingham, A.H., Cotrufo, M.F., 2016. Organic nitrogen storage in mineral soil: Implications for policy and management. *Science of The Total Environment* 551, 116-126.
- Carter, M.R., 1995. Analysis of Soil Organic Matter Storage in Agroecosystems, In: Carter, M.R., Stewart, B.A. (Eds.), *Structure and Organic Matter Storage in Agricultural Soils*. Taylor & Francis, Boca Raton, FL, pp. 3-11.
- Christensen, B.T., 1992. Physical fractionation of soil and organic matter in primary particle size and density separates, In: Stewart, B.A. (Ed.), *Advances in Soil Science*. Springer New York, pp. 1-90.
- Cotrufo, M.F., Wallenstein, M.D., Boot, C.M., Denef, K., Paul, E., 2013. The Microbial Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition with soil organic matter stabilization: do labile plant inputs form stable soil organic matter? *Global Change Biology* 19, 988-995.
- De Nobili, M., Contin, M., Mahieu, N., Randall, E.W., Brookes, P.C., 2008. Assessment of chemical and biochemical stabilization of organic C in soils from the long-term experiments at Rothamsted (UK). *Waste Management* 28, 723-733.
- Denef, K., Six, J., 2005. Clay mineralogy determines the importance of biological versus abiotic processes for macroaggregate formation and stabilization. *European Journal of Soil Science* 56, 469-479.
- Goebel, M.-O., Woche, S.K., Bachmann, J., 2009. Do soil aggregates really protect encapsulated organic matter against microbial decomposition? *Biologia* 64, 443-448.
- Golchin, A., Oades, J., Skjemstad, J., Clarke, P., 1994. Soil structure and carbon cycling. *Soil Research* 32, 1043-1068.
- Goldberg, S., 1989. Interaction of aluminum and iron oxides and clay minerals and their effect on soil physical properties: A review. *Communications in Soil Science and Plant Analysis* 20, 1181-1207.
- Hassink, J., 1997. The capacity of soils to preserve organic C and N by their association with clay and silt particles. *Plant and Soil* 191, 77-87.
- Heggenmann, T., Welp, G., Amelung, W., Angst, G., Franz, S.O., Koszinski, S., Schmidt, K., Pätzold, S., 2017. Proximal gamma-ray spectrometry for site-independent in situ prediction of soil texture on ten heterogeneous fields in Germany using support vector machines. *Soil and Tillage Research* 168, 99-109.

Horn, R., 1990. Aggregate characterization as compared to soil bulk properties. *Soil and Tillage Research* 17, 265-289.

Imhoff, S., da Silva, A.P., Dexter, A., 2002. Factors Contributing to the Tensile Strength and Friability of Oxisols. *Soil Science Society of America Journal* 66, 1656-1661.

Kahle, M., Kleber, M., Jahn, R., 2002. Carbon storage in loess derived surface soils from Central Germany: Influence of mineral phase variables. *Journal of Plant Nutrition and Soil Science* 165, 141-149.

Kaiser, M., Berhe, A.A., 2014. How does sonication affect the mineral and organic constituents of soil aggregates?—A review. *Journal of Plant Nutrition and Soil Science* 177, 479-495.

Kaiser, M., Berhe, A.A., Sommer, M., Kleber, M., 2012. Application of ultrasound to disperse soil aggregates of high mechanical stability. *Journal of Plant Nutrition and Soil Science* 175, 521-526.

Karchegani, P.M., Ayoubi, S., Mosaddeghi, M.R., Honarjoo, N., 2012. Soil organic carbon pools in particle-size fractions as affected by slope gradient and land use change in hilly regions, western Iran. *Journal of Mountain Science* 9, 87-95.

Kennedy, M.J., Pevear, D.R., Hill, R.J., 2002. Mineral Surface Control of Organic Carbon in Black Shale. *Science* 295, 657-660.

Kiem, R., Knicker, H., Kögel-Knabner, I., 2002. Refractory organic carbon in particle-size fractions of arable soils I: distribution of refractory carbon between the size fractions. *Organic Geochemistry* 33, 1683-1697.

Kiem, R., Kögel-Knabner, I., 2002. Refractory organic carbon in particle-size fractions of arable soils II: organic carbon in relation to mineral surface area and iron oxides in fractions <6 µm. *Organic Geochemistry* 33, 1699-1713.

Kleber, M., Sollins, P., Sutton, R., 2007. A conceptual model of organo-mineral interactions in soils: self-assembly of organic molecular fragments into zonal structures on mineral surfaces. *Biogeochemistry* 85, 9-24.

Kögel-Knabner, I., Guggenberger, G., Kleber, M., Kandeler, E., Kalbitz, K., Scheu, S., Eusterhues, K., Leinweber, P., 2008. Organo-mineral associations in temperate soils: Integrating biology, mineralogy, and organic matter chemistry. *Journal of Plant Nutrition and Soil Science* 171, 61-82.

Kölbl, A., Kögel-Knabner, I., 2004. Content and composition of free and occluded particulate organic matter in a differently textured arable Cambisol as revealed by solid-state ¹³C NMR spectroscopy. *Journal of Plant Nutrition and Soil Science* 167, 45-53.

Kölbl, A., Leifeld, J., Kögel-Knabner, I., 2005. A comparison of two methods for the isolation of free and occluded particulate organic matter. *Journal of Plant Nutrition and Soil Science* 168, 660-667.

Kösters, R., Preger, A.C., Du Preez, C.C., Amelung, W., 2013. Re-aggregation dynamics of degraded cropland soils with prolonged secondary pasture management in the South African Highveld. *Geoderma* 192, 173-181.

Krull, E.S., Baldock, J.A., Skjemstad, J.O., 2003. Importance of mechanisms and processes of the stabilisation of soil organic matter for modelling carbon turnover. *Functional plant biology* 30, 207-222.

Ladd, J.N., Foster, R.C., Skjemstad, J.O., 1993. Soil structure: carbon and nitrogen metabolism. *Geoderma* 56, 401-434.

Lynch, J.M., Bragg, E., 1985. Microorganisms and Soil Aggregate Stability, In: Stewart, B.A. (Ed.), *Advances in Soil Science*. Springer New York, New York, NY, pp. 133-171.

Mehra, O.P., Jackson, M.L., 1958. Iron Oxide Removal from Soils and Clays by a Dithionite-Citrate System Buffered with Sodium Bicarbonate. *Clays and Clay Minerals* 7, 317-327.

Mueller, C.W., Schlund, S., Prietzel, J., Kögel-Knabner, I., Gutsch, M., 2012. Soil Aggregate Destruction by Ultrasonication Increases Soil Organic Matter Mineralization and Mobility. *Soil Science Society of America Journal* 76, 1634-1643.

Oades, J., Waters, A., 1991. Aggregate hierarchy in soils. *Soil Research* 29, 815-828.

Oades, J.M., 1993. The role of biology in the formation, stabilization and degradation of soil structure. *Geoderma* 56, 377-400.

Rodionov, A., Amelung, W., Urusevskaja, I., Zech, W., 2001. Origin of the enriched labile fraction (ELF) in Russian Chernozems with different site history. *Geoderma* 102, 299-315.

Schöning, I., Morgenroth, G., Kögel-Knabner, I., 2005. O/N-alkyl and alkyl C are stabilised in fine particle size fractions of forest soils. *Biogeochemistry* 73, 475-497.

Schröder, P., Huber, B., Olazábal, U., Kämmerer, A., Munch, J.C., 2002. Land use and sustainability: FAM Research Network on Agroecosystems. *Geoderma* 105, 155-166.

Schweizer, S.A., Fischer, H., Häring, V., Stahr, K., 2017. Soil structure breakdown following land use change from forest to maize in Northwest Vietnam. *Soil and Tillage Research* 166, 10-17.

Six, J., Bossuyt, H., Degryze, S., Denef, K., 2004. A history of research on the link between (micro)aggregates, soil biota, and soil organic matter dynamics. *Soil and Tillage Research* 79, 7-31.

Six, J., Conant, R.T., Paul, E.A., Paustian, K., 2002. Stabilization mechanisms of soil organic matter: Implications for C-saturation of soils. *Plant and Soil* 241, 155-176.

Six, J., Elliott, E.T., Paustian, K., 2000. Soil macroaggregate turnover and microaggregate formation: a mechanism for C sequestration under no-tillage agriculture. *Soil Biology and Biochemistry* 32, 2099-2103.

Tisdall, J.M., 1996. Formation of soil aggregates and accumulation of soil organic matter, In: Carter, M.R., Stewart, B.A. (Eds.), *Structure and organic matter storage in agricultural soils*. Taylor & Francis, Boca Raton, FL, pp. 57-87.

Tisdall, J.M., Oades, J.M., 1982. Organic matter and water-stable aggregates in soils. *Journal of Soil Science* 33, 141-163.

Totsche, K.U., Amelung, W., Gerzabek, M.H., Guggenberger, G., Klumpp, E., Knief, C., Lehdorff, E., Mikutta, R., Peth, S., Prechtel, A., Ray, N., Kögel-Knabner, I., 2017. Microaggregates in soils. *Journal of Plant Nutrition and Soil Science*, n/a-n/a.

Virto, I., Barré, P., Chenu, C., 2008. Microaggregation and organic matter storage at the silt-size scale. *Geoderma* 146, 326-335.

Wagner, S., Cattle, S.R., Scholten, T., 2007. Soil-aggregate formation as influenced by clay content and organic-matter amendment. *Journal of Plant Nutrition and Soil Science* 170, 173-180.

Wilding, L.G., 1985. Spatial variability: Its documentation, accommodation and implication to soil survey, In: Nielsen, D.R., J., B. (Eds.), *Soil spatial variability: proceedings of a workshop of the ISSS and the SSSA*, Las Vegas, USA, Wageningen, Netherlands, pp. 166-187.

Yang, X., Lee, J., Barker, D.E., Wang, X., Zhang, Y., 2012. Comparison of six particle size distribution models on the goodness-of-fit to particulate matter sampled from animal buildings. *Journal of the Air & Waste Management Association* 62, 725-735.

Yu, H., Ding, W., Chen, Z., Zhang, H., Luo, J., Bolan, N., 2015. Accumulation of organic C components in soil and aggregates. *Scientific Reports* 5, 13804.

Zhang, X., Amelung, W., Yuan, Y., Samson-Liebig, S., Brown, L., Zech, W., 1999. Land-use effects on amino sugars in particle size fractions of an Argiudoll. *Applied Soil Ecology* 11, 271-275.




Fig. 1. Overview about the fractionation process using wet sieving and pressure filtration. According to the different size fractions the samples were divided into macroaggregates, large, and small SMA. Mechanical disaggregation via ultrasonication allowed a differentiation between occluded (ultrasonic treatment, red arrow) and free (no treatment, blue arrow) SMA.




Fig. 2. Texture of H_2O_2 oxidized soil samples obtained from the toposequence at an agricultural site.

485




Fig. 3. Mass percentage of macroaggregates and SMA in bulk soil (a) and within SMA (b) of soils with different clay content (small and large).

486




Fig. 4. Mass percentage of free and occluded small SMA; significant differences are indicated with asterisks.

487




Fig. 5. Volume based size distribution (a, c) and cumulative percentage (b, d) of free (blue/dotted) and occluded (red/straight) SMA for soils with small (a, b) and large (b, c) clay content.

488




Fig. 6. SOC content in dependence of soil clay content for all SMA (a) and their corresponding fractions including small (b) and large (c) SMA. The proportion of the free form is indicated in black and the occluded form in grey. Significant differences are indicated with asterisks.