

**Influence of dexamethasone on
O-(2-[¹⁸F]-fluoroethyl)-L-tyrosine uptake in the human brain and
quantification of tumor uptake**

Carina Stegmayr¹, Gabriele Stoffels¹, Elena Rota Kops¹, Philipp Lohmann¹,
Norbert Galldiks^{1,2,3}, Nadim J. Shah^{1,4,5}, Bernd Neumaier¹ and Karl-Josef
Langen^{1,4,6}

¹ Institute of Neuroscience and Medicine (INM-3, INM-4; INM-5; INM-11),
Forschungszentrum Jülich, Jülich, Germany

² Dept. of Neurology, University of Cologne, Cologne Germany

³ Center of Integrated Oncology (CIO), Universities of Cologne and
Bonn, Cologne, Germany

⁴ Jülich-Aachen Research Alliance (JARA) – Section JARA-Brain, Aachen,
Germany

⁵ Dept. of Neurology, RWTH University Aachen, Aachen, Germany

⁶ Dept. of Nuclear Medicine, RWTH University Hospital Aachen,
Aachen, Germany

Word count: 4385

Address for correspondence:

Karl-Josef Langen, M.D.
Institute of Neuroscience and
Medicine
Forschungszentrum Jülich
D-52425 Jülich, Germany
Phone: 0049-2461-61-5900
E-Mail: k.j.langen@fz-juelich.de Fax:
0049-2461-61-8261

First author:

Carina Stegmayr, Ph.D.
Institute of Neuroscience and
Medicine
Forschungszentrum Jülich
D-52425 Jülich, Germany
Phone: 0049-2461-61-9796
E-Mail: c.stegmayr@fz-juelich.de

Running title: Influence of corticoids on FET uptake

ABSTRACT

Purpose: O-(2-¹⁸F-fluoroethyl)-L-tyrosine ([¹⁸F]FET) is an established PET tracer for brain tumor imaging. This study explores the influence of dexamethasone therapy on [¹⁸F]FET uptake in the normal brain and its influence on the maximum and mean tumor-to-brain ratio (TBR).

Methods: [¹⁸F]FET PET scans of 160 brain tumor patients were evaluated (80 dexamethasone treated, 80 untreated; each group with 40 men/40 women). The standardized uptake value of [¹⁸F]FET uptake in the normal brain (SUV_{brain}) in the different groups was compared. Nine patients were examined repeatedly with and without dexamethasone therapy.

Results: SUV_{brain} of [¹⁸F]FET uptake was significantly higher in dexamethasone treated patients than in untreated patients (SUV_{brain} 1.33 ± 0.1 versus 1.06 ± 0.16 in male and 1.45 ± 0.25 versus 1.31 ± 0.28 in female patients). Similar results were observed in patients with serial PET scans. Furthermore, compared to men, a significantly higher SUV_{brain} was found in women, both with and without dexamethasone treatment. There were no significant differences between the different groups for TBR_{max} and TBR_{mean}, which could have been masked by the high standard deviation.

In a patient with a stable brain metastasis investigated twice with and without dexamethasone, the TBR_{max} and the biological tumor volume (BTV) decreased considerably after dexamethasone due to an increased SUV_{brain}.

Conclusion: Dexamethasone treatment appears to increase the ¹⁸F-FET uptake in the normal brain. An effect on TBR_{max}, TBR_{mean} and BTV cannot be

Running title: Influence of corticoids on FET uptake

excluded which should be considered especially for treatment monitoring and the estimation of BTV using [^{18}F]FET PET.

Key Words: PET; brain tumors; amino acids; ^{18}F -Fluoroethyltyrosine; FET; dexamethasone; tumor-to-brain ratio

INTRODUCTION

Brain tumor diagnostics with PET using radiolabeled amino acids such as O-(2- ^{18}F -fluoroethyl)-L-tyrosine ([^{18}F]FET) is gaining increasing interest and has recently been recommended by the Response Assessment in Neuro-Oncology (RANO) working group at every stage of management of patients with brain tumors [1-2]. Amino acid PET using [^{18}F]FET provides important additional information to conventional MRI especially with respect to delineation of tumor extent, the differentiation of treatment-related effects from tumor progression and for treatment monitoring [3-14].

Treatment with dexamethasone is the current standard therapy for patients suffering from brain edema, as it rapidly restores brain homeostasis and the integrity of the blood brain barrier (BBB), and thus reduces symptoms [15-16]. Recently, we were able to show that dexamethasone treatment reduces the permeability of the BBB in rat gliomas but does not change [^{18}F]FET uptake in the tumors [17]. Furthermore, we observed that restoration of the BBB after antiangiogenic therapy of rat gliomas with bevacizumab did not influence [^{18}F]FET uptake in the tumors [18]. These observations indicate that [^{18}F]FET uptake in

Running title: Influence of corticoids on FET uptake

brain tumors is widely independent of the permeability of the BBB and that viability of tumor tissue during therapy is more reliably reflected by [^{18}F]FET uptake than by contrast-enhanced MRI.

In another experimental study, however, we found some evidence that the tumor-to-brain ratio (TBR) decreased after dexamethasone treatment caused by an increased tracer uptake in normal brain tissue and not by reduced [^{18}F]FET uptake in the tumor [19]. Since it can happen that a patient during chemotherapy is once treated with dexamethasone and once not, changes of TBR of [^{18}F]FET uptake could be misinterpreted and interfere with the assessment of response to chemotherapy. Moreover, the biological tumor volume (BTV) is determined by a predefined threshold above background and could be influenced by an increased standardized uptake value in normal brain tissue ($\text{SUV}_{\text{brain}}$) after dexamethasone treatment.

The aim of this study was to explore the effect of dexamethasone treatment on [^{18}F]FET uptake in normal brain of patients with gliomas, and, consequently, to estimate the possible impact on TBR_{mean} , TBR_{max} , and BTV.

PATIENTS AND METHODS

Patients

Out of 1170 patients who were referred to our institution from November 2014 to August 2016 for tumor diagnostics using [^{18}F]FET PET, 160 patients were selected consecutively according to the following criteria: A) no dexamethasone prior to PET scans for at least two weeks (80 patients) and B) dexamethasone treatment for at least two weeks prior to the PET scans in doses ranging from 0.5 mg/day to 24 mg/day. The groups contained a balanced sex distribution, i.e. 40 men and 40 women per subgroup. Further information on mean age, dose of dexamethasone, histological or suspected diagnosis and, if appropriate, specific pretreatment of the different groups are provided in Table 1. Nine patients were scanned multiple times (two to four times) with and without dexamethasone therapy. The ethics committee of the University of Aachen approved the retrospective data evaluation. There was no conflict with the Declaration of Helsinki. All subjects gave prior written informed consent for the PET investigation and the use of the data for scientific evaluations.

PET imaging with ^{18}F -FET

The tracer was produced in-house as described elsewhere [20]. All patients fasted for at least 4 h before PET scanning, according to the German guidelines for brain tumor imaging using radiolabeled amino acid

Running title: Influence of corticoids on FET uptake

analogues [21] and were asked to take their drugs as usual, including dexamethasone, on the day of the PET scan. PET imaging was performed in 80 patients (20 per each group) on an ECAT Exact HR+ PET scanner (Siemens Medical Systems, Erlangen, Germany) in 3-dimensional mode (axial field of view, 15.5 cm; image resolution, 6 mm) and in 80 patients (20 per each group) simultaneously with MR imaging using a BrainPET insert (Siemens Medical Systems, Erlangen, Germany). The BrainPET is a compact cylinder that fits in the bore of the Magnetom Trio MR scanner (Siemens Medical Systems, Erlangen, Germany) (axial field of view, 19.2 cm; optimum image resolution, 3 mm) [22]. The dynamic PET studies were acquired up to 50 min after intravenous injection of approximately 3 MBq/kg body weight [^{18}F]FET. Reconstruction parameters were as similar as possible considering the different reconstruction environment of the two PET scanners. For the Exact HR+ scanner iterative reconstruction with 16 subsets and 6 iterations without any filtering was applied, while for the BrainPET an iterative reconstruction with 32 iterations and 2 subsets with a final image Gaussian filtering of 2.5 mm^3 was used; attenuation correction for the ECAT HR+ PET scan was based on a transmission scan, and for the BrainPET scan on a template-based approach [23]. The data were reconstructed dynamically with different frame lengths: 5 x 60 s, 5 x 180 s, and 6 x 300 s covering the 50 min whole scan time and allowing a time activity curve for the analysis.

PET data analysis

The volume-of-interest (VOI) analysis and the calculation of the BTV were based on the averaged PET data from 20–40 minutes post-injection using PMOD (Version 3.505, PMOD Technologies Ltd.). [^{18}F]FET uptake in the tissue was expressed as standardized uptake value (SUV) by dividing the radioactivity concentration (kBq/ml) in the tissue by the radioactivity amount (kBq) injected per gram of body weight. To determine the $\text{SUV}_{\text{brain}}$, a three-dimensional elliptic VOI of 15.5 was placed contralateral to the tumor in normal appearing perisylvian cortex representing almost exclusively grey matter. Another 2D ROI (6.7cm^2) was placed in the centrum semiovale representing exclusively white matter, to test for different effects of dexamethasone in grey and white matter ($\text{SUV}_{\text{white}}$). Furthermore, the area under curve (AUC) of the time activity curve of FET uptake in the brain VOI was determined. The maximum tumor SUV was determined by a 2 ml VOI centered on the maximal tumor uptake, mean tumor SUV and BTV by a 3-dimensional auto-contouring process using a tumor-to-brain ratio (TBR) of 1.6 as described previously [24]. TBR_{mean} and TBR_{max} were calculated by dividing the mean and maximum SUV in the tumor VOIs by the mean SUV of normal brain.

Statistics

Descriptive statistics are provided as mean and standard deviation (SD). A t-test was performed to test a difference between the two scanners.

Running title: Influence of corticoids on FET uptake

Distribution of patients in the different subgroups was evaluated by Fisher exact tests for 2×2 contingency tables. Two Way (TW) ANOVAs with All Pairwise Multiple Comparison Procedures (Holm-Sidak method) were performed to detect differences between dexamethasone treated and untreated patients as well as between men and women without correction for multiple testing. Data without normal distribution were tested with the Mann-Whitney Rank Sum test (TBR_{mean} and TBR_{max}). *P* values of 0.05 or less were considered significant. Pearson Product Correlations were performed for testing correlations between two variables. Statistical analysis was performed using SigmaPlot for Windows, Version 12.5.

RESULTS

SUV_{brain} of [^{18}F]FET uptake was significantly higher in dexamethasone treated patients than in untreated patients (SUV_{brain} 1.33 ± 0.10 versus 1.06 ± 0.16 in male and 1.45 ± 0.25 versus 1.31 ± 0.28 in female patients; both $P < 0.001$). The effect was likewise significant in white matter (Table 1). Furthermore, a significantly higher SUV_{brain} was found in women than in men, independently of dexamethasone treatment (Table 1, Fig. 1). Similar results were obtained when using AUC as a quantitative parameter for comparison. SUV_{brain} and AUC_{brain} values did not differ significantly between the ECAT HR+ PET scanner and the BrainPET scanner (t-test: $P = 0.234$ and 0.233 , respectively).

Running title: Influence of corticoids on FET uptake

In the nine patients who were scanned multiple times (two to four times) with and without dexamethasone therapy the SUV_{brain} values were highest in all but one case during dexamethasone therapy (Tab. 2).

All Pairwise Multiple Comparison Procedures found significant differences between subgroups, that is, men without dexamethasone had the lowest $[^{18}\text{F}]\text{FET}$ uptake in normal brain tissue and women with dexamethasone showed the highest uptake (Fig. 3), with a difference of 37% between the two groups.

A slight correlation was found between SUV_{brain} and dexamethasone dose in men ($P < 0.01$; $r = 0.44$), but not in women (Fig. 4A). There was no correlation between SUV_{brain} and body mass index (BMI) (Fig. 4B), dose of $[^{18}\text{F}]\text{FET}$, or age (data not shown) in men or women.

The influence of changing SUV_{brain} of $[^{18}\text{F}]\text{FET}$ uptake after dexamethasone treatment is further illustrated in one male patient with brain metastasis who underwent a $[^{18}\text{F}]\text{FET}$ PET scan during dexamethasone therapy (4 mg/d) and nine weeks later a second scan without dexamethasone (Tab. 3). No other treatment or medication was administered during this time. During follow-up, the tumor size remained unchanged on MR images, clinical symptoms remained stable (Fig. 5) and the maximum SUV of FET uptake in the tumor did merely change (-3%). In contrast, the SUV_{brain} was 19% higher during dexamethasone therapy leading to a drop of the TBR_{max} by 18%. Calculation of BTV using the threshold based method yielded a 70

% smaller BTV with dexamethasone than without, leading to a 14 % higher SUV_{mean} in the corresponding tumor VOI (Tab. 3). Thus, the higher SUV_{brain} during dexamethasone treatment was compensated by a higher SUV_{mean} in the tumor VOI and the TBR_{mean} was less dependent on dexamethasone therapy (-4%) than the TBR_{max} (-18 %).

DISCUSSION

The aim of this retrospective study was to evaluate whether dexamethasone treatment changes [^{18}F]FET uptake in normal brain tissue and might influence PET parameters such as tumor $SUV_{mean/max}$, tumor volume, and $TBR_{mean/max}$, which are important for diagnosis, grading, treatment planning and control, and follow-up.

We observed that dexamethasone treatment increased [^{18}F]FET uptake in normal brain tissue in both men and women (+24% and 9%, respectively). This effect was identical for grey and white matter. The difference in SUV_{brain} could be demonstrated not only in the mean values of the different subgroups, but it was also clearly detectable in eight out of nine patients, who underwent repeated [^{18}F]FET PET with and without dexamethasone treatment. There was no correlation between SUV_{brain} and the dose of dexamethasone in women and a very poor correlation in men (Fig. 4a). This is not unexpected because the Cushing threshold for dexamethasone is only 1-2 mg/day and high metabolic effects can be expected at low doses.

Running title: Influence of corticoids on FET uptake

Thus, an all or nothing effect on SUV_{brain} can be expected at doses used for brain tumor patients. Hence, the increase of $[^{18}\text{F}]\text{FET}$ uptake in normal brain tissue under dexamethasone therapy is a reproducible phenomenon which might interfere with the quantification of $[^{18}\text{F}]\text{FET}$ uptake in brain tumors especially for treatment monitoring of chemotherapy if dexamethasone therapy is initiated or interrupted in the meantime.

The key question in this context is whether $[^{18}\text{F}]\text{FET}$ uptake in the brain and the tumor is affected by the drug in the same way. As mentioned above, our data from animal experiments suggest that $[^{18}\text{F}]\text{FET}$ uptake in the tumor is not increased by dexamethasone treatment in contrast to brain tissue and that the TBR thereby changes [19]. For humans, this question cannot easily be answered by comparing $[^{18}\text{F}]\text{FET}$ uptake in the brain tumors of different patients with and without dexamethasone treatment since the tracer uptake varies largely between different types of brain tumors, causing high standard deviation when calculating statistics (Tab. 1). Repeated measurements in the same patients with and without dexamethasone in a short time interval with stable disease would be required, which we have not performed for ethical reasons. Nevertheless, the data of one patient with a brain metastasis and a short time interval between two $[^{18}\text{F}]\text{FET}$ PET scans with and without dexamethasone medication confirmed the hypothesis of a different influence of dexamethasone therapy on tracer uptake in the tumor and normal brain tissue. Maximum SUV of $[^{18}\text{F}]\text{FET}$ uptake in the tumor remained constant (-

Running title: Influence of corticoids on FET uptake

3%) while SUV_{brain} was 19% higher during dexamethasone therapy leading to a drop of the TBR_{max} by 18% (Tab. 3, Fig 5). In addition, the threshold-based determination of BTV using a value of 1.6 above background resulted in a change in volume of 70%.

Interestingly, Herholz and colleagues [25] reported a lower TBR_{max} of the amino acid tracer ^{11}C -L-methionine (MET) uptake in high grade gliomas after corticoid therapy but no SUV data are reported. It is tempting to speculate that the lower TBR of amino acid uptake during corticosteroid treatment was caused by higher $[^{11}\text{C}]\text{MET}$ uptake in the brain tissue, rather than lower $[^{11}\text{C}]\text{MET}$ uptake in the tumor.

Based on the results from our previous animal experiments we tested the short term effect of dexamethasone on $[^{18}\text{F}]\text{FET}$ uptake in a group of six healthy male rats which yielded the same increase of $[^{18}\text{F}]\text{FET}$ uptake of approx. 20% after 48 h of dexamethasone treatment (see supplementary table 1). This small time span argues against the involvement of long-term effects due to corticoid treatment and cross-effects with other medication.

The cause of the phenomenon is still unknown. It has been shown that glucocorticoids induce increased protein breakdown and consumption of leucine [26-27], while the transport system LAT1 itself has been shown to be unaffected in rat kidneys by glucocorticoids [28]. $[^{18}\text{F}]\text{FET}$, like leucine, is a substrate of the LAT transporter and uptake might be upregulated due to

Running title: Influence of corticoids on FET uptake

increased amino acid consumption and oxidation. This effect might be different in tumor cells, which already have an increased metabolism.

Furthermore, we observed higher [^{18}F]FET uptake in the healthy brain tissue of women than of men. Sex-specific differences in metabolism is a well-known phenomenon and leads to discussions in pre-clinical as well as in clinical studies [29-31]. Sex-specific metabolism might also be the reason for different increase of brain SUV after dexamethasone therapy in men in women. As there was no correlation for brain SUV and BMI (Fig. 4B) and a poor correlation for brain SUV and BMI in another patient population evaluated in our group [32], we assume that the high sex-specific differences are more likely due to differences in metabolism than to different volume of distribution caused by a higher percentage of muscle mass in men which shows increased FET uptake [33].

Our study is limited by its retrospective nature and the fact that the effects of dexamethasone treatment on TBR and BTV could be directly demonstrated in a single patient only. Therefore, a prospective study in the same patients is needed to confirm these effects. Nevertheless, the effect of dexamethasone treatment on TBR and BTV has been clearly demonstrated in animal experiments which in conjunction with this study represents an important information for clinicians in this field.

CONCLUSION

Running title: Influence of corticoids on FET uptake

In conclusion, dexamethasone treatment increases [^{18}F]FET uptake in the normal brain. An effect on TBRmax, TBRmean and BTV cannot be excluded which should be considered in particular during treatment monitoring with [^{18}F]FET PET and if BTV is used for treatment planning.

CONFLICT OF INTEREST STATEMENT

The authors declare that they have no conflict of interest.

ACKNOWLEDGEMENTS

The authors wish to thank Erika Wabbals, Silke Grafmüller and Sascha Rehbein for technical assistance in radiosynthesis of [^{18}F]FET and Silke Frensch, Suzanne Schaden, and Kornelia Frey for technical assistance in performing the PET measurements.

REFERENCES

1. Langen KJ, Watts C (2016) Neuro-oncology: Amino acid PET for brain tumours - ready for the clinic? *Nature reviews Neurology* 12:375-376.
2. Albert NL, Weller M, Suchorska B, et al. (2016) Response Assessment in Neuro-Oncology working group and European Association for Neuro-Oncology recommendations for the clinical use of PET imaging in gliomas. *Neuro-oncology*.
3. Galldiks N, Langen K, Holy R, et al. (2012) Assessment of treatment response in patients with glioblastoma using [18F]Fluoroethyl-L-Tyrosine PET in comparison to MRI. *Journal of nuclear medicine : official publication, Society of Nuclear Medicine* 53:1048-1057.
4. Pauleit D, Stoffels G, Bachofner A, et al. (2009) Comparison of (18)F-FET and (18)F-FDG PET in brain tumors. *Nuclear medicine and biology* 36:779-787.
5. Pichler R, Dunzinger A, Wurm G, et al. (2010) Is there a place for FET PET in the initial evaluation of brain lesions with unknown significance? *European journal of nuclear medicine and molecular imaging* 37:1521-1528.
6. Popperl G, Gotz C, Rachinger W, Gildehaus FJ, Tonn JC, Tatsch K (2004) Value of O-(2-[18F]fluoroethyl)- L-tyrosine PET for the diagnosis of recurrent glioma. *European journal of nuclear medicine and molecular imaging* 31:1464-1470.
7. Floeth FW, Pauleit D, Sabel M, et al. (2007) Prognostic value of O-(2-18F-fluoroethyl)-L-tyrosine PET and MRI in low-grade glioma. *Journal of nuclear medicine : official publication, Society of Nuclear Medicine* 48:519-527.
8. Jansen NL, Suchorska B, Wenter V, et al. (2015) Prognostic Significance of Dynamic 18F-FET PET in Newly Diagnosed Astrocytic High-Grade Glioma. *Journal of nuclear medicine : official publication, Society of Nuclear Medicine* 56:9-15.
9. Weckesser M, Langen KJ, Rickert CH, et al. (2005) O-(2-[18F]fluorethyl)-L-tyrosine PET in the clinical evaluation of primary brain tumours. *European journal of nuclear medicine and molecular imaging* 32:422-429.
10. Calcagni ML, Galli G, Giordano A, et al. (2011) Dynamic O-(2-[18F]fluoroethyl)-L-tyrosine (F-18 FET) PET for glioma grading: assessment of individual probability of malignancy. *Clinical nuclear medicine* 36:841-847.
11. Galldiks N, Langen KJ (2015) Applications of PET imaging of neurological tumors with radiolabeled amino acids. *The quarterly journal of nuclear medicine and molecular imaging : official publication of the Italian Association of Nuclear Medicine* 59:70-82.
12. Herholz K, Langen KJ, Schiepers C, Mountz JM (2012) Brain tumors. *Seminars in nuclear medicine* 42:356-370.

Running title: Influence of corticoids on FET uptake

13. Langen KJ, Hamacher K, Weckesser M, et al. (2006) O-(2-[18F]fluoroethyl)-L-tyrosine: uptake mechanisms and clinical applications. *Nuclear medicine and biology* 33:287-294.
14. Wester HJ, Herz M, Weber W, et al. (1999) Synthesis and radiopharmacology of O-(2-[18F]fluoroethyl)-L-tyrosine for tumor imaging. *Journal of nuclear medicine : official publication, Society of Nuclear Medicine* 40:205-212.
15. Kotsarini C, Griffiths PD, Wilkinson ID, Hoggard N (2010) A systematic review of the literature on the effects of dexamethasone on the brain from in vivo human-based studies: implications for physiological brain imaging of patients with intracranial tumors. *Neurosurgery* 67:1799-1815; discussion 1815.
16. Sarin R, Murthy V (2003) Medical decompressive therapy for primary and metastatic intracranial tumours. *Lancet neurology* 2:357-365.
17. Stegmayer C, Bandelow U, Oliveira D, et al. (2017) Influence of blood-brain barrier permeability on O-(2-18F-fluoroethyl)-L-tyrosine uptake in rat gliomas. *European journal of nuclear medicine and molecular imaging* 44:408-416.
18. Stegmayer C, Oliveira D, Niemietz N, et al. (2017) Influence of bevacizumab on blood-brain barrier permeability and O-(2-18F-fluoroethyl)-L-tyrosine uptake in rat gliomas. *Journal of nuclear medicine : official publication, Society of Nuclear Medicine*.
19. Stegmayer C, Schoneck M, Oliveira D, et al. (2016) Reproducibility of O-(2-(18)F-fluoroethyl)-L-tyrosine uptake kinetics in brain tumors and influence of corticoid therapy: an experimental study in rat gliomas. *European journal of nuclear medicine and molecular imaging* 43:1115-1123.
20. Hamacher K, Coenen HH (2002) Efficient routine production of the 18F-labelled amino acid O-2-18F fluoroethyl-L-tyrosine. *Applied radiation and isotopes : including data, instrumentation and methods for use in agriculture, industry and medicine* 57:853-856.
21. Langen KJ, Bartenstein P, Boecker H, et al. (2011) [German guidelines for brain tumour imaging by PET and SPECT using labelled amino acids]. *Nuklearmedizin Nuclear medicine* 50:167-173.
22. Herzog H, Langen KJ, Weirich C, et al. (2011) High resolution BrainPET combined with simultaneous MRI. *Nuklearmedizin Nuclear medicine* 50:74-82.
23. Rota Kops E, Hautzel H, Herzog H, Antoch G, Shah NJ (2015) Comparison of Template-Based Versus CT-Based Attenuation Correction for Hybrid MR/PET Scanners. *IEEE Transactions on Nuclear Science* 62:2115-2121.
24. Rapp M, Heinzl A, Galldiks N, et al. (2013) Diagnostic performance of 18F-FET PET in newly diagnosed cerebral lesions suggestive of glioma. *Journal of nuclear medicine : official publication, Society of Nuclear Medicine* 54:229-235.
25. Herholz K, Holzer T, Bauer B, et al. (1998) 11C-methionine PET for differential diagnosis of low-grade gliomas. *Neurology* 50:1316-1322.

26. Beaufrere B, Horber FF, Schwenk WF, et al. (1989) Glucocorticosteroids increase leucine oxidation and impair leucine balance in humans. *The American journal of physiology* 257:E712-721.
27. Garrel DR, Moussali R, De Oliveira A, Lesiege D, Lariviere F (1995) RU 486 prevents the acute effects of cortisol on glucose and leucine metabolism. *J Clin Endocrinol Metab* 80:379-385.
28. Schwertfeger M, Pissowotzki K, Fleck C, Taylor PM (2003) Regulation of L-leucine transport in rat kidney by dexamethasone and triiodothyronine. *Amino acids* 25:75-83.
29. Varlamov O, Bethea CL, Roberts CT, Jr. (2014) Sex-specific differences in lipid and glucose metabolism. *Front Endocrinol (Lausanne)* 5:241.
30. Clayton JA, Collins FS (2014) Policy: NIH to balance sex in cell and animal studies. *Nature* 509:282-283.
31. Fuller CM, Insel PA (2014) I don't know the question, but sex is definitely the answer! Focus on "In pursuit of scientific excellence: sex matters" and "Do you know the sex of your cells?". *American journal of physiology Cell physiology* 306:C1-2.
32. Verger A, Stegmayr C, Galldiks N, et al. (2018) Evaluation of factors influencing ¹⁸F-FET uptake in the brain. *NeuroImage Clinical* 17:491-497.
33. Pauleit D, Floeth F, Herzog H, et al. (2003) Whole-body distribution and dosimetry of O-(2-[¹⁸F]fluoroethyl)-L-tyrosine. *European journal of nuclear medicine and molecular imaging* 30:519-524.

Tab. 1

Patient population and distribution of diseases.

* *P*-values for male vs. female patients and dexamethasone treatment vs no dexamethasone treatment (w/o). # For post-hoc/subgroup *P*-values see Fig. 3.

	Men w/o	Women w/o	Men under dexamethasone	Women under dexamethasone	<i>P</i> -value
n	40	40	40	40	
Age	52.5 ± 10.9	48.7 ± 12.7	51.8 ± 10.5	53.1 ± 11.2	n.s.
Dexamethasone (mg)	-	-	9.0 ± 6.8	7.3 ± 5.2	n.s.
HGG	28	18	25	21	n.s.
LGG	8	14	3	5	n.s.
Metastasis	2	2	10	11	n.s.
Unclear/other	2	6	2	2	n.s.

Pretreatment					
Surgery	26	20	26	25	n.s.
Radiotherapy	21	17	30	32	n.s.
Chemotherapy	19	18	22	22	n.s.

SUV _{cortex}	1.06 ± 0.16	1.33 ± 0.19	1.31 ± 0.28	1.45 ± 0.25	< 0.001* [#]
SUV _{wm}	0.85 ± 0.12	1.08 ± 0.17	1.02 ± 0.22	1.21 ± 0.19	≤ 0.005*
AUC _{cortex}	24.82 ± 3.56	30.78 ± 4.33	30.58 ± 6.46	33.59 ± 5.62	< 0.001*
TBR _{mean}	2.02 ± 0.54	1.92 ± 0.53	2.08 ± 0.44	1.93 ± 0.33	n.s.
TBR _{max}	2.67 ± 1.07	2.40 ± 1.13	2.68 ± 1.17	2.38 ± 0.78	n.s.

Tab. 2

SUV brain of nine patients with and without dexamethasone scanned multiple times. Except of the last patient, the SUV during dexamethasone therapy was always higher than the SUV without therapy.


Patient, sex	FET PET	SUV _{brain}	Dexamethasone
1, m	1	1.33	Yes
	2	1.12	No
2, f	1	1.61	No
	2	1.28	No
	3	1.62	Yes
	4	1.93	Yes
3, m	1	1.57	Yes
	2	1.48	No
	3	1.23	No
4, f	1	1.38	No
	2	1.95	Yes
5, m	1	1.20	No
	2	1.72	Yes
6, f	1	1.36	No
	2	1.89	Yes
	3	1.38	No
7, m	1	1.32	No
	2	1.12	No
	3	1.49	Yes
8, f	1	1.13	No
	2	1.43	Yes
9, f	1	1.35	No
	2	1.28	Yes

Tab. 3

a, [^{18}F]FET PET parameters of a patient with a brain metastasis investigated twice, during dexamethasone treatment (Sep/15) and without (Nov/15).


a	Dexamethasone 09/15	No dexamethasone 11/15	Difference under dexamethasone treatment
SUV _{brain}	1.33	1.12	+ 19 %
Tumor volume (ml) (Threshold based)	2.72	8.99	- 70 %
Tumor SUV _{mean} (Threshold based)	2.35	2.06	+ 14 %
Tumor SUV _{max}	2.07	2.14	- 3 %
TBR _{mean}	1.77	1.84	- 4 %
TBR _{max}	1.56	1.91	- 18 %

Fig. 1




Time-activity-curves (mean SUV with SD) of dexamethasone-treated (Dex) and non-treated (w/o) men (a) and women (b). Under dexamethasone therapy, the mean SUV increases in men more than in women.

Fig. 2



Time-activity-curves (mean SUV_{brain} with SD) of female (f) and male (m) patients without dexamethasone therapy (a) and with therapy (b). The mean SUV of female patients is higher than the mean SUV of male patients, regardless of therapy.


Fig. 3



The boxplot diagram shows significant differences between the subgroups. Men not receiving dexamethasone (m w/o) show lowest SUV in healthy brain tissue, women under dexamethasone therapy (f dexamethasone) have highest uptake.


* $P \leq 0.05$; ** $P \leq 0.01$; *** $P \leq 0.001$

Fig. 4



Correlation plots of SUV_{brain} vs. (a) dose of dexamethasone per day, and (b) body mass index (BMI), separately for men and women. SUV_{brain} vs. dose of dexamethasone correlated in men ($P < 0.01$; $r = 0.44$), but not in women.

Fig. 5



Contrast enhanced MR images (left) and corresponding $[^{18}\text{F}]\text{FET}$ PET SUV images of a male patient with brain metastasis at an interval of 10 weeks. Visually, no significant tumor growth could be observed in MR or PET images. At the time of first PET scan (a), the patient received dexamethasone. Nine weeks later (b), dexamethasone was already discontinued and background activity was visually lower, resulting in higher tumor volume due to threshold based VOI definition (Tab. 2).

Supplementary data

Table 1

[¹⁸F]FET SUV_{brain} of six male healthy rats, measured with and without dexamethasone.

Rat No.	Baseline SUV _{brain}	Dexamethasone SUV _{brain}	Difference
1	0.67	0.80	+20 %
2	0.66	0.81	+24 %
3	0.71	0.81	+14 %
4	0.80	0.86	+7 %
5	0.79	0.80	+2 %
6	0.77	0.75	-2 %
t-test	0.75 ± 0.06	0.81 ± 0.04	P=0.016