

Introduction

- **Cognitive sex differences** well documented in **behavioral and functional brain imaging** (fMRI).
- Structural MRI has identified a **structural sexual dimorphism** of the human brain (3).
- **Sex differences** are also found in **resting state (RS) brain connectivity** (e.g. 8, 11).

Aims of the present study:

1. Employ a **machine learning approach** on RS data to address generalizability of previous findings to independent samples.
2. Delineate **regionally specific brain networks** underlying successful classification of novel subjects' sex.
3. Further understanding of a possible **sexual dimorphism of the RS connectome**.

Methods

Samples:

- Two mutually exclusive samples of unrelated subjects constructed of Human Connectome Project data (HCP S1200 release, (7)).
- Sample 1: 434 subjects (217 males, age range: 22-37, mean age: 28.6 years),
- Sample 2: 310 subjects (155 males, age range: 22-36, mean age: 28.5 years).
- Males and females matched for age, twin-status and education within each sample.

Functional imaging data:

- Resting state (RS): 1200 volumes per subject.
- Siemens Skyra 3T scanner (TR=720ms).
- Standard realignment and normalization.
- FSL-FIX denoising (5).
- Individual RS connectomes extracted for 436 ROIs based on (6).

Sex Classification:

- Linear SVM (LibSVM toolbox, (1)) model for classification of subjects' sex from RS connectome.
- Nested optimization of cost parameter.
- 10 repetitions of a 10-fold cross-validation.
- Across sample classification: fitting of the model on sample 1 and testing it on sample 2.

Whole brain vs. ROI based classification:


- (1) Whole brain connectome.
- (2) Each individual ROI's connectivity profile (436 parcels).

Result Summary:


- ROI based analyses performed separately for each sample and conservatively characterized by minimum across the two samples.

Results


1 ROI based minimum classification accuracy across both samples



2 ROI based classification accuracy for between-sample classification



3 Classification accuracy for the whole connectome analysis and the ten ROIs with highest classification accuracy across sample 1 and sample 2



Whole Brain Connectome

- 10-fold cross-validation performance for whole brain connectome:
 - Sample 1: 79.3%
 - Sample 2: 78.8%
- Across sample classification performance:
 - 81.4%.
 (possibly due to larger training set)

Whole brain RS connectome allows for the prediction of an unknown subject's sex at ~ 80% accuracy!

Regional Connectivity

- ROI based analyses identified regions for which the connectivity profile differentiated most strongly between the sexes.
- Highest regional accuracies:
 - Medial brain regions in anterior cingulate and cingulate gyrus.
 - Left lateralized inferior frontal gyrus and inferior temporal gyrus
- Regions displaying top classification accuracies highly similar for within-sample and between-sample classification.

Classification accuracies for top ROIs are only marginally lower than whole connectome analyses!

Discussion

- Both within- and between-sample cross-validation allowed reliable classification of unknown subjects' sex from RS connectivity profiles → **robust sexual dimorphism of the resting state connectome**.
- Predictive power of local brain connectivity almost as high as whole brain connectivity → **regionally specific effects**.
- Within- and between-sample prediction based on highly similar brain regions → **reliability of findings**.
- Regions with top prediction power are mainly

- located along (**anterior**) **cingulate cortex**.
- Cortical regions with top prediction accuracies are **left lateralized**.
- Left inferior frontal **inter-hemispheric connectivity** has been shown to **vary across the menstrual cycle in women**, but to remain **stable in men** (9).
- Similar frontal regions reported in relation to **differing cognitive strategies between the sexes** (10).
- Sex differences in cingulate cortex reported in connection with **emotional reactivity and**

- cognitive control of emotion** (2,4).
- Regions most clearly differentiating between the sexes are related to **cognitive control of behaviour and emotion**.
- Findings might help explain why **sex differences** are mostly found in **cognitive strategies** employed by men and women, but not in behavioural performance per se.
- Results substantiate a **sexual dimorphism in RS connectivity** → male and females differ not only in brain structure, but also in functional brain organization.

1) Chang, C.C., Lin, C.J. (2011). ACM Transactions on Intelligent Systems and Technology, 2, 27:1-27.
2) Domes, G., Schulze, L., Böttger, M., Grossmann, A., Hauenstein, K., Wirtz, P.H., Heinrichs, M., Herpertz, S.C. (2010). Hum Brain Mapp, 31(5):758-69.
3) Goldstein, J.M., Seidman, L.J., Horton, N.J., Makris, N., Kennedy, D.N., Caviness, V.S. Jr, Faraone, S.V., Tsuang, M.T. (2001). Cereb Cortex, 11(6):490-497.
4) Koch, K., Pauly, K., Kellermann, T., Seifert, N.Y., Reske, M., Backes, V., Stöcker, T., Shah, N.J., Amunts, K., Kircher, T., Scheider, F., Habel, U. (2007). Neuropsychologia, 45(12):2744-54.
5) Salimi-Khorshidi, G., Douaud, G., Beckmann, C.F., Glasser, M.F., Griffanti, L., Smith S.M. (2014). NeuroImage, 90: 449-468.
6) Schaefer, A., Kong, R., Gordon, E.M., Laumann, T.O., Zuo, X.N., Holmes, A.J., Eickhoff, S.B., Yeo, B.T.T. (2017). Cerebral Cortex, 18:1-20.
7) Van Essen, D.C. (2012). NeuroImage, 62: 2222-2231.
8) Weis, S., Hodgetts, S., Hausmann, M. (2017). Brain Cogn., in press.
9) Weis, S., Hausmann, M., Stoffers, B., Vohn, R., Kellermann, T., Sturm, W. (2008). J Neurosci., 28(50):13401-13410.
10) Weiss, E., Siedentopf, C.M., Hofer, A., Deisenhammer, E.A., Hoptman, M.J., Kremser, C., Golaszewski, S., Felber, S., Fleischhacker, W.W., Delazer, M. (2003). Neurosci Lett., 344(3):169-172.
11) Zhang, C., Dougherty, C.C., Baum, S.A., White, T., Michael, A.M. (2018). Hum Brain Mapp. 39(4):1765-1776.