



Aircraft-based observations of isoprene epoxydiol-derived secondary organic aerosol (IEPOX-SOA) in the tropical upper troposphere over the Amazon region

Christiane Schulz¹, Johannes Schneider¹, Bruna Amorim Holanda¹, Oliver Appel¹, Anja Costa², Suzane S. de Sá³, Volker Dreiling⁴, Daniel Fütterer⁵, Tina Jurkat-Witschas⁵, Thomas Klimach¹, Martina Krämer², Scot T. Martin^{3,7}, Stephan Mertes⁸, Mira L. Pöhlker¹, Daniel Sauer⁵, Christiane Voigt^{5,6}, Bernadett Weinzierl^{5,a}, Helmut Ziereis⁵, Martin Zöger⁴, Meinrat O. Andreae^{1,9}, Paulo Artaxo¹⁰, Luiz A. T. Machado¹¹, Ulrich Pöschl¹, Manfred Wendisch¹², and Stephan Borrmann^{1,6}

¹Particle Chemistry, Biogeochemistry and Multiphase Chemistry Departments, Max Planck Institute for Chemistry, Mainz, Germany

²Institut für Energie- und Klimaforschung (IEK7), Forschungszentrum Jülich GmbH, Jülich, Germany

³School of Engineering and Applied Sciences, Harvard University, Cambridge, Massachusetts, USA

⁴Flight Experiments, German Aerospace Center (Deutsches Zentrum für Luft- und Raumfahrt), Oberpfaffenhofen, Germany

⁵Institute for Atmospheric Physics, German Aerospace Center (Deutsches Zentrum für Luft- und Raumfahrt), Oberpfaffenhofen, Germany

⁶Institute for Physics of the Atmosphere, Johannes Gutenberg University, Mainz, Germany

⁷Department of Earth and Planetary Sciences, Harvard University, Cambridge, Massachusetts, USA

⁸Leibniz Institute for Tropospheric Research, Leipzig, Germany

⁹Scripps Institution of Oceanography, University of California San Diego, La Jolla, California, USA

¹⁰Instituto de Física, Universidade de São Paulo, São Paulo, Brazil

¹¹Instituto Nacional de Pesquisas Espaciais (INPE), Centro de Previsão de Tempo e Estudos Climáticos, São José dos Campos, Brazil

¹²Leipzig Institute for Meteorology, University of Leipzig, Leipzig, Germany

^anow at: Faculty of Physics, University of Vienna, Vienna, Austria

Correspondence: Johannes Schneider (johannes.schneider@mpic.de)

Abstract. During the ACRIDICON-CHUVA field project (September - October 2014; based in Manaus, Brazil) aircraft-based in-situ measurements of aerosol chemical composition were conducted in the tropical troposphere over the Amazon using the High Altitude and Long Range Research Aircraft (HALO), covering altitudes from the boundary layer height up to 14.4 km. The submicron non-refractory aerosol was characterized by flash-vaporization/electron impact-ionization aerosol particle mass spectrometry. The results show that significant secondary organic aerosol (SOA) formation by isoprene oxidation products occurs in the upper troposphere, leading to increased organic aerosol mass concentrations above 10 km altitude. The median organic mass concentrations in the upper troposphere above 10 km range between 1.0 and 2.1 $\mu\text{g m}^{-3}$ (referring to standard temperature and pressure; STP) with interquartile ranges of 0.6 to 3.0 $\mu\text{g m}^{-3}$ (STP), representing 70 % of the total submicron non-refractory aerosol particle mass. The presence of isoprene epoxydiol-derived isoprene secondary organic aerosol (IEPOX-SOA) was confirmed by marker peaks in the mass spectra. We estimate the contribution of IEPOX-SOA to the total organic aerosol in the upper troposphere to be about 20 %. After isoprene emission from vegetation, oxidation processes



occur at low altitudes and/or during transport to higher altitudes, which may lead to the formation of IEPOX (one oxidation product of isoprene). Reactive uptake or condensation of IEPOX on pre-existing particles leads to IEPOX-SOA formation and subsequently increasing organic mass in the upper troposphere. This organic mass increase was accompanied by an increase of the nitrate mass concentrations, most likely due to NO_x production by lightning. We further found that the ammonium
 5 contained in the aerosol particles is not sufficient to neutralize the particulate sulfate and nitrate. Analysis of the ion ratio of NO^+ to NO_2^+ indicated that nitrate in the upper troposphere exists mainly in the form of organic nitrate. IEPOX-SOA and organic nitrates are coincident with each other, indicating that IEPOX-SOA forms in the upper troposphere either on acidic nitrate particles forming organic nitrates derived from IEPOX or on already neutralized organic nitrate aerosol particles.

1 Introduction

10 Volatile organic compounds (VOCs) emitted by vegetation can lead to the formation of secondary organic aerosol (SOA) through atmospheric oxidation and further chemical processes (e.g., Zhang et al., 2007; Jimenez et al., 2009; Hallquist et al., 2009; Shrivastava et al., 2017). One important VOC is isoprene (C_5H_8 , 2-methyl-1,3-butadiene), the most abundant non-methane hydrocarbon with a global emission rate of $\sim 500 \text{ Tg y}^{-1}$, with a large contribution coming from the Amazon rain forest (Guenther et al., 2012). Also the conditions for photooxidative reactions are favoured in this tropical region. Isoprene is
 15 a short-lived atmospheric gas, which is oxidized in the atmosphere by reactions with the hydroxyl radical (OH), nitrate radical (NO_3), or ozone (O_3). OH-initiated oxidation leads to isoprene peroxide radicals (ISOPOO). Depending on the NO_x and HO_x concentrations, ISOPOO can react further through different pathways.

For HO_2 dominant conditions, meaning conditions with low amounts ($< 1 \text{ ppb}$) of NO (Wennberg, 2013), ISOPOO will mainly react with hydroperoxyl radicals (HO_2) to the intermediate oligomer hydroxyhydroperoxides (ISOPOOH). Further oxidation
 20 of ISOPOOH may lead to isoprene epoxydiols (IEPOX), which then can partition into the particle phase by condensation or reactive uptake and, thus, SOA derived from IEPOX (IEPOX-SOA) can be formed (e.g., Claeys et al., 2004; Carlton et al., 2009; Paulot et al., 2009; Surratt et al., 2010; Lin et al., 2012; Budisulistiorini et al., 2013; Worton et al., 2013; St. Clair et al., 2016; Liu et al., 2016b). Laboratory studies demonstrate that around 50 % of isoprene-derived particulate matter is associated with IEPOX production and uptake through the HO_2 pathway and in the presence of acidic aerosol particles (Liu et al.,
 25 2015). The existence of aerosol as seed particles seems to be necessary (Surratt et al., 2010; Lin et al., 2012), but also the acidity of aerosol can influence the formation yield of IEPOX-SOA. Laboratory and field studies found a correlation between IEPOX-SOA and sulfate, which is related to the acidity of aerosol (Surratt et al., 2007; Budisulistiorini et al., 2013). Although the IEPOX pathway is considered as the dominant one, further laboratory studies show that also other gas-phase reactions of ISOPOOH with multifunctional hydroperoxides contribute to the formation of isoprene-derived SOA (e.g., Krechmer et al.,
 30 2015; Liu et al., 2016b; Riva et al., 2016).

A different oxidation pathway occurs at increased NO concentrations ($> 1 \text{ ppb}$) in NO dominant conditions (Wennberg, 2013). Reaction pathways for ISOPOO change towards reactions with NO instead of HO_2 , leading to hydroxynitrates and/or hydrox-yalkoxy radicals, which decompose further to methacrolein (MACR) and methyl vinyl ketone (MVK). Whereas MACR is a



precursor for isoprene-derived SOA, MVK has almost no SOA contribution (Surratt et al., 2006; Liu et al., 2016a).

In the central Amazon region, both HO₂ and NO dominated conditions were observed dependent on the time and location of measurements (Kuhn et al., 2010; Andreae et al., 2015; Martin et al., 2016). Thus, isoprene can undergo different gas-phase reactions depending on several conditions, leading either to IEPOX-SOA or to other types of isoprene-derived SOA.

- 5 Several field studies for investigating IEPOX-SOA were conducted in different regions. The extensive study by Hu et al. (2015) summarizes those with a focus on IEPOX-SOA measured by aerosol mass spectrometry. The tracer ion at mass to charge ratio (m/z) 82 (C₅H₆O) was identified to be related to IEPOX-SOA in combination with the ion at m/z 53 (C₄H₅) (Robinson et al., 2011; Allan et al., 2014; Hu et al., 2015). In order to quantify the amount of IEPOX-SOA measured by aerosol mass spectrometry, the ratio of the signal at m/z 82 to the whole organic signal was introduced and is defined as f_{82} (e.g., Budisulistiorini et al., 2013; Allan et al., 2014; Hu et al., 2015). Background values of f_{82} were calculated on the basis of different field studies
 10 worldwide. In areas with predominant monoterpene emissions, the background value is $3.1 \pm 0.6 \%$. Areas with a strong biomass burning and urban influence a background value of $1.7 \pm 0.1 \%$ is reported, whereas ambient organic aerosol with a strong isoprene influence under low NO conditions shows an increased value of $6.5 \pm 2.2 \%$ (Hu et al., 2015).

- The AMAZE-08 campaign focused on SOA production mechanisms at a pristine continental site in the Amazon basin during
 15 the wet season (Martin et al., 2010). The submicron aerosol particles were found to be dominated by secondary organic material (Chen et al., 2009). From positive matrix factorization (PMF) analysis one factor was associated with the reactive uptake of IEPOX to acidic haze, fog or cloud droplets (Chen et al., 2015). Another field campaign that was conducted in the Amazon basin is the GoAmazon2014/5 field campaign (Martin et al., 2016). One focus of GoAmazon2014/5 is the study of aerosol sources and SOA formation and aging comparing the dry and wet season and the influence of urban pollution (Shilling et al.,
 20 2018). Another focus is the investigation of parameters influencing the pathways for isoprene oxidation. Shifts in the prevailing regime of NO or HO₂ pathways for isoprene photooxidation in the central region of Amazonia were studied (Liu et al., 2016b). Also, shifts in the production of IEPOX-SOA with changing concentrations of sulfate and NO_x in the boundary layer of central Amazon region were studied as part of GoAmazon2014/5 (de Sá et al., 2017). Increased NO concentrations suppress IEPOX-SOA production. Despite the enhancing effect of increased sulfate concentrations for IEPOX-SOA production, the NO effect
 25 is dominating (de Sá et al., 2017). During another airborne measurement campaign in the Amazon rain forest (SAMBBA), the highest f_{82} values were around 9 % measured at the top of the boundary layer with a maximum flight altitude of 5 km (Allan et al., 2014). A recent paper by Mei et al. (2018) compares aircraft measurements from the GoAmazon2014/5 campaign with data collected during the ACRIDICON-CHUVA campaign and with measurements taken at a ground station in the Amazon (T3). The results show good agreement for many measured atmospheric parameters and gives the opportunity to validate the
 30 data quality for several measurements conducted on different sampling platforms (Mei et al., 2018).

- Very few measurements at altitudes higher than the boundary layer were reported. From single particle mass spectrometric measurements IEPOX sulfate esters were identified in 80 % and 50 % of the analyzed particles measured at altitudes of 5 and 10 km in the tropical free troposphere, respectively (Froyd et al., 2010). In the boundary layer, IEPOX sulfate ester in aerosol particles did not occur. This is explained by a low abundance of acidic aerosol particles acting as seed particles and with a
 35 relatively short time since the emission of isoprene (Froyd et al., 2010). However, with lofting of isoprene and its derivatives



above the boundary layer, it is suggested that IEPOX can partition more efficiently to acidic aerosol particles. The reactive uptake of IEPOX contributed 1-20 % of the tropospheric aerosol mass in the tropics where continental convection was active (Froyd et al., 2010).

The presence of SOA in the tropical upper troposphere can have different effects. For example, it can be entrained into the tropical transition layer (TTL) from where further slow, radiatively-driven lifting could transport them into the lower stratosphere (Fueglistaler et al., 2009). There, and in the TTL region of enhanced new particle formation near the tropopause, the SOA could become part of the global tropical layer of elevated submicron particle abundances (Borrmann et al., 2010; Weigel et al., 2011). It has also been suggested that aerosol formation in the upper troposphere can provide a source for cloud condensation nuclei for lower altitudes in the Amazon region (Wang et al., 2016; Andreae et al., 2018).

Another interesting aspect is the presence of organic nitrates. Many studies reported on the difficulties to measure organic nitrates quantitatively, but suggested also a possibility to estimate the amount or at least the presence of organic nitrates based on the measured ion ratios of NO^+ to NO_2^+ using aerosol mass spectrometry (e.g., Bruns et al., 2010; Farmer et al., 2010; Rollins et al., 2010; Fry et al., 2013; Ayres et al., 2015; Kiendler-Scharr et al., 2016; Ng et al., 2017; Schneider et al., 2017). Oxidation of VOCs with the highly reactive nitrate radical NO_3 can lead to different nitrogen-containing oxidation products that can partition to the aerosol-phase (e.g., Kroll and Seinfeld, 2008; Shrivastava et al., 2017). The nitrate radical oxidation of VOCs can contribute up to 20 % of the global VOC oxidation and is supposed to increase the aerosol mass significantly (Boyd et al., 2015).

Field measurements have shown that the major aerosol-phase product of monoterpene oxidation with nitrate radical is likely a hydroperoxy nitrate ($\text{C}_{10}\text{H}_{17}\text{NO}_5$) whereas the analogous isoprene oxidation product has a contribution of less than 1 % of the total organic nitrate and occurs more in gas phase (Ayres et al., 2015). Laboratory studies suggested that isoprene-derived organic nitrates are formed from SOA reactions, but undergo substitution reactions in which nitrate is substituted by sulfate (Darer et al., 2011). Studies from the south eastern US showed, that organic nitrate aerosol particles from monoterpenes are strongly influenced by anthropogenic pollutants and may contribute to 19-34 % of the total organic aerosol content (Xu et al., 2015). Polluted urban regions are often dominated by inorganic nitrates (Farmer et al., 2010). However, in rural forested areas a dominance of particulate organic nitrates formed from oxidation of monoterpenes was reported (Fry et al., 2013). A recent study from the Amazon showed with measurements at ground level that up to 87 % of the total nitrate can be attributed to organic nitrate (de Sá et al., 2018).

This study presents submicron aerosol chemical composition measurements and focuses on the presence of IEPOX-SOA at different altitudes above the Amazon. The analysis herein uses data from the ACRIDICON-CHUVA campaign, which was conducted in September 2014 (Wendisch et al., 2016). The study gives insights into the photooxidative state of organic aerosols, the presence of IEPOX-SOA, and also the presence of particulate organic nitrates. A comparison of these parameters for different altitudes is presented.



2 The ACRIDICON-CHUVA campaign and instrumentation on HALO

The aircraft campaign ACRIDICON-CHUVA aimed at the investigation of convective cloud systems to better understand and to quantify aerosol-cloud-interactions and radiative effects of convective clouds. ACRIDICON is the acronym for “Aerosol, Cloud, Precipitation, and Radiation Interactions and Dynamics of Convective Cloud Systems”; CHUVA stands for “Cloud Processes of the Main Precipitation Systems in Brazil: A Contribution to Cloud Resolving Modeling and to the GPM (Global Precipitation Measurement)”. This campaign was performed with HALO (High Altitude and Long Range Research Aircraft), which is operated by the German Aerospace Center (DLR). During September and October 2014 (dry season (Andreae et al., 2015)), 14 flights were conducted in the region around Manaus, Brazil, with a radius of 1300 km above the Amazon rain forest and to the Atlantic coast. Altitudes up to 14.4 km were reached. HALO was equipped with instruments for measuring basic meteorological parameters, atmospheric radiation, trace gases, and aerosol properties, such as size, number, and mass concentration and chemical composition. An overview on the campaign, its objectives and the instrumentation properties and uncertainties can be found in Wendisch et al. (2016), Voigt et al. (2017) and Machado et al. (2017).

2.1 In-situ aircraft instrumentation

2.1.1 Basic meteorological data

Basic meteorological data were obtained from the BAsic HALO Measurement And Sensor System (BAHAMAS) at 1 s time resolution. BAHAMAS acquires data from air flow and thermodynamic sensors as well as from the aircraft avionics and a high precision inertial reference system to derive basic meteorological parameters like pressure, temperature and the 3-D wind vector as well as aircraft position and altitude. Water vapour mixing ratio and further derived humidity parameters are measured by SHARC (Sophisticated Hygrometer for Atmospheric ResearCh) based on direct absorption measurement by a tunable diode laser (TDL) system. Typical absolute accuracy of the basic meteorological data is 0.5 K for temperature, 0.3 hPa for pressure, 0.4-0.6 m s⁻¹ for wind and (5 % ± 1 ppmv) for water vapour volume mixing ratio.

2.1.2 Aerosol number concentration measurements

Aerosol particle number concentrations were measured using the Aerosol Measurement System (AMETYST). It includes four butanol-based condensation particle counters (modified model 5.410 by Grimm Aerosol Technik, Ainring, Germany) with flow rates of 0.6 and 0.3 l min⁻¹, configured with different nominal lower cutoff diameters at 4 nm and 10 nm (set via the temperature difference between saturator and condenser). AMETYST samples behind the HALO Aerosol Submicrometer Inlet (HASI) mounted on top of the fuselage, which provides near-isokinetic sampling of aerosol particles up to diameters of 2-3 µm. During this campaign, sampling line losses and inlet transmission limited the lower size cutoff to about 20 nm in the upper troposphere. As cloud hydrometeors are found to cause artefacts in the detected number concentrations, cloud passages have been removed from the data set. For details see Wendisch et al. (2016) and Andreae et al. (2018).



2.1.3 Submicron particle chemical composition measurements

During the ACRIDICON-CHUVA campaign we operated a Compact Time-of-Flight Aerosol Mass Spectrometer (C-ToF-AMS) (Drewnick et al., 2005; Canagaratna et al., 2007; Schmale et al., 2010) to investigate background aerosol composition around deep convective cloud systems. The C-ToF-AMS was connected to the HASI for sampling aerosol particles. The instrument was connected via a 6.25 mm (1/4-inch) stainless steel tubing. Aerosol particles enter the C-ToF-AMS via a constant pressure inlet controlling the volumetric flow into the instrument. In contrast to the classical approach (Bahreini et al., 2008) this custom-made (at the Max Planck Institute for Chemistry) device consists of two plates with a flexible orifice in between and is connected to a rotor. Squeezing or relaxing the orifice leads to a change of the volumetric flow rate into the instrument such that the pressure in the aerodynamic lens can be maintained constant. Thus, a constant mass flow into the instrument is achieved. Therefore, all data reported here represent conditions during the calibration on ground with a pressure of 995 hPa and temperatures of 300 K. These conditions define our standard temperature and pressure (STP). Further details of the constant pressure inlet are subject of a separate publication (Molleker, 2018). The C-ToF-AMS was operated with a time resolution of 30 s, which corresponds to roughly 6 km flight path. An overall accuracy of about 30 % has been reported in previous studies (Canagaratna et al., 2007; Middlebrook et al., 2012).

The refractory black carbon (rBC) particles were measured with a Single Particle Soot Photometer (SP2, Droplet Measurement Techniques, Longmont, CO, USA). The instrument uses a laser-induced incandescence technique to quantify the mass of rBC in individual aerosol particles (Stephens et al., 2003; Schwarz et al., 2006). Calibrations of the incandescent signal were conducted before, during and after the campaign using size-selected fullerene soot particles. The scattering signal was calibrated using either spherical polystyrene latex size standards or ammonium sulfate particles of different diameters selected by a differential mobility analyzer.

2.1.4 NO and NO_y measurements

Measurements of nitric oxide (NO) and total reactive nitrogen (NO_y) were conducted by a dual-channel chemiluminescence detector (CLD-SR, Eco Physics). Ambient air is sampled via a standard HALO trace gas inlet with a Teflon tube. For the NO_y channel, the chemi-luminescence detector is combined with a custom-built gold converter that reduces all oxidized reactive nitrogen species to NO (Ziereis et al., 2000). NO from ambient air reacts with O₃ produced from an ozone generator in a chamber resulting in excited NO₂. The emitted luminescence signal is detected. The detector channel is equipped with a pre-reaction chamber for determining cross-reactions of other compounds in ambient air reacting with O₃. The time resolution is 1 s. The precision and accuracy of the measurements depend on the ambient concentrations, with typical values of 5 % and 7 % (NO) and 10 % and 15 % (NO_y), respectively.



Table 1. Relative ionisation efficiencies (RIEs) used for data analysis. RIE_{NH_4} and RIE_{SO_4} are determined from calibrations with ammonium nitrate and sulfate before, during and after the campaign. RIE_{NO_3} and RIE_{Org} are literature values (Alfarra et al., 2004; Canagaratna et al., 2007).

RIE_{NO_3}	RIE_{NH_4}	RIE_{SO_4}	RIE_{Org}
1.1	3.77	0.89	1.4

3 Data analysis

3.1 General C-ToF-AMS data approaches

The C-ToF-AMS was calibrated with monodisperse ammonium nitrate and sulfate before, during and after the campaign in order to estimate relative ionisation efficiencies (RIE) for nitrate and sulfate. The resulting calibration values can be found in Tab.1.

A value of 0.5 for the collection efficiency (CE) was used for all flights (Matthew et al., 2008; Middlebrook et al., 2012). The main fraction of the measured aerosol mass consists of organic matter (see Sect. 4.2), which has no clear effect on the CE. Furthermore, large amounts of nitrate that would lead to a higher CE were not encountered during the campaign (Middlebrook et al., 2012).

Detection limits were derived as three times the standard deviation from the background signal according to Reitz (2011). Here, a time-dependent cubic spline function was used to determine a detection limit for each data point. Averaged over a single flight, the detection limit is around $0.11 \mu\text{g m}^{-3}$ for organics, $0.02 \mu\text{g m}^{-3}$ for nitrate, $0.03 \mu\text{g m}^{-3}$ for sulfate and $0.09 \mu\text{g m}^{-3}$ for ammonium. It should be noted here that ammonium can experience additional uncertainty due to cross-relations with water in the fragmentation table (Allan et al., 2004). A vertical profile of the averaged detection limits for all species can be found in the supplement (see Fig. S1).

3.2 Oxidation state of the organic aerosol

The oxidation state of the organic aerosol indicates the degree of photochemical aging. It can be determined using the correlation between the ratio of the signal at m/z 44 (mostly CO_2^+) to the total organic signal, f_{44} , and the ratio of the signal at m/z 43 (mostly $\text{C}_2\text{H}_3\text{O}^+$) to the total organic signal, f_{43} (Ng et al., 2010). These two ions are important tools to identify the photochemical aging of organic aerosol components in the atmosphere. Organic aerosol can be grouped into oxygenated organic aerosol (OOA) and hydrocarbon-like organic aerosol (HOA), whereby OOA is further classified into low-volatile OOA (LV-OOA) and semi-volatile OOA (SV-OOA) (e.g., Alfarra et al., 2004; Zhang et al., 2005; Canagaratna et al., 2007; Zhang et al., 2007; Jimenez et al., 2009; Ng et al., 2011; Ortega et al., 2016). The difference between these two sub-components is represented by the two ions at m/z 43 and m/z 44. These two ions, or more specifically f_{43} and f_{44} , change during photochemical aging of organic aerosol, leading to higher f_{44} and lower f_{43} values for LV-OOA than SV-OOA. With increasing



photochemical aging, organic aerosol of different origins become more similar in terms of chemistry (Jimenez et al., 2009). We use the ratio $R_{44/43}$ (f_{44} divided by f_{43}) to show the effect of photochemical aging on organic aerosol particles dependent on the altitude.

3.3 Tracer for isoprene epoxydiol-derived secondary organic aerosol (IEPOX-SOA)

- 5 The tracer ion at m/z 82 ($C_5H_6O^+$) is attributed to methylfuran and was identified to be related to IEPOX-SOA (Robinson et al., 2011). Due to thermal vaporisation and subsequent electron impact ionization in the C-ToF-AMS, isoprene photooxidation products in the aerosol are decomposed and lead to an increased signal at m/z 82 (Robinson et al., 2011; Lin et al., 2012). A strong correlation is found with the ion at m/z 53, which corresponds to $C_4H_5^+$ (Robinson et al., 2011; Lin et al., 2012). Calculating the ratio of the signal at m/z 82 to the whole organic signal leads to the fraction f_{82} . This fraction is identified as a tracer
- 10 just for IEPOX-SOA, not for other isoprene-derived SOA from different reaction pathways (e.g., Robinson et al., 2011; Allan et al., 2014; Hu et al., 2015). A study by Hu et al. (2015) found background values of $f_{C_5H_6O^+}$ (ratio of $C_5H_6O^+$ to the total organic signal) for different regions worldwide. For urban and biomass-burning influenced regions an averaged background value of $1.7 \pm 0.1 \%$ is found. Areas strongly impacted by monoterpene emissions show higher averaged background values of $3.1 \pm 0.6 \%$. For ambient organic aerosol particles that are influenced by enhanced isoprene emissions and no extensive NO
- 15 emissions with presence of HO_2 , values of $6.5 \pm 2.2 \%$ have been reported (Hu et al., 2015). This enhanced value indicates a strong IEPOX-SOA influence. All the mentioned values are derived from high resolution ToF-AMS (HR-ToF-AMS) data. A C-ToF-AMS with unit mass resolution might show some interferences from other ions at m/z 82, but nevertheless the tracer f_{82} is also seen as useful for identifying IEPOX-SOA (Hu et al., 2015).

Figure 1 shows the normalized, measured mass spectrum for a period with high f_{82} during flight AC13 (for the flight notation, see Wendisch et al. (2016)) conducted on 19 September 2014 at an altitude of 12.6 km. The averaging time for this spectrum was one minute (15:54-15:55 UTC). The two peaks for determining IEPOX-SOA, m/z 82 and m/z 53, are labelled and clearly visible in the spectrum. For m/z 82 and m/z 53 the same RIE as for organics is applied (Allan et al., 2014). The calculated f_{82} from this spectrum is 27.4 %.

A method for the estimation of the mass concentration of IEPOX-SOA is reported by Hu et al. (2015), see Eq. 1. For this calculation the mass concentration at m/z 82 [$C(m/z$ 82)], total organic mass concentration [$C(Org)$], a reference f_{82} value for IEPOX-SOA ($f_{82}^{IEPOXSOA}$), and a background value (f_{82}^{Bg}) are taken into account. The reference value for $f_{82}^{IEPOXSOA}$ is set to 22 % (Hu et al., 2015). The background value f_{82}^{Bg} can be determined with an empirical equation and depends on the influence of urban, biomass burning or strong monoterpene emissions (Hu et al., 2015). For our data set we assume strong monoterpene emission influence and use Eq. 2 to determine f_{82}^{Bg} according to Hu et al. (2015).

$$30 \quad C(IEPOX-SOA) = \frac{C(m/z \ 82) - C(Org) \cdot f_{82}^{Bg}}{f_{82}^{IEPOXSOA} - f_{82}^{Bg}} \quad (1)$$

$$f_{82}^{Bg} = 0.0077 - 0.019 \cdot f_{44} \quad (2)$$




Figure 1. Normalized mass spectrum of IEPOX-SOA during flight AC13 conducted on 19 September 2014 at an altitude of 12.6 km. Averaging time for this spectrum was one minute (15:54-15:55 UTC). The two distinct IEPOX-SOA tracers m/z 82 and m/z 53 are labelled. The calculated f_{82} from this mass spectrum is 27.4 %.

In Eq. 1, $C(m/z\ 82)$, and $C(\text{Org})$ designate mass concentrations in units of $\mu\text{g m}^{-3}$.

3.4 Particulate organic nitrates

Several studies show that qualitative measurements and quantification of organic nitrates are of major interest (e.g., Fry et al., 2009; Bruns et al., 2010; Farmer et al., 2010; Ayres et al., 2015; Kiendler-Scharr et al., 2016; Liu et al., 2017; Schneider et al., 2017). Organic nitrates are decomposed during the evaporation and/or ionisation processes in the C-ToF-AMS and, therefore, are divided into an organic and a nitrate signal (Farmer et al., 2010). The following use of RONO_2 refers to the nitrate content of organic nitrates, as the organic content cannot be estimated with the described methods. Due to observations during the measurement campaign, the analysis of organic nitrates is described as one part of this study. Four methods are applied to find out whether organic nitrates have been present during the measurements.

- A first estimation of organic nitrates can be derived from the ratio of the nitrate-related ions at m/z 30 (NO^+) and m/z 46 (NO_2^+). The signal at m/z 30 is mostly from NO^+ , but also the organic ion CH_2O^+ can contribute with a small amount (Allan et al., 2014). Such interferences at m/z 30 with CH_2O^+ are corrected in the evaluation software by the fragmentation table (Allan et al., 2004), but it is not possible to distinguish unambiguously between the NO^+ and the CH_2O^+ ions with a C-ToF-AMS. The signal at m/z 46 is usually dominated by NO_2^+ ions (Jimenez et al., 2003; Allan et al., 2004). The ratio of NO^+ to NO_2^+ is different for inorganic ammonium nitrate and organic nitrate. The ratio for inorganic nitrate is known from the ionisation efficiency calibration with pure ammonium nitrate. For our instrument its value lies between 1.49 and 1.56 with a mean and standard deviation value of 1.52 ± 0.03 and is derived from calibration measurements during the campaign. For organic nitrates the literature presents a range of ratios of NO^+ to NO_2^+ that are higher than the ratio for inorganic nitrates and lie between 5 and 12.5 (e.g., Farmer et al., 2010; Fry et al., 2009, 2011; Rollins et al., 2009; Bruns et al., 2010; Boyd et al., 2015).



- Second, a range of possible mass concentrations of particulate organic nitrate (pRONO_2) can be determined by using C-ToF-AMS data. This range is defined with an upper and lower limit. We calculate the amount of inorganic nitrate from neutralization with ammonium and subtract this value from the measured nitrate. For the upper limit we assume full neutralization of sulfate by ammonium and allow only the remaining excess ammonium to be available for neutralization of nitrate (see Eq. 3). Resulting negative values in the first step (neutralisation of sulfate) mean that not enough ammonium for full neutralisation of sulfate was available and were set to zero. In this case, nitrate could exist as organic nitrate such that the upper limit for organic nitrate equals total measured nitrate. For the lower limit we assumed that nitrate was neutralized by the available amount of ammonium (see Eq.4). Resulting negative values were taken as due to statistical variation. The remaining nitrate is the lowest possible amount of organic nitrate, assuming that particulate nitric acid is not present.

$$\text{Upper Limit: } \text{pRONO}_2^{\text{up}} = C_{\text{NO}_3} - \left[\left(C_{\text{NH}_4} - \frac{36}{96} \times C_{\text{SO}_4} \right) \times \frac{62}{18} \right] \quad (3)$$

$$\text{Lower Limit: } \text{pRONO}_2^{\text{low}} = C_{\text{NO}_3} - \left[C_{\text{NH}_4} \times \frac{62}{18} \right] \quad (4)$$

In Eq. 3 and Eq. 4 $\text{pRONO}_2^{\text{up}}$, $\text{pRONO}_2^{\text{low}}$, C_{NO_3} , C_{NH_4} , and C_{SO_4} designate mass concentrations.

- The third possibility to estimate the mass concentration of the nitrate content of organic nitrates comes from recent studies (Farmer et al., 2010; Kiendler-Scharr et al., 2016) (see Eq. 5). Besides the measured ratio of the mass concentrations for NO^+ to NO_2^+ (R_{ambient}), the ratio of NO^+ to NO_2^+ derived from calibrations with ammonium nitrate (R_{cal}), and a fixed value for the ratio from organic nitrates (R_{RONO_2}) are used. In our analyses a value of 10 for R_{RONO_2} was taken as described in Kiendler-Scharr et al. (2016). For a detection limit, Bruns et al. (2010) took $0.1 \mu\text{g m}^{-3}$ for a conservative data evaluation of $\text{pRONO}_2^{\text{Farmer}}$.

$$\text{pRONO}_2^{\text{Farmer}} = \text{NO}_3 \times \left[\frac{(1 + R_{\text{RONO}_2}) \times (R_{\text{ambient}} - R_{\text{cal}})}{(1 + R_{\text{ambient}}) \times (R_{\text{RONO}_2} - R_{\text{cal}})} \right] \quad (5)$$

- The fourth estimation method is described by Fry et al. (2013) and links the ratios determined for inorganic and organic nitrate. The ratios of NO^+ to NO_2^+ calculated for inorganic and organic nitrate are instrument specific, but a proportional co-variation is observed. Thus, a ‘ratio of the ratios’, χ , is proposed and a value for χ of 2.25 ± 0.35 is reported.

$$\chi = \frac{R_{\text{RONO}_2}}{R_{\text{cal}}} \quad (6)$$

With this χ and our calibration value R_{cal} , a value for R_{RONO_2} is calculated using Eq. 6. The next step puts the derived ratio for organic nitrates into Eq. 5 to determine $\text{pRONO}_2^{\text{Fry}}$.

It should be emphasized again, that only the mass concentration of the nitrate content of organic nitrate is determined by all of the described methods.



4 Results and discussion

Aerosol data obtained during 13 flights of the ACRIDICON-CHUVA campaign were evaluated. Data collected during take-off and landing were removed, and cloud passages were not considered. In the following section, vertical profiles of median and interquartile ranges of different parameters are shown. They are calculated for 500 m altitude bins. Data below 100 m were not considered to eliminate the influence of the airport at take-off and landing. Data above 14 km were not used in the binned vertical profiles due to the low amount of available data points.

4.1 Meteorological conditions - boundary layer

The boundary layer (BL) height was determined with the help of the ambient temperature (T_{amb}), the virtual potential temperature (Θ_v), relative humidity with respect to water RH_w , and aerosol number concentration for particle diameters larger than 20 nm ($N_{d>20\text{nm}}$) (Fig. 2). During the whole campaign the vertical profiles of T_{amb} and Θ_v showed almost no deviation, indicating stable conditions for the measurement period. The daily evolution of the BL height was noticeable during the flights, which lasted typically seven hours. The maximum heights of the BL varied between 1.2 km and 2.3 km, depending on the flight time. The highest BL heights were measured later in the afternoon. Over the whole measurement period, a mean height of the BL was found to be at 1.7 ± 0.2 km (Fig. 2, horizontal dashed line; see also Fig. S2). For the binned vertical profile shown in Fig. 2 the BL height is not so obvious due to smoothing effects while averaging over all flights. The observed BL height is consistent with previous studies in the Amazonian dry season (Andreae et al., 2018). Above the BL is the convective cloud layer, which reached altitudes of about 4–5 km during the campaign (Andreae et al., 2018). The thermal tropopause was at an altitude of 16.9 ± 0.6 km (mean and standard deviation) (Andreae et al., 2018). Therefore, all flights of the ACRIDICON-CHUVA campaign were performed in the troposphere.

Figure 2 shows the vertical profiles of median relative humidity with respect to water (RH_w) and median aerosol number concentration for particle diameters larger than 20 nm ($N_{d>20\text{nm}}$). RH_w decreases above the BL showing a minimum with constant median values between 5 and 9 km. At higher altitudes, RH_w increases again with altitude. The median values of aerosol number concentration are constant in the BL, decrease at middle altitudes with a minimum at 4 km and rise strongly at altitudes above 7 km. This increase was interpreted as evidence for new particle formation at altitudes above 7 km by Andreae et al. (2018).

4.2 Aerosol mass concentration

In Fig. 3, the vertical profiles of median mass concentrations of organics, nitrate, sulfate, ammonium, and black carbon given in $\mu\text{g m}^{-3}$ (STP) are shown, with STP calculated for $T = 300$ K and $p = 995$ hPa from calibration measurements at the ground. At all altitudes, the main fraction of the submicron particulate mass consists of organic matter. The highest aerosol mass concentration, in terms of total aerosol mass as well as median values for all species, are observed at lower altitudes between 0.1 and 4.5 km. This includes the BL, see Sect. 4.1. An exception from that is nitrate, showing maximum median values at




Figure 2. Vertical profiles of (a) ambient temperature (T_{amb}), (b) virtual potential temperature (Θ_v), (c) relative humidity with respect to water (RH_w), and (d) aerosol number concentration for particle diameters larger than 20 nm ($N_{d>20\text{nm}}$) for all flights during the ACRIDICON-CHUVA campaign. Here medians (connected dots) with interquartile ranges (shaded area) are shown for each plot. The horizontal dashed line shows the mean height of the top of the boundary layer.

altitudes above 10 km. At middle altitudes (between 5 and 8 km) the mass concentrations of all shown species decrease rapidly. A different behaviour of the species is observed at high altitudes between 8 and 14 km. The mass concentrations increase again with increasing altitude for organics and nitrate. For this altitude range, nitrate shows the highest median values. Although the median values decrease between 13 and 14 km, the 75 percentile range still indicates that high mass concentrations were encountered. In Sect. 4.5 the increase in nitrate mass concentration is discussed with respect to the potential existence of organic nitrates. In contrast to this, the mass concentrations of sulfate and black carbon are highest at lower altitudes, decline above 4.5 km and stay constant at middle and high altitudes. Just between 13 and 14 km, sulfate median values show a slight increase again. At lower altitudes, the mass concentrations of ammonium are highest, above 4.5 km decreasing and staying constantly low for the rest of the altitude range.

- 10 The vertical distribution of aerosol mass concentration allows the classification into three different regions. These are the lower troposphere (LT) ranging from 0.1 to 4.5 km, the middle troposphere (MT) covering altitudes between 4.5 and 8 km and the upper troposphere (UT), which includes altitudes between 8 and 14 km (see Fig. 3).




Figure 3. Vertical profiles of (a) organics (green), (b) nitrate (blue), (c) sulfate (red), (d) ammonium (yellow), (e) black carbon (grey), and (f) total aerosol (black) median mass concentration and interquartile ranges (in 500 m bins) for 13 flights of the ACRIDICON-CHUVA campaign. Horizontal dashed lines indicate the division into lower, middle and upper troposphere.

4.3 Oxidation state of the organic aerosol

As described in Sect. 3.2, we calculated the correlation between f_{43} and f_{44} . Figure 4, Panel (a) presents the data from all flights for the two different altitude regimes LT (0.1 to 4 km) and UT (8 to 14.4 km) averaged over two minutes. The dashed lines represent the region where previous boundary layer aerosol data lie (Ng et al., 2010). The arrow illustrates the direction (upper left corner) in which data points are ‘moving’ when photochemical aging occurs. The two different colours indicate the altitude dependency. The dark green markers represent data sampled in the LT, whereas the light green markers show data from the UT. Although there is an overlapping region of both, there is a difference between LT and UT. Most of the data sampled in the LT are located towards the upper left corner of the triangle, meaning that they are more oxidized. In comparison to this, data from the UT show different properties. Lower f_{44} and at the same time increased f_{43} values show a lower oxidation level of the organic aerosol. Also presented in Fig. 4, Panel (a) are the median values with the interquartile ranges for LT and UT, respectively. The median value for the UT has a higher f_{43} and a lower f_{44} than that for the LT. That means organic aerosol measured in the UT is significantly less photooxidized than in the LT. Thus, the organic aerosol particles in LT and UT are different from each other. Some of the LT organic aerosol may be transported to higher altitudes, but most of the organic aerosol measured in the UT must have a different source that is not located in the BL or in the LT. The possibility that substantial amounts of aerosol are transported from the BL into the UT has been ruled out by the study of Andreae et al. (2018), based on the absence of detectable amounts of black carbon in the UT (also see Fig. 3) and other differences in the properties of aerosol in the LT and UT.




Figure 4. Scatter (triangle) plot of f_{44} against f_{43} for (a) the lower troposphere (dark green, LT) and upper troposphere (light green, UT) for two minutes averaged data and (b) a comparison between different field campaigns performed in the Amazon region. Dashed lines indicate triangular area according to the criteria introduced by Ng et al. (2010). Cubic coloured markers and boxes show median values and interquartile ranges for LT and UT, respectively.

Previous field measurements that have been performed in the Amazon allow a comparison of the presented data set with measurements taken at the ground at two different stations (T3 and T0t). Station T3 is an open field ca. 70 km west of Manaus and has frequent pollution influence from this city, whereas T0t is in a near-pristine rain forest ca. 60 km northwest of Manaus. Thus, they provide anthropogenically influenced or natural measurement conditions, respectively. For further information on the research stations see Martin et al. (2010) and Martin et al. (2016).

The data were collected during the AMAZE-08 and the GoAmazon2014/5 campaigns during the wet season (AMAZE-08: Feb/Mar 2008; GoAmazon2014/5: Feb/Mar 2014) and represent ground measurements (Chen et al., 2009; de Sá et al., 2018). Figure 4, Panel (b) illustrates the median and interquartile ranges for the different data sets. Median values from the GoAmazon2014/5 campaign are similar to the median values derived from the ACRIDICON-CHUVA campaign sampled in the LT, showing that the organic aerosol is oxidized. In comparison, the AMAZE-08 campaign data differ, showing that the organic aerosol during AMAZE-08 was less oxidized. Our data for the UT differ from both these data sets, indicating again a different source for organic aerosol than in the LT.

Figure 5 shows the vertical profiles of median and interquartile ranges for $R_{44/43}$, f_{82} , calculated IEPOX-SOA mass concentration, and relative contribution of IEPOX-SOA to the organic mass concentration, $R_{\text{IEPOX-SOA}/\text{Org}}$, respectively. Panel (a) illustrates the changes of $R_{44/43}$ with altitude. In the BL $R_{44/43}$ is constant, demonstrating that this layer is well mixed. With increasing altitudes this ratio decreases significantly to much lower values than in the LT. The lowest median values are observed in the UT.

This raises the question about the source of the observed organics in the UT. There are three different possibilities. First, horizontal long range transport and a subsequent mixing of these air masses with convectively lofted air could occur. Second,




Figure 5. Vertical profiles with medians and interquartile ranges of (a) $R_{44/43}$, (b) f_{82} , (c) $C(\text{IEPOX-SOA})$, and (d) $R_{\text{IEPOX-SOA/Org}}$. Horizontal dashed lines indicate divisions into LT with BL, MT and UT. Vertical dashed line in Panel (b) presents the derived value for areas with strong IEPOX-SOA influence (see Sect. 3.3) reported by Hu et al. (2015).

particles from the near or distant boundary layer might be transported aloft. A third possibility would be in-situ secondary organic aerosol formation (SOA) in the UT.

A horizontal long range transport of air masses can be excluded due to the less photochemically aged organics in the UT. The organic aerosol particles in the UT show a lower $R_{44/43}$ ratio meaning that they did not experience much photooxidation, as it would be expected from aerosol influenced by long range transport.

The second possibility, representing the fast convective vertical transport of boundary layer particles, can also be ruled out. The aerosol number concentrations differ considerably (see Fig. 2, Panel d) between the LT and UT with strongly increased values in the UT. Furthermore, the sulfate and black carbon aerosol mass concentrations show the highest values in the LT and decrease at higher altitudes (see Fig. 3, Panel c and d). In case of a fast convective vertical transport of boundary layer particles, the sulfate and black carbon aerosol mass concentrations would show similar values in the LT and UT. The decrease at altitudes above 4.5 km indicates that the aerosol particles have been efficiently removed (e.g. scavenging) during vertical transport (Andreae et al., 2018).

This leads to the conclusion that the third possibility, in-situ SOA formation with subsequent growth of the aerosol particles to large enough sizes that they can be detected by the C-ToF-AMS, is the dominant process in the UT.

In the following we present evidence that the formation of IEPOX-SOA in the UT can partly explain this observation.



4.4 Observations of isoprene epoxydiol-derived secondary organic aerosol (IEPOX-SOA)

Figure 5 presents the vertical profile of the IEPOX-SOA tracer f_{82} (Panel (b)). In the LT, f_{82} shows constant median values around 5 %, slightly increasing in the interquartile ranges in the upper part of the LT. These values are similar to values for areas with strong IEPOX-SOA influence (see Sect. 3.3 and vertical dashed line in Fig. 5, Panel (b)). In the UT, the values of f_{82} are again quite constant over the altitude range, but with increased median values around 8 %. The values in the UT are even higher than the ones in the LT, which already show a strong influence of IEPOX-SOA. This indicates that IEPOX-SOA can have an important impact also in the UT.

Although the characteristics of the MT are not the focus here due to the overall low organic mass concentrations, it should be mentioned that the highest median values of f_{82} were observed in the MT between 4.5 and 8 km. The interquartile ranges are extended here, but the median values reach up to 12 %.

Figure 5, Panel (c) depicts the vertical profile of the median IEPOX-SOA mass concentrations, which are calculated using Eq. 1. The lowest values can be found in the LT and MT. However, in the UT, especially at altitudes above 10 km, a strong increase of IEPOX-SOA is observed. Here IEPOX-SOA contributes up to 20 % of the organic mass concentration in the UT (see Fig. 5, Panel (d)). The highest contribution of IEPOX-SOA to the organic aerosol mass is observed in the MT, whereas in the LT up to 10 % can be attributed to IEPOX-SOA. This suggests that in the MT as well as in the UT IEPOX-SOA formation can occur. The correlation of f_{82} with $R_{44/43}$ is presented in Fig. 6 using data averaged over two minutes. Again, the two different green colours refer to the LT (dark green) and the UT (light green). In the LT, high $R_{44/43}$ but low f_{82} values are observed. The f_{82} values are similar to reported ones that describe a strong IEPOX-SOA influence (see Sect. 3.3). Interestingly, the values with low $R_{44/43}$, indicating less photooxidized organic aerosol, are correlated with even higher f_{82} .

This suggests that IEPOX-SOA is contributing to in-situ SOA formation in the UT. In general, SOA formation can occur either through new particle formation (NPF) with subsequent growth or by condensation or reactive uptake on pre-existing particles without NPF. However, at the time that NPF occurs, the aerosol particles would be too small to be measurable with the C-ToF-AMS, implying that growth of these newly formed particles is necessary.

The observed enhanced NO mixing ratios in the UT would likely change the reaction pathway of isoprene to a non-IEPOX route and subsequently no IEPOX-SOA formation would occur. Based on the observed IEPOX-SOA in the UT, the oxidation of isoprene to IEPOX must occur before reaching these high altitudes.

In previous laboratory studies it was found that acidic aerosol is needed for reactive uptake or condensation of gaseous IEPOX onto particles. In the laboratory, the acidic conditions are often realised by using sulfate seed particles (e.g., Surratt et al., 2010; Darer et al., 2011; Lin et al., 2012, 2013). Also in field studies a correlation between IEPOX-SOA and sulfate aerosol was proposed (e.g., Lin et al., 2013; Allan et al., 2014; Hu et al., 2015; Xu et al., 2015; Marais et al., 2016; de Sá et al., 2017).

Figure 7 shows the scatter plot of IEPOX-SOA against nitrate (Panel (a)) and sulfate (Panel (b)) mass concentrations for LT and UT, respectively. The data are averaged over two minutes. In the LT, no correlation between IEPOX-SOA and nitrate is found. However, in the UT a stronger correlation between IEPOX-SOA and nitrate can be seen (see Fig. 7, Panel (a)). For sulfate, the correlation is very low both in the LT and UT (see Fig. 7, Panel (b)). This indicates that sulfate might not be necessary for the




Figure 6. Scatter plot of f_{82} against $R_{44/43}$ for the lower troposphere (dark green, LT) and upper troposphere (light green, UT). Data are averaged over two minutes. Red markers and boxes show median values and interquartile ranges for LT and UT, respectively.

formation of IEPOX-SOA, but nitrate could be an important, possibly sufficient component in the UT.

For our data, taking only the inorganic species (nitrate and sulfate) into account for acidity calculations, the measured ammonium would not be sufficient to neutralize all of the inorganic species in the UT such that the aerosol would then be regarded as acidic aerosol (see Fig. S3). Although the correlation between IEPOX-SOA and nitrate is weak, it could indicate that not only sulfate, but also nitrate could provide the acidic conditions for the partitioning of IEPOX to IEPOX-SOA. The partitioning of IEPOX on nitrate aerosol results in organic nitrates. The formation and the estimated amount of organic nitrates are discussed in Sect. 4.5.

4.5 Particulate organic nitrate

The vertical profiles of NO and NO_y mixing ratios are shown in Fig. 8, Panel (a) and (b). The mixing ratios of both species are highest in the LT and decrease towards the MT before reaching increased values in the UT again. In the LT, increased NO_x mixing ratios arise from anthropogenic emissions coming from Manaus (Kuhn et al., 2010) and other pollution sources. A likely source for NO_x in the UT is the production by lightning (Schumann and Huntrieser, 2007). Due to the relatively low O_3 values in the UT, NO_x exists mainly in form of NO . After conversion to NO_y (mainly HNO_3), NO_x can act here also as a source for (organic) nitrate aerosol and explain the increase in nitrate aerosol mass concentration (see Sect. 4.2).

As already mentioned in Sect. 4.4, the detected ammonium in the presented data set did not suffice to fully neutralize the measured nitrate and sulfate. However, also organics can react with inorganic species forming organic nitrates and sulfates. In the following, the four approaches to estimate the presence of organic nitrates as described in Sect. 3.4 are discussed.

A first estimation of the particulate nitrate content from organic nitrates is the ratio of NO^+ to NO_2^+ . From calibration measure-




Figure 7. Scatter plot of $C(\text{IEPOX-SOA})$ against nitrate (Panel (a)) and sulfate (Panel (b)) mass concentrations for the lower troposphere (dark green, LT) and upper troposphere (light green, UT). Data are averaged over two minutes and presented together with the values for Pearson's r^2 for the correlation between $C(\text{IEPOX-SOA})$ and nitrate (Panel (a)) and sulfate (Panel (b)) mass concentration for LT and UT, respectively. In Panel (a) linear regression for the correlation between $C(\text{IEPOX-SOA})$ and nitrate mass concentration in the UT is presented.




Figure 8. Vertical profiles of (a) NO mixing ratio, (b) reactive nitrogen NO_y mixing ratio, (c) NO^+ (m/z 30, light blue) and NO_2^+ (m/z 46, dark blue) mass signal and (d) ratio of NO^+ (m/z 30) to NO_2^+ (m/z 46). Horizontal dashed lines indicate divisions into LT with BL, MT and UT. The vertical dashed line in Panel (d) presents the ratio of NO^+ to NO_2^+ derived during calibration measurements.




Figure 9. Vertical profiles of (a) calculated lower and upper limits of organic nitrate mass concentration (approach Sect. 3.4, this study), (b) measured nitrate mass concentration and organic nitrate mass concentration calculated according to Farmer et al. (2010), and (c) measured nitrate mass concentration and organic nitrate mass concentration calculated according to Fry et al. (2013). Horizontal dashed lines give divisions into LT with BL, MT and UT. The vertical dashed line in Panel (b) shows the conservative detection limit of $0.1 \mu\text{g m}^{-3}$ after Bruns et al. (2010).

ments with ammonium nitrate during the campaign this ratio is known and was found to be in the range between 1.49 and 1.56 with a mean and standard deviation value of 1.52 ± 0.03 . Vertical profiles of median values of m/z 30 and m/z 46 and of the ratio NO^+ to NO_2^+ are depicted in Fig. 8, Panel (c) and (d), respectively. During the whole vertical profile the two lines behave similarly, except for the altitude range between 2 and 6 km, where the distance between them becomes smaller. Compared to the values derived from calibrations with ammonium nitrate, the measured ratios of NO^+ to NO_2^+ during the flights are much higher and the median values range between 4 and 6. As described in Sect. 3.4, higher NO^+ to NO_2^+ ratios are linked to organic nitrates and the observed ratio profile can be seen as a first evidence for the presence of organic nitrates.

The second estimation provides a range with a lower and upper limit of nitrate mass concentration of organic nitrates according to Eq. 3 and 4 (Sect. 3.4). Figure 9, Panel (a) shows the estimated lower and upper limits as a vertical profile. In the LT the derived values are below zero such that a presence of organic nitrates is unlikely. However, with increasing altitude, the lower and upper limits are also increasing. Especially at altitudes higher than 10 km, both parameters are above zero and the presence of organic nitrates becomes likely. The weakness of this method lies in its dependence on ammonium measurements (see Eq. 3 and 4). Interferences of water in the fragmentation table can lead to a biased estimation of ammonium concentrations (Allan et al., 2004). Biased ammonium concentrations can be one reason for the derived negative values for upper and lower limits of pRONO_2 .

The third approach uses Eq. 5 to estimate the amount of nitrate mass concentration of organic nitrates. As described in Kiendler-



Scharr et al. (2016), a fixed value of 10 for R_{RONO_2} is used here (see Sect. 3.4). In Fig. 9, Panel (b) the calculated particulate nitrate mass concentration of organic nitrates ($\text{pRONO}_2^{\text{Farmer}}$) is presented. The measured nitrate mass concentration is also shown. Values for $\text{pRONO}_2^{\text{Farmer}}$ that are below the detection limit are shown as empty triangles, the ones above in filled triangles. The detection limit is set to $0.1 \mu\text{g m}^{-3}$, which was reported by Bruns et al. (2010) as a conservative approach. Calculated $\text{pRONO}_2^{\text{Farmer}}$ are quite similar to the measured nitrate in the vertical profile, although they are slightly smaller. Interestingly, at altitudes above 10 km the derived values are above the detection limit. This points again to the presence of organic nitrates, especially in the UT at altitudes above 10 km, but the presence of organic nitrates at altitudes below 10 km cannot be excluded. The fourth approach is based on Eq. 6 and addresses an estimation of the ratio of organic nitrates from the ratio for inorganic nitrates determined from the usual calibration measurements (see Sect. 3.4). Using our value of $R_{\text{cal}} = 1.53 \pm 0.03$ results in $R_{\text{RONO}_2} = 3.44 \pm 0.05$ (mean and standard deviation). This value is lower than the ratios reported above from standards and also lower than the ratio that is used for the estimation of Kiendler-Scharr et al. (2016). The vertical profile of $\text{pRONO}_2^{\text{Fry}}$ is shown in Fig. 9, Panel (c), again with the measured nitrate mass concentration in the back. In accord with the lower ratio for organic nitrates, the vertical profile of $\text{pRONO}_2^{\text{Fry}}$ shows slightly higher values than the measured nitrate and also than the estimated $\text{pRONO}_2^{\text{Farmer}}$. At altitudes above 10 km the highest values are reached here also, suggesting the likely presence of organic nitrates. Using the estimation by Fry et al. (2013) yields higher values than measured, which shows an overestimation of $\text{pRONO}_2^{\text{Fry}}$.

In Fig. 10, a comparison between the measured nitrate mass concentration and the calculated pRONO_2 after Farmer et al. (2010) (Panel (a)) and Fry et al. (2013) (Panel (b)) is presented. The data are divided into LT (dark blue) and UT (light blue). The 1:1 line implies, that all nitrate is present as organic nitrate with a relative fraction of 100 %. The slope of the linear regression that lies below the 1:1 line describes the relative content of nitrate that is present as organic nitrate less than 100 %. The linear regression in Fig. 10 (Panel (b)) lies above the 1:1 line, representing an overestimation of the used method. Based on the estimation after Farmer et al. (2010) around 80 % of the measured nitrate data in the UT and 50 % in the LT could be explained with organic nitrates (see Fig. 10, Panel (a)). For the comparison with Fry et al. (2013), an overestimation of the amount of organic nitrates results for the UT, whereas for the LT almost 90 % of the measured nitrate could be explained by organic nitrates (see Fig. 10, Panel (b)). As a qualitative result both comparisons show that in the UT a higher organic nitrate fraction is observed than in the LT.

The quantification of the nitrate content of organic nitrates remains difficult, but a qualitative statement can be made. Methods one, three and four agree with each other that at all altitudes organic nitrates are likely present. In the MT, the calculated values lie below the detection limit introduced by Bruns et al. (2010). However, in the UT above 10 km the presence of organic nitrates is supported by the results of all four methods.

Organic sulfates can also be formed in the absence of sufficient ammonium. Farmer et al. (2010) reported that organic sulfates cannot be quantified using C-ToF-AMS data due to a missing tracer ion. Organic sulfates would lead to similar fragmentation patterns as inorganic sulfates (Farmer et al., 2010). Facing the low mass concentration of sulfate, especially in the UT, the presence of organic nitrates likely plays a larger role than organic sulfates.

The study by Darer et al. (2011) about formation and stability of organic nitrates and sulfates reports that nitrate and sulfates




Figure 10. Scatter plots of measured nitrate mass concentration against organic nitrate mass concentration calculated according to (a) Kiendler-Scharr et al. (2016) and (b) Fry et al. (2013) for the lower troposphere (dark blue, LT) and upper troposphere (light blue, UT), respectively. The horizontal dashed line shows the conservative detection limit of $0.1 \mu\text{g m}^{-3}$ according to Bruns et al. (2010). The 1:1 line indicates that all particulate nitrate is present as organic nitrate. Also shown are the two fit functions (red) for UT and LT in both plots, respectively.

have similar kinetic properties regarding the reaction with tertiary epoxides. According to their study, organic nitrates have shorter lifetimes than organic sulfates and are stable only for short time periods (a few days) before they undergo substitution reactions of nitrate by sulfate or water. Unfortunately, no statement of the temperature dependency of the reactions is given (Darer et al., 2011). The temperatures measured in the UT are low (210-240 K) and could slow down chemical reaction processes and possibly shift the reactions in favour of organic nitrates for a longer time period. In field experiments it was shown that low temperatures shift the chemistry to the formation of organic nitrates (Lee et al., 2014). The condensation of organic nitrates may be important for the growth of newly formed particles in the atmosphere (Berkemeier et al., 2016). Also the phase state of the particles may be important for condensational growth. Over the Amazon basin, SOA particles are predicted to be solid at altitudes above 5 km (Shiraiwa et al., 2017).

5 Summary and Conclusion

We presented results from airborne aerosol measurements with a C-ToF-AMS conducted during the ACRIDICON-CHUVA campaign in September and October 2014 in the tropical lower, middle and upper troposphere over the Amazon region. Vertical profiles of the aerosol mass concentrations for organics, nitrate, sulfate, ammonium and black carbon show an overall decrease of the mass concentrations above the lower troposphere. For organics and nitrate the mass concentrations increase again with increasing altitude in the upper troposphere. The characteristics of the organic aerosol were analysed. The photooxidation state of the organics shows a well-mixed lower troposphere with mainly oxidized organics, whereas in the upper troposphere less



oxidized organics were observed. Fast vertical transport from the boundary layer or horizontal long-range transport can be excluded as an explanation for this feature. Thus, SOA formation in the upper troposphere is proposed as the most likely process explaining the less oxidized organics. Furthermore, IEPOX-SOA was identified at all altitudes indicating a strong influence on the organic aerosol composition. Previous measurements that reported the enhanced IEPOX-SOA influence in the boundary layer were confirmed (e.g., Allan et al., 2014; Hu et al., 2015; de Sá et al., 2017). In the upper troposphere, the IEPOX-SOA mass concentration increases and is associated with less oxidized organics. This suggests that after emission of isoprene by vegetation, oxidation of isoprene by HO_2 proceeds at low altitudes and/or during vertical transport to higher altitudes. The oxidation product, IEPOX, must have been formed before reaching the upper troposphere, where different conditions for the oxidation pathway of isoprene are observed (increased NO and NO_y mixing ratios). In the upper troposphere IEPOX can then partition on pre-existing aerosol particles and thus IEPOX-SOA is formed.

Furthermore, the increase of IEPOX-SOA in the upper troposphere is most likely associated with organic nitrate formation. An increase of nitrate mass concentration was observed in the upper troposphere. Four different methods to estimate the presence and the nitrate mass of organic nitrates were applied and the results support the fact that organic nitrates are present at altitudes above 10 km.

These findings suggest that the formation of IEPOX-SOA and organic nitrates are combined with each other. Two processes could explain the partitioning of IEPOX to the aerosol-phase. The first possibility is that nitrate provides the required acidic conditions and IEPOX is uptaken into the pre-existing aerosol containing organic nitrates. The second possibility is that IEPOX partitions on already neutralized organic nitrates. In this case, the pre-existing aerosol would not need to be acidic.

The formation of IEPOX-SOA in the upper troposphere is an important source for organic aerosol particles at these high altitudes, contributing about 20 % to the total organic aerosol. Further vertical transport of the aerosol particles may lead to entrainment of upper tropospheric aerosol into the tropical tropopause layer (TTL), the "gate to the stratosphere" (Fueglistaler et al., 2009). By this process, organic aerosol particles of tropospheric origin could enter the stratosphere. Also the downward transport of these particles could lead to a regular influence on the aerosol composition at lower altitudes and especially in the boundary layer (Wang et al., 2016; Andreae et al., 2018). This would have effects on cloud condensation nuclei production and thus on cloud properties, and also on the radiative budget.

Emissions from the rainforest influence the tropical atmosphere in several ways and up to high altitudes. The processes described in this study may provide further understanding of the mechanisms that occurred in the pre-industrial tropical atmosphere.

6 Data availability

The measured mass concentration data collected with the C-ToF-AMS are available on the HALO data base (HALO-DB). The link is <https://halo-db.pa.op.dlr.de/>.



Author contributions. MOA, PA, LAM, UP and MW designed the research project. VD and MZ provided BAHAMAS data. BW, DS and DF performed aerosol number concentration measurements. BAH and MP provided black carbon data. HZ provided NO and NO_y data. AC, MK, TJW and CV provided cloud data. CS and JS conducted the C-ToF-AMS measurements during the campaign. CS analysed the data with the help of JS and wrote the manuscript. All co-authors commented on the manuscript.

- 5 *Acknowledgements.* Special thanks to the whole ACRIDICON-CHUVA team for the successful campaign and fruitful data meetings and discussions. Logistics were handled by DLR-FX; a special thanks for the great support and organisation before, during and after the campaign. Also a special thank to the pilots for realisation of the specific flight patterns. This work was supported by BMBF, grant No. 01LG1205E (ROMIC-SPITFIRE) and by DFG (HALO-SPP 1294, SCHN1138/1-2). Significant funding for the instrument integration, certification, and operation was provided by the support of the Max Planck Society for HALO, as well as from internal resources of the Particle Chemistry
- 10 Department at Max Planck Institute for Chemistry. P. Artaxo acknowledges FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo) grants 2013/05014-0 and FAPESP 2017/17047-0. We thank the LBA program runned by INPA (Brazilian National Institute for Amazonian Research). We also thank CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) for the Expedition Licence 00254/2013-9. For technical support in the laboratory thanks to Thomas Böttger and Florian Rubach. Also to Franziska Köllner a special thanks for the helpful discussions.



References

- Alfarra, M. R., Coe, H., Allan, J. D., Bower, K. N., Boudries, H., Canagaratna, M. R., Jimenez, J. L., Jayne, J. T., Garforth, A. A., Li, S. M., and Worsnop, D. R.: Characterization of urban and rural organic particulate in the lower Fraser valley using two Aerodyne Aerosol Mass Spectrometers, *Atmos. Environ.*, 38, 5745–5758, <https://doi.org/10.1016/j.atmosenv.2004.01.054>, 2004.
- 5 Allan, J. D., Delia, A. E., Coe, H., Bower, K. N., Alfarra, M. R., Jimenez, J. L., Middlebrook, A. M., Drewnick, F., Onasch, T. B., Canagaratna, M. R., Jayne, J. T., and Worsnop, D. R.: A generalised method for the extraction of chemically resolved mass spectra from Aerodyne Aerosol Mass Spectrometer data, *Journal of Aerosol Science*, 35, 909–922, <https://doi.org/10.1016/j.jaerosci.2004.02.007>, 2004.
- Allan, J. D., Morgan, W. T., Darbyshire, E., Flynn, M. J., Williams, P. I., Oram, D. E., Artaxo, P., Brito, J., Lee, J. D., and Coe, H.: Airborne observations of IEPOX-derived isoprene SOA in the Amazon during SAMBBA, *Atmos. Chem. Phys.*, 14, 11 393–11 407, <https://doi.org/10.5194/acp-14-11393-2014>, 2014.
- 10 Andreae, M. O., Acevedo, O. C., Araujo, A., Artaxo, P., Barbosa, C. G. G., Barbosa, H. M. J., Brito, J., Carbone, S., Chi, X., Cintra, B. B. L., da Silva, N. F., Dias, N. L., Dias, C. Q., Ditas, F., Ditz, R., Godoi, A. F. L., Godoi, R. H. M., Heimann, M., Hoffmann, T., Kesselmeier, J., Konemann, T., Kruger, M. L., Lavric, J. V., Manzi, A. O., Lopes, A. P., Martins, D. L., Mikhailov, E. F., Moran-Zuloaga, D., Nelson, B. W., Nolscher, A. C., Nogueira, D. S., Piedade, M. T. F., Pohlker, C., Poschl, U., Quesada, C. A., Rizzo, L. V., Ro, C. U., Ruckteschler, N., Sa, L. D. A., Sa, M. D., Sales, C. B., dos Santos, R. M. N., Saturno, J., Schongart, J., Sorgel, M., de Souza, C. M., de Souza, R. A. F., Su, H., Targhetta, N., Tota, J., Trebs, I., Trumbore, S., van Eijck, A., Walter, D., Wang, Z., Weber, B., Williams, J., Winderlich, J., Wittmann, F., Wolff, S., and Yanez-Serrano, A. M.: The Amazon Tall Tower Observatory (ATTO): overview of pilot measurements on ecosystem ecology, meteorology, trace gases, and aerosols, *Atmos. Chem. Phys.*, 15, 10 723–10 776, <https://doi.org/10.5194/acp-15-10723-2015>, 2015.
- 20 Andreae, M. O., Afchine, A., Albrecht, R., Holanda, B. A., Artaxo, P., Barbosa, H. M. J., Borrmann, S., Cecchini, M. A., Costa, A., Dollner, M., Fütterer, D., Järvinen, E., Jurkat, T., Klimach, T., Konemann, T., Knote, C., Krämer, M., Krisna, T., Machado, L. A. T., Mertes, S., Minikin, A., Pöhlker, C., Pöhlker, M. L., Pöschl, U., Rosenfeld, D., Sauer, D., Schlager, H., Schnaiter, M., Schneider, J., Schulz, C., Spanu, A., Sperling, V. B., Voigt, C., Walser, A., Wang, J., Weinzierl, B., Wendisch, M., and Ziereis, H.: Aerosol characteristics and particle production in the upper troposphere over the Amazon Basin, *Atmos. Chem. Phys.*, 18, 921–961, <https://doi.org/10.5194/acp-18-921-2018>, <https://www.atmos-chem-phys.net/18/921/2018/>, 2018.
- 25 Ayres, B. R., Allen, H. M., Draper, D. C., Brown, S. S., Wild, R. J., Jimenez, J. L., Day, D. A., Campuzano-Jost, P., Hu, W., de Gouw, J., Koss, A., Cohen, R. C., Duffey, K. C., Romer, P., Baumann, K., Edgerton, E., Takahama, S., Thornton, J. A., Lee, B. H., Lopez-Hilfiker, F. D., Mohr, C., Wennberg, P. O., Nguyen, T. B., Teng, A., Goldstein, A. H., Olson, K., and Fry, J. L.: Organic nitrate aerosol formation via NO₃ and biogenic volatile organic compounds in the southeastern United States, *Atmos. Chem. Phys.*, 15, 13 377–13 392, <https://doi.org/10.5194/acp-15-13377-2015>, 2015.
- 30 Bahreini, R., Dunlea, E. J., Matthew, B. M., Simons, C., Docherty, K. S., DeCarlo, P. F., Jimenez, J. L., Brock, C. A., and Middlebrook, A. M.: Design and operation of a pressure-controlled inlet for airborne sampling with an aerodynamic aerosol lens, *Aerosol. Sci. Technol.*, 42, 465–471, <https://doi.org/10.1080/02786820802178514>, 2008.
- Berkemeier, T., Ammann, M., Mentel, T. F., Poschl, U., and Shiraiwa, M.: Organic Nitrate Contribution to New Particle Formation and Growth in Secondary Organic Aerosols from alpha-Pinene Ozonolysis, *Environ Sci Technol*, 50, 6334–42, <https://doi.org/10.1021/acs.est.6b00961>, <https://www.ncbi.nlm.nih.gov/pubmed/27219077>, 2016.



- Borrmann, S., Kunkel, D., Weigel, R., Minikin, A., Deshler, T., Wilson, J. C., Curtius, J., Volk, C. M., Homan, C. D., Ulanovsky, A., Ravegnani, F., Viciani, S., Shur, G. N., Belyaev, G. V., Law, K. S., and Cairo, F.: Aerosols in the tropical and subtropical UT/LS: in-situ measurements of submicron particle abundance and volatility, *Atmos. Chem. Phys.*, 10, 5573–5592, <https://doi.org/10.5194/acp-10-5573-2010>, 2010.
- 5 Boyd, C. M., Sanchez, J., Xu, L., Eugene, A. J., Nah, T., Tuet, W. Y., Guzman, M. I., and Ng, N. L.: Secondary organic aerosol formation from the β -pinene+NO₃ system: effect of humidity and peroxy radical fate, *Atmos. Chem. Phys.*, 15, 7497–7522, <https://doi.org/10.5194/acp-15-7497-2015>, <https://www.atmos-chem-phys.net/15/7497/2015/>, 2015.
- Bruns, E. A., Perraud, V., Zelenyuk, A., Ezell, M. J., Johnson, S. N., Yu, Y., Imre, D., Finlayson-Pitts, B. J., and Alexander, M. L.: Comparison of FTIR and particle mass spectrometry for the measurement of particulate organic nitrates, *Environ Sci Technol*, 44, 1056–61, <https://doi.org/10.1021/es9029864>, <https://www.ncbi.nlm.nih.gov/pubmed/20058917>, 2010.
- 10 Budisulistiorini, S. H., Canagaratna, M. R., Croteau, P. L., Marth, W. J., Baumann, K., Edgerton, E. S., Shaw, S. L., Knipping, E. M., Worsnop, D. R., Jayne, J. T., Gold, A., and Surratt, J. D.: Real-time continuous characterization of secondary organic aerosol derived from isoprene epoxydiols in downtown Atlanta, Georgia, using the Aerodyne Aerosol Chemical Speciation Monitor, *Environ Sci Technol*, 47, 5686–94, <https://doi.org/10.1021/es400023n>, <https://www.ncbi.nlm.nih.gov/pubmed/23638946>, 2013.
- 15 Canagaratna, M. R., Jayne, J. T., Jimenez, J. L., Allan, J. D., Alfarra, M. R., Zhang, Q., Onasch, T. B., Drewnick, F., Coe, H., Middlebrook, A., Delia, A., Williams, L. R., Trimborn, A. M., Northway, M. J., DeCarlo, P. F., Kolb, C. E., Davidovits, P., and Worsnop, D. R.: Chemical and microphysical characterization of ambient aerosols with the Aerodyne Aerosol Mass Spectrometer, *Mass Spectrom Rev*, 26, 185–222, <https://doi.org/10.1002/mas.20115>, <https://www.ncbi.nlm.nih.gov/pubmed/17230437>, 2007.
- Carlton, A. G., Wiedinmyer, C., and Kroll, J. H.: A review of Secondary Organic Aerosol (SOA) formation from isoprene, *Atmos. Chem. Phys.*, 9, 4987–5005, 2009.
- 20 Chen, Q., Farmer, D. K., Schneider, J., Zorn, S. R., Heald, C. L., Karl, T. G., Guenther, A., Allan, J. D., Robinson, N., Coe, H., Kimmel, J. R., Pauliquevis, T., Borrmann, S., Poschl, U., Andreae, M. O., Artaxo, P., Jimenez, J. L., and Martin, S. T.: Mass spectral characterization of submicron biogenic organic particles in the Amazon Basin, *Geophys. Res. Lett.*, 36, <https://doi.org/10.1029/2009gl039880>, 2009.
- 25 Chen, Q., Farmer, D. K., Rizzo, L. V., Pauliquevis, T., Kuwata, M., Karl, T. G., Guenther, A., Allan, J. D., Coe, H., Andreae, M. O., Poschl, U., Jimenez, J. L., Artaxo, P., and Martin, S. T.: Submicron particle mass concentrations and sources in the Amazonian wet season (AMAZE-08), *Atmos. Chem. Phys.*, 15, 3687–3701, <https://doi.org/10.5194/acp-15-3687-2015>, 2015.
- Claeys, M., Graham, B., Vas, G., Wang, W., Vermeylen, R., Pashynska, V., Cafmeyer, J., Guyon, P., Andreae, M. O., Artaxo, P., and Maenhaut, W.: Formation of secondary organic aerosols through photooxidation of isoprene, *Science*, 303, 1173–1176, <https://doi.org/10.1126/science.1092805>, 2004.
- 30 Darer, A. I., Cole-Filipiak, N. C., O'Connor, A. E., and Elrod, M. J.: Formation and Stability of Atmospherically Relevant Isoprene-Derived Organosulfates and Organonitrates, *Environ. Sci. Technol.*, 45, 1895–1902, <https://doi.org/10.1021/es103797z>, <http://dx.doi.org/10.1021/es103797z>, 2011.
- de Sá, S. S., Palm, B. B., Campuzano-Jost, P., Day, D. A., Newburn, M. K., Hu, W., Isaacman-VanWertz, G., Yee, L. D., Thalman, R., Brito, J., Carbone, S., Artaxo, P., Goldstein, A. H., Manzi, A. O., Souza, R. A. F., Mei, F., Shilling, J. E., Springston, S. R., Wang, J., Surratt, J. D., Alexander, M. L., Jimenez, J. L., and Martin, S. T.: Influence of urban pollution on the production of organic particulate matter from isoprene epoxydiols in central Amazonia, *Atmos. Chem. Phys.*, 17, 6611–6629, <https://doi.org/10.5194/acp-17-6611-2017>, 2017.



- de Sá, S. S., Palm, B. B., Campuzano-Jost, P., Day, D. A., Hu, W., Isaacman-VanWertz, G., Yee, L. D., Brito, J., Carbone, S., Ribeiro, I. O., Cirino, G. G., Liu, Y. J., Thalman, R., Sedlacek, A., Funk, A., Schumacher, C., Shilling, J. E., Schneider, J., Artaxo, P., Goldstein, A. H., Souza, R. A. F., Wang, J., McKinney, K. A., Barbosa, H., Alexander, M. L., Jimenez, J. L., and Martin, S. T.: Urban influence on the concentration and composition of submicron particulate matter in central Amazonia, *Atmos. Chem. Phys. Discuss.*, 2018, 1–56, <https://doi.org/10.5194/acp-2018-172>, <https://www.atmos-chem-phys-discuss.net/acp-2018-172/>, 2018.
- 5 Drennick, F., Hings, S. S., DeCarlo, P., Jayne, J. T., Gonin, M., Fuhrer, K., Weimer, S., Jimenez, J. L., Demerjian, K. L., Borrmann, S., and Worsnop, D. R.: A new time-of-flight aerosol mass spectrometer (TOF-AMS) - Instrument description and first field deployment, *Aerosol. Sci. Technol.*, 39, 637–658, <https://doi.org/10.1080/02786820500182040>, 2005.
- Farmer, D. K., Matsunaga, A., Docherty, K. S., Surratt, J. D., Seinfeld, J. H., Ziemann, P. J., and Jimenez, J. L.: Response of an aerosol mass spectrometer to organonitrates and organosulfates and implications for atmospheric chemistry, *Proc Natl Acad Sci USA*, 107, 6670–6675, <https://doi.org/10.1073/pnas.0912340107>, <http://www.pnas.org/content/107/15/6670.abstract>, 2010.
- 10 Froyd, K. D., Murphy, S. M., Murphy, D. M., de Gouw, J. A., Eddingsaas, N. C., and Wennberg, P. O.: Contribution of isoprene-derived organosulfates to free tropospheric aerosol mass, *Proc Natl Acad Sci USA*, 107, 21 360–21 365, <https://doi.org/10.1073/pnas.1012561107>, <http://www.pnas.org/content/107/50/21360.abstract>, 2010.
- 15 Fry, J. L., Kiendler-Scharr, A., Rollins, A. W., Wooldridge, P. J., Brown, S. S., Fuchs, H., Dubé, W., Mensah, A., dal Maso, M., Tillmann, R., Dorn, H.-P., Brauers, T., and Cohen, R. C.: Organic nitrate and secondary organic aerosol yield from NO₃ oxidation of β -pinene evaluated using a gas-phase kinetics/aerosol partitioning model, *Atmos. Chem. Phys.*, 9, 1431–1449, <https://doi.org/10.5194/acp-9-1431-2009>, <https://www.atmos-chem-phys.net/9/1431/2009/>, 2009.
- Fry, J. L., Kiendler-Scharr, A., Rollins, A. W., Brauers, T., Brown, S. S., Dorn, H.-P., Dubé, W. P., Fuchs, H., Mensah, A., Rohrer, F., Tillmann, R., Wahner, A., Wooldridge, P. J., and Cohen, R. C.: SOA from limonene: role of NO₃ in its generation and degradation, *Atmos. Chem. Phys.*, 11, 3879–3894, <https://doi.org/10.5194/acp-11-3879-2011>, <https://www.atmos-chem-phys.net/11/3879/2011/>, 2011.
- 20 Fry, J. L., Draper, D. C., Zarzana, K. J., Campuzano-Jost, P., Day, D. A., Jimenez, J. L., Brown, S. S., Cohen, R. C., Kaser, L., Hansel, A., Cappellin, L., Karl, T., Roux, A. H., Turnipseed, A., Cantrell, C., Lefer, B. L., and Grossberg, N.: Observations of gas- and aerosol-phase organic nitrates at BEACHON-RoMBAS 2011, *Atmos. Chem. Phys.*, 13, 8585–8605, <https://doi.org/10.5194/acp-13-8585-2013>, 2013.
- 25 Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical Tropopause Layer, *Reviews of Geophysics*, 47, <https://doi.org/ArtId1004> 10.1029/2008rg000267, 2009.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geoscientific Model Development*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012.
- 30 Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, T. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155–5236, <https://doi.org/10.5194/acp-9-5155-2009>, <https://www.atmos-chem-phys.net/9/5155/2009/>, 2009.
- 35 Hu, W. W., Campuzano-Jost, P., Palm, B. B., Day, D. A., Ortega, A. M., Hayes, P. L., Krechmer, J. E., Chen, Q., Kuwata, M., Liu, Y. J., de Sa, S. S., McKinney, K., Martin, S. T., Hu, M., Budisulistiorini, S. H., Riva, M., Surratt, J. D., St Clair, J. M., Isaacman-Van Wertz, G., Yee, L. D., Goldstein, A. H., Carbone, S., Brito, J., Artaxo, P., de Gouw, J. A., Koss, A., Wisthaler, A., Mikoviny, T., Karl, T., Kaser, L., Jud, W., Hansel, A., Docherty, K. S., Alexander, M. L., Robinson, N. H., Coe, H., Allan, J. D., Canagaratna, M. R., Paulot, F., and Jimenez,



- J. L.: Characterization of a real-time tracer for isoprene epoxydiols-derived secondary organic aerosol (IEPOX-SOA) from aerosol mass spectrometer measurements, *Atmos. Chem. Phys.*, 15, 11 807–11 833, <https://doi.org/10.5194/acp-15-11807-2015>, 2015.
- Jimenez, J. L., Jayne, J. T., Shi, Q., Kolb, C. E., Worsnop, D. R., Yourshaw, I., Seinfeld, J. H., Flagan, R. C., Zhang, X. F., Smith, K. A., Morris, J. W., and Davidovits, P.: Ambient aerosol sampling using the Aerodyne Aerosol Mass Spectrometer, *J. Geophys. Res.-Atmos.*, 5 108, <https://doi.org/10.1029/2001jd001213>, 2003.
- Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, E. J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, 10 J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimo, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525–9, <https://doi.org/10.1126/science.1180353>, <https://www.ncbi.nlm.nih.gov/pubmed/20007897>, 2009.
- Kiendler-Scharr, A., Mensah, A. A., Friese, E., Topping, D., Nemitz, E., Prevot, A. S. H., Aijala, M., Allan, J., Canonaco, F., Canagaratna, 15 M., Carbone, S., Crippa, M., Dall'Osto, M., Day, D. A., De Carlo, P., Di Marco, C. F., Elbern, H., Eriksson, A., Freney, E., Hao, L., Herrmann, H., Hildebrandt, L., Hillamo, R., Jimenez, J. L., Laaksonen, A., McFiggans, G., Mohr, C., O'Dowd, C., Otjes, R., Ovadnevaite, J., Pandis, S. N., Poulain, L., Schlag, P., Sellegri, K., Swietlicki, E., Tiitta, P., Vermeulen, A., Wahner, A., Worsnop, D., and Wu, H. C.: Ubiquity of organic nitrates from nighttime chemistry in the European submicron aerosol, *Geophys. Res. Lett.*, 43, 7735–7744, <https://doi.org/10.1002/2016gl069239>, 2016.
- Krechmer, J. E., Coggon, M. M., Massoli, P., Nguyen, T. B., Crounse, J. D., Hu, W., Day, D. A., Tyndall, G. S., Henze, D. K., Rivera-Rios, J. C., Nowak, J. B., Kimmel, J. R., Mauldin, R. L., Stark, H., Jayne, J. T., Sipilä, M., Junninen, H., Clair, J. M. S., Zhang, X., Feiner, P. A., Zhang, L., Miller, D. O., Brune, W. H., Keutsch, F. N., Wennberg, P. O., Seinfeld, J. H., Worsnop, D. R., Jimenez, J. L., and Canagaratna, M. R.: Formation of Low Volatility Organic Compounds and Secondary Organic Aerosol from Isoprene Hydroxyhydroperoxide Low-NO 20 Oxidation, *Environ. Sci. Technol.*, 49, 10 330–10 339, <https://doi.org/10.1021/acs.est.5b02031>, <http://dx.doi.org/10.1021/acs.est.5b02031>, 25 2015.
- Kroll, J. H. and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and evolution of low-volatility organics in the atmosphere, *Atmos. Environ.*, 42, 3593 – 3624, <https://doi.org/10.1016/j.atmosenv.2008.01.003>, <http://www.sciencedirect.com/science/article/pii/S1352231008000253>, 2008.
- Kuhn, U., Ganzeveld, L., Thielmann, A., Dindorf, T., Schebeske, G., Welling, M., Sciare, J., Roberts, G., Meixner, F. X., Kesselmeier, 30 J., Lelieveld, J., Kolle, O., Ciccioli, P., Lloyd, J., Trentmann, J., Artaxo, P., and Andreae, M. O.: Impact of Manaus City on the Amazon Green Ocean atmosphere: ozone production, precursor sensitivity and aerosol load, *Atmos. Chem. Phys.*, 10, 9251–9282, <https://doi.org/10.5194/acp-10-9251-2010>, 2010.
- Lee, L., Wooldridge, P. J., Gilman, J. B., Warneke, C., de Gouw, J., and Cohen, R. C.: Low temperatures enhance organic nitrate formation: evidence from observations in the 2012 Uintah Basin Winter Ozone Study, *Atmos. Chem. Phys.*, 14, 12 441–12 454, 35 <https://doi.org/10.5194/acp-14-12441-2014>, 2014.
- Lin, Y.-H., Zhang, Z., Docherty, K. S., Zhang, H., Budisulistiorini, S. H., Rubitschun, C. L., Shaw, S. L., Knipping, E. M., Edgerton, E. S., Kleindienst, T. E., Gold, A., and Surratt, J. D.: Isoprene Epoxydiols as Precursors to Secondary Organic Aerosol Formation: Acid-



- Catalyzed Reactive Uptake Studies with Authentic Compounds, *Environ. Sci. Technol.*, 46, 250–258, <https://doi.org/10.1021/es202554c>, <http://dx.doi.org/10.1021/es202554c>, 2012.
- Lin, Y.-H., Knipping, E. M., Edgerton, E. S., Shaw, S. L., and Surratt, J. D.: Investigating the influences of SO₂ and NH₃ levels on isoprene-derived secondary organic aerosol formation using conditional sampling approaches, *Atmos. Chem. Phys.*, 13, 8457–8470, <https://doi.org/10.5194/acp-13-8457-2013>, <https://www.atmos-chem-phys.net/13/8457/2013/>, 2013.
- Liu, J., D'Ambro, E. L., Lee, B. H., Lopez-Hilfiker, F. D., Zaveri, R. A., Rivera-Rios, J. C., Keutsch, F. N., Iyer, S., Kurten, T., Zhang, Z., Gold, A., Surratt, J. D., Shilling, J. E., and Thornton, J. A.: Efficient Isoprene Secondary Organic Aerosol Formation from a Non-IEPOX Pathway, *Environ. Sci. Technol.*, 50, 9872–9880, <https://doi.org/10.1021/acs.est.6b01872>, <http://dx.doi.org/10.1021/acs.est.6b01872>, 2016a.
- Liu, J., Russell, L. M., Lee, A. K. Y., McKinney, K. A., Surratt, J. D., and Ziemann, P. J.: Observational evidence for pollution-influenced selective uptake contributing to biogenic secondary organic aerosols in the southeastern U.S., *Geophys. Res. Lett.*, <https://doi.org/10.1002/2017GL074665>, <http://dx.doi.org/10.1002/2017GL074665>, 2017.
- Liu, Y., Kuwata, M., Strick, B. F., Geiger, F. M., Thomson, R. J., McKinney, K. A., and Martin, S. T.: Uptake of Epoxydiol Isomers Accounts for Half of the Particle-Phase Material Produced from Isoprene Photooxidation via the HO₂ Pathway, *Environ. Sci. Technol.*, 49, 250–258, <https://doi.org/10.1021/es5034298>, <http://dx.doi.org/10.1021/es5034298>, 2015.
- Liu, Y., Brito, J., Dorris, M. R., Rivera-Rios, J. C., Seco, R., Bates, K. H., Artaxo, P., Duvoisin, S., J., Keutsch, F. N., Kim, S., Goldstein, A. H., Guenther, A. B., Manzi, A. O., Souza, R. A., Springston, S. R., Watson, T. B., McKinney, K. A., and Martin, S. T.: Isoprene photochemistry over the Amazon rainforest, *Proc Natl Acad Sci USA*, 113, 6125–30, <https://doi.org/10.1073/pnas.1524136113>, <https://www.ncbi.nlm.nih.gov/pubmed/27185928>, 2016b.
- Machado, L. A. T., Calheiros, A. J. P., Biscaro, T., Giangrande, S., Silva Dias, M. A. F., Cecchini, M. A., Albrecht, R., Andreae, M. O., Araujo, W. F., Artaxo, P., Borrmann, S., Braga, R., Burleyson, C., Eichholz, C. W., Fan, J., Feng, Z., Fisch, G. F., Jensen, M. P., Martin, S. T., Pöschl, U., Pöhlker, C., Pöhlker, M. L., Ribaud, J.-F., Rosenfeld, D., Saraiva, J. M. B., Schumacher, C., Thalman, R., Walter, D., and Wendisch, M.: Overview: Precipitation Characteristics and Sensitivities to the Environmental Conditions during GoAmazon2014/5 and ACRIDICON-CHUVA, *Atmospheric Chemistry and Physics Discussions*, 2017, 1–41, <https://doi.org/10.5194/acp-2017-990>, <https://www.atmos-chem-phys-discuss.net/acp-2017-990/>, 2017.
- Marais, E. A., Jacob, D. J., Jimenez, J. L., Campuzano-Jost, P., Day, D. A., Hu, W., Krechmer, J., Zhu, L., Kim, P. S., Miller, C. C., Fisher, J. A., Travis, K., Yu, K., Hanisco, T. F., Wolfe, G. M., Arkinson, H. L., Pye, H. O. T., Froyd, K. D., Liao, J., and McNeill, V. F.: Aqueous-phase mechanism for secondary organic aerosol formation from isoprene: application to the southeast United States and co-benefit of SO₂ emission controls, *Atmos. Chem. Phys.*, 16, 1603–1618, <https://doi.org/10.5194/acp-16-1603-2016>, 2016.
- Martin, S. T., Andreae, M. O., Althausen, D., Artaxo, P., Baars, H., Borrmann, S., Chen, Q., Farmer, D. K., Guenther, A., Gunthe, S. S., Jimenez, J. L., Karl, T., Longo, K., Manzi, A., Müller, T., Pauliquevis, T., Petters, M. D., Prenni, A. J., Pöschl, U., Rizzo, L. V., Schneider, J., Smith, J. N., Swietlicki, E., Tota, J., Wang, J., Wiedensohler, A., and Zorn, S. R.: An overview of the Amazonian Aerosol Characterization Experiment 2008 (AMAZE-08), *Atmos. Chem. Phys.*, 10, 11 415–11 438, <https://doi.org/10.5194/acp-10-11415-2010>, 2010.
- Martin, S. T., Artaxo, P., Machado, L. A. T., Manzi, A. O., Souza, R. A. F., Schumacher, C., Wang, J., Andreae, M. O., Barbosa, H. M. J., Fan, J., Fisch, G., Goldstein, A. H., Guenther, A., Jimenez, J. L., Pöschl, U., Silva Dias, M. A., Smith, J. N., and Wendisch, M.: Introduction: Observations and Modeling of the Green Ocean Amazon (GoAmazon2014/5), *Atmos. Chem. Phys.*, 16, 4785–4797, <https://doi.org/10.5194/acp-16-4785-2016>, <https://www.atmos-chem-phys.net/16/4785/2016/>, 2016.



- Matthew, B. M., Middlebrook, A. M., and Onasch, T. B.: Collection Efficiencies in an Aerodyne Aerosol Mass Spectrometer as a Function of Particle Phase for Laboratory Generated Aerosols, *Aerosol. Sci. Technol.*, 42, 884–898, <https://doi.org/10.1080/02786820802356797>, <http://dx.doi.org/10.1080/02786820802356797>, 2008.
- Mei, F., Comstock, J. M., Wang, J., Pekour, M., Shilling, J., Long, C., Tomlinson, J., Schmid, B., Hubbe, J., Krisna, T., Giez, A., Wendisch, M., Weinzierl, B., Zoeger, M., Schulz, C., Schlager, H., Schneider, J., de Sa, S. S., Springston, S., Cecchini, M. A., Machado, L. A. T., and Martin, S.: Inter-comparison of aircraft measurements during GoAmazon14/5, in prep. for *Atmos. Meas. Tech.*, 2018.
- Middlebrook, A. M., Bahreini, R., Jimenez, J. L., and Canagaratna, M. R.: Evaluation of Composition-Dependent Collection Efficiencies for the Aerodyne Aerosol Mass Spectrometer using Field Data, *Aerosol. Sci. Technol.*, 46, 258–271, <https://doi.org/10.1080/02786826.2011.620041>, 2012.
- Molleker, S.: in prep., 2018.
- Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H., Docherty, K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J. H., Hildebrandt, L., Donahue, N. M., DeCarlo, P. F., Lanz, V. A., Prevot, A. S. H., Dinar, E., Rudich, Y., and Worsnop, D. R.: Organic aerosol components observed in Northern Hemispheric datasets from Aerosol Mass Spectrometry, *Atmos. Chem. Phys.*, 10, 4625–4641, <https://doi.org/10.5194/acp-10-4625-2010>, 2010.
- Ng, N. L., Canagaratna, M. R., Jimenez, J. L., Zhang, Q., Ulbrich, I. M., and Worsnop, D. R.: Real-time methods for estimating organic component mass concentrations from aerosol mass spectrometer data, *Environ. Sci. Technol.*, 45, 910–6, <https://doi.org/10.1021/es102951k>, <https://www.ncbi.nlm.nih.gov/pubmed/21186814>, 2011.
- Ng, N. L., Brown, S. S., Archibald, A. T., Atlas, E., Cohen, R. C., Crowley, J. N., Day, D. A., Donahue, N. M., Fry, J. L., Fuchs, H., Griffin, R. J., Guzman, M. I., Herrmann, H., Hodzic, A., Iinuma, Y., Jimenez, J. L., Kiendler-Scharr, A., Lee, B. H., Luecken, D. J., Mao, J. Q., McLaren, R., Mutzel, A., Osthoff, H. D., Ouyang, B., Picquet-Varrault, B., Platt, U., Pye, H. O. T., Rudich, Y., Schwantes, R. H., Shiraiwa, M., Stutz, J., Thornton, J. A., Tilgner, A., Williams, B. J., and Zaveri, R. A.: Nitrate radicals and biogenic volatile organic compounds: oxidation, mechanisms, and organic aerosol, *Atmos. Chem. Phys.*, 17, 2103–2162, <https://doi.org/10.5194/acp-17-2103-2017>, 2017.
- Ortega, A. M., Hayes, P. L., Peng, Z., Palm, B. B., Hu, W., Day, D. A., Li, R., Cubison, M. J., Brune, W. H., Graus, M., Warneke, C., Gilman, J. B., Kuster, W. C., de Gouw, J., Gutiérrez-Montes, C., and Jimenez, J. L.: Real-time measurements of secondary organic aerosol formation and aging from ambient air in an oxidation flow reactor in the Los Angeles area, *Atmos. Chem. Phys.*, 16, 7411–7433, <https://doi.org/10.5194/acp-16-7411-2016>, <https://www.atmos-chem-phys.net/16/7411/2016/>, 2016.
- Paulot, F., Crounse, J. D., Kjaergaard, H. G., Kürten, A., St. Clair, J. M., Seinfeld, J. H., and Wennberg, P. O.: Unexpected Epoxide Formation in the Gas-Phase Photooxidation of Isoprene, *Science*, 325, 730–733, <https://doi.org/10.1126/science.1172910>, <http://science.sciencemag.org/content/325/5941/730>, 2009.
- Reitz, P.: Chemical composition measurements of cloud condensation nuclei and ice nuclei by aerosol mass spectrometry, Ph.D. thesis, University Mainz, <http://nbn-resolving.org/urn:nbn:de:hebis:77-28691>, 2011.
- Riva, M., Budisulistiorini, S. H., Chen, Y., Zhang, Z., D'Ambro, E. L., Zhang, X., Gold, A., Turpin, B. J., Thornton, J. A., Canagaratna, M. R., and Surratt, J. D.: Chemical Characterization of Secondary Organic Aerosol from Oxidation of Isoprene Hydroxyhydroperoxides, *Environ. Sci. Technol.*, 50, 9889–9899, <https://doi.org/10.1021/acs.est.6b02511>, <http://dx.doi.org/10.1021/acs.est.6b02511>, 2016.
- Robinson, N. H., Hamilton, J. F., Allan, J. D., Langford, B., Oram, D. E., Chen, Q., Docherty, K., Farmer, D. K., Jimenez, J. L., Ward, M. W., Hewitt, C. N., Barley, M. H., Jenkin, M. E., Rickard, A. R., Martin, S. T., McFiggans, G., and Coe, H.: Evidence for a signif-



- icant proportion of Secondary Organic Aerosol from isoprene above a maritime tropical forest, *Atmos. Chem. Phys.*, 11, 1039–1050, <https://doi.org/10.5194/acp-11-1039-2011>, 2011.
- Rollins, A. W., Kiendler-Scharr, A., Fry, J. L., Brauers, T., Brown, S. S., Dorn, H.-P., Dubé, W. P., Fuchs, H., Mensah, A., Mentel, T. F., Rohrer, F., Tillmann, R., Wegener, R., Wooldridge, P. J., and Cohen, R. C.: Isoprene oxidation by nitrate radical: alkyl nitrate and secondary organic aerosol yields, *Atmos. Chem. Phys.*, 9, 6685–6703, <https://doi.org/10.5194/acp-9-6685-2009>, <https://www.atmos-chem-phys.net/9/6685/2009/>, 2009.
- Rollins, A. W., Fry, J. L., Hunter, J. F., Kroll, J. H., Worsnop, D. R., Singaram, S. W., and Cohen, R. C.: Elemental analysis of aerosol organic nitrates with electron ionization high-resolution mass spectrometry, *Atmos. Meas. Tech.*, 3, 301–310, <https://doi.org/10.5194/amt-3-301-2010>, 2010.
- 10 Schmale, J., Schneider, J., Jurkat, T., Voigt, C., Kalesse, H., Rautenhaus, M., Lichtenstern, M., Schlager, H., Ancellet, G., Arnold, F., Gerding, M., Mattis, I., Wendisch, M., and Borrmann, S.: Aerosol layers from the 2008 eruptions of Mount Okmok and Mount Kasatochi: In situ upper troposphere and lower stratosphere measurements of sulfate and organics over Europe, *J. Geophys. Res.-Atmos.*, 115, <https://doi.org/ArtId0010710.1029/2009jd013628>, 2010.
- Schneider, J., Mertes, S., van Pinxteren, D., Herrmann, H., and Borrmann, S.: Uptake of nitric acid, ammonia, and organics in orographic clouds: mass spectrometric analyses of droplet residual and interstitial aerosol particles, *Atmos. Chem. Phys.*, 17, 1571–1593, <https://doi.org/10.5194/acp-17-1571-2017>, 2017.
- 15 Schumann, U. and Huntrieser, H.: The global lightning-induced nitrogen oxides source, *Atmos. Chem. Phys.*, 7, 3823–3907, <https://doi.org/10.5194/acp-7-3823-2007>, <https://www.atmos-chem-phys.net/7/3823/2007/>, 2007.
- Schwarz, J. P., Gao, R. S., Fahey, D. W., Thomson, D. S., Watts, L. A., Wilson, J. C., Reeves, J. M., Darbeheshti, M., Baumgardner, D. G., Kok, G. L., Chung, S. H., Schulz, M., Hendricks, J., Lauer, A., Kärcher, B., Slowik, J. G., Rosenlof, K. H., Thompson, T. L., Langford, A. O., Loewenstein, M., and Aikin, K. C.: Single-particle measurements of midlatitude black carbon and light-scattering aerosols from the boundary layer to the lower stratosphere, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2006JD007076>, <http://dx.doi.org/10.1029/2006JD007076>, 2006.
- 20 Shilling, J. E., Pekour, M. S., Fortner, E., Artaxo, P., de Sá, S. S., Hubbe, J. M., Longo, K. M., Machado, L. A. T., Martin, S. T., Springston, S. R., Tomlinson, J., and Wang, J.: Particle-phase chemical composition measurements onboard the G-1 research aircraft during the GoAmazon2014/5 campaign, in prep. for *Atmos. Chem. Phys.*, 2018.
- Shiraiwa, M., Li, Y., Tsimpidi, A. P., Karydis, V. A., Berkemeier, T., Pandis, S. N., Lelieveld, J., Koop, T., and Pöschl, U.: Global distribution of particle phase state in atmospheric secondary organic aerosols, *Nature Communications*, 8, 15002, <https://doi.org/10.1038/ncomms15002>, <http://dx.doi.org/10.1038/ncomms15002>, 2017.
- 30 Shrivastava, M., Cappa, C. D., Fan, J., Goldstein, A. H., Guenther, A. B., Jimenez, J. L., Kuang, C., Laskin, A., Martin, S. T., Ng, N. L., Petaja, T., Pierce, J. R., Rasch, P. J., Roldin, P., Seinfeld, J. H., Shilling, J., Smith, J. N., Thornton, J. A., Volkamer, R., Wang, J., Worsnop, D. R., Zaveri, R. A., Zelenyuk, A., and Zhang, Q.: Recent advances in understanding secondary organic aerosol: Implications for global climate forcing, *Reviews of Geophysics*, 55, 509–559, <https://doi.org/10.1002/2016RG000540>, <http://dx.doi.org/10.1002/2016RG000540>, 2017.
- 35 St. Clair, J. M., Rivera-Rios, J. C., Crounse, J. D., Knap, H. C., Bates, K. H., Teng, A. P., Jørgensen, S., Kjaergaard, H. G., Keutsch, F. N., and Wennberg, P. O.: Kinetics and Products of the Reaction of the First-Generation Isoprene Hydroxy Hydroperoxide (ISOPOOH) with OH, *J. Phys. Chem. A*, 120, 1441–1451, <https://doi.org/10.1021/acs.jpca.5b06532>, <http://dx.doi.org/10.1021/acs.jpca.5b06532>, 2016.



- Stephens, M., Turner, N., and Sandberg, J.: Particle identification by laser-induced incandescence in a solid-state laser cavity, *Appl. Opt.*, 42, 3726–3736, <https://doi.org/10.1364/AO.42.003726>, <http://ao.osa.org/abstract.cfm?URI=ao-42-19-3726>, 2003.
- Surratt, J. D., Murphy, S. M., Kroll, J. H., Ng, N. L., Hildebrandt, L., Sorooshian, A., Szmigielski, R., Vermeylen, R., Maenhaut, W., Claeys, M., Flagan, R. C., and Seinfeld, J. H.: Chemical composition of secondary organic aerosol formed from the photooxidation of isoprene, *Journal of Physical Chemistry A*, 110, 9665–90, <https://doi.org/10.1021/jp061734m>, <https://www.ncbi.nlm.nih.gov/pubmed/16884200>, 2006.
- Surratt, J. D., Kroll, J. H., Kleindienst, T. E., Edney, E. O., Claeys, M., Sorooshian, A., Ng, N. L., Offenberg, J. H., Lewandowski, M., Jaoui, M., Flagan, R. C., and Seinfeld, J. H.: Evidence for Organosulfates in Secondary Organic Aerosol, *Environ. Sci. Technol.*, 41, 517–527, <https://doi.org/10.1021/es062081q>, <http://dx.doi.org/10.1021/es062081q>, 2007.
- 10 Surratt, J. D., Chan, A. W., Eddingsaas, N. C., Chan, M., Loza, C. L., Kwan, A. J., Hersey, S. P., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Reactive intermediates revealed in secondary organic aerosol formation from isoprene, *Proc Natl Acad Sci USA*, 107, 6640–5, <https://doi.org/10.1073/pnas.0911114107>, <https://www.ncbi.nlm.nih.gov/pubmed/20080572>, 2010.
- Voigt, C., Schumann, U., Minikin, A., Abdelmonem, A., Afchine, A., Borrmann, S., Boettcher, M., Buchholz, B., Bugliaro, L., Costa, A., Curtius, J., Dollner, M., Dörnbrack, A., Dreiling, V., Ebert, V., Ehrlich, A., Fix, A., Forster, L., Frank, F., Fütterer, D., Giez, A., Graf, K., Groß, J.-U., Groß, S., Heimerl, K., Heinold, B., Hüneke, T., Järvinen, E., Jurkat, T., Kaufmann, S., Kenntner, M., Klingebiel, M., Klimach, T., Kohl, R., Krämer, M., Krisna, T. C., Luebke, A., Mayer, B., Mertes, S., Molleker, S., Petzold, A., Pfeilsticker, K., Port, M., Rapp, M., Reutter, P., Rolf, C., Rose, D., Sauer, D., Schäfler, A., Schlage, R., Schnaiter, M., Schneider, J., Spelten, N., Spichtinger, P., Stock, P., Walser, A., Weigel, R., Weinzierl, B., Wendisch, M., Werner, F., Wernli, H., Wirth, M., Zahn, A., Ziereis, H., and Zöger, M.: ML-CIRRUS: The Airborne Experiment on Natural Cirrus and Contrail Cirrus with the High-Altitude Long-Range Research Aircraft
- 20 HALO, *Bulletin of the American Meteorological Society*, 98, 271–288, <https://doi.org/10.1175/BAMS-D-15-00213.1>, <https://doi.org/10.1175/BAMS-D-15-00213.1>, 2017.
- Wang, J., Krejci, R., Giangrande, S., Kuang, C., Barbosa, H. M. J., Brito, J., Carbone, S., Chi, X., Comstock, J., Ditas, F., Lavric, J., Manninen, H. E., Mei, F., Moran-Zuloaga, D., Pöhlker, C., Pöhlker, M. L., Saturno, J., Schmid, B., Souza, R. A. F., Springston, S. R., Tomlinson, J. M., Toto, T., Walter, D., Wimmer, D., Smith, J. N., Kulmala, M., Machado, L. A. T., Artaxo, P., Andreae, M. O., Petäjä, T., and Martin, S. T.: Amazon boundary layer aerosol concentration sustained by vertical transport during rainfall, *Nature*, 539, 416, <https://doi.org/10.1038/nature19819>, <http://dx.doi.org/10.1038/nature19819>, 2016.
- Weigel, R., Borrmann, S., Kazil, J., Minikin, A., Stohl, A., Wilson, J. C., Reeves, J. M., Kunkel, D., de Reus, M., Frey, W., Lovejoy, E. R., Volk, C. M., Viciani, S., D’Amato, F., Schiller, C., Peter, T., Schlager, H., Cairo, F., Law, K. S., Shur, G. N., Belyaev, G. V., and Curtius, J.: In situ observations of new particle formation in the tropical upper troposphere: the role of clouds and the nucleation mechanism, *Atmos. Chem. Phys.*, 11, 9983–10 010, <https://doi.org/10.5194/acp-11-9983-2011>, 2011.
- 30 Wendisch, M., Poschl, U., Andreae, M. O., Machado, L. A. T., Albrecht, R., Schlager, H., Rosenfeld, D., Martin, S. T., Abdelmonem, A., Afchine, A., Araujo, A. C., Artaxo, P., Aufmhoff, H., Barbosa, H. M. J., Borrmann, S., Braga, R., Buchholz, B., Cecchini, M. A., Costa, A., Curtius, J., Dollner, M., Dorf, M., Dreiling, V., Ebert, V., Ehrlich, A., Ewald, F., Fisch, G., Fix, A., Frank, F., Fütterer, D., Heckl, C., Heidelberg, F., Hüneke, T., Jakel, E., Järvinen, E., Jurkat, T., Kanter, S., Kastner, U., Kenntner, M., Kesselmeier, J., Klimach, T., Knecht, M., Kohl, R., Kolling, T., Kramer, M., Kruger, M., Krisna, T. C., Lavric, J. V., Longo, K., Mahnke, C., Manzi, A. O., Mayer, B., Mertes, S., Minikin, A., Molleker, S., Munch, S., Nillius, B., Pfeilsticker, K., Pöhlker, C., Roiger, A., Rose, D., Rosenowow, D., Sauer, D., Schnaiter, M., Schneider, J., Schulz, C., de Souza, R. A. F., Spanu, A., Stock, P., Vila, D., Voigt, C., Walser, A., Walter, D., Weigel, R., Weinzierl, B., Werner, F., Yamasoe, M. A., Ziereis, H., Zinner, T., and Zöger, M.: ACRIDICON-CHUVA CAMPAIGN Studying Tropical



- Deep Convective Clouds and Precipitation over Amazonia Using the New German Research Aircraft HALO, B. Am. Meteorol. Soc., 97, 1885–1908, <https://doi.org/10.1175/Bams-D-14-00255.1>, 2016.
- Wennberg, P.: Let's abandon the "high NO_x" and "low NO_x" terminology, IGAC News, 50, 3–4, 2013.
- Worton, D. R., Surratt, J. D., Lafranchi, B. W., Chan, A. W., Zhao, Y., Weber, R. J., Park, J. H., Gilman, J. B., de Gouw, J., Park, C., Schade, G., Beaver, M., Clair, J. M., Crounse, J., Wennberg, P., Wolfe, G. M., Harrold, S., Thornton, J. A., Farmer, D. K., Docherty, K. S., Cubison, M. J., Jimenez, J. L., Frossard, A. A., Russell, L. M., Kristensen, K., Glasius, M., Mao, J., Ren, X., Brune, W., Browne, E. C., Pusede, S. E., Cohen, R. C., Seinfeld, J. H., and Goldstein, A. H.: Observational insights into aerosol formation from isoprene, Environ. Sci. Technol., 47, 11 403–13, <https://doi.org/10.1021/es4011064>, <https://www.ncbi.nlm.nih.gov/pubmed/24004194>, 2013.
- 5 Xu, L., Guo, H., Boyd, C. M., Klein, M., Bougiatioti, A., Cerully, K. M., Hite, J. R., Isaacman-VanWertz, G., Kreisberg, N. M., Knote, C., Olson, K., Koss, A., Goldstein, A. H., Hering, S. V., de Gouw, J., Baumann, K., Lee, S. H., Nenes, A., Weber, R. J., and Ng, N. L.: Effects of anthropogenic emissions on aerosol formation from isoprene and monoterpenes in the southeastern United States, Proc Natl Acad Sci USA, 112, 37–42, <https://doi.org/10.1073/pnas.1417609112>, <https://www.ncbi.nlm.nih.gov/pubmed/25535345>, 2015.
- 10 Zhang, Q., Alfarra, M. R., Worsnop, D. R., Allan, J. D., Coe, H., Canagaratna, M. R., and Jimenez, J. L.: Deconvolution and quantification of hydrocarbon-like and oxygenated organic aerosols based on aerosol mass spectrometry, Environ. Sci. Technol., 39, 4938–52, <https://doi.org/10.1021/es048568l>, <https://www.ncbi.nlm.nih.gov/pubmed/16053095>, 2005.
- 15 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimono, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, Geophys. Res. Lett., 34, L13 801, <https://doi.org/10.1029/2007gl029979>, 2007.
- 20 Ziereis, H., Schlager, H., Schulte, P., van Velthoven, P. F. J., and Slemr, F.: Distributions of NO, NO_x, and NO_y in the upper troposphere and lower stratosphere between 28° and 61°N during POLINAT 2, Journal of Geophysical Research, 105, 3653–3664, <https://doi.org/10.1029/1999JD900870>, <http://dx.doi.org/10.1029/1999JD900870>, 2000.