




THERMODIFFUSION AS A PROBE FOR PROTEIN HYDRATION


11.09.2018 | DOREEN NIETHER, MONA SARTER, ANDREAS STADLER AND SIMONE WIEGAND

13th International Meeting on Thermodiffusion


MOTIVATION

Microscale thermophoresis (MST)


- method to determine kinetic constant of binding reactions
- protein's response to thermal gradient changes when ligand binds
- detected through change in fluorescence intensity during titration
- change due to modification of hydration shell



thermodiffusion changes



equilibration constants of biochemical reactions




quantitative information about hydration shell?

titration curve:
 $\Delta I_{\text{fluor}} \propto \Delta S_T$

Hydration has strong influence on thermophoretic response.

- Can this connection be quantified?
- Can it be used to gain information about change in hydration shell upon complex formation?




OPEN QUESTIONS

CONTRIBUTIONS

Aqueous systems

$$S_T \approx S_T^i + S_T^{chem}$$

- aqueous systems
- S_T influenced by hydrogen bonds
 - HB network of water
 - HB between solute and water
- T -dependence of HB $\Leftrightarrow$ thermodiffusion





RESULTS

Correlation with $\log P$


$$\Delta S_T = S_T(50^\circ\text{C}) - S_T(20^\circ\text{C})$$

- ΔS_T is measure for temperature dependence $\rightarrow$ proper chemical contribution
- ΔS_T correlates with $\log P$
- connection between hydration and thermodiffusion

$\rightarrow$ Simone Wiegand
Wednesday, 11:25



$c = 1 - 10 \text{ wt}\%$






saccharides: P. Blanco et al., J. Phys. Chem. B 114, 2807 (2010)
urea: D. Niether et al., PCCP., 20, 1012 (2018).
formamide: D. Niether et al., PNAS, 113, 4272 (2016).

UREA + WATER

NEMD-simulations – microscopic understanding




- non-equilibrium molecular dynamics simulations
- T -dependence of S_T decreases with rising concentration




UREA + WATER

NEMD-simulations – microscopic understanding

- NEMD simulations show that interactions between **urea** and **water** decrease



w.f. = 0.10








w.f. = 0.50

- increasing U-W interactions: slope goes from negative to positive

STREPTAVIDIN + BIOTIN


Protein-ligand system



streptavidin tetramer

PROTEIN LIGAND INTERACTIONS

Influence of hydration layer



ΔH enthalpy: forces in the protein

VdW interactions
H-bonding
screened charges

ΔS entropy: number of accessible states

$$\Delta S = \Delta S_{\text{protein}} + \Delta S_{\text{hydration}}$$

< 0 ?

ligand binding determined by change of free energy


$$\Delta G = G^{\text{bound}} - G^{\text{free}} = \Delta H - T\Delta S$$

QUASI-ELASTIC NEUTRON SCATTERING


Determination of protein dynamics in solution

Andreas Stadler and Mona Sarter,
ICS-1, JCNS and RWTH Aachen

EISF SPHERES




EISF –
elastic
incoherent
structure
factor $\rightarrow$
amplitudes
of motion



$$A_0(q) = \exp(-\langle x^2 \rangle * q^2)$$

measured in D_2O : only dynamic of **protein**




Quasi-elastic scattering:

- small energy exchange between neutron and particle
- processes with distribution of energies (translations, rotations, ...)

CONFORMATIONAL ENTROPY

Reduction of protein conformational entropy due to biotin binding

$$\Delta S_{conf} = 3R \cdot \ln \left[\sqrt{\frac{\langle x^2 \rangle_{bound}}{\langle x^2 \rangle_{free}}} \right]$$





- streptavidin is more mobile than streptavidin+biotin complex
- fast motions attributed to side-chains
- conformational entropy negative on ps to ns time-scale
- entropy change free protein → complex:

$$\Delta S^{QENS} = -2.0 \pm 0.2 \text{ kJ mol}^{-1} \text{ K}^{-1}$$

Time-scale measured	$\Delta S_{conf} \left[\frac{\text{kJ}}{\text{mol K}} \right]$
ps	-1.8 ± 0.1
100 ps	-1.4 ± 0.5
ns	-2.2 ± 0.3


ITC

Isothermal titration calorimetry




measures free energy change upon complex formation (**protein + hydration shell**)

$$\Delta S^{ITC} = -0.1 \pm 0.1 \text{ kJ mol}^{-1} \text{ K}^{-1}$$



RESULTS

Streptavidin + biotin





QENS	ITC	Difference
$\Delta S^{QENS} =$ $-1.8 \pm 0.2 \text{ kJ mol}^{-1} \text{ K}^{-1}$	$\Delta S^{ITC} =$ $-0.1 \pm 0.1 \text{ kJ mol}^{-1} \text{ K}^{-1}$	$\Delta S^{hs} = \Delta S^{ITC} - \Delta S^{QENS} =$ $1.9 \pm 0.3 \text{ kJ mol}^{-1} \text{ K}^{-1}$
entropy change of protein	entropy change of protein + hydration shell	→ entropy of hydration shell increases when biotin binds

RESULTS


Streptavidin + biotin

- reduced ΔS_T for strep + biotin
→ surface of complex less hydrophilic
- indicates breaking of HB
- results from neutron scattering data:
 - protein-complex more rigid
 - entropy increase in hydration shell



SUMMARY

- Influence of hydration on S_T was investigated
- Conclusion on complex formation of streptavidin: disordered hydration layer compensates less flexible protein complex
- Clear change in ΔS_T when biotin binds on streptavidin, connection to entropy change in hydration shell



ACKNOWLEDGEMENT

Simone Wiegand, Jan K.G. Dhont, and the ICS-3

collaborators

Silvia Di Lecce and Fernando Bresme,
Imperial College London, UK – NEMD simulations



Mona Sarter, Andreas Stadler, Bernd König, and Jörg Fitter,
FZJ and RWTH Aachen – streptavidin QENS

Thank you for your attention!

SETUP

Infra-red Thermal Diffusion Forced Rayleigh Scattering (IR-TDFRS)

homogeneous
temperature
and particle
distribution




laser grating




$\Delta T = 20-100 \mu\text{K}$

temperature
grating



↓
refractive index
grating




thermal diffusion

↓
concentration
grating




Measured quantity:
Intensity of the diffracted beam





SETUP

Signal


Measured quantity:
Intensity of the diffracted beam



Molecules/colloides with higher refractive index moves to...
...cold side



...warm side



$$\zeta_{\text{het}} = \left(\begin{matrix} 1 & 4 & 2 & 4 & 3 \\ 1 & 4 & 2 & 4 & 3 \end{matrix} \right) - \frac{(\partial n / \partial c)_{p,T} \begin{matrix} D_T \\ D \end{matrix}}{(\partial n / \partial T)_{p,c} \begin{matrix} D_T \\ D \end{matrix}} c(1-c) \frac{1}{\tau - \tau_{\text{th}}} \left[\tau \left(1 - e^{-t/\tau} \right) - \tau_{\text{th}} \left(1 - e^{-t/\tau_{\text{th}}} \right) \right]$$

temperature

A

S_T