



The composition of the Asian monsoon UTLS as observed by GLORIA during StratoClim

Michael Höpfner (1), Sören Johansson (1), Jörn Ungermann (2), Felix Friedl-Vallon (1), Peter Preusse (2), Johannes Orphal (1), Bernard Legras (3), Roland Ruhnke (1), Fred Stroh (2), Carmen Ullwer (1), Bärbel Vogel (2), Heike Vogel (1), Ingo Wohltmann (4), and the GLORIA-StratoClim-Team

(1) Karlsruhe Institute of Technology, Institute of Meteorology and Climate Research, Karlsruhe, Germany (michael.hoepfner@kit.edu), (2) Institut für Energie und Klimaforschung, Stratosphäre, IEK-7, Forschungszentrum Jülich, Jülich, Germany, (3) Laboratoire de Météorologie Dynamique, UMR8539, IPSL, UPMC/ENS/CNRS/Ecole Polytechnique, Paris, France, (4) Alfred Wegener Institute for Polar and Marine Research, Potsdam, Germany

The GLORIA (Gimballed Limb Observer for Radiance Imaging of the Atmosphere) infrared limb-imaging spectrometer has been operated on board the Geophysica high-altitude aircraft during the StratoClim field campaign from Kathmandu, Nepal in July/August 2017. During several flights over Nepal and India two dimensional cross-sections of various trace gases (e.g. ozone, water vapour, nitric acid, ethane, HCFC-22, ammonia) and aerosol/cloud cover along the flight path have been retrieved from the limb spectra covering an altitude range between about 10 and 20 km with a vertical resolution of 0.5-1 km. In this contribution we will present the detailed observational data together with a discussion supported by trajectory analyses. Further, the GLORIA measurements will be studied by comparison to simulations performed with the atmospheric model systems CLaMS and ICON-ART (ICOsahedral Nonhydrostatic model - Aerosols and Reactive Trace gases).