

Higher order fluctuations and correlations of conserved charges from lattice QCD

Szabolcs Borsanyi,^a Zoltan Fodor,^{a,b,c} Jana N. Guenther^{a,d} Sandor. K. Katz,^d K. K. Szabó,^{a,c} Attila Pasztor,^a Israel Portillo,^e Claudia Ratti^e

^a*University of Wuppertal, Department of Physics, Wuppertal D-42097, Germany*

^b*Eötvös University, Budapest 1117, Hungary*

^c*Jülich Supercomputing Centre, Jülich D-52425, Germany*

^d*University of Regensburg, Department of Physics, Regensburg D-93053, Germany*

^e*Department of Physics, University of Houston, Houston, TX 77204, USA*

E-mail: borsanyi@uni-wuppertal.de, fodor@bodri.elte.hu,
Jana.Guenther@t-online.de, katz@bodri.elte.hu,
szaboka@general.elte.hu, apasztor@bodri.elte.hu,
iportillovazquez@gmail.com, cratti@uh.edu

ABSTRACT: We calculate several diagonal and non-diagonal fluctuations of conserved charges in a system of 2+1+1 quark flavors with physical masses, on a lattice with size $48^3 \times 12$. Higher order fluctuations at $\mu_B = 0$ are obtained as derivatives of the lower order ones, simulated at imaginary chemical potential. From these correlations and fluctuations we construct ratios of net-baryon number cumulants as functions of temperature and chemical potential, which satisfy the experimental conditions of strangeness neutrality and proton/baryon ratio. Our results qualitatively explain the behavior of the measured cumulant ratios by the STAR collaboration.

Contents

1	Introduction	1
2	Fluctuations and imaginary chemical potentials	2
3	Analysis details	4
3.1	Lattice setup	4
3.2	Correlated fit with priors	5
3.3	Cross-correlators	6
3.4	Error Analysis	8
4	Phenomenology at finite chemical potential	9
5	Conclusions and outlook	12
A	Results for the correlators	14
B	Statistics and lattice details	14

1 Introduction

One of the most challenging goals in the study of Quantum Chromodynamics (QCD) is a precise mapping of the phase diagram of strongly interacting matter. First principle, lattice QCD simulations predict that the transition from hadrons to deconfined quarks and gluons is a smooth crossover [1–6], taking place in the temperature range $T \simeq 145 - 165$ MeV. Lattice simulations cannot presently be performed at finite density due to the sign problem, thus leading to the fact that the QCD phase diagram is still vastly unexplored when the asymmetry between matter and antimatter becomes large.

With the advent of the second Beam Energy Scan (BES-II) at the Relativistic Heavy Ion Collider (RHIC), scheduled for 2019-2020, there is a renewed interest in the heavy ion community towards the phases of QCD at moderate-to-large densities. A rich theoretical effort is being developed in support of the experimental program; several observables are being calculated, in order to constrain the existence and location of the QCD critical point and to observe it experimentally.

Fluctuations of conserved charges (electric charge Q , baryon number B and strangeness S) are among the most relevant observables for the finite-density program for several reasons. One possible way to extend lattice results to finite density is to perform Taylor expansions of the thermodynamic observables around chemical potential $\mu_B = 0$ [7–11]. Fluctuations of conserved charges are directly related to the Taylor expansion coefficients of such observables, thus, they are needed to extend first principle approaches to the regions

of the phase diagram relevant to RHIC. An other popular method to extend observables to finite density is the analytical continuation from imaginary chemical potentials [12–16]. The agreement between the analytical continuation and Taylor expansion was shown for the transition temperature by Bonati et al in Ref. [17].

Fluctuations can also be measured directly, and a comparison between theoretical and experimental results allows to extract the chemical freeze-out temperature T_f and chemical potential μ_{Bf} as functions of the collision energy [18–22]. Such fluctuations have been recently calculated and extrapolated using the Taylor method in Ref. [23]. Finally, higher order fluctuations of conserved charges are proportional to powers of the correlation length and are expected to diverge at the critical point, thus providing an important signature for its experimental detection [9, 24, 25].

In this paper, we calculate several diagonal and non-diagonal fluctuations of conserved charges up to sixth-order and give estimates for higher orders, in the temperature range $135 \text{ MeV} \leq T \leq 220 \text{ MeV}$, for a system of 2+1+1 dynamical quarks with physical masses and lattice size $48^3 \times 12$. We simulate the lower-order fluctuations at imaginary chemical potential and extract the higher order fluctuations as derivatives of the lower order ones at $\mu_B = 0$. This method has been successfully used in the past and proved to lead to a more precise determination of the higher order fluctuations, compared to their direct calculation [26, 27]. The direct method (see e.g. [7]) requires the evaluation of several terms and is affected by a signal-to-noise ratio which is decreasing as a power law of the spatial volume V , with an exponent that grows with the order of the susceptibility.

We also construct combinations of these diagonal and non-diagonal fluctuations in order to study the ratio of the cumulants of the net-baryon number distribution as functions of temperature and chemical potential by means of their Taylor expansion in powers of μ_B/T . We discuss their qualitative comparison with the experimental results from the STAR collaboration, as well as the validity of the truncation of the Taylor series.

The paper is organized as follows: we first discuss the use of imaginary chemical potentials in Section 2. Section 3 gives details on the lattice setup, on the fitting procedure, on its generalization for cross-correlators, and finally on the error estimation. The phenomenological results for the ratios of kurtosis, skewness and variance of the baryon number are presented in Section 4. Conclusions and outlook are discussed in Section 5, while in the Appendix we present all diagonal and non-diagonal fluctuations needed to construct the cumulant ratios shown in Section 4, and give additional technical details.

2 Fluctuations and imaginary chemical potentials

The chemical potentials are implemented on a flavor-by-flavor basis, their relation to the phenomenological baryon (B), electric charge (Q) and strangeness (S) chemical potentials

are given by

$$\begin{aligned}\mu_u &= \frac{1}{3}\mu_B + \frac{2}{3}\mu_Q \\ \mu_d &= \frac{1}{3}\mu_B - \frac{1}{3}\mu_Q \\ \mu_s &= \frac{1}{3}\mu_B - \frac{1}{3}\mu_Q - \mu_S.\end{aligned}\tag{2.1}$$

The observables we are looking at are the derivatives of the free energy with respect to the chemical potentials. Since the free energy is proportional to the pressure, we can write:

$$\chi_{i,j,k}^{B,Q,S} = \frac{\partial^{i+j+k}(p/T^4)}{(\partial\hat{\mu}_B)^i(\partial\hat{\mu}_Q)^j(\partial\hat{\mu}_S)^k},\tag{2.2}$$

with

$$\hat{\mu}_i = \frac{\mu_i}{T}.\tag{2.3}$$

These are the generalized fluctuations we calculated around $\mu = 0$ in our previous work [28].

The fermion determinant $\det M(\mu)$ is complex for real chemical potentials, prohibiting the use of traditional simulation algorithms. For imaginary μ , however, the determinant stays real. The chemical potential is introduced through weighted temporal links in the staggered formalism:

$$U_0(\mu) = e^\mu U_0, \quad U_0^\dagger(\mu) = e^{-\mu} U_0^\dagger\tag{2.4}$$

Thus, an imaginary μ translates into a phase factor for the antiperiodic boundary condition in the Dirac operator. Due to the $Z(3)$ symmetry of the gauge sector, there is a non-trivial periodicity in the imaginary quark chemical potential $\mu_q \rightarrow \mu_q + i(2\pi/3)T$, which translates to the baryochemical potential as $\mu_B \rightarrow \mu_B + i2\pi T$, the Roberge-Weiss symmetry. This is independent of the charge conjugation symmetry $\mu_B \leftrightarrow -\mu_B$. As a result, e.g. for the imaginary part of the baryon density:

$$\langle B \rangle|_{\mu_B/T=i\pi-\epsilon} = -\langle B \rangle|_{\mu_B/T=i\pi+\epsilon}\tag{2.5}$$

At $\mu_B = i\pi T$ there is a first order phase transition at all temperatures above the Roberge-Weiss critical end point T_{RW} [29]. When μ_B crosses $i\pi T$ in the imaginary direction, the imaginary baryon density is discontinuous. This behaviour is illustrated in Fig. 1, where the imaginary baryon density as a function of the imaginary chemical potential is shown. At low temperature the Hadron Resonance Gas model predicts $\langle B \rangle \sim \sinh(\mu_B/T)$, thus for imaginary values we expect a sine function below T_c : $\text{Im}\langle B \rangle \sim \sin(\text{Im}\mu_B/T)$. At temperatures slightly above T_c , we observe that further Fourier components appear in addition to $\sin(\text{Im}\mu_B/T)$ with alternating coefficients, these are consistent with a repulsive interaction between baryons [30]. At very high temperatures, on the other hand, $\langle B \rangle$ is a polynomial of μ_B since the diagrams contributing to its $\sim \mu_B^5$ and higher order components are suppressed by asymptotic freedom [31, 32]. The Stefan-Boltzmann limit is non-vanishing




Figure 1. Cartoon for the imaginary baryon number ($\text{Im } \chi_1^B$) as a function of the imaginary chemical potential. T_{RW} is the temperature of the Roberge-Weiss critical point.

only for two Taylor coefficients of $\text{Im } \langle B \rangle$, giving $\text{Im } \langle B \rangle|_{\mu_B/T=i\pi-\epsilon} = 8\pi/27$. At finite temperatures above T_{RW} this expectation value is smaller but positive. By Eq. (2.5), it implies a first order transition at $\mu_B = i\pi T$.

The order of the transition at T_{RW} heavily depends on the quark masses [33, 34]. For physical quark masses one obtains $T_{RW} = 208(5)$ MeV, and the scaling around the end-point is consistent with the Ising exponents [35]. This implies that, for physical parameters, the transition is limited to $\mu_B = i\pi T$ without any other structures between the imaginary interval $[0, i\pi)$ [33].

Thus, we have only the range $\mu/T \in [0, i\pi)$ to explore the μ -dependence of the observables. Recent simulations in this range include the determination of the transition line, where the slope was determined on the negative side of the $T - \mu_B^2$ phase diagram. Using analyticity arguments, this coefficient gives the curvature of the transition line on the real $T - \mu_B$ phase diagram [36–38]. Apart from the transition temperature, we used imaginary chemical potentials also to extrapolate the equation of state to real μ_B [26], which serves as an alternative approach to the Taylor extrapolation [39]. In a recent study D’Elia et al. have used the low order fluctuations at imaginary chemical potentials to calculate generalized quark number susceptibilities [27].

3 Analysis details

3.1 Lattice setup

In this work we calculate high order fluctuations by studying the imaginary chemical potential dependence of various generalized quark number susceptibilities.

We use a tree-level Symanzik improved gauge action, with four times stout smeared ($\rho = 0.125$) staggered fermions. We simulate $2 + 1 + 1$ dynamical quarks, where the light flavors are tuned in a way to reproduce the physical pion and kaon masses and we set $\frac{m_c}{m_s} = 11.85$ [40]. For the zero-temperature runs that we used for the determination of the

bare masses and the coupling, the volumes satisfy $Lm_\pi > 4$. The scale is determined via f_π . More details on the scale setting and lattice setup can be found in [28].

Our lattice ensembles are generated at eighteen temperatures in the temperature range 135...220 MeV. We simulate at eight different values of imaginary μ_B given as: $\mu_B^{(j)} = iT \frac{j\pi}{8}$ for $j \in \{0, 1, 2, 3, 4, 5, 6, 7\}$. In this work the analysis is done purely on a $48^3 \times 12$ lattice, we leave the continuum extrapolation for future work.

In terms of quark chemical potentials we generate ensembles with $\mu_u = \mu_d = \mu_s = \mu_B/3$. In each simulation point we calculate all derivatives in Eq. (2.2) up to fourth order. Thanks to our scan in $\text{Im } \hat{\mu}_B$, we can calculate additional μ_B derivatives. Ref. [27] uses various “trajectories” in the $\mu_B - \mu_Q - \mu_S$ space, allowing the numerical determination of higher e.g. μ_Q and μ_S derivatives. We find relatively good signal for the μ_Q and μ_S derivatives by directly evaluating Eq. (2.2) within one simulation. We recently summarized the details of the direct calculation in Ref. [28].

3.2 Correlated fit with priors

We start with the analysis for $\chi_2^B(T)$, $\chi_4^B(T)$ and $\chi_6^B(T)$. Our goal is to calculate these quantities at zero chemical potential, using the imaginary chemical potential data up to $\chi_B^4(T, \hat{\mu}_B)$. In this work we extract these derivatives at a fixed temperature. Results for different temperatures are obtained completely independently, an interpolation in temperature is not necessary at any point. Thus, the error bars in our results plot will be independent. The errors between the quantities $\chi_2^B(T)$, $\chi_4^B(T)$ and $\chi_6^B(T)$ will be highly correlated, though, since these are extracted through the same set of ensembles at the given temperature. This correlation will be taken into account when combined quantities are calculated, or when an extrapolation to real chemical potential is undertaken.

Thus we consider the ensembles at a fixed temperature T . For each value of imaginary $\mu_B \neq 0$ we determine χ_1^B , χ_2^B , χ_3^B and χ_4^B from simulation, while for $\mu_B = 0$ only χ_2^B and χ_4^B can be used, since χ_1^B and χ_3^B are odd functions of μ_B and therefore equal to zero.

We make the ansatz for the pressure:

$$\chi_0^B(\hat{\mu}_B) = c_0 + c_2 \hat{\mu}_B^2 + c_4 \hat{\mu}_B^4 + c_6 \hat{\mu}_B^6 + c_8 \hat{\mu}_B^8 + c_{10} \hat{\mu}_B^{10}, \quad (3.1)$$

where the Taylor expansion coefficients c_n are related to the baryon number fluctuations χ_n^B by: $n!c_n = \chi_n^B$. Our data do not allow for an independent determination of c_8 and c_{10} . Nevertheless, in order to have some control over these terms we make the assumption

$$|\chi_8^B| \lesssim \chi_4^B \quad (3.2)$$

$$|\chi_{10}^B| \lesssim \chi_4^B \quad (3.3)$$

or in terms of the c_n coefficients

$$\begin{aligned} 8!c_8 &\lesssim 4!c_4 \\ 10!c_{10} &\lesssim 4!c_4. \end{aligned}$$

We can then rewrite our ansatz as

$$\chi_0^B(\hat{\mu}_B) = c_0 + c_2 \hat{\mu}_B^2 + c_4 \hat{\mu}_B^4 + c_6 \hat{\mu}_B^6 + \frac{4!}{8!} c_4 \epsilon_1 \hat{\mu}_B^8 + \frac{4!}{10!} c_4 \epsilon_2 \hat{\mu}_B^{10}. \quad (3.4)$$

where ϵ_1 and ϵ_2 are drawn randomly from a normal distribution with mean -1.25 and variance 2.75. We use the same distribution for all temperatures.

In effect, our c_8 and c_{10} coefficients are stochastic variables. There is sufficient probability to obtain coefficients that slightly break Eq. (3.3), should the data prefer larger than expected fluctuations, e.g. due to a nearby critical end point. The used distribution for $\epsilon_{1,2}$ actually implements a prior for χ_8^B and χ_{10}^B . In the HRG model we know that $\chi_2^B = \chi_4^B = \chi_6^B = \chi_8^B = \chi_{10}^B$, which is well represented by the prior distribution. At high temperatures χ_B^6 and higher coefficients quickly approach zero, as obtained in Hard Thermal Loop results [41]. In the transition regime, the higher moments of the baryon fluctuations are dominated by the fact that the transition line is μ_B -dependent. Starting from $\mu_B = 0$ at a fixed temperature between T_c and T_{RW} , a crossover line is developed as the imaginary chemical potential is introduced. The magnitude of higher order fluctuations in the transition regime can be estimated by a very simple observation. The behaviour of the quark density $\chi_1^B(T, \hat{\mu}_B)$ is reasonably approximated by $\mu_B \chi_2^B(T + T_c \kappa \hat{\mu}_B^2, \hat{\mu}_B = 0)$, where κ is the curvature of the transition line in the $\mu_B - T$ phase diagram. In this approximation, the only source of μ_B -dependence is coming from the curvature of the transition line. Calculating the μ_B derivatives gives a basic estimate for χ_8^B , which we used to tune the prior distribution.

For this ansatz we calculate the following derivatives, which are the actually simulated lattice observables:

$$\chi_1^B(\hat{\mu}_B) = 2c_2\hat{\mu}_B + 4c_4\hat{\mu}_B^3 + 6c_6\hat{\mu}_B^5 + \frac{4!}{7!}c_4\epsilon_1\hat{\mu}_B^7 + \frac{4!}{9!}c_4\epsilon_2\hat{\mu}_B^9 \quad (3.5)$$

$$\chi_2^B(\hat{\mu}_B) = 2c_2 + 12c_4\hat{\mu}_B^2 + 30c_6\hat{\mu}_B^4 + \frac{4!}{6!}c_4\epsilon_1\hat{\mu}_B^6 + \frac{4!}{8!}c_4\epsilon_2\hat{\mu}_B^8 \quad (3.6)$$

$$\chi_3^B(\hat{\mu}_B) = 24c_4\hat{\mu}_B + 120c_6\hat{\mu}_B^3 + \frac{4!}{5!}c_4\epsilon_1\hat{\mu}_B^5 + \frac{4!}{7!}c_4\epsilon_2\hat{\mu}_B^7 \quad (3.7)$$

$$\chi_4^B(\hat{\mu}_B) = 24c_4 + 360c_6\hat{\mu}_B^2 + c_4\epsilon_1\hat{\mu}_B^4 + \frac{4!}{6!}c_4\epsilon_2\hat{\mu}_B^6. \quad (3.8)$$

We perform a correlated fit for the four measured observables, thus obtaining the values of c_2 , c_4 and c_6 for each temperature, and the corresponding χ_2^B , χ_4^B and χ_6^B . We repeat the fit for 1000 random draws for ϵ_1 and ϵ_2 . The result is weighted using the Akaike Information Criterion [42]. Through these weights we get a posterior distribution from the prior distribution. Our final estimate for χ_8^B represents this posterior distribution. We do not show the posterior for χ_{10}^B , which is mostly noise.

These results are shown in Fig. 2, together with an estimate of χ_8^B , related to χ_4^B by Eq. (3.4).

3.3 Cross-correlators

So far we only considered derivatives with respect to the baryonic chemical potential. In our previous, direct analysis in Ref. [28], the μ_B -derivatives had larger errors than μ_Q - or μ_S -derivatives. For μ_Q , the most noisy disconnected contributions come with smaller prefactors, while for μ_S the disconnected contributions are small due to the heavier strange mass. Our approach was designed to improve the μ_B -derivatives only. Therefore, the μ_S




Figure 2. Results for χ_2^B , χ_4^B , χ_6^B and an estimate for χ_8^B as functions of the temperature, obtained from the single-temperature analysis. We plot χ_8^B in green to point out that its determination is guided by a prior, which is linked to the χ_4^B observable by Eq. (3.4). The red curve in each panel corresponds to the Hadron Resonance Gas (HRG) model result.

and μ_Q derivatives have to be simulated directly and without the support from the fit that we used in the μ_B direction. Our result on χ_{jk}^{QS} improved only due to the increase in the statistics since [28].

On the other hand, baryon-strange and baryon-charge mixed derivatives do benefit from the imaginary μ_B data. We simulate various $\chi_{i,j,k}^{B,Q,S}$ with the appropriate values of j and k and all possible values of i so that $i + j + k \leq 4$. For each group of fluctuations with the same j and k we perform a fit analogous to the procedure described in Section 3.2.

Let's take the example of $j = 1, k = 0$. Our ansatz for cross-correlators is analogous to Eqs. (3.5)-(3.8):

$$\chi_{01}^{BS}(\hat{\mu}_B) = \chi_{11}^{BS} \hat{\mu}_B + \frac{1}{3!} \chi_{31}^{BS} \hat{\mu}_B^3 + \frac{1}{5!} \chi_{51}^{BS} \hat{\mu}_B^5 + \frac{1}{7!} \chi_{71}^{BS} \hat{\mu}_B^7 + \frac{1}{9!} \chi_{91}^{BS} \hat{\mu}_B^9 \quad (3.9)$$

We truncated the expression at tenth order. The priors assume $|\chi_{71}^{BS}| \lesssim |\chi_{31}^{BS}|$ and $|\chi_{91}^{BS}| \lesssim |\chi_{31}^{BS}|$, as it is certainly true at high temperature and within the HRG model. The

prior distribution is wider than 1, we used the same mean and variance as in the channel with no μ_S derivative.

When we use Eq. (3.9) we take χ_1^S , χ_{11}^{BS} , χ_{21}^{BS} and χ_{31}^{BS} as correlated quartets for each imaginary chemical potential and determine the three free coefficients of Eq. (3.9). This fitting procedure is repeated 1000 times with random $\chi_{71}^{BS}/\chi_{31}^{BS}$ and $\chi_{91}^{BS}/\chi_{31}^{BS}$ coefficients. Again, using the Akaike weights we constrain the prior distribution. The resulting estimate for χ_{71}^{BS} along with the fit coefficients are shown in Fig. 3. The posterior for χ_{91}^{BS} is not only noisy, but it is probably heavily contaminated by the higher orders that we did not account for.




Figure 3. χ_{11}^{BS} , χ_{31}^{BS} , χ_{51}^{BS} and an estimate for χ_{71}^{BS} as functions of the temperature. The red curves are the HRG model results.

The other channels with higher μ_S or μ_Q derivatives are obtained analogously. These are plotted in Appendix A.

3.4 Error Analysis

For a reliable comparison between experimental measurements and theoretical calculations, the error estimate is an important ingredient. Our statistical error is estimated through the jackknife method. For our systematic error there are several sources. We determine our systematic error by the histogram method described in [43], where each analysis is weighted with the Akaike information criteria. We include the influence of the number of points in the μ_B direction, by either including or ignoring the data from our highest value of μ_B . A very important source for our systematic error is the influence of the higher order

contributions in μ_B . This effect was estimated by adding the higher order terms with pre-factors ϵ_1 and ϵ_2 as described in Section 3.2. We consider 1000 different ϵ pairs and add the different analyses to our histogram. The width of the histogram using Akaike weights corresponding to the fit quality gives the systematic errors for the fit coefficients, and from the same histogram we obtain the posterior distributions for ϵ_1 . The physical quantities that are constrained only by the posterior distribution are plotted with green symbols.

These histograms are built independently for each number (j and k) of μ_S and μ_Q derivatives. When calculating the systematics for the cumulant ratios (Section 4) we need to calculate different combinations of diagonal and non-diagonal fluctuations from the available analyses. Though these fits (corresponding to the same temperature) are carried out separately we keep track of the statistical correlation, by maintaining the jackknife ensembles throughout the analysis. The correct propagation of systematic errors is a more elaborate procedure. When χ_{ijk}^{BSQ} coefficients are combined with different j, k pairs, different histograms have to be combined. If we had only two variables to combine, each of the 2000 first fit variants should be combined with each of the 2000 second fit variants and use the product of the respective probability weights. Instead, we combine the fit results by drawing ‘good’ fits by importance sampling from each histogram independently. In this way, $\mathcal{O}(100)$ random combinations of χ_{ijk}^{BSQ} results already give convergence for each discussed quantity and its error bar. For the results in this paper we used 1000 such random combinations. This procedure assumes that between different j, k pairs the prior distribution is uncorrelated.

4 Phenomenology at finite chemical potential

For a comparison with heavy ion collision experiments, the cumulants of the net-baryon distribution are very useful observables. The first four cumulants are the mean M_B , the variance σ_B^2 , the skewness S_B and the kurtosis κ_B . By forming appropriate ratios, we can cancel out explicit volume factors. However, the measured distributions themselves may still depend on the volume, which one should take into account when comparing to experiments.

Heavy ion collisions involving lead or gold atoms at $\mu_B > 0$ correspond to the following situation

$$\langle n_S \rangle = 0 \quad \langle n_Q \rangle = 0.4 \langle n_B \rangle. \quad (4.1)$$

For each T and μ_B pair, we have to first calculate μ_Q and μ_S that satisfy this condition. The resulting $\mu_Q(\mu_B)$ and $\mu_S(\mu_B)$ functions, too, can be Taylor expanded [19, 20], introducing

$$q_j = \frac{1}{j!} \left. \frac{d^j \hat{\mu}_Q}{(d\hat{\mu}_B)^j} \right|_{\mu_B=0} \quad (4.2)$$

$$s_j = \frac{1}{j!} \left. \frac{d^j \hat{\mu}_S}{(d\hat{\mu}_B)^j} \right|_{\mu_B=0}. \quad (4.3)$$

We investigate three different ratios of cumulants:

$$\frac{M_B}{\sigma_B^2} = \frac{\chi_1^B(T, \hat{\mu}_B)}{\chi_2^B(T, \hat{\mu}_B)} = \hat{\mu}_B r_{12}^{B,1} + \hat{\mu}_B^3 r_{12}^{B,3} + \dots \quad (4.4)$$

$$\frac{S_B \sigma_B^3}{M_B} = \frac{\chi_3^B(T, \hat{\mu}_B)}{\chi_1^B(T, \hat{\mu}_B)} = r_{31}^{B,0} + \hat{\mu}_B^2 r_{31}^{B,2} + \dots \quad (4.5)$$

$$\kappa_B \sigma_B^2 = \frac{\chi_4^B(T, \hat{\mu}_B)}{\chi_2^B(T, \hat{\mu}_B)} = r_{42}^{B,0} + \hat{\mu}_B^2 r_{42}^{B,2} + \hat{\mu}_B^4 r_{42}^{B,4} + \dots \quad (4.6)$$

The μ_B -dependence of the $\chi_i^B(T, \hat{\mu}_B)$ can again be written as a Taylor series:

$$\begin{aligned} \chi_{i,j,k}^{BQS}(\hat{\mu}_B) &= \chi_{i,j,k}^{BQS}(0) + \hat{\mu}_B \left[\chi_{i+1,j,k}^{BQS}(0) + q_1 \chi_{i,j+1,k}^{BQS}(0) + s_1 \chi_{i,j,k+1}^{BQS}(0) \right] \\ &+ \frac{1}{2} \hat{\mu}_B^2 \left[\chi_{i+2,j,k}^{BQS}(0) + q_1^2 \chi_{i,j+2,k}^{BQS}(0) + s_1^2 \chi_{i,j,k+2}^{BQS}(0) \right. \\ &\left. + 2q_1 s_1 \chi_{i,j+1,k+1}^{BQS}(0) + 2q_1 \chi_{i+1,j+1,k}^{BQS}(0) + 2s_1 \chi_{i+1,j,k+1}^{BQS}(0) \right] + \dots \end{aligned} \quad (4.7)$$

The χ coefficients that we determined in Section 3 include derivatives up to sixth order, and we have estimates for the eighth order, too. The fit coefficients corresponding to the tenth order are likely to be contaminated by higher orders, that we did not include into the ansatz. These χ_{ijk}^{BQS} coefficients, however, are given for $j+k \leq 4$, which is the highest order that we used in μ_Q and μ_S .

This list of coefficients allows us to calculate the $r_{ij}^{B,k}$ coefficients from Equations (4.4), (4.5) and (4.6). The results for the $r_{ij}^{B,k}$ coefficients are shown in Figures 4, 5 and 6. We confirm the observation from Ref. [23] that the coefficient $r_{42}^{B,2}$ has a similar temperature dependence as $r_{31}^{B,2}$ but it is ~ 3 times larger in magnitude.

For higher order coefficients, higher order derivatives in μ_S and μ_Q are needed. The direct simulations have a rapidly increasing error with the order of the derivative, and very large statistics would be needed to improve our calculations at this point. Another possibility would be to simulate new ensembles with finite μ_S and μ_Q and do a similar fit as for the μ_B direction. This approach has been used in [27].

After calculating the Taylor coefficients for $S_B \sigma_B^3 / M_B$ and $\kappa_B \sigma_B^2$, we use these results to extrapolate these quantities to finite chemical potential. They are shown in Figure 7. In the left panel, $S_B \sigma_B^3 / M_B$ is shown as a function of the chemical potential for different temperatures. The Taylor expansion for this quantity is truncated at $\mathcal{O}(\hat{\mu}_B^2)$. The black points in the figure are the experimental results from the STAR collaboration from an analysis of cumulant ratios measured at mid-rapidity, $|y| \leq 0.5$, including protons and anti-protons with transverse momenta $0.4 \text{ GeV} \leq p_t \leq 2.0 \text{ GeV}$ [44, 45]. The beam energies were translated to chemical potentials using the fitted formula of Ref. [46]. Even if we do not quantitatively compare the lattice bands to the measurements to extract the freeze-out parameters, as experimental higher order fluctuations might be affected by several effects of non-thermal origin and our lattice results are not continuum extrapolated, we notice that the trend of the data with increasing μ_B can be understood in terms of our Taylor expansion.

In the right panel, we show $\kappa_B \sigma_B^2$ as a function of μ_B / T for different temperatures. The darker bands correspond to the extrapolation up to $\mathcal{O}(\hat{\mu}_B^2)$, whereas the lighter bands also include the $\mathcal{O}(\hat{\mu}_B^4)$ term. Also in this case, the black points are the experimental results from




Figure 4. Taylor expansion coefficients for $\frac{M_B}{\sigma_B} = \frac{\chi_1^B(T, \hat{\mu}_B)}{\chi_2^B(T, \hat{\mu}_B)}$ as functions of the temperature: $r_{12}^{B,1}$ (left panel) and $r_{12}^{B,3}$ (right panel).




Figure 5. Taylor expansion coefficients for $\frac{S_B \sigma_B^3}{M_B} = \frac{\chi_3^B(T, \hat{\mu}_B)}{\chi_1^B(T, \hat{\mu}_B)}$ as functions of the temperature: $r_{31}^{B,0}$ (left panel) and $r_{31}^{B,2}$ (right panel).

the STAR collaboration with transverse momentum cut $0.4 \text{ GeV} \leq p_t \leq 2.0 \text{ GeV}$ [44, 45]. Notice that, due to the fact that the $r_{42}^{B,4}$ is positive in the range $160 \text{ MeV} \leq T \leq 195 \text{ MeV}$, we observe a non-monotonic behavior in $\kappa_B \sigma_B^2$ for $T = 160 \text{ MeV}$ at large chemical potentials. By comparing the two different truncations of the Taylor series we can conclude that, as we increase the temperature, the range of applicability of our Taylor series decreases: while at $T = 150 \text{ MeV}$ the two orders agree in the whole μ_B/T range shown in the figure, at $T = 160 \text{ MeV}$ the central line of the next-to-next-to-leading order bends upwards and is not contained in the next-to-leading order band. To make the NLO prediction precise substantially more computer time would be needed.




Figure 6. Taylor expansion coefficients for $\kappa_B \sigma_B^2 = \frac{\chi_4^B(T, \hat{\mu}_B)}{\chi_2^B(T, \hat{\mu}_B)}$ as functions of the temperature: $r_{42}^{B,0}$ (left panel) $r_{42}^{B,2}$ (middle panel), $r_{42}^{B,4}$ (right panel). The latter is not obtained independently, but by means of the prior ansatz (see text): for this reason, we plot it in green.




Figure 7. $S_B \sigma_B^3 / M_B$ (left panel) and $\kappa_B \sigma_B^2$ (right panel) extrapolated to finite chemical potential. The left panel is extrapolated up to $\mathcal{O}(\hat{\mu}_B^2)$. In the right panel, the darker bands correspond to the extrapolation up to $\mathcal{O}(\hat{\mu}_B^2)$, whereas the lighter bands also include the $\mathcal{O}(\hat{\mu}_B^4)$ term.

5 Conclusions and outlook

In this manuscript, we have calculated several diagonal and non-diagonal fluctuations of electric charge, baryon number and strangeness up to sixth-order, in a system of 2+1+1 quark flavors with physical quark masses, on a lattice with size $48^3 \times 12$. The analysis has been performed simulating the lower order fluctuations at zero and imaginary chemical potential μ_B , and extracting the higher order fluctuations as derivatives of the lower order ones at $\mu_B = 0$. The chemical potentials for electric charge and strangeness have both been set to zero in the simulations. From these fluctuations, we have constructed ratios of baryon number cumulants as functions of T and μ_B , by means of a Taylor series which takes into account the experimental constraints $\langle n_S \rangle = 0$ and $\langle n_Q \rangle = 0.4 \langle n_B \rangle$. These ratios qualitatively explain the behavior observed in the experimental measurements by the STAR collaboration as functions of the collision energy.

We focused on observables (baryon distribution, ratios of cumulants) that are less sensitive to lattice artefacts. An obvious extension of our work will be the use of finer lattices and a continuum extrapolation. The other extension is to use a two- or even three-dimensional mapping of the space of the imaginary chemical potentials using non-vanishing μ_S and μ_Q . That would not only improve the μ_S - and μ_Q -derivatives, but would allow us to study the melting of states with various strangeness and electric charge quantum numbers. Our first study in this direction using strangeness chemical potentials was published in Ref. [47].

Acknowledgements

This project was funded by the DFG grant SFB/TR55. This work was supported by the Hungarian National Research, Development and Innovation Office, NKFIH grants KKP126769 and K113034. An award of computer time was provided by the INCITE program. This research used resources of the Argonne Leadership Computing Facility, which is a DOE Office of Science User Facility supported under Contract DE-AC02-06CH11357. The authors gratefully acknowledge the Gauss Centre for Supercomputing e.V. (www.gauss-centre.eu) for funding this project by providing computing time on the GCS Supercomputer JUQUEEN[48] at Jülich Supercomputing Centre (JSC) as well as on HAZELHEN at HLRS Stuttgart, Germany. This material is based upon work supported by the National Science Foundation under grants no. PHY-1654219 and OAC-1531814 and by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, within the framework of the Beam Energy Scan Theory (BEST) Topical Collaboration. C.R. also acknowledges the support from the Center of Advanced Computing and Data Systems at the University of Houston.

A Results for the correlators

In this Appendix we present the non-diagonal fluctuations of conserved charges needed to construct the cumulant ratios at finite chemical potential μ_B , satisfying the constraints $\langle n_S \rangle = 0$ and $\langle n_Q \rangle = 0.4\langle n_B \rangle$.

Like we did for the diagonal χ_i^B , we simulate lower order fluctuations at finite imaginary chemical potential and extract the higher order fluctuations as derivatives of the lower order ones at $\mu_B = 0$: in particular, we simulate various $\chi_{i,j,k}^{B,Q,S}$ with the appropriate values of j and k and all possible values for i so that

$$i + j + k \leq 4, \quad (\text{A.1})$$

and extract the corresponding $\chi_{i,j,k}^{B,Q,S}$ with $i + j + k \leq 6$ and an estimate for $i + j + k = 8$ and sometimes even $i + j + k = 10$. By estimate (shown in green) we mean the posterior distribution that we get for the two highest orders when using priors, as discussed in the main text. In total we need 15 channels to obtain all the necessary terms.

In the following plots we show these results organized by the number of charge derivatives (j) in Figs. 8-12. It is notoriously difficult to calculate charge correlators using staggered fermions [28]. Correlators that are not protected by a baryon derivative are affected by significant discretization errors. It is understood in the HRG model context that discretization errors mostly affect the contributions from pions and kaons. Staggered lattice effects introduce the highest relative errors for the lightest mesons. Luckily, however, quantities with such discretization effects come with a small pre-factor into the final formulas of Eqs. (4.4)-(4.6). If we had a complete isospin symmetry (factor 0.5 between $\langle n_Q \rangle$ and $\langle n_B \rangle$ in Eq. (4.1)) then electric charge correlators would play no role at all in the extrapolation of baryon fluctuations.

B Statistics and lattice details

In Table 1 we give the number of analyzed configurations per ensemble. The simulation parameters and the details of the analysis are given in Ref. [28].

The determination of the μ derivatives follows the lines of Ref. [7, 28]. We calculate four quantities per configuration and per quark mass

$$A_j = \frac{d}{d\mu_j} \log(\det M_j)^{1/4} = \frac{1}{4} \text{tr} M_j^{-1} M_j', \quad (\text{B.1})$$

$$B_j = \frac{d^2}{(d\mu_j)^2} \log(\det M_j)^{1/4} = \frac{1}{4} \text{tr} \left(M_j'' M_j^{-1} - M_j' M_j^{-1} M_j' M_j^{-1} \right), \quad (\text{B.2})$$

$$C_j = \frac{d^3}{(d\mu_j)^3} \log(\det M_j)^{1/4} = \frac{1}{4} \text{tr} \left(M_j' M_j^{-1} - 3 M_j'' M_j^{-1} M_j' M_j^{-1} + 2 M_j' M_j^{-1} M_j' M_j^{-1} M_j' M_j^{-1} \right), \quad (\text{B.3})$$

$$D_j = \frac{d^4}{(d\mu_j)^4} \log(\det M_j)^{1/4} = \frac{1}{4} \text{tr} \left(M_j'' M_j^{-1} - 4 M_j' M_j^{-1} M_j' M_j^{-1} - 3 M_j'' M_j^{-1} M_j'' M_j^{-1} + 12 M_j'' M_j^{-1} M_j' M_j^{-1} M_j' M_j^{-1} - 6 M_j' M_j^{-1} M_j' M_j^{-1} M_j' M_j^{-1} M_j' M_j^{-1} \right), \quad (\text{B.4})$$




Figure 8. Results containing no electric charge derivative on the various correlators on our $48^3 \times 12$ lattice as functions of the temperature. Green data points denote our estimates for the high orders, these were fitted using a prior distribution. The red curves are the HRG model results.

Here M_j is the fermion matrix corresponding to the j -th quark mass in the system. M' and M'' indicate the first and higher order derivatives with respect to the quark chemical potential. For this simple staggered action higher order derivatives are equal to lower order ones, $M''' = M'$ and $M'''' = M''$ by construction. These traces are calculated using the standard stochastic method, by calculating the effect of the matrices on random sources. At finite (imaginary) chemical potentials we used 4×256 Gaussian random sources for the light quarks and 4×128 sources for the strange quarks. The analysis was accelerated by calculating 256 eigenvectors of the Dirac operator first. These eigenvectors were then fed into an Eig-CG algorithm.

Using the isospin symmetry ($m_u = m_d$), the $ABCD$ traces can be used to calculate the χ^{uds} derivatives with the following formulas:

$$\chi_{200}^{uds} = +\langle B_u \rangle + \langle A_u^2 \rangle - \langle A_u \rangle^2 \quad (\text{B.5})$$

$$\chi_{110}^{uds} = +\langle A_u^2 \rangle - \langle A_u \rangle^2 \quad (\text{B.6})$$

$$\chi_{101}^{uds} = +\langle A_u A_s \rangle - \langle A_s \rangle \langle A_u \rangle \quad (\text{B.7})$$




Figure 9. Results containing one electric charge derivative on the various correlators on our $48^3 \times 12$ lattice as functions of the temperature. Green data points denote our estimates for the high orders, these were fitted using a prior distribution. The red curves are the HRG model results.

$$\chi_{300}^{uds} = +\langle C_u \rangle + 3\langle A_u B_u \rangle + \langle A_u^3 \rangle - 3\langle B_u \rangle \langle A_u \rangle - 3\langle A_u \rangle \langle A_u^2 \rangle + 2\langle A_u \rangle^3 \quad (\text{B.8})$$

$$\chi_{210}^{uds} = +\langle A_u B_u \rangle + \langle A_u^3 \rangle - \langle B_u \rangle \langle A_u \rangle - 3\langle A_u \rangle \langle A_u^3 \rangle + 2\langle A_u \rangle^3 \quad (\text{B.9})$$

$$\chi_{120}^{uds} = +\langle A_u B_u \rangle + \langle A_u^3 \rangle - \langle B_u \rangle \langle A_u \rangle - 3\langle A_u \rangle \langle A_u^2 \rangle + 2\langle A_u \rangle^3 \quad (\text{B.10})$$

$$\chi_{111}^{uds} = +\langle A_u A_u A_s \rangle - \langle A_s \rangle \langle A_u^2 \rangle - 2\langle A_u \rangle \langle A_u A_s \rangle + 2\langle A_s \rangle \langle A_u \rangle^2 \quad (\text{B.11})$$




Figure 10. Results containing two electric charge derivatives on the various correlators on our $48^3 \times 12$ lattice as functions of the temperature. Green data points denote our estimates for the high orders, these were fitted using a prior distribution. The red curves are the HRG model results. Charge correlators without baryon derivative (here χ_{22}^{QS}) are expected to have significant discretization errors.

$$\begin{aligned}
\chi_{400}^{uds} = & \langle D_u \rangle + 3\langle B_u B_u \rangle + 4\langle A_u C_u \rangle + 6\langle A_u^2 B_u \rangle + \langle A_u^4 \rangle \\
& - 4\langle C_u \rangle \langle A_u \rangle - 3\langle B_u \rangle^2 - 6\langle B_u \rangle \langle A_u^2 \rangle - 12\langle A_u \rangle \langle A_u B_u \rangle \\
& - 4\langle A_u \rangle \langle A_u^3 \rangle - 3\langle A_u A_u \rangle \langle A_u^2 \rangle + 12\langle B_u \rangle \langle A_u \rangle^2 \\
& + 12\langle A_u \rangle^2 \langle A_u^2 \rangle - 6\langle A_u \rangle^4
\end{aligned} \tag{B.12}$$

$$\begin{aligned}
\chi_{310}^{uds} = & \langle A_u C_u \rangle + 3\langle A_u^2 B_u \rangle + \langle A_u^4 \rangle - \langle C_u \rangle \langle A_u \rangle - 3\langle B_u \rangle \langle A_u^2 \rangle \\
& - 6\langle A_u \rangle \langle A_u B_u \rangle - 4\langle A_u \rangle \langle A_u^3 \rangle - 3\langle A_u^2 \rangle \langle A_u^2 \rangle \\
& + 6\langle B_u \rangle \langle A_u \rangle^2 + 12\langle A_u \rangle \langle A_u \rangle \langle A_u^2 \rangle - 6\langle A_u \rangle^4
\end{aligned} \tag{B.13}$$

$$\begin{aligned}
\chi_{220}^{uds} = & \langle B_u^2 \rangle + 2\langle A_u^2 B_u \rangle + \langle A_u^4 \rangle - \langle B_u \rangle^2 - 2\langle B_u \rangle \langle A_u^2 \rangle \\
& - 4\langle A_u \rangle \langle A_u B_u \rangle - 4\langle A_u \rangle \langle A_u^3 \rangle - 3\langle A_u^2 \rangle \langle A_u^2 \rangle \\
& + 4\langle B_u \rangle \langle A_u \rangle \langle A_u \rangle + 12\langle A_u \rangle \langle A_u \rangle \langle A_u^2 \rangle - 6\langle A_u \rangle^4
\end{aligned} \tag{B.14}$$

$$\begin{aligned}
\chi_{211}^{uds} = & \langle A_u B_u A_s \rangle + \langle A_u^3 A_s \rangle - \langle A_s \rangle \langle A_u B_u \rangle - \langle A_s \rangle \langle A_u^3 \rangle - \langle B_u \rangle \langle A_u A_s \rangle - \langle B_u A_s \rangle \langle A_u \rangle \\
& - 3\langle A_u \rangle \langle A_u^2 A_s \rangle - 3\langle A_u A_s \rangle \langle A_u^2 \rangle + 2\langle A_s \rangle \langle B_u \rangle \langle A_u \rangle + 6\langle A_s \rangle \langle A_u \rangle \langle A_u^2 \rangle \\
& + 6\langle A_u \rangle^2 \langle A_u A_s \rangle - 6\langle A_s \rangle \langle A_u \rangle^3
\end{aligned} \tag{B.15}$$




Figure 11. Results containing three electric charge derivatives on the various correlators on our $48^3 \times 12$ lattice as functions of the temperature. Green data points denote our estimates for the high orders, these were fitted using a prior distribution. The red curves are the HRG model results. Charge correlators without baryon derivative (here χ_{31}^{QS}) are expected to have significant discretization errors.




Figure 12. χ_4^Q , χ_{24}^{BQ} and estimate for χ_{44}^{BQ} as functions of the temperature. The quantity χ_4^Q has severe cut-off effects on this lattice [28]. The red curves are the HRG model results.

If the listed products of the A, B, C, D traces are calculated as products of the stochastic estimators, a bias could be introduced. Thus, in products different random vectors have to be used in each factor. Alternatively, the expectation value of the bias has to be subtracted. The last step is to express the derivatives in terms of μ_B , μ_Q and μ_S in Eq. (2.2) using Eqs. (2.1), which is a straightforward exercise.

T [MeV]	$\hat{\mu}_B^I = 0$	$\hat{\mu}_B^I = 0.4$	$\hat{\mu}_B^I = 0.8$	$\hat{\mu}_B^I = 1.2$	$\hat{\mu}_B^I = 1.6$	$\hat{\mu}_B^I = 2.0$	$\hat{\mu}_B^I = 2.4$	$\hat{\mu}_B^I = 2.7$
135	17871	1647	2680	4377	2375	3449	2622	2008
140	22624	1625	3583	2975	3499	5321	3129	3211
145	17195	2439	5255	4468	3191	2846	4959	4117
150	18429	2048	3404	10115	6450	5665	3211	3254
155	17494	1624	4735	4938	3911	7813	3670	3485
160	12688	1607	4459	4831	3382	3917	4831	4990
165	18472	1935	4976	8113	8466	4984	5235	4321
170	14417	1987	2704	8820	8053	8023	5916	3273
175	12018	2034	2006	4748	3878	11330	6178	5583
180	12446	2104	2089	5424	4514	6057	5910	4466
185	14184	2151	2138	3112	3086	5934	7733	3767
190	13741	1693	3395	4395	8140	10410	4201	3844
195	15013	1758	3643	5334	8420	5707	3884	4003
200	14974	2300	2262	5999	10709	5033	5496	4203
205	7788	2126	2125	5951	5873	8294	3087	4333
210	4014	1957	1949	12174	6649	3543	2999	3146
215	2506	1783	7056	2268	2244	1711	1674	2090
220	9172	1810	3548	4264	5498	1754	1717	2163

Table 1. Statistics of our simulations on the $48^3 \times 12$ lattice. We list the number of stored and analyzed gauge configurations. These configurations were separated by ten Rational Hybrid Monte Carlo updates.

References

- [1] Y. Aoki, G. Endrodi, Z. Fodor, S. Katz and K. Szabo, *The Order of the quantum chromodynamics transition predicted by the standard model of particle physics*, *Nature* **443** (2006) 675 [[hep-lat/0611014](#)].
- [2] Y. Aoki, Z. Fodor, S. Katz and K. Szabo, *The QCD transition temperature: Results with physical masses in the continuum limit*, *Phys.Lett.* **B643** (2006) 46 [[hep-lat/0609068](#)].
- [3] Y. Aoki, S. Borsanyi, S. Durr, Z. Fodor, S. D. Katz et al., *The QCD transition temperature: results with physical masses in the continuum limit II.*, *JHEP* **0906** (2009) 088 [[0903.4155](#)].
- [4] WUPPERTAL-BUDAPEST COLLABORATION collaboration, S. Borsanyi et al., *Is there still any T_c mystery in lattice QCD? Results with physical masses in the continuum limit III*, *JHEP* **1009** (2010) 073 [[1005.3508](#)].
- [5] T. Bhattacharya, M. I. Buchoff, N. H. Christ, H. T. Ding, R. Gupta et al., *The QCD phase transition with physical-mass, chiral quarks*, [1402.5175](#).
- [6] A. Bazavov, T. Bhattacharya, M. Cheng, C. DeTar, H. Ding et al., *The chiral and deconfinement aspects of the QCD transition*, *Phys.Rev.* **D85** (2012) 054503 [[1111.1710](#)].
- [7] C. Allton, S. Ejiri, S. Hands, O. Kaczmarek, F. Karsch et al., *The QCD thermal phase transition in the presence of a small chemical potential*, *Phys.Rev.* **D66** (2002) 074507 [[hep-lat/0204010](#)].
- [8] C. Allton, M. Doring, S. Ejiri, S. Hands, O. Kaczmarek et al., *Thermodynamics of two flavor QCD to sixth order in quark chemical potential*, *Phys.Rev.* **D71** (2005) 054508 [[hep-lat/0501030](#)].
- [9] R. V. Gavai and S. Gupta, *QCD at finite chemical potential with six time slices*, *Phys. Rev.* **D78** (2008) 114503 [[0806.2233](#)].
- [10] MILC collaboration, S. Basak et al., *QCD equation of state at non-zero chemical potential*, *PoS LATTICE2008* (2008) 171 [[0910.0276](#)].
- [11] O. Kaczmarek, F. Karsch, E. Laermann, C. Miao, S. Mukherjee et al., *Phase boundary for the chiral transition in (2+1)-flavor QCD at small values of the chemical potential*, *Phys.Rev.* **D83** (2011) 014504 [[1011.3130](#)].
- [12] Z. Fodor and S. Katz, *A New method to study lattice QCD at finite temperature and chemical potential*, *Phys.Lett.* **B534** (2002) 87 [[hep-lat/0104001](#)].
- [13] P. de Forcrand and O. Philipsen, *The QCD phase diagram for small densities from imaginary chemical potential*, *Nucl. Phys.* **B642** (2002) 290 [[hep-lat/0205016](#)].
- [14] M. D’Elia and M.-P. Lombardo, *Finite density QCD via imaginary chemical potential*, *Phys. Rev.* **D67** (2003) 014505 [[hep-lat/0209146](#)].
- [15] Z. Fodor and S. Katz, *Lattice determination of the critical point of QCD at finite T and μ* , *JHEP* **0203** (2002) 014 [[hep-lat/0106002](#)].
- [16] Z. Fodor and S. Katz, *Critical point of QCD at finite T and μ , lattice results for physical quark masses*, *JHEP* **0404** (2004) 050 [[hep-lat/0402006](#)].
- [17] C. Bonati, M. D’Elia, F. Negro, F. Sanfilippo and K. Zambello, *Curvature of the pseudocritical line in QCD: Taylor expansion matches analytic continuation*, [1805.02960](#).

- [18] F. Karsch, *Determination of Freeze-out Conditions from Lattice QCD Calculations*, *Central Eur.J.Phys.* **10** (2012) 1234 [[1202.4173](#)].
- [19] A. Bazavov, H. Ding, P. Hegde, O. Kaczmarek, F. Karsch et al., *Freeze-out Conditions in Heavy Ion Collisions from QCD Thermodynamics*, *Phys.Rev.Lett.* **109** (2012) 192302 [[1208.1220](#)].
- [20] S. Borsanyi, Z. Fodor, S. Katz, S. Krieg, C. Ratti et al., *Freeze-out parameters: lattice meets experiment*, *Phys.Rev.Lett.* **111** (2013) 062005 [[1305.5161](#)].
- [21] S. Borsanyi, Z. Fodor, S. Katz, S. Krieg, C. Ratti et al., *Freeze-out parameters from electric charge and baryon number fluctuations: is there consistency?*, *Phys.Rev.Lett.* **113** (2014) 052301 [[1403.4576](#)].
- [22] C. Ratti, *Lattice QCD and heavy ion collisions: a review of recent progress*, [1804.07810](#).
- [23] HOTQCD collaboration, A. Bazavov et al., *Skewness and kurtosis of net baryon-number distributions at small values of the baryon chemical potential*, [1708.04897](#).
- [24] M. A. Stephanov, K. Rajagopal and E. V. Shuryak, *Event-by-event fluctuations in heavy ion collisions and the QCD critical point*, *Phys.Rev.* **D60** (1999) 114028 [[hep-ph/9903292](#)].
- [25] M. Cheng, N. Christ, S. Datta, J. van der Heide, C. Jung et al., *The QCD equation of state with almost physical quark masses*, *Phys.Rev.* **D77** (2008) 014511 [[0710.0354](#)].
- [26] J. Gunther, R. Bellwied, S. Borsanyi, Z. Fodor, S. D. Katz, A. Pasztor et al., *The QCD equation of state at finite density from analytical continuation*, *EPJ Web Conf.* **137** (2017) 07008 [[1607.02493](#)].
- [27] M. D’Elia, G. Gagliardi and F. Sanfilippo, *Higher order quark number fluctuations via imaginary chemical potentials in $N_f = 2 + 1$ QCD*, *Phys. Rev.* **D95** (2017) 094503 [[1611.08285](#)].
- [28] R. Bellwied, S. Borsanyi, Z. Fodor, S. D. Katz, A. Pasztor, C. Ratti et al., *Fluctuations and correlations in high temperature QCD*, *Phys. Rev.* **D92** (2015) 114505 [[1507.04627](#)].
- [29] A. Roberge and N. Weiss, *Gauge Theories With Imaginary Chemical Potential and the Phases of QCD*, *Nucl.Phys.* **B275** (1986) 734.
- [30] V. Vovchenko, A. Pasztor, Z. Fodor, S. D. Katz and H. Stoecker, *Repulsive baryonic interactions and lattice QCD observables at imaginary chemical potential*, *Phys. Lett.* **B775** (2017) 71 [[1708.02852](#)].
- [31] J. I. Kapusta and C. Gale, *Finite-Temperature Field Theory*. Cambridge University Press, second ed., 2006.
- [32] A. Vuorinen, *Quark number susceptibilities of hot QCD up to $g^{*6} \ln g$* , *Phys.Rev.* **D67** (2003) 074032 [[hep-ph/0212283](#)].
- [33] O. Philipsen and C. Pinke, *The nature of the Roberge-Weiss transition in $N_f = 2$ QCD with Wilson fermions*, *Phys. Rev.* **D89** (2014) 094504 [[1402.0838](#)].
- [34] F. Cuteri, C. Pinke, A. Sciarra, C. Czaban and O. Philipsen, *Roberge-Weiss transition in $N_f = 2$ QCD with Wilson fermions and $N_\tau = 6$* , *Phys. Rev. D* (2015) 054507 [[1512.07180](#)].
- [35] C. Bonati, M. D’Elia, M. Mariti, M. Mesiti, F. Negro and F. Sanfilippo, *Roberge-Weiss endpoint at the physical point of $N_f = 2 + 1$ QCD*, *Phys. Rev.* **D93** (2016) 074504 [[1602.01426](#)].

- [36] C. Bonati, M. D'Elia, M. Mariti, M. Mesiti, F. Negro and F. Sanfilippo, *Curvature of the chiral pseudocritical line in QCD: Continuum extrapolated results*, *Phys. Rev.* **D92** (2015) 054503 [[1507.03571](#)].
- [37] R. Bellwied, S. Borsanyi, Z. Fodor, J. Günther, S. D. Katz, C. Ratti et al., *The QCD phase diagram from analytic continuation*, *Phys. Lett.* **B751** (2015) 559 [[1507.07510](#)].
- [38] P. Cea, L. Cosmai and A. Papa, *Critical line of 2+1 flavor QCD: Toward the continuum limit*, *Phys. Rev.* **D93** (2016) 014507 [[1508.07599](#)].
- [39] A. Bazavov et al., *The QCD Equation of State to $\mathcal{O}(\mu_B^6)$ from Lattice QCD*, [1701.04325](#).
- [40] C. McNeile, C. Davies, E. Follana, K. Hornbostel and G. Lepage, *High-Precision c and b Masses, and QCD Coupling from Current-Current Correlators in Lattice and Continuum QCD*, *Phys.Rev.* **D82** (2010) 034512 [[1004.4285](#)].
- [41] N. Haque, J. O. Andersen, M. G. Mustafa, M. Strickland and N. Su, *Three-loop HTLpt Pressure and Susceptibilities at Finite Temperature and Density*, *Phys.Rev.* **D89** (2014) 061701 [[1309.3968](#)].
- [42] H. Akaike *IEEE Transactions on Automatic Control* **19** (1974) 716.
- [43] S. Durr, Z. Fodor, J. Frison, C. Hoelbling, R. Hoffmann et al., *Ab-Initio Determination of Light Hadron Masses*, *Science* **322** (2008) 1224 [[0906.3599](#)].
- [44] STAR collaboration, X. Luo, *Energy Dependence of Moments of Net-Proton and Net-Charge Multiplicity Distributions at STAR*, *PoS CPOD2014* (2014) 019 [[1503.02558](#)].
- [45] STAR collaboration, J. Thäder, *Higher Moments of Net-Particle Multiplicity Distributions*, [1601.00951](#).
- [46] A. Andronic, P. Braun-Munzinger and J. Stachel, *Hadron production in central nucleus-nucleus collisions at chemical freeze-out*, *Nucl.Phys.* **A772** (2006) 167 [[nucl-th/0511071](#)].
- [47] P. Alba et al., *Constraining the hadronic spectrum through QCD thermodynamics on the lattice*, *Phys. Rev.* **D96** (2017) 034517 [[1702.01113](#)].
- [48] *Juqueen: Ibm blue gene/q supercomputer system at the jülich supercomputing centre*, Tech. Rep. 1 A1, Jülich Supercomputing Centre, <http://dx.doi.org/10.17815/jlsrf-1-18>, 2015.