

# ANOMALOUS DIFFUSION, DILATION, AND EROSION IN IMAGE PROCESSING

joint work with Sophia Vorderwülbecke & Bernhard Burgeth

SIAM CSE 2019 (MS 62) | February 25, 2019 | Andreas Kleefeld | Jülich Supercomputing Centre, Germany

# TABLE OF CONTENTS

- Part 1: Introduction & motivation
- Part 2: Anomalous diffusion
- Part 3: Modified dilation & erosion
- Part 4: Numerical results
- Part 5: Summary & outlook

# Part I: Introduction & motivation

# INTRODUCTION & MOTIVATION

## General idea

- Time-dependent partial differential equations (PDEs) arise naturally in image processing.
- For example: convolution of image with Gaussian kernel which is equivalent to solving a linear diffusion equation.
- Other PDEs: dilation/erosion (evolution equations).
- Can serve as building blocks for higher morphological operations (opening, closing, gradients) or deblurring filters.

# INTRODUCTION & MOTIVATION

## What is new?

- Different type of generalization of an evolution equation.
- Temporal derivative of fractional order  $\alpha$ :  $\frac{\partial^\alpha}{\partial t^\alpha}$  with  $\alpha \in (0, 2)$ .
- Definition of the fractional derivative as an extension of integration concatenated with regular differentiation (Caputo).
- Global information are considered.
- Also interesting for other applications.
- Up to now this approach was only considered for specific fractional orders as  $\alpha = 1/2$  and not for morphological operations.

## Part II: Anomalous diffusion

# ANOMALOUS DIFFUSION

## Mathematical model

- Diffusion equation:

$$\frac{{}^c\partial^\alpha}{\partial t^\alpha} u = \operatorname{div}(\kappa \operatorname{grad} u),$$

where  $\kappa$  is a constant.

- Caputo fractional derivative:

$$\frac{{}^c\partial^\alpha}{\partial t^\alpha} u = \frac{1}{\Gamma(m+1-\alpha)} \int_0^t \frac{u^{(m+1)}(\tau)}{(t-\tau)^{\alpha-m}} d\tau,$$

where  $m = \lfloor \alpha \rfloor$  and  $0 < \alpha < 1$  or  $1 < \alpha < 2$ .

- Initial condition(s): given gray-value image and in case of super-diffusion we need a second initial condition.
- Boundary condition: homogeneous Neumann.

# ANOMALOUS DIFFUSION

## Space discretization

- 2D-grid with  $h = 1$  and  $M \times N$  grid points.
- Approximation of Laplace operator with centered differences for interior nodes:  $\kappa(u_{i+1,j} + u_{i-1,j} + u_{i,j+1} + u_{i,j-1} - 4u_{i,j})$ .
- Homogeneous Neumann boundaries for exterior nodes.



# ANOMALOUS DIFFUSION

## Time discretization

- Grid of the form  $t_k = k\Delta t$ ,  $k = 0, \dots, P$  with grid size  $\Delta t = T/P$ .
- Approximation of Caputo derivative by Grünwald-Letnikov formula:

$$\left. \frac{{}_C \partial^\alpha u}{\partial t^\alpha} \right|_{\mathbf{x}=(x_i, y_j)}^{t=t_{k+1}} \approx \sum_{\ell=0}^{k+1} c_\ell^{(\alpha)} u_{i,j}^{k+1-\ell} - \sum_{n=0}^m \frac{(t_{k+1})^{n-\alpha}}{\Gamma(n-\alpha+1)} u^{(m)}(x_i, y_j),$$

where

$$c_0^{(\alpha)} = (\Delta t)^{-\alpha}, \quad c_k^{(\alpha)} = \left(1 - \frac{1+\alpha}{k}\right) c_{k-1}^{(\alpha)}.$$

# ANOMALOUS DIFFUSION

## Numerical schemes

Explicit:

$$\begin{aligned} & \sum_{\ell=0}^{k+1} c_{\ell}^{(\alpha)} u_{i,j}^{k+1-\ell} - \sum_{n=0}^m \frac{(t_{k+1})^{n-\alpha}}{\Gamma(n-\alpha+1)} u^{(m)}(x_i, y_j) \\ & = \kappa \left( u_{i+1,j}^k + u_{i-1,j}^k + u_{i,j+1}^k + u_{i,j-1}^k - 4u_{i,j}^k \right). \end{aligned}$$

Implicit:

$$\begin{aligned} & \sum_{\ell=0}^{k+1} c_{\ell}^{(\alpha)} u_{i,j}^{k+1-\ell} - \sum_{n=0}^m \frac{(t_{k+1})^{n-\alpha}}{\Gamma(n-\alpha+1)} u^{(m)}(x_i, y_j) \\ & = \kappa \left( u_{i+1,j}^{k+1} + u_{i-1,j}^{k+1} + u_{i,j+1}^{k+1} + u_{i,j-1}^{k+1} - 4u_{i,j}^{k+1} \right). \end{aligned}$$

# ANOMALOUS DIFFUSION

## Numerical schemes

Explicit:

$$\mathbf{u}^{k+1} = A \mathbf{u}^k - \mathbf{b}_{ex} \text{ with } A = \alpha I_{MN} + (\Delta t)^\alpha \kappa \cdot D_2.$$

Implicit:

$$B \mathbf{u}^{k+1} = \mathbf{b}_{im} \text{ with } B = -(\Delta t)^{-\alpha} I_{MN} + \kappa \cdot D_2.$$

$D_2$  is the 2D-Laplacian and  $\mathbf{b}_{ex}$  and  $\mathbf{b}_{im}$  are given by

$$\mathbf{b}_{ex} = (\Delta t)^\alpha \left( \sum_{\ell=2}^{k+1} c_\ell^{(\alpha)} \mathbf{u}^{k+1-\ell} - \sum_{n=0}^m \frac{(t_{k+1})^{n-\alpha}}{\Gamma(n-\alpha+1)} u^{(m)}(x_i, y_j) \right),$$

$$\mathbf{b}_{im} = -\alpha(\Delta t)^{-\alpha} \mathbf{u}^k + \left( \sum_{\ell=2}^{k+1} c_\ell^{(\alpha)} \mathbf{u}^{k+1-\ell} - \sum_{n=0}^m \frac{(t_{k+1})^{n-\alpha}}{\Gamma(n-\alpha+1)} u^{(m)}(x_i, y_j) \right).$$

## Part III: Modified dilation & erosion

# MODIFIED DILATION & EROSION

## Mathematical model & discretization

- Dilation & erosion equation:

$$\frac{{}^c\partial^\alpha}{\partial t^\alpha} u = \pm \sqrt{\left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial u}{\partial y}\right)^2}.$$

- Approximation of Caputo fractional derivative as before.
- Approximation in space by first-order finite difference scheme of Rouy-Tourin:

$$\left[ \max(-u_{i,j} + u_{i-1,j}, u_{i+1,j} - u_{i,j}, 0)^2 + \max(-u_{i,j} + u_{i,j-1}, u_{i,j+1} - u_{i,j}, 0)^2 \right]^{1/2}.$$

# MODIFIED DILATION & EROSION

## Numerical schemes

As before, we obtain an iterative scheme of the form

$$\mathbf{u}^{k+1} = \alpha \mathbf{u}^k + (\Delta t)^\alpha \mathbf{b}_{dt} \pm (\Delta t)^\alpha \sqrt{\mathbf{b}_{dx}^2 + \mathbf{b}_{dy}^2},$$

where

$$\mathbf{b}_{dt} = - \sum_{l=2}^{k+1} c_l^{(\alpha)} \mathbf{u}^{k+1-l} + \frac{t_{k+1}^{-\alpha}}{\Gamma(1-\alpha)} \mathbf{u}^0$$

and the  $i$ -th,  $j$ -th entry of  $\mathbf{b}_{dx}$  is given by

$$\max(-u_{i,j}^k + u_{i-1,j}^k, u_{i+1,j}^k - u_{i,j}^k, 0)$$

and  $\mathbf{b}_{dy}$  analogously.

# Part IV: Numerical results

# NUMERICAL RESULTS

## Stability

- Linear test problem:


$$\begin{aligned}\frac{{}^c\partial^\alpha u(t)}{\partial t^\alpha} &= \lambda u(t), \quad \lambda \in \mathbb{C}, \\ u(0) &= u_0 \text{ for } 0 < \alpha \leq 1, \\ \text{and additionally } u'(0) &= u_1 \text{ for } 1 < \alpha < 2.\end{aligned}$$

- Explicit method:  $\mathbb{C} \setminus \{(1 - z)^\alpha / z : |z| \leq 1\}$ .
- Implicit method:  $\mathbb{C} \setminus \{(1 - z)^\alpha : |z| \leq 1\}$ .



# NUMERICAL RESULTS


## Stability



**Figure:** Stability regions for explicit Euler method using parameters  $\alpha = 0.4$ ,  $\alpha = 0.6$ , and  $\alpha = 0.8$  (first row) and  $\alpha = 1.0$ ,  $\alpha = 1.2$ , and  $\alpha = 1.4$  (second row).

# NUMERICAL RESULTS

## Stability



**Figure:** Stability regions for implicit Euler method using parameters  $\alpha = 0.4$ ,  $\alpha = 0.6$ , and  $\alpha = 0.8$  (first row) and  $\alpha = 1.0$ ,  $\alpha = 1.2$ , and  $\alpha = 1.4$  (second row).

# NUMERICAL RESULTS

## Stability

- Interval of stability is  $(-2^\alpha, 0)$ .
- Implicit Euler method is  $A$ -stable for  $0 < \alpha \leq 1$  whereas we loose this property for  $1 < \alpha < 2$ .
- Could investigate  $A(\theta)$  stability, where  $\theta \leq \pi/2$  will depend on  $\alpha$ .
- We obtain the  $\theta$  angles (in degrees  $^\circ$ ) 90, 81, 72, 63, 54, 45, 36, 27, 18, and 9 for the parameters  $\alpha = 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8$ , and 1.9, respectively.
- Hence, it appears to be that  $\theta$  is given by  $(2 - \alpha) \cdot 90^\circ$  for  $1 \leq \alpha < 2$  (the proof remains open).

# NUMERICAL RESULTS

## Convergence

- Homogeneous initial conditions: convergence order 1
- Non-homogenous initial conditions: convergence order depends on  $\alpha$
- Calculation of error:

$$\frac{{}^c\partial^\alpha u(t)}{\partial t^\alpha} = t^2, \quad u(0) = 0, \quad 0 \leq t \leq 1, \quad 1 < \alpha \leq 2$$

with exact solution

$$u(t) = \frac{\Gamma(3 + \alpha)}{\Gamma(3)} t^{2+\alpha}.$$

Estimated convergence order (EOC):

$$\text{EOC} = \frac{\log(E_{\Delta t}/E_{\Delta t/2})}{\log(2)}, \text{ where } E_{\Delta t} = |u(1) - \tilde{u}_{\Delta t}(1)|.$$

# NUMERICAL RESULTS

## Convergence

	$\alpha = 0.4$		$\alpha = 0.8$		$\alpha = 1.0$		$\alpha = 1.2$	
$\Delta t$	$E_{\Delta t}$	EOC	$E_{\Delta t}$	EOC	$E_{\Delta t}$	EOC	$E_{\Delta t}$	EOC
1/10	0.1220		0.0685		0.0483		0.0324	
1/20	0.0627	0.96	0.0350	0.97	0.0246	0.98	0.0164	0.99
1/40	0.0318	0.98	0.0177	0.98	0.0124	0.99	0.0082	1.00
1/80	0.0160	0.99	0.0089	0.99	0.0062	0.99	0.0041	1.00
1/160	0.0080	1.00	0.0045	1.00	0.0031	1.00	0.0021	1.00
1/320	0.0040	1.00	0.0022	1.00	0.0016	1.00	0.0010	1.00
1/640	0.0020	1.00	0.0011	1.00	0.0008	1.00	0.0005	1.00

**Table:** Estimated order of convergence for the explicit Euler method using the parameters  $\alpha = 0.4, 0.8, 1.0$ , and  $1.2$ .

# NUMERICAL RESULTS


## Convergence

	$\alpha = 0.4$		$\alpha = 0.8$		$\alpha = 1.0$		$\alpha = 1.2$	
$\Delta t$	$E_{\Delta t}$	EOC	$E_{\Delta t}$	EOC	$E_{\Delta t}$	EOC	$E_{\Delta t}$	EOC
1/10	0.0323		0.0487		0.0517		0.0519	
1/20	0.0161	1.00	0.0241	1.01	0.0254	1.02	0.0253	1.03
1/40	0.0081	1.00	0.0120	1.01	0.0126	1.01	0.0125	1.02
1/80	0.0040	1.00	0.0060	1.00	0.0063	1.00	0.0062	1.01
1/160	0.0020	1.00	0.0030	1.00	0.0031	1.00	0.0031	1.00
1/320	0.0010	1.00	0.0015	1.00	0.0016	1.00	0.0015	1.00
1/640	0.0005	1.00	0.0007	1.00	0.0008	1.00	0.0008	1.00


**Table:** Estimated order of convergence for the implicit Euler method using the parameters  $\alpha = 0.4, 0.8, 1.0$ , and  $1.2$ .

# NUMERICAL RESULTS


## Anomalous diffusion




(a)  $T = 1, \alpha = 1/2$




(b)  $T = 1, \alpha = 3/4$




(c)  $T = 1, \alpha = 1$



(d)  $T = 10, \alpha = 1/2$



(e)  $T = 10, \alpha = 3/4$




(f)  $T = 10, \alpha = 1$


**Figure:** Anomalous sub-diffusion with  $T = 1, 10$  and  $\alpha = 1/2, 3/4, 1$  for the Lena image.

# NUMERICAL RESULTS


## Modified dilation




(a)  $T = 1, \alpha = 1/2$




(b)  $T = 1, \alpha = 3/4$



(c)  $T = 1, \alpha = 1$



(d)  $T = 10, \alpha = 1/2$



(e)  $T = 10, \alpha = 3/4$



(f)  $T = 10, \alpha = 1$

**Figure:** Modified dilation with  $T = 1, 10$  and  $\alpha = 1/2, 3/4, 1$  for the Lena image.



# Part V: Summary & outlook

# SUMMARY & OUTLOOK

- Modified standard diffusion as well as dilation & erosion for gray-valued images.
- Treated numerically by explicit and implicit Euler method.
- Showed convergence and stability.
  
- Consider second-order approximation of the Caputo fractional derivative.
- Multistep methods (BDF, Adams-Moulton, and Adams-Bashforth methods).
- Consider corresponding inverse problems (denoising).
- Extension for higher morphological operations.
- Extending the approach to color images.

# REFERENCE



A. KLEEFELD, S. VORDERWÜLBECKE, & B. BURGETH, *Anomalous diffusion, dilation, and erosion in image processing*, International Journal of Computer Mathematics 95 (6–7), 1375–1393 (2018), special issue: “Advances on Computational Fractional Partial Differential Equations”.