

1 **Immediate and carry-over effects of increased soil frost on soil respiration**
2 **and microbial activity in a spruce forest**

3 Kaijun Yang^{1, 2}, Changhui Peng^{3, 4}, Josep Peñuelas^{2, 5}, Paul Kardol⁶, Zhijie Li^{1, 7}, Li Zhang¹,
4 Xiangyin Ni¹, Kai Yue¹, Bo Tan¹, Rui Yin⁸, Zhenfeng Xu^{1*}

5 1. Institute of Ecology and Forest, Sichuan Agricultural University, Chengdu, China

6 2. Global Ecology Unit CREAF-CSIC-UAB, CSIC, Barcelona, Catalonia, Spain

7 3. Department of Biological Science, Institute of Environment Sciences, University of
8 Quebec at Montreal, Montreal, Canada;

9 4. Center for Ecological Forecasting and Global Change, College of Forestry, Northwest
10 Agriculture & Forest University, Yangling, Shaanxi, China;

11 5. CREAF, Barcelona, Spain

12 6. Department of Forest Ecology and Management, Swedish University of Agricultural
13 Science, 90183, Umeå, Sweden

14 7. Forschungszentrum Jülich GmbH, Agrosphere (IBG-3), Jülich, Germany

15 8. Helmholtz-Centre for Environmental Research-UFZ, Department of Community Ecology,
16 Theodor-Lieser-Strasse 4, 06110 Halle (Saale), Germany

17

18 **Correspondence:** Dr. Zhenfeng Xu, Email: xuzf@sicau.edu.cn, Tel: +86 28 86290957,
19 No.211, Huimin Road, Wenjiang, Chengdu, Sichuan, 611130 China

20 **Key words:** Snow exclusion, soil enzyme, microbial biomass, nitrogen availability, soil
21 aggregate, fine root

22 **Running title:** Frost effect on soil carbon cycling

Abstract

Increased soil frost associated with winter climate change could have immediate and carry-over effects on biological processes in high-altitude forest soils, but the nature of these processes remain poorly understood. We conducted a snow-exclusion experiment to investigate the immediate and cross-seasonal effects of increased soil frost on soil CO₂ efflux and biological activity in a subalpine spruce forest on the eastern Tibetan Plateau, China. The increased frost reduced soil CO₂ efflux by ~15 and ~19% in the winters of 2015/2016 and 2016/2017, respectively. Increased frost tended to decrease soil basal respiration, the amount of microbial phospholipid fatty acids and the activities of enzymes involved in soil carbon cycling during the winters. Winter soil nitrogen availabilities were higher in the snow-exclusion treatment than in the control plots. However, these effects did not carry over to the following growing season. Our results suggest that increased frost reduces winter soil respiration by direct environmental effects (e.g. soil temperature) and indirect biological processes (e.g. microbial biomass and activity), whereas increased frost did not have any cross-seasonal effects. These findings underscore the ecological importance of seasonal snowpack and microbe-associated carbon processes in subalpine forests where winter snowfall is decreasing substantially.

1. Introduction

Seasonal snow cover is a major control of biogeochemical cycling in cold environments (Jusselme et al., 2016). Many snowy areas at high latitudes and altitudes have experienced substantial climate change in recent decades, and this trend is predicted to continue in this century (IPCC, 2013). Climate-induced changes are particularly rapid in high latitude and alpine ecosystems, where rising temperatures have profound effects on winter conditions, such as snowfall, soil frost and extreme climatic events (Liu et al., 2012; IPCC, 2013). Winter precipitation in these regions is more likely to occur in the form of rain rather than snow due to winter warming (Wang et al., 2016). The lack of insulating snow cover could consequently increase soil frost (Groffman et al., 2001a; Bokhorst et al., 2013), which could in turn have complex and large impacts on soil microbiological and biochemical processes in cold forests.

Winter biological processes and their controls are not as well understood as growing-season processes, despite the importance of winter warming and biological activity in cold systems (Sanders-DeMott and Templer, 2017). Recent studies have found that soil biological processes are sensitive to warming-induced changes in winter conditions, especially snow cover and soil frost (Haei and Laudon, 2015; De Long et al. 2016; Li et al., 2016). Snow removal has negative or neutral influences on winter soil respiration in boreal and temperate forests (Groffman et al., 2006; Aanderud et al., 2013; Bokhorst et al., 2013), but changes in snow cover can also alter biological processes in snow-free periods (Muhr et al., 2009; Wubs et al., 2018). Snow exclusion can suppress soil respiration in the snow-free season in high-latitude ecosystems (Öquist and Laudon, 2008; Zhao et al. 2017). To our knowledge, however, soil biological responses to changing soil frost have rarely been investigated in both snow-covered and snow-free periods in the same experiment. A better understanding of the impacts of intensified soil frost on the biotic and abiotic controls over the dynamics of soil C in both snow-covered and snow-free seasons is thus essential for accurately modeling and predicting potential C feedbacks in a warmer world.

Altered soil frost may directly and indirectly affect soil C cycling, such as by affecting soil temperature and moisture (Aanderud et al., 2013; Song et al., 2017), soil microbial biomass and activity (Monson et al., 2006b; Sorensen et al., 2016) and substrate quality and quantity (Brooks et al., 2004; Steinweg et al., 2008; Comerford et al., 2013). The direction and magnitude of biological responses to increased frost may be determined by the combined effect of these processes. Diverse techniques have provided insight in recent years into the impacts of winter climate change on soil C cycling as the importance of winter processes has increased (Li et al., 2016a). Most field-manipulation studies have focused mainly on high-latitude systems, including peatlands and boreal forests (Sanders-DeMott and Templer, 2017). Soil biological responses from low-latitude cold systems with unique winter conditions, such as Tibetan subalpine forests, however, remain unknown.

The Tibetan Plateau, the Earth's 'Third Pole', has warmed substantially, especially in winter (Chen et al., 2013). Winter snowfall has decreased at a rate of 0.6 mm y⁻¹ in recent decades (Wang et al., 2016; Xu et al., 2017). Seasonal snowpack in this region has unique characteristics, such as shorter duration and shallower depth relative to high latitudes. Winter soil temperature is also near the physical melting point and is sensitive to changes in snow cover (Li et al., 2017). The subalpine forests of southwestern China contain a large amount of soil organic C (Zhang et al., 2013), but most studies of global-change biology have only focused on responses during the growing season (e.g. Xu et al., 2012; 2015; Yin et al., 2013), even though warming is extremely pronounced and microbial activity is unexpectedly high during winter (Wang et al., 2016; Wang et al. 2012; Tan et al., 2014). Future soil frost will also likely affect the biological and environmental controls of soil C cycling in these forests, but the underlying mechanisms of such processes remain unknown.

We conducted a snow-manipulation experiment to investigate the immediate and carry-over effects of increased winter frost on soil C cycling in a spruce forest on the eastern Tibetan Plateau. Specifically, we hypothesized that (1) more intense frost in the soil as a result of

exclusion of snow would decrease microbial activity and soil respiration in winter; (2) frost-reduced biological processes would carry over into the subsequent snow-free growing season.

2. Materials and methods

2.1 Site description

The field manipulation experiment was conducted in a dragon spruce (*Picea asperata* Mast.) stand at the Long-term Research Station of Alpine Forest Ecosystems of Sichuan Agricultural University on the eastern Tibetan Plateau of China (31°15'N, 102°53'E; 3021 m a.s.l.). The mean annual precipitation and temperature are 850 mm and 3.0 °C, respectively. Snow generally begins to accumulate in late November and melts in late March the following year. The soil is classified as a Cambic Umbrisol (IUSS Working Group WRB, 2007). The soil (0-15 cm) contains 88.5 g kg⁻¹ organic C and 5.4 g kg⁻¹ nitrogen (N) and has a pH of 6.4 (Li et al., 2017).

2.2 Experimental design

Winter snowfall was excluded using shelters to intensify soil frost. Shelters are considered to be a useful tool for studying the responses of soil processes to winter climate change because they can effectively reduce snow cover and minimize the changes in other unwanted environmental conditions (Li et al., 2016a). In early November 2015, six wooden roofs were set up in the spruce forest to prevent the accumulation of snow on the ground. One control plot was established in the vicinity of each roof. The roofs were 2 m in height with a ground area of 3 × 3 m. The snow manipulation began in mid-November and ended in late March the following year.

2.3 Soil sampling

Soil samples were collected from the topsoil (0-15 cm) in the frost period (FP, late January), early thawing period (ETP, early April) and the middle of the growing season (MGS, mid-August) in the year of 2016 and 2017, respectively. Three cores (5 cm in diameter, 15 cm in depth) were collected in each plot at each sampling. The three cores from each plot were combined to form one composite sample. Each composite sample was passed through a 2-mm sieve, and any visible living plant material was manually removed. The sieved soil was used for biochemical analysis.

2.4 Soil CO₂ efflux

Two PVC collars (20 cm in diameter, 12 cm in height) were permanently installed in each plot for measuring soil respiration. Soil CO₂ efflux was measured using a portable infrared gas analyzer (Li-8100, Li-Cor Inc., Lincoln, USA) between 10:00 and 14:00 (Beijing time, China Standard Time) approximately every two weeks during the experimental period. The measurements were repeated twice for each collar to minimize measurement errors. Soil temperature and moisture at a depth of 5 cm were also measured near each collar using thermocouple and Theta probes, respectively, connected to the LI-8100. Small red flags were attached to the PVC collars in the control plots to minimize disturbance during the period of snow cover. During the winters of 2015/2016 (four times) and 2016/2017 (once), the surface snow was removed carefully from the top of the collars when the snowpack was thicker than the height of the collar. We then waited 5 min to allow the system to equilibrate before measuring the CO₂ efflux. The removed snow was gently backfilled after the measurements. We measured CO₂ efflux under the natural snowpack (~10 cm) in the winter of 2015/2016 in the same forest stand adjacent to the snow manipulation site. CO₂ efflux was also measured after removing the snow within and around the collars. CO₂ efflux did not differ significantly

before and after snow removal (unpublished data). Snow removal therefore likely negligibly affected the quantification of immediate CO₂ efflux, at least within an interval of a few minutes.

2.5 Microclimate, extractable N and microbial respiration

Air temperature 2 m above the ground in the forest stand was measured using Thermochron iButton DS1923–F5 Recorders (Maxim Dallas Semiconductor Corp., USA) every 2 h during the experimental period. Meanwhile, soil temperatures 5 cm below the surface were recorded in the snow-exclusion and control plots, respectively. Snow depth in the control plots was measured by a metal ruler approximately every two weeks during winter.

Soil extractable N (nitrate, NO₃⁻-N, and ammonium, NH₄⁺-N) was extracted with 2 M KCl (1:5 soil:solution). The extracts were shaken for 1 h and filtered with a filter paper. The concentrations of NO₃⁻-N and NH₄⁺-N in the extracts were determined by colorimetry (Li et al., 2017).

The rate of soil microbial respiration was estimated using alkali absorption (Anderson et al., 1982). Soil samples (50 g) were incubated in 1-L jars at 20 °C for 2 weeks. Empty jars without soil were used as controls. The CO₂ produced was captured with 0.5 M NaOH in a beaker suspended inside each jar. The NaOH solution was removed and titrated with 0.25 M HCL solution to determine the amount of CO₂ produced. Microbial respiration was reported as mg CO₂ kg⁻¹ soil d⁻¹.

2.6 Aggregate fraction and fine-root biomass

Aggregates were isolated as described by Kristiansen et al. (2006). Two soil cores from each plot were collected from the 0-15 cm layer using an auger 10 cm in diameter in the early thawing periods of 2015/2016 and 2016/2017 winters. Soil samples were air-dried to optimal

moisture (~10-15%) that would allow limited mechanical stress to maximize brittle failure along natural planes of weakness, and the samples were then gently manually crumbled to <8 mm. The recovered samples were transferred to a nest of sieves (2 and 0.25 mm) and shaken at 100 min⁻¹ for 2 min. All visible roots and stones were removed, and aggregates >2 mm (large macroaggregates) were collected. The same procedure was used for the material retained on the 0.25 mm sieve, isolating an aggregate size class 0.25-2 mm (small macroaggregates). The remaining material passing through the 0.25 mm sieve was identified as aggregate class <0.25 mm (microaggregates).

Two soil cores were collected from each plot using an auger (15 cm long and 10 cm in diameter) in the ETPs of 2015/2016 and 2016/2017 winters. Root samples were washed in the laboratory on sieves (mesh size 0.1 mm) and dried to constant weight at 65 °C. Fine roots (<2 mm in diameter) were separated into live and dead components based on their color and mechanical consistency.

2.7 Assays of soil phospholipid fatty acids and enzymes

Microbial biomass was estimated as the total extractable phospholipid fatty acids (PLFAs) with a modified method described by White et al., (1996). Lipids from 2 g of fresh soil were extracted in a chloroform-methanol-phosphate buffer mixture (1:2:0.8). The phospholipids in the extracts were transformed by alkaline methanolysis into fatty acid methyl esters (FAMES), which were identified by gas chromatography/mass spectrometry (GC/MS-QP2010 Series, Shimadzu, Japan). Fatty acids were quantified by comparisons of the peak areas from the sample with the peak areas of internal standards at 19:0 (nonadecanoic methyl ester) of the known concentration. The areas were used to estimate the abundance of PLFA markers, which were expressed as nmole g⁻¹ dry soil.

We assessed the activities of four enzymes involved in soil C cycling: two hydrolytic enzymes, β -glucosidase (BG) that catalyzes one of the later steps of cellulose degradation and

β -N-acetyl-glucosaminidase (NAG) involved in the breakdown of chitin and fungal cell walls, and two oxidases, polyphenol oxidase (PPO) that breaks down recalcitrant polymers such as lignin and humic compounds and peroxidase (POD), a nonspecific enzyme that oxidizes and depolymerizes lignin. The activities were measured using assay techniques described by Allison and Jastrow (2006). Substrate solutions were 5 mM pNP- β -glucopyranoside for BG, 50 mM pyrogallol and 50 mM EDTA for PPO, 2 mM pNP- β -N-acetylglucosaminide for NAG and 5 mM L-DOPA and 10 μ L of 0.3% H₂O₂ for POD. Activities were determined using a microplate spectrophotometer and expressed as μ mol of substrate produced or consumed h⁻¹ g⁻¹ dry soil.

Data analysis

A repeated-measures ANOVA was used to test the effects of treatment, sampling date and their interactions on all response variables. A Bonferroni post hoc test was used to examine the treatment effect on the variables on a given sampling date when the interaction of treatment and sampling date was significant ($P < 0.05$), and a paired t -test was used when the interaction was not significant. All data were tested for the assumptions of an ANOVA before analysis. Heterogeneous data were ln-transformed before analysis. An exponential regression model was used to describe the relationship between CO₂ efflux and soil temperature during specific periods (winter, growing season and entire year). All data from two winters or growing seasons were used for the analyses due to the limited number of measurements. Winter was defined as the period between the first day in autumn and the last day in spring when soil temperature was continually below 5 °C for 5 d in the control plots. The temperature sensitivity (Q_{10}) of soil respiration was estimated using van't Hoff equation (Van's Hoff, 1898). $R = \alpha \times e^{\beta \times T}$, Where R is the soil respiration rate (μ mol m⁻² s⁻¹), T is the soil temperature at 5 cm (°C), α and β are parameters. The Q_{10} values are calculated as: $Q_{10} = e^{10 \times \beta}$. All statistical tests were performed using the Software Statistical Package for the Social

Sciences (SPSS) version 17.0 (IBM SPSS Statistics Inc., Chicago, IL, USA).

3. Results

3.1 Treatment effect on winter soil conditions

The mean and minimum air temperatures were -2.1 and -14.1 °C during the winter of 2015/2016 and -0.9 °C and -6.4 °C during the winter of 2016/2017, respectively (Figure 1 a). The maximum snow depth was 40 cm in the winter of 2015/2016 but only 23 cm in the winter of 2016/2017. The mean air temperature in the winter of 2015/2016 (-2.1 °C) was comparable to the seven-year average of -2.4 °C for 2010-2016. The mean air temperature in the winter of 2016/2017 (-0.9 °C), however, was the highest in the last seven winters and 1.5 °C higher than the mean.

The snow-exclusion treatment successfully created a more intense frost regime in both winters (Figure 1 a). The minimum daily mean soil temperatures were -2.2 °C (2015/2016) and -2.4 °C (2016/2017) in the snow-exclusion plots but were only -0.5 °C (2015/2016) and -1.3 °C (2016/2017) in the control plots. The numbers of days with differences in soil temperature ≥ 0.5 °C between the treatment and control plots were 42 and 56 for the winters of 2015/2016 and 2016/2017, respectively. Such differences were mainly during mid- and late winter when snow cover was >10 cm. Soil temperature fluctuated more in the treatment than the control plots. Volumetric soil moisture was similar between the treatment and control plots across the two years ($F=3.364$, $P = 0.116$, Figure 1 b).

3.2 Soil CO₂ efflux

The snow exclusion lowered CO₂ efflux early in the winter of 2015/2016 and in mid-winter of 2016/2017. The snow exclusion reduced CO₂ efflux by averages of 15% and 19% in the winters of 2015/2016 and 2016/2017, respectively, and these reductions were

statistically significant ($F = 11.13$, $P < 0.01$ for the winter of 2015/2016; $F = 9.143$, $P < 0.05$ for the winter of 2016/2017). The snow-exclusion manipulation, however, did not affect CO₂ efflux during the snow-free growing seasons ($F = 1.065$, $P = 0.323$ for 2016; $F = 1.354$, $P = 0.305$ for 2017). Mean CO₂ efflux differed marginally between the frost regimes in the winter of 2015/2016 ($t = 2.006$, $P = 0.076$; Table 1), but differed significantly between the regimes in the winter of 2016/2017 ($t = 3.909$, $P < 0.01$). Mean CO₂ efflux nevertheless did not differ significantly between the treatment and control plots in either growing season ($t = -1.584$, $P = 0.335$ for 2016; $t = -0.285$, $P = 0.465$ for 2017).

Soil CO₂ efflux increased exponentially with soil temperature throughout the study period (Figure 3a-c). Soil temperature explained 82-83% of the variation in CO₂ effluxes during the growing seasons (Figure 3b) but explained only 52-53% of the variations in the winters (Figure 3a). Soil temperature explained 90-91% of the variance in CO₂ effluxes when the data for the two years were pooled (Figure 3c). The temperature sensitivity (Q_{10}) of the CO₂ efflux was 23.3, 3.2 and 4.4 in the treatment plots and 22.6, 3.3 and 4.7 in the control plots for winter, growing season and the entire year, respectively. Q_{10} did not differ significantly between the frost regimes for each period modeled (all $P > 0.05$).

3.3 Soil PLFAs and microbial respiration

The intensified frost tended to decrease the soil PLFAs biomarkers. PLFA content was lower in the snow-exclusion treatment than the control plots in the FPs of 2015/2016 ($t = -2.072$, $P < 0.05$) and 2016/2017 ($t = -3.686$, $P < 0.05$; Figure 4) but did not differ significantly in the MGSs of 2016 ($t = 1.368$, $P = 0.245$) or 2017 ($t = 0.035$, $P = 0.895$).

Microbial activity, measured as basal respiration without roots, was estimated by determining CO₂ emission. The intensified frost tended to decrease soil microbial respiration in the winter. The snow-exclusion treatment negatively affected soil microbial respiration in the FP of 2015/2016 ($t = -0.918$, $P < 0.05$; Figure 5) and in the ETP of 2016/2017 ($t = -5.821$,

$P < 0.01$) but had no effect in the MGSs of 2016 and 2017 (both $P > 0.05$)

3.4 Soil enzymes

The activities of the soil enzymes varied significantly with sampling date (all $P < 0.01$, Figure 6a-d). The snow-exclusion treatment tended to inhibit activities in the winter. Activity was significantly lower in the treatment than in the control plots for BG in the ETP of 2015/2016 ($t = -1.975$, $P < 0.05$; Figure 6a) and for PPO in the ETP of 2016/2017 ($t = -2.643$, $P < 0.05$; Figure 6b). The intensified frost decreased POD activity in the FPs of 2016 and 2017 (both $P < 0.05$, Figure 6c) and decreased NAG activity in the ETPs of 2016 and 2017 (both $P < 0.05$, Figure 6d) but did not significantly affect the activities of the enzymes in the MGSs of 2016 or 2017.

3.5 Soil extractable N

Frost treatment, sampling date and their interaction all significantly affected soil $\text{NH}_4^+\text{-N}$ concentration (all $P < 0.05$, Figure 7a). The snow-exclusion treatment increased $\text{NH}_4^+\text{-N}$ concentrations in the FP and ETP of 2015/2016 (all $P < 0.01$) but not in the winter of 2016/2017 (both $P > 0.05$). Likewise, the intensified frost increased $\text{NO}_3^-\text{-N}$ concentrations in both winters ($F = 16.575$, $P < 0.01$; Figure 7b). $\text{NO}_3^-\text{-N}$ concentrations were significantly higher in the treatment than the control plots in the ETP of 2015/2016 ($t = 2.309$, $P < 0.05$) and in the FP of 2016/2017 ($t = 5.017$, $P < 0.01$). Neither $\text{NH}_4^+\text{-N}$ nor $\text{NO}_3^-\text{-N}$ concentration, however, differed between the frost regimes in the MGS of 2016 and 2017 (both $P > 0.05$).

3.6 Aggregate fraction and fine-root biomass

The relative distribution of the aggregate-size classes of the bulk soil was in the order small macroaggregates (0.25-2 mm) > large macroaggregates (>2 mm) > microaggregates (<0.25 mm) irrespective of frost regime ($F = 221.75$, $P < 0.001$; Table 2). The snow-exclusion treatment did not affect the distribution of aggregates in the size classes ($F =$

0.159, $P = 0.897$), the live fine-root biomass ($F = 0.202$, $P = 0.663$; Table 3) or the dead fine-root biomass ($F = 0.171$, $P = 0.688$) in the ETP.

4. Discussion

We investigated the impact of intensified soil frost on soil C cycling in a Tibetan subalpine spruce forest using a field experiment manipulating frost. Our main objective was to determine whether differences in winter frost conditions induced immediate and carry-over effects on soil CO₂ efflux. In line with the first hypothesis, snow exclusion resulted in more intensive soil frost, which decreased soil respiration in the winter season. However, contrary to what we expected, increased winter frost did not have carry-over effects on soil respiration in the subsequent growing season. Several potential mechanisms have been tested to account for the underlying the changes of CO₂ efflux during the snow-covered and snow-free seasons.

Firstly, frost manipulation could decrease winter CO₂ efflux, in part, by the direct effect of temperature, because snow exclusion during the winter decreased soil temperature. A slight reduction in soil temperature over the lower range is especially important to CO₂ production, because the temperature sensitivity of microbial processes is extremely high at low temperatures (Davidson and Janssens, 2006; Schütt et al., 2014). Soil temperature in our study accounted for only 52-53% of the variance in winter CO₂ efflux. CO₂ efflux was very sensitive to small changes in soil temperature throughout the winter, particularly near 0 °C. Q_{10} was much higher in the winter than in the growing season regardless of frost regime. A several studies have described a surprisingly higher temperature sensitivities for winter soil respiration in temperate and boreal forests (Monson et al., 2006a; Muhr et al., 2009; Wang et al., 2010). Q_{10} , however, was estimated over a narrow temperature span (~5-6 °C), and the temperature-CO₂ relationship was as weak as in temperate forests (Schindlbacher et al., 2007; Wang et al., 2010; Schindlbacher et al., 2014).

Other factors co-varying with soil temperature may also likely regulate winter CO₂ efflux. The intensified frost may have produced a stronger ‘freezing drought’, which would likely limit microbial activity and the extracellular diffusion of substrates (Rivkina et al., 2000). Soil moisture did not differ significantly between the treatment and control plots during the winters, suggesting that soil moisture was not likely responsible for the decreased winter CO₂ efflux. The lack of significant differences in CO₂ efflux during the snow-free seasons was likewise partially attributed to the lack of significant differences in both soil temperature and moisture between the treatment and control plots during the growing season.

Secondly, root activity is extremely low during dormant seasons, and winter soil respiration in cold ecosystems is primarily derived from microbial decomposition (Muhr et al., 2009; Wang et al., 2010). Winter soil respiration is thus largely determined by the biomass and activity of soil microbes (Lipson et al., 2002; Moorhead et al., 2014). Soil microbes are very susceptible to soil frost (Monson et al., 2006b; Aanderud et al., 2013), which can kill a substantial proportion of the organisms by the rupture of cell membranes by ice crystals (Sulkava and Huhta, 2003; Jusselme et al., 2016; Gavazov et al., 2017). We also found that the increased frost significantly reduced soil microbial PLFAs, implying a lowered potential for the microbial community to metabolize soil C in the winters. A significant decline in the cross-winter microbial PLFAs also implied that seasonal frost would kill the soil microbes in the spruce forest irrespective of frost manipulation. Our measurements of microbial basal respiration, excluding plant roots, also indicated a similar decline with winter frost, consistent with *in situ* soil CO₂ efflux. Frost-induced decreases in winter soil respiration may thus largely be attributed to the lower microbial biomass and activity. Soil PLFAs and basal respiration in the middle of the growing season nevertheless did not differ significantly between frost regimes. These observations may partially account for the neutral effect of increased frost on CO₂ efflux during the snow-free growing seasons.

Soil enzymes play very important roles in the cycling of soil C and nutrients. Little

attention has been paid to enzymatic activities in studies of winter climate change, despite the importance of soil enzymes in soil C cycling. A recent study found that enzymatic activities were negatively correlated with the intensity of soil frost in mixed-hardwood forests (Sorensen et al., 2016), and another experiment also found that snow removal decreased the activity of soil invertase in an alpine spruce-fir forest (Tan et al., 2014). We assayed the activities of four enzymes involved in soil C cycling to further assess the functional capacity of soil. The intensified frost tended to reduce soil enzymatic activities. Soil enzymatic activities are strongly temperature-dependent (Tabatabai, 1982), so a decrease in soil temperature caused by snow-exclusion may, to some extent, reduce soil enzymatic activities directly. The lower activities may also partly be attributed to the smaller population size of the microbes, which are an important source of enzyme synthesis. Soil enzymes, as proximate agents of the decomposition of soil organic C, can break down plant and microbial cell walls and catabolize macromolecules into soluble substrates for microbial assimilation (Sinsabaugh et al., 2008). Frost-induced decreases in enzymatic activities may thus constrain this decomposition, which could also partly account for the lower winter CO₂ efflux. Conversely, intensified frost did not affect activities in the snow-free growing seasons, which may account for the lack of significant responses during the subsequent growing season.

Thirdly, soil frost may also have affected the decomposition of soil C in winter by altering nutrient availability. Intensified frost can increase the mortality of roots and microbes (Henry, 2007; Repo et al., 2014; Blume-Werry et al., 2016), which are important substrates for soil microbial metabolism during winter (Schimel et al., 2004). Dead roots and microbes are also main N sources during winter in cold systems (Chapin III et al., 1988; Tierney et al., 2001). In an earlier study we observed that soil at -5 °C could release considerable extractable N in the soils of this spruce forest, possibly due to the effect of freezing on microbial mortality (Xu et al., 2014). The snow-exclusion treatment in the present study stimulated the production of soil extractable N in the two winters, likely due mainly to the increased microbial mortality. An increase in N availability but a decrease in soil PLFAs throughout the

winter, irrespective of the frost regime, may also support this conclusion. Live and dead fine-root biomass also notably did not differ significantly between frost regimes later in the winter, further suggesting that the increased N availability was mainly attributed to microbial mortality rather than root injury. An increase in N availability coincided with a decrease in CO₂ efflux, implying that the cycling of soil C could be decoupled from N availability during winter under intensified frost.

In addition to microbial and root mortality, substrate availability could have been affected by the physically disruptive effects of frost on soil aggregates (Chai et al., 2014). Freezing can break down macroaggregates into microaggregates (Oztas and Fayetorbay, 2003). Microaggregates with a larger surface area have more contact points, which can potentially increase the amount of substrate decomposed by microorganisms (Grogan et al., 2004). Snow removal increased the fraction of microaggregates in a northern hardwood forest, implying that soil substrate could become more accessible to soil microorganisms (Steinweg et al., 2008). Our observations, however, did not provide further evidence that more intense frost could disrupt aggregates in the soil of this Tibetan spruce forest. The intensified frost did not affect the distribution of aggregates among the size classes, suggesting that frost-associated changes to aggregates may not importantly affect soil respiration in the spruce forest during the winter and growing season.

Lastly, the flux of CO₂ derived from decaying litter accounts for a considerable part of total soil respiration during winter (Uchida et al., 2005). CO₂ flux derived from aboveground litter accounts for an average of 14.2% of total soil respiration in this spruce-forest stand (Xiong et al., 2015). In a previous study we also found that the mass loss of spruce needles over the winter constituted 18.3-28.8% of the net loss rates for the entire year (Xu et al., 2016). The lack of snow cover at this experimental site decreased the temperature of the surface soil by an average of 1.4 °C during the winter (Li et al., 2017), implying that litter decomposition was most likely inhibited by the lower temperatures. A growing number of

studies have documented that thick snow covers can provide relatively stable conditions for biological activity, favoring the decomposition of plant litter (Christenson et al., 2010; Bokhorst et al., 2013; Saccone et al., 2013). The rates of decomposition of litter from subalpine tree species in this area similarly decrease with decreasing snow depth (Ni et al., 2014; He et al., 2015). The lower rate of litter decomposition due to the lack of snow cover may therefore also have contributed to the lower winter soil respiration in the snow-exclusion plots, but further supporting evidence is needed.

5. Implications

The climate on the Tibetan Plateau has changed considerably in recent decades, especially in winter (Wang et al., 2016). Winter snowfall has tended to decrease substantially due to strong winter warming (Xu et al., 2017). The decrease or absence of insulating snow cover associated with climate change may thus increase the duration and intensity of soil frost in the future in this special region. The importance of soil frost, large storage of soil C and sensitivity of snow cover to winter warming indicate that understanding the potential effects of projected frost increases on soil C cycling in the subalpine forests of western China is essential. To our knowledge, our study is the first to identify the effects of changes in soil frost on soil C cycling in a Tibetan forest. Our results generally indicate that more intense soil frost decreases winter soil respiration and biological activities. Winter soil CO₂ emission was lower in the snow-exclusion than in the control plots during the two winters of the study. Intensified soil frost, however, did not affect soil CO₂ efflux and biological activities during the subsequent growing season, suggesting that a short-term change in snow cover does not produce large carry-over effects in snow-free periods. If the observed effects apply to natural conditions, intensified soil frost would decrease the amount of soil C released to the atmosphere from subalpine forests during winter, but additional supporting evidence is

needed.

This study was conducted during two contrasting winters (cold winter and thick snow cover in 2015/2016 and mild winter and thin snow cover in 2016/2017) so offered a good opportunity for determining the effect of the lack of snow cover in winters with different weather on soil C cycling in the Tibetan spruce forest. The decrease in soil respiration due to frost in the first winter occurred early but then disappeared, suggesting that soil biological processes may begin to acclimate to the frost late in the winter. Soil respiration early in the mild winter of 2016/2017 did not differ significantly between the treatment and control plots, mainly due to the absence of an insulating snow cover. CO₂ effluxes, however, were lower in the treatment plots after the formation of a steady snow cover (>10 cm). In addition to variable snowfall, extreme winter events (e.g. warm weather and snow storms) may become more frequent and likely under scenarios of future climate, indicating the complexity and uncertainty of winter climate change in this specific region. The comparably strong climate change and variable winter snowfall on the Tibetan Plateau bring great challenges and opportunities for studying winter climate change and its impacts on the structure and function of Tibetan ecosystems. Long-term monitoring is strongly needed for exploring the natural winter variations and underlying mechanisms of the observed phenomena to help models for providing more realistic predictions of future winter conditions.

The frost intensity due to the lack of snow cover was low at our experimental forest site, unlike in temperate and boreal forests (e.g. Groffman et al., 2006; Muhr et al., 2009; Sorensen et al., 2016), but the difference in temperature minima was nearly 2 °C, likely due to the site-specific characteristics, such as winter snowfall, air temperature, properties of soil heat transfer and albedo. Such soil frost, however, had large impacts on soil respiration, microbial PLAFs, enzymatic activity and N availability, suggesting that Tibetan forest soils will be sensitive to changing soil frost in the future. The direction and magnitude of the response of soil respiration to intensified soil frost may largely depend on the interaction between less

snowfall and warmer temperature in winter. Winter warming may offset the negative effects induced by frost to some extent. Seasonal snow cover in cold regions plays a key role in decoupling soil from cold winter weather, but soil temperature is often insensitive to a small change in air temperature. Changes in snow cover will thus likely have a stronger influence on soil biological processes than winter warming itself (Groffman et al., 2001a), and long-term changes in soil frost may also have carry-over effects on soil C dynamics during the subsequent growing season in cold systems (Zhao et al., 2017). More research is warranted to integrate potential factors and separate their relative importance for a better understanding and ability to predict potential C feedbacks in snowy regions under a warmer future.

6. Conclusions

This study explored the immediate and carry-over effects of intensified soil frost on soil C cycling in a subalpine spruce forest on the Tibetan Plateau of China. Our results suggested that the lack of snow cover increased the intensity of soil frost, which decreased soil respiration in the winters but did not affect it during the subsequent growing seasons. Frost decreased microbial biomass and activities in the winters but not in the snow-free growing seasons. More intense soil frost did not affect the size distribution of soil aggregates or the fine-root biomass. Predicted soil frost due to winter climate change may thus decrease winter soil respiration by direct environmental effects (e.g. soil temperatures) and indirect biological processes (e.g. microbial biomass and activities) in the subalpine forests on the Tibetan Plateau. Intensified soil frost did not cause cross-seasonal effects on soil CO₂ efflux or biological activities in the subsequent growing seasons. Our observations underscore the ecological importance of seasonal snowpack and microbe-associated C processes in subalpine

forest soils. These findings improve our understanding of the response of soil C dynamics to winter climate change in this region experiencing large decreases in winter snowfall.

Acknowledgements

We are very grateful to Fan Yang, Liyan Zhuang, Yu Tan and Rui Cao for their great support with field and lab. We also thank Martin Schädler, Prof. Dr. Wanqin Yang and Pro. Dr. Fuzhong Wu for their helpful suggestions on the statistical analysis. We also thank the experimental platform—Long-term Research Station of Alpine Forest Ecosystems. This work was supported by the National Natural Science Foundation of China (31570601 and 31700542) and the National Key Research and Development Program of China (2016YFC0502505 and 2017YFC0505003). Kaijun Yang appreciates the financial support from the China Scholarship Council (CSC).

Competing interests

The authors declare no competing financial interests.

References

- Aanderud, Z.T., Jones, S.E., Schoolmaster Jr, D.R., Fierer, N., Lennon, J.T., 2013. Sensitivity of soil respiration and microbial communities to altered snowfall. *Soil Biology & Biochemistry* 57, 217–227.
- Anderson, J.P.E., 1982. Soil Respiration. In: Page, A.L., Miller, R.H., Keeney, D.R. (Eds.), *Methods of Soil Analysis, Part 2: Chemical and Microbiological Properties*. American Society of Agronomy-Soil Science Society of America, Madison, pp. 831-871.
- Allison, S.D., Jastrow, J.D., 2006. Activities of extracellular enzymes in physically isolated fractions of restored grassland soils. *Soil Biology & Biochemistry* 38, 3245–3256.
- Blume-Werry, G., Kreyling, J., Laudon, H., Milbau, A., 2016. Short-term climate change manipulation effects do not scale up to long-term legacies: effects of an absent snow cover on boreal forest plants. *Journal of Ecology* 104, 1638–1648.
- Bokhorst, S., Metcalfe, D.B., Wardle, D.A., 2013. Reduction in snow depth negatively affects decomposers but impact on decomposition rates is substrate dependent. *Soil Biology & Biochemistry* 62, 157–164.
- Brooks, P.D., Mcknight, D., Elder, K., 2004. Carbon limitation of soil respiration under winter snowpacks: potential feedbacks between growing season and winter carbon fluxes. *Global Change Biology* 11, 231–238.
- Chai, Y.J., Zeng, X.B., S, Z.E., Bai, L.Y., Su, S.M., Huang, T., 2014. Effects of freeze-thaw on aggregate stability and the organic carbon and nitrogen enrichment ratios in aggregate fractions. *Soil Use and Management* 30, 507–516.
- Chapin III, F.S., Fetcher, N., Kielland, K., Everett, K.R., Linkins, A.E., 1988. Productivity and nutrient cycling of Alaskan tundra: enhancement by flowing soil water. *Ecology* 69, 693–702.
- Chen, H., Zhu, Q.A., Peng, C.H., Wu, N., Wang Y.F., Fang, X.Q., Gao, Y.H., Zhu, D., Yang, G., Tian, J.Q., Kang, X., Piao, S.L., Ouyang, H., Xiang, W.H., Luo, Z.B., Jiang, H.,

519 Song, X.Z., Zhang, Y., Yu, G.R., Zhao, X.Q., Gong, P., Yao, T.D., Wu, J.H., 2013. The
520 impacts of climate change and human activities on biogeochemical cycles on the
521 Qinghai-Tibetan Plateau. *Global Change Biology* 19, 2940–2955.

522 Christenson, L.M., Mitchell, M.J., Groffman, P.M., Lovett, G.M., 2010. Winter climate
523 change implications for decomposition in northeastern forests: comparisons of sugar
524 maple litter with herbivore fecal inputs. *Global Change Biology* 16, 2589–2601.

525 Comerford, D.P., Schaberg, P.G., Templer, P.H., Soggi, A.M., Campbell, J.L., Wallin K.F.,
526 2013. Influence of experimental snow removal on root and canopy physiology of sugar
527 maple trees in a northern hardwood forest. *Oecologia* 171, 261–269.

528 Davidson, E.A., Janssens, I.A., 2006. Temperature sensitivity of soil carbon decomposition
529 and feedbacks to climate change. *Nature* 440, 165–173.

530 De Long, J.R., Laudon, H, Blume-Werry, G, Kardol, P. 2016. Nematode community resistant
531 to deep soil frost in boreal forest soils. *Pedobiologia* 59, 243–251.

532 Du, E., Zhou, Z., Li, P., Jiang, L., Hu, X., Fang, J., 2013. Winter soil respiration during
533 soil-freezing process in a boreal forest in Northeast China. *Journal of Plant Ecology* 6,
534 800 349–357.

535 Gavazov, K., Ingrisch, J., Hasibeder, R., Mills, R.T.E., Buttler, A., Gleixner, G., Pumpanen, J.,
536 Bahn, M., 2017. Winter ecology of a subalpine grassland: Effects of snow removal on
537 soil respiration, microbial structure and function. *Science of the Total Environment* 590,
538 316-324.

539 Groffman, P.M., Driscoll, C.T., Fahey, T.J., Hardy, J.P., Fitzhugh, R.D., Tierney, G.L., 2001a.
540 Colder soils in a warmer world: A snow manipulation study in a northern hardwood
541 forest ecosystem. *Biogeochemistry* 56, 135–150.

542 Groffman, P.M., Driscoll, C.T., Fahey, T.J., Hardy, J.P., Fitzhugh, R.D., Tierney, G.L., 2001b.
543 Effects of mild winter freezing on soil nitrogen and carbon dynamics in a northern
544 hardwood forest. *Biogeochemistry* 56, 191–213.

545 Groffman, P.M. , Hardy, J.P., Driscoll, C.T., Fahey, T.J., 2006. Snow depth, soil freezing, and

546 fluxes of carbon dioxide, nitrous oxide and methane in a northern hardwood forest.
547 *Global Change Biology* 12, 1748–1760.

548 Grogan, P., Michelsen, A., Ambus, P., Jonasson, S., 2004. Freeze-thaw regime effects on
549 carbon and nitrogen dynamics in sub-arctic heath tundra mesocosms. *Soil Biology &*
550 *Biochemistry* 36, 641–654.

551 Haei, M., Laudon, H., 2015. Carbon dynamics and changing winter conditions: a review of
552 current understanding and future research directions. *Biogeosciences Discussion* 12,
553 15763–15808.

554 Henry, H.A.L., 2007. Soil freeze-thaw cycle experiments: trends, methodological weaknesses
555 and suggested improvements. *Soil Biology & Biochemistry* 39, 977–986

556 He, W., Wu, F.Z., Zhang, D.J., Yang, W.Q., Tan, B., Zhao, Y.Y., Wu, Q.Q., 2015. The
557 effects of forest gaps on cellulose degradation in the foliar litter of two shrub species in
558 an alpine fir forest. *Plant and Soil* 393, 109–122.

559 IPCC (2013) *Climate Change 2013: The Physical Science Basis. Contribution of Working*
560 *Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*
561 *Change*. Cambridge University Press, Cambridge, UK and New York, NY.

562 Janssens, I.A., Pilegaard, K., 2003. Large seasonal changes in Q_{10} of soil respiration in a
563 beech forest. *Global Change Biology* 9, 911–918.

564 Jusselme, M.D., Saccone, P., Zinger, L., Faure, M., Le Roux X., Guillaumaud, N., Bernard, L.,
565 Clement, J. C., Poly, F., 2016. Variations in snow depth modify N-related soil microbial
566 abundances and functioning during winter in subalpine grassland. *Soil Biology &*
567 *Biochemistry* 92, 27–37.

568 Kristiansen, S.M., Schjønning, P., Thomsen, I.K., Olesen, J.E., Kristensen, K., Christensen,
569 B.T., 2006. Similarity of differently sized macro-aggregates in arable soils of different
570 texture. *Geoderma* 137, 147–154.

571 Lipson, D.A., Schadt, C.W., Schmidt, S.K., 2002. Changes in soil microbial community
572 structure and function in an alpine dry meadow following spring snow melt. *Microbial*

573 Ecology 43, 307–314.

574 Liu, J.P., Curry, J. A., Wang, H.J., Song, M.R., Horton, R. M., 2012. Impact of declining
575 Arctic sea ice on winter snowfall. *Proceedings of the National Academy of Sciences* 109,
576 6781–6783.

577 Li, W.B., Wu, J.B., Bai, E., Jin, C.J., Wang, A.Z., Yuan, F.H., Guan, D.X., 2016a. Response
578 of terrestrial carbon dynamics to snow cover change: A meta-analysis of experimental
579 manipulation (II). *Soil Biology & Biochemistry* 103, 388–393.

580 Li, Z.J., Yang, W.Q., Yue, K., Justine, M.F., He, R.Y., Yang, K.J., Zhuang, L.Y., Wu, F.Z., Tan,
581 B., Zhang, L., Xu, Z.F., 2017. Effects of snow absence on winter soil nitrogen dynamics in
582 a subalpine spruce forest of southwestern China. *Geoderma*, 307, 107–113.

583 Monson, R.K., Lipson, D.L., Burns, S.P., Turnipseed, A.A., Delany, A.C., Williams, M.W.,
584 Schmidt, S.K., 2006b. Winter forest soil respiration controlled by climate and microbial
585 community composition. *Nature* 439, 711–714.

586 Moorhead, D., Lashermes, G., Recous, S., Bertrand, I., 2014. Interacting microbe and litter
587 quality controls on litter decomposition: a modeling analysis. *PLOS ONE* 9: e108769.

588 Muhr, J., Borken, W., Matzner, E., 2009. Effects of soil frost on soil respiration and its
589 radiocarbon signature in a Norway spruce forest soil. *Global Change Biology* 15, 782–
590 793.

591 Ni, X.Y., Yang, W.Q., Li, H., Xu, L.Y., He, J., Tan, B., Wu, F.Z., 2014. The responses of
592 early foliar litter humification to reduced snow cover during winter in an alpine forest.
593 *Canadian Journal of Soil Science* 94, 453–461.

594 Oquist, M.G., Laudon, H., 2008. Winter soil frost conditions in boreal forests control growing
595 season soil CO₂ concentration and its atmospheric exchange. *Global Change Biology* 14,
596 2839–2847.

597 Oztas, T., Fayetorbay, F., 2003. Effect of freezing and thawing processes on soil aggregate
598 stability. *Catena* 52, 1–8.

599 Repo, T., Sirki, S., Heinonen, J., Lavigne, A., Roitto, M., Koljonen, E., Koljonen, E., Sutinen,

600 S., Finer, L., 2014. Effects of frozen soil on growth and longevity of fine roots of
 601 Norway spruce. *Forest Ecology and Management* 313, 112–122.

602 Rivkina, E.M., Friedmann, E.I., McKay, C.P., Gilichinsky, D.A., 2000. Metabolic activity of
 603 permafrost bacteria below the freezing point. *Applied and Environmental Microbiology*
 604 66, 3230–3233.

605 Saccone, P., Morin, S., Baptist, F., Bonneville, J.M., Colace, M.P., Domine, F., Faure, M.,
 606 Geremia, R., Lochet, J., Poly, F., Lavorel, S., Clement, J.C., 2013. The effects of
 607 snowpack properties and plant strategies on litter decomposition during winter in
 608 subalpine meadows. *Plant and Soil* 363, 215–229.

609 Sanders-DeMott, R., Templer, P.H., 2017. What about winter? Integrating the missing season
 610 into climate change experiments in seasonally snow covered ecosystems. *Methods in*
 611 *Ecology and Evolution* 8, 1183–1191.

612 Schimel, J.P., Bilbrough, C., Welker, J. A., 2004. Increased snow depth affects microbial
 613 activity and nitrogen mineralization in two Arctic tundra communities. *Soil Biology &*
 614 *Biochemistry* 36, 217–227.

615 Schindlbacher, A., Jandl, R., Schindlbacher, S., 2014. Natural variations in snow cover do not
 616 affect the annual soil CO₂ efflux from a mid-elevation temperate forest. *Global Change*
 617 *Biology* 20, 622–632.

618 Schindlbacher, A., Zechmeister-Boltenstern, S., Glatzel, G., Jandl, R., 2007. Winter soil
 619 respiration from an Austrian mountain forest. *Agricultural and Forest Meteorology* 146,
 620 205–215.

621 Schutt, M., Borken, W., Spott, O., Stange, C.F., Matzner, E., 2014. Temperature sensitivity of
 622 C and N mineralization in temperate forest soils at low temperatures. *Soil Biology &*
 623 *Biochemistry* 69, 320–327.

624 Sinsabaugh, R.L., Lauber, C.L., Weintraub, M.N., Ahmed, B., Allison, S.D., Crenshaw, C.,
 625 Contosta, A.R., Cusack, D., Frey, S., Gallo M.E., 2008. Stoichiometry of soil enzyme
 626 activity at global scale. *Ecology Letters* 11, 1252–1264.

627 Song, Y., Zou, Y.C., Wang, G.P., Yu, X.F., 2017. Altered soil carbon and nitrogen cycles due
628 to the freeze-thaw effect: A meta-analysis. *Soil Biology & Biochemistry* 109, 35–49.

629 Sorensen, P.O., Templer, P.H., Finzi, A.C., 2016. Contrasting effects of winter snowpack and
630 soil frost on growing season microbial biomass and enzyme activity in two
631 mixed-hardwood forests. *Biogeochemistry* 128, 141–154.

632 Steinweg, J.M., Fisk, M.C., McAlexander, B., Groffman, P.M., Hardy, J.P., 2008.
633 Experimental snowpack reduction alters organic matter and net N mineralization
634 potential of soil macroaggregates in a northern hardwood forest. *Biology and Fertility of*
635 *Soils* 45, 1–10

636 Sulkava, P., Huhta, V., 2003. Effects of hard frost and freeze–thaw cycles on decomposer
637 communities and N mineralisation in boreal forest soil. *Applied Soil Ecology* 22, 225–
638 239.

639 Tan, B., Wu, F.Z., Yang, W.Q., He, X.H., 2014. Snow removal alters soil microbial biomass
640 and enzyme activity in a Tibetan alpine forest. *Applied Soil Ecology* 76, 34–41.

641 Tierney, G.L., Fahey, T.J., Groffman, P.M., Hardy, J.P., Fitzhugh, R.D., Driscoll, C.T., 2001.
642 Soil freezing alters fine root dynamics in a northern hardwood forest. *Biogeochemistry*
643 56, 175–190.

644 Uchidaa, M., Mo, W., Nakatsubo, T., Tsuchiya, Y., Horikoshi, T., Koizumi, H., 2005.
645 Microbial activity and litter decomposition under snow cover in a cool-temperate
646 broad-leaved deciduous forest. *Agricultural and Forest Meteorology* 134, 102–109.

647 Van's Hoff, J.H., 1898. *Lectures on Theoretical and Physical Chemistry. Part 1: Chemical*
648 *Dynamics*. Edward Arnold, London, pp. 224–229.

649 Wang, A., Wu, F.Z., Yang, W.Q., Wu, Z.C., Wang, X.X., Tan, B., 2012. Abundance and
650 composition dynamics of ammonia-oxidizing archaea in alpine fir forest on the eastern
651 Tibetan Plateau of China. *Canadian Journal of Microbiology* 58, 572–580.

652 Wang, C., Han, Y., Chen, J., Wang, X., Zhang, Q., Bond-Lamberty, B., 2013. Seasonality of
653 soil CO₂ efflux in a temperate forest: Biophysical effects of snowpack and spring freeze–

929 thaw cycles. *Agricultural and Forest Meteorology* 177, 83–92.

Wang, J., Zhang, M.J., Wang, S.J., Ren, Z.G., Che, Y.J., Qiang, F., Qu, D.Y., 2016. Decrease in snowfall/rainfall ratio in the Tibetan Plateau from 1961 to 2013. *Journal of Geographphy Sciences* 26, 1277–1288.

Wang, W., Peng, S.S., Wang, T., Fang, J.Y., 2010. Winter soil CO₂ efflux and its contribution to annual soil respiration in different ecosystems of a forest-steppe ecotone, north China. *Soil Biology & Biochemistry* 42, 451–458.

White, D.C., Stair, J.O., Ringelberg, D.B., 1996. Quantitative comparisons of in situ microbial biodiversity by signature biomarker analysis. *Journal of Industrial Microbiology* 17, 185–196.

Wubs, E.R.J., Woodin, S.J., Stutter, M.I., Wipf, S., Sommerkorn, M., Wal, R.V.D., 2018. Two decades of altered snow cover does not affect soil microbial ability to catabolize carbon compounds in an oceanic alpine heath. *Soil Biology & Biochemistry* 124, 101–104.

Xiong, L., Xu, Z.F., Yang, W.Q., Yin, R., Tang, S.S., Wang, B., Xu, L.Y., Chang, C.H., 2015. Aboveground litter contribution to soil respiration in a subalpine dragon spruce plantation of western Siehuan. *Acta Ecologica Sinica*, 35, 4678–4686, in Chinese with an English abstract.

Xu, M., Qi, Y., 2001. Spatial and seasonal variations of Q_{10} determined by soil respiration measurements at a Sierra Nevadan forest. *Global Biogeochemistry Cycles* 15, 687–696.

Xu, W.F., Ma, L.J., Ma, M.N., Zhang, H.C., Yuan, W.P., 2017. Spatial–temporal variability of snow cover and depth in the Qinghai-Tibetan Plateau. *Journal of Climate* 30, 1521–1533.

Xu, Z.F., Liu, Q., Yin, H.J., 2014. Effects of temperature on soil net nitrogen mineralisation in two contrasting forests on the eastern Tibetan Plateau, China. *Soil Research* 52, 562–567.

Xu, Z.F., Pu, X.Z., Yin, H.J., Zhao, C.Z., Liu, Q., Wu, F.Z., 2012. Warming effects on the early decomposition of three litter types, Eastern Tibetan Plateau, China. *European Journal of Soil Science* 63, 360–367.

- Xu, Z.F., Wan, C.A., Xiong, P., Tang, Z., Hu, R., Cao, G., Liu, Q., 2010. Initial responses of soil CO₂ efflux and C, N pools to experimental warming in two contrasting forest ecosystems, Eastern Tibetan Plateau, China. *Plant and Soil* 336, 183–195.
- Xu, Z.F., Zhao, C.Z., Yin, H.J., Liu, Q., 2015. Warming and forest management interactively affect the decomposition of subalpine forests on the eastern Tibetan Plateau: A four-year experiment. *Geoderma* 239, 223–228.
- Xu, Z.F., Zhu, J.X., Wu, F.Z., Liu, Y., Tan, B., Yang, W.Q., 2016. Effects of litter quality and climate change along an elevational gradient on litter decomposition of subalpine forests, Eastern Tibetan Plateau, China. *Journal of Forestry Research* 27, 505–511.
- Yin, H.J., Li, Y.F., Xiao, J., Xu, Z.F., Cheng, X.Y., Liu, Q., 2013. Enhanced root exudation stimulates soil nitrogen transformations in a subalpine coniferous forest under experimental warming. *Global Change Biology* 19, 2158–2167.
- Zhang, Y.D., Gu, F.X., Liu, S.R., Liu, Y.C., Li, C., 2013. Variations of carbon stock with forest types in subalpine region of southwestern China. *Forest Ecology and Management* 300, 88–95.
- Zhao, J.B., Peichl, M., Nilsson, M.B., 2017. Long-term enhanced winter soil frost alters growing season CO₂ fluxes through its impact on vegetation development in a boreal peatland. *Global Change Biology* 23, 3139–3153.

Table 1 Mean soil CO₂ efflux ($\mu\text{mol m}^{-2} \text{s}^{-1}$, means \pm SEs) during the winter and growing season.

Year	Period	Treatment	Control
2016	Winter	0.47 ± 0.08	0.55 ± 0.08
	Growing season	2.07 ± 0.28	2.19 ± 0.29
2017	Winter	0.43 ± 0.06	0.53 ± 0.07
	Growing season	2.23 ± 0.26	2.25 ± 0.30

Table 2. Relative distribution of the aggregate size classes (% , means \pm SEs) in the snow-exclusion treatment and control plots in the early thawing period of 2016 and 2017.

Year	Size	Treatment	Control
2016	>2 mm	34.9 ± 5.7	32.9 ± 6.9
	0.25-2 mm	59.9 ± 4.7	58.6 ± 4.2
	<0.25 mm	5.2 ± 2.0	8.5 ± 2.8
2017	>2 mm	35.3 ± 6.4	36.7 ± 4.5
	0.25-2 mm	53.9 ± 3.8	50.7 ± 2.2
	<0.25 mm	10.8 ± 3.6	12.6 ± 2.8

712

713 **Table 3.** Content of live and dead fine roots (g m⁻², means \pm SEs) up to a depth of 15 cm in
 714 the snow-exclusion treatment and control plots in the early thawing period of 2016 and 2017.

Year	Fine root pool	Treatment	Control
2016	Live	243.7 \pm 45.6	251.7 \pm 35.6
	Dead	92.8 \pm 14.4	79.4 \pm 26.3
2017	Live	221.2 \pm 38.5	215.6 \pm 28.9
	Dead	94.3 \pm 22.6	88.1 \pm 16.4

715

716

717

718

719

720

721

722

723

Figure legends

Figure 1. Air temperature, soil temperature and snow depth (a) and soil moisture (b) in the snow-exclusion treatment and control plots during the experimental period.

Figure 2. Soil CO₂ efflux (means \pm SEs) in the snow-exclusion treatment and control plots during the experimental period. Significant differences between the control and treatment on a given date are indicated by asterisks ($P < 0.05$).

Figure 3. Exponential relationships between soil CO₂ efflux and soil temperature (means \pm SEs) in the treatment and control plots for winter, growing season and entire year, respectively.


Figure 4. Total soil phospholipid fatty acids (means \pm SEs) in the snow-exclusion treatment and control plots. Significant differences between the control and treatment on a given date are indicated by asterisks ($P < 0.05$). FP, frost period; ETP, early thawing period; MGS, middle of the growing season.

Figure 5. Soil microbial respiration (means \pm SEs) in the snow-exclusion treatment and control plots. Significant differences between the control and treatment on a given date are indicated by asterisks ($P < 0.05$). FP, frost period; ETP, early thawing period; MGS, middle of the growing season.

Figure 6. Activities of (a) β -glucosidase, (b) polyphenol oxidase, (c) peroxidase and (d) β -N-acetyl-glucosaminidase (means \pm SEs) in the snow-exclusion treatment and control plots. Significant differences between the control and treatment on a given date are indicated by

asterisks ($P < 0.05$). FP, frost period; ETP, early thawing period; MGS, middle of the growing season.

Figure 7. Soil ammonium (a) and nitrate (b) concentrations (means \pm SEs) in the snow-exclusion treatment and control plots. Significant differences between the control and treatment on a given date are indicated by asterisks ($P < 0.05$). FP, frost period; ETP, early thawing period; MGS, middle of the growing season.




763

764

765

Figure 2



783 **Figure 3**






Figure 4




Figure 5




Figure 6




Figure 7

