

# Exposure to air pollution and noise is associated with cognition and brain structure in older adults

Presented During: Poster Session

Tuesday, June 11, 2019: 12:45 PM - 02:45 PM

## Poster No:

T377

## Submission Type:

Abstract Submission

## Authors:

Christiane Jockwitz<sup>1,2,3</sup>, René Nussbaum<sup>1,4,5</sup>, Sarah Lucht<sup>5,6</sup>, Susanne Moebus<sup>7</sup>, Barbara Hoffmann<sup>5</sup>, Svenja Caspers<sup>1,3,4</sup>

## Institutions:

<sup>1</sup>Institute of Neuroscience and Medicine (INM- 1), Research Center Juelich, Juelich, Germany, Juelich, Germany, <sup>2</sup>Department of Psychiatry, Psychotherapy and Psychosomatics, RWTH Aachen University, Medical Faculty, Aachen, Germany, <sup>3</sup>JARA-BRAIN, Juelich-Aachen Research Alliance, Juelich, Germany, <sup>4</sup>Institute of Anatomy I, Medical Faculty, Heinrich Heine University Duesseldorf, Duesseldorf, Germany, <sup>5</sup>Environmental Epidemiology Group, Institute of Occupational, Social and Environmental Medicine, Centre for Health and Society, Medical Faculty, Heinrich Heine University Duesseldorf, Germany, <sup>6</sup>Institute for Medical Statistics, Medical Faculty, Heinrich Heine University Düsseldorf, Duesseldorf, Germany, <sup>7</sup>Institute of Medical Informatics, Biometry and Epidemiology, University of Duisburg-Essen, Essen, Germany

## Introduction:

Aging is associated with decreases in brain structure and cognitive performance [1,2]. However, during older ages, high inter-individual variability has been reported, with age explaining only minor parts of this variance [3]. Thus, other factors come into focus, such as environmental exposures to air pollution (AP) and noise. Previous studies have shown adverse associations of long-term AP and noise exposure with global cognitive abilities in older adults [4], implying a relation between the two exposures, cognitive abilities and brain structure. Therefore, the current study aimed at identifying exposure-related differences of AP and noise on cognitive abilities and structural atrophy in cognitive brain networks in older adults.

## Methods:

The 615 older adults who were included in the current analysis (mean age:  $61.5 \pm 6.7$ , 271 f), were recruited as part of the population-based 1000BRAINS study [5], an extension of the preceding Heinz

Nixdorf Recall study [6]. Residential long-term exposures to AP included particulate matter (PM<sub>10</sub>, PM<sub>2.5</sub>, PM<sub>2.5abs</sub>), accumulation mode particle number (PNAM), nitrogen oxides (NO<sub>x</sub>, NO<sub>2</sub>) and distance to the nearest major road. Residential exposure to noise was modeled for weighted 24h (LDEN) and nighttime (Lnight) noise (for details about AP/ noise modelling, see [4]). Linear regressions were used to discover relations between AP/noise exposure and cognitive performance in five cognitive domains (Attention, Executive Function, Episodic Memory, Working Memory, Language). Based on the resulting associations, cognitive brain networks were extracted from functional resting state images using independent component analysis (voxel size: 2.4mm<sup>3</sup>; FSL [7]). Structural brain atrophy was measured using the local gyrification index (LGI; FreeSurfer [8]) on T1 weighted structural brain images (available n=590; 3 Tesla MR-Scanner, voxel size: 1mm<sup>3</sup>). Afterwards, AP/noise exposure was related to LGI values within the extracted networks (covariates: age, sex, education, neighborhood unemployment, alcohol consumption, smoking, physical activity; results were significant at p < .05).

### Results:

We found negative associations between AP (PM<sub>2.5</sub>, PM<sub>2.5abs</sub>, NO<sub>x</sub>, PNAM) and language, as well as noise (LDEN, Lnight) and working memory performance. Based on these results, LGI values within the frontoparietal network were extracted (bilateral: dorsolateral prefrontal cortex [DLPFC], posterior cingulate cortex [PCC]/precuneus and inferior parietal lobule [IPL]). AP (PM<sub>10</sub>, PM<sub>2.5</sub>, NO<sub>2</sub>, and NO<sub>x</sub>) was negatively associated with LGI values in the right PCC/precuneus and IPL. Distance to major roads showed negative associations with LGI values in right DLPFC and PCC/precuneus. Further, PNAM showed negative associations with the right DLPFC. In contrast, noise (LDEN, Lnight) was positively associated with LGI values in the right DLPFC.

### Conclusions:

The current study investigated associations between environmental exposure to AP/ noise and cognitive abilities and structural atrophy in the frontoparietal network. Long-term exposure to AP was adversely associated with language functions, suggesting an accelerated cognitive decline in older adults who are more strongly exposed to AP. These associations were accompanied by a more pronounced right hemispheric structural brain atrophy, especially in posterior parts of the frontoparietal network, thereby providing structural correlates for two major functionally based aging theories, i.e. posterior to anterior shift in aging [9] and the right hemi aging theory [10]. Notably, noise exposure, modeled at the outdoor façade, was associated with a preservation of brain structure, which could be explained by protective behaviors in those exposed to more noise, e.g. window closing, and needs further investigation. Our results suggest that exposure to AP may be one important factor that contributes to the high inter-individual variability of brain structure and cognitive decline in older adults.

### Imaging Methods:

Anatomical MRI <sup>2</sup>

### Lifespan Development:

Aging <sup>1</sup>

### Keywords:

Aging  
Cognition

<sup>1|2</sup>Indicates the priority used for review

**My abstract is being submitted as a Software Demonstration.**

No

**Please indicate below if your study was a "resting state" or "task-activation" study.**

Resting state

Other

**Healthy subjects only or patients (note that patient studies may also involve healthy subjects):**

Healthy subjects

**Was any human subjects research approved by the relevant Institutional Review Board or ethics panel? NOTE: Any human subjects studies without IRB approval will be automatically rejected.**

Yes

**Was any animal research approved by the relevant IACUC or other animal research panel? NOTE: Any animal studies without IACUC approval will be automatically rejected.**

Not applicable

**Please indicate which methods were used in your research:**

Functional MRI

Structural MRI

Neuropsychological testing

**For human MRI, what field strength scanner do you use?**

3.0T

**Which processing packages did you use for your study?**

FSL

Free Surfer

**Provide references using author date format**

[1] Hedden, & Gabrieli (2004). Insights into the ageing mind: a view from cognitive neuroscience.

Nat Rev Neurosci, 5(2): p. 87-96.

[2] Sowell, et al. (2003). Mapping cortical change across the human life span. Nature Neuroscience, 6(3): p. 309-15.

[3] Jockwitz, et al. (2017). Age- and function-related regional changes in cortical folding of the default mode network in older adults. Brain Struct Funct, 222(1): p. 83-99.

- [4] Tzivian, et al. (2016). Long-term air pollution and traffic noise exposures and mild cognitive impairment in older adults: a cross-sectional analysis of the Heinz Nixdorf recall study. *Environmental health perspectives*, 124(9), 1361.
- [5] Caspers, et al. (2014). Studying variability in human brain aging in a population-based German cohort-rationale and design of 1000BRAINS. *Front Aging Neurosci*, 6: p. 149.
- [6] Schmermund, et al. (2002). Assessment of clinically silent atherosclerotic disease and established and novel risk factors for predicting myocardial infarction and cardiac death in healthy middle-aged subjects: rationale and design of the Heinz Nixdorf RECALL Study. *Risk Factors, Evaluation of Coronary Calcium and Lifestyle. Am. Heart J.* 144, 212–218.
- [7] Beckmann, et al. (2005). Investigations into resting-state connectivity using independent component analysis. *Philos. Trans. R. Soc. B Biol. Sci.* 360, 1001–1013.
- [8] Schaer, et al. (2008). A surface-based approach to quantify local cortical gyrification. *Medical Imaging, IEEE Transactions on*, 27(2), 161-170.
- [9] Davis, et al. (2007). Que PASA? The posterior–anterior shift in aging. *Cerebral cortex*, 18.
- [10] Dolcos, et al. (2002). "Hemispheric asymmetry and aging: right hemisphere decline or asymmetry reduction." *Neuroscience & Biobehavioral Reviews*, 26.7: 819-825.