

Anatomy and Function of Four New Cytoarchitectonic Areas in the Human Lateral Orbitofrontal Cortex

Magdalena Wojtasik¹, Sebastian Bludau², Simon B. Eickhoff^{3,4}, Hartmut Mohlberg², Svenja Caspers^{2,5}, Katrin Amunts^{1, 2}

¹C. & O. Vogt – Institute for Brain Research, Heinrich-Heine-University Hospital Düsseldorf, Germany, ²Institute of Neuroscience and Medicine 1 (INM-1), Research Center Jülich, Germany, ³Institute of Neuroscience and Medicine 7 (INM-7), Research Center Jülich, Germany, ⁴Institute for Systems Neuroscience, Heinrich-Heine-University Düsseldorf, Germany, ⁵Institute of Anatomy I, University Hospital Düsseldorf, Germany

BACKGROUND

The lateral orbitofrontal cortex (IOFC) has been cytoarchitecturally mapped by different authors, resulting in a different number of areas and nomenclature (e.g., BA47¹, Area *F*² or Area 47/12³). Most of the maps are two-dimensional, and therefore cannot be directly compared with neuroimaging data.



Functional neuroimaging and co-activational analysis revealed IOFC involvement in memory and reward processing⁴. It receives neuronal input from the sensory systems, the amygdala and the hypothalamus⁵.

The present study was designed:

- to investigate the cytoarchitecture of the IOFC using image analysis and a statistically reproducible approach for detecting cytoarchitectonic borders in a sample of ten *post mortem* brains,
- to generate maps which reflect the anatomical intersubject variability in a common reference space,
- to provide cytoarchitectonic probabilistic maps in stereotaxic space as a prerequisite for comparison with *in vivo* findings,
- to use meta-analysis as an approach to identify functional correlates of the identified areas.


METHODS

- Microstructural analysis of 10 human *post mortem* brains⁶
- Quantitative and statistically reproducible detection of cytoarchitectonic borders in 2D⁷
- Analysis of cytoarchitectonic similarity based on a hierarchical cluster analysis
- Macroanatomical analysis of sulci and gyri
- Calculation of cytoarchitectonic probability maps in the MNI reference space Colin27 and maximum probability maps⁸
- Analysis of co-activational patterns of every IOFC area with meta-analytic connectivity modelling (MACM)⁹
- Functional decoding of all detected IOFC areas with the BrainMap database^{10,11}




RESULTS

Cytoarchitecture of four new areas in the IOFC




Cytoarchitecture of four identified areas in the lateral OFC. Roman letters indicate respective layers and sublayers. I = molecular layer, II = external granular layer, III = external pyramidal layer (with sublayers a-c), IV = internal granular layer, V = internal pyramidal layer (with sublayers a & b), VI = multimodal layer. Scale bar = 500 µm.

Hierarchical cluster analysis



High structural similarity between Fo5 & Fo7, also Fo4 & Fo6 revealing lateral clustering of the four new IOFC areas, based on higher granularity in Fo5 & Fo7 as well as a more striate appearance in Fo4 & Fo6. Frontal pole area Fp1, medial OFC area Fo3 and Broca area 45 show an increasing structural dissimilarity compared to the IOFC areas.

Section sequence of identified areas




Extent of the areas in serial brain sections of an individual brain. Fo4 follows Fo5 on the anterior orbital gyrus. Fo4 has a larger volume than Fo5. Fo5 starts on the most rostral part of the inferior frontal gyrus, at the frontomarginal sulcus, entering the anterior orbital gyrus only shortly. Fo6 occupies the lateral orbital gyrus, separating Fo7 and from adjoining areas. Fo7 lies buried in the lateral orbital sulcus between Fo4 and Fo6, terminating in the transverse orbital sulcus.


AOG = anterior orbital gyrus, FMS = frontomarginal sulcus, IOS = intermediate orbital sulcus, LOG = lateral orbital gyrus, LOS = lateral orbital sulcus, MOG = medial orbital gyrus, MOS = medial orbital sulcus, POG = posterior orbital gyrus, TOS = transverse orbital sulcus.

Probability maps in stereotaxic MNI space


Continuous probability maps of areas Fo4, Fo5, Fo6 & Fo7 revealing their location and extent in the stereotaxic MNI reference brain 'Colin27', ranging from high (red = 100% = 10 out of 10 brains) to low (blue = 10% = 1 out of 10 brains) probable appearance in ten analyzed brains according to the interindividual variability of area extents.



Interindividual variability of sulcal patterns in the OFC



Maximum probability map & Functional decoding



Right areas	Function
Fo4	Reward processing
	Working memory
Fo5	Explicit memory
	Perception of physical pain
Fo6	Disgust, Anger
	Gustational perception
	Perception of physical pain
	Action inhibition
Fo7	Emotional processing
	Gustational perception

Left areas	Function
Fo4	Gustational perception
	Working memory
Fo5	Reward processing
	Language processing, Phonology
	Working & Explicit memory
Fo6	Language processing, Semantics
	Gustational perception
Fo7	Olfactory perception
	Language processing, Orthography, Semantics
	Memory

Maximum probability map of identified areas Fo4, Fo5, Fo6 & Fo7 superimposed onto the MNI reference brain 'Colin27'. Threshold = 0.4. Areas were smoothed prior to final image creation. Additional functional decoding of every IOFC area in both hemispheres was calculated using the BrainMap database^{10,11}.

CONCLUSIONS

- Microanatomy:** Identification of our new cortical areas Fo4, Fo5, Fo6 & Fo7 based on differences in the cytoarchitecture. Partial overlap with a region previously being described as BA47 by Brodmann¹.
- Predominantly high cell density in the deep parts of layer III, weak layer IV and separation of layer V into two sublayers V_a and V_b according to cytoarchitectonic analysis.
- Clustering of areas Fo4 + Fo6 versus Fo5 + Fo7 based on structural differences in granularity, lamination and greater pyramidal cells in layer IIIC in areas Fo5 + Fo7.
- Macroanatomy:** Detection of three different sulcal patterns, with diversely distributed medial, lateral and transverse orbital sulci and therefore changing arrangements of the four new areas.
- Mapping:** Calculation of cytoarchitectonic probabilistic maps and the Maximum Probability Map in the MNI reference brain 'Colin27' as well as MNI152.
- The maps are publicly available in the atlas of the Human Brain Project and the JuBrain Cytoarchitectonic atlas (QR codes).
- Meta-Analysis:** Detection of co-activational clustering according to MACM analysis, with areas of the inferior parietal lobule, the frontal pole and medial orbital cortex as well as Broca areas 44 and 45, centromedial and laterobasal amygdaloid nuclei and hippocampal area CA1.
- All areas are involved in cognitive processing and emotional evaluation of sensory input. Areas Fo5, Fo6 & Fo7 participate in language processing in the left hemisphere.

REFERENCES & ACKNOWLEDGEMENTS

- Brodmann, K. (1909). Vergleichende Lokalisationslehre der Großhirnrinde. Verlag von Johann Ambrosius Barth, Leipzig.
- Triarhou, L.C. (2007). A Proposed Number System for the 107 Cortical Areas of Economo and Koskinas, and Brodmann Area Correlations. *Stereotactic and Functional Neurosurgery* 85, 204–215.
- Öngür, D., Ferry, A.T., and Price, J.L. (2003). Architectonic subdivision of the human orbital and medial prefrontal cortex. *The Journal of Comparative Neurology* 460, 425–449.
- Rolls, E.T. (2000). The Orbitofrontal Cortex and Reward. *Cerebral Cortex* 10, 284–294.
- Rolls, E.T. (2004). The functions of the orbitofrontal cortex. *Brain and Cognition* 55, 11–29.
- Amunts, K., Schleicher, A., Bürgel, U., Mohlberg, H., Uylings, H.B.M., and Zilles, K. (1999). Broca's region revisited: Cytoarchitecture and intersubject variability. *The Journal of Comparative Neurology* 412, 319–341.
- Schleicher, A., Amunts, K., Geyer, S., Morosan, P., and Zilles, K. (1999). Observer-Independent Method for Microstructural Parcellation of Cerebral Cortex: A Quantitative Approach to Cytoarchitectonics. *NeuroImage* 9, 165–177.
- Amunts, K., and Zilles, K. (2015). Architectonic Mapping of the Human Brain beyond Brodmann. *Neuron* 88, 1086–1107.
- Eickhoff, S.B., Laird, A.R., Grefkes, C., Wang, L.E., Zilles, K., and Fox, P.T. (2009). Coordinate-based activation likelihood estimation meta-analysis of neuroimaging data: A random-effects approach based on empirical estimates of spatial uncertainty. *Human Brain Mapping* 30, 2907–2926.
- Laird, A.R. (2009). ALE meta-analysis workflows via the BrainMap database: Progress towards a probabilistic functional brain atlas. *Frontiers in Neuroinformatics* 3.
- Eickhoff, S.B., Paus, T., Caspers, S., Grosbras, M.-H., Evans, A.C., Zilles, K., and Amunts, K. (2007). Assignment of functional activations to probabilistic cytoarchitectonic areas revisited. *NeuroImage* 36, 511–521.
- Caspers, S., Eickhoff, S.B., Zilles, K., and Amunts, K. (2013). Microstructural grey matter parcellation and its relevance for connectome analyses. *NeuroImage* 80, 16–26.
- Eickhoff, S.B., Stephan, K.E., Mohlberg, H., Grefkes, C., Fink, G.R., Amunts, K., and Zilles, K. (2005). A new SPM toolbox for combining probabilistic cytoarchitectonic maps and functional imaging data. *NeuroImage* 25, 1325–1335.
- Ono, M., Kubik, S., Abernathy, C.D. (1990). *Atlas of the Cerebral Sulci*. Thieme Medical Publishers. New York.
- Chiavaras, M.M., Petrides M. (2000). Orbitofrontal sulci of the human and macaque monkey brain. *J Comp Neurol* 422, 35–54.

This project has received funding from the European Union's Horizon 2020 Research and Innovation Programme under Grant Agreement No. 785907 (HBP SGA2).

