

Towards a Storage Ring Electric Dipole Moment Measurement

J. Pretz

RWTH Aachen & FZ Jülich
on behalf of the JEDI & CPEDM collaboration



Aachen, DPG meeting, March 2019

Outline

- Introduction & Motivation

What are EDMs?, What do we know about EDMs?,
Why are EDMs interesting?


- Experimental Methods

How to measure **charged** particle EDMs?

- Strategy towards a storage ring EDM measurement

Introduction & Motivation

Electric Dipole Moments (EDM)




- permanent separation of positive and negative charge
- fundamental property of particles (like magnetic moment, mass, charge)
- existence of EDM only possible via violation of time reversal $\mathcal{T} \stackrel{\mathcal{CP}\mathcal{T}}{=} \mathcal{CP}$ and parity $\mathcal{P}$ symmetry
- close connection to “matter-antimatter” asymmetry
- **axion** field leads to oscillating EDM


$\mathcal{CP}$ –Violation & connection to EDMs

Standard Model	
Weak interaction CKM matrix	→ unobservably small EDMs
Strong interaction θ_{QCD}	→ best limit from neutron EDM
beyond Standard Model	
e.g. SUSY	→ accessible by EDM measurements


EDM in SM and SUSY




EDM in SM and SUSY



SUSY



EDM: Current Upper Limits



storage rings: EDMs of **charged** hadrons: $p, d, {}^3\text{He}$


Experimental Method

Experimental Method: Generic Idea

For **all** EDM experiments (neutron, proton, atoms, ...):

Interaction of $\vec{d}$ with electric field $\vec{E}$

For charged particles: apply electric field in a storage ring:



$$\frac{d\vec{S}}{dt} \propto \vec{d} \times \vec{S}$$

In general:

$$\frac{d\vec{S}}{dt} = \vec{\Omega} \times \vec{S}$$


build-up of vertical polarization $s_{\perp} \propto \vec{d}$, if $\vec{s}_{horz} \parallel \vec{p}$ (**frozen spin**)

Experimental Method: Generic Idea

For **all** EDM experiments (neutron, proton, atoms, ...):

Interaction of $\vec{d}$ with electric field $\vec{E}$

For charged particles: apply electric field in a storage ring:



$$\frac{d\vec{S}}{dt} \propto \textcolor{red}{d}\vec{E} \times \vec{S}$$

In general:

$$\frac{d\vec{s}}{dt} = \vec{\Omega} \times \vec{s}$$

build-up of vertical polarization $s_{\perp} \propto \mathbf{d}$, if $\vec{s}_{\text{horz}} \parallel \vec{p}$ (frozen spin)

Spin Precession: Thomas-BMT Equation

$$\frac{d\vec{s}}{dt} = \vec{\Omega} \times \vec{s} = \frac{-q}{m} \left[\underbrace{G\vec{B} + \left(G - \frac{1}{\gamma^2 - 1}\right) \vec{v} \times \vec{E}}_{= \vec{\Omega}_{\text{MDM}}} + \underbrace{\frac{\eta}{2}(\vec{E} + \vec{v} \times \vec{B})}_{= \vec{\Omega}_{\text{EDM}}} \right] \times \vec{s}$$

electric dipole moment (EDM): $\vec{d} = \eta \frac{q\hbar}{2mc} \vec{s}$,

magnetic dipole moment (MDM): $\vec{\mu} = 2(G + 1) \frac{q\hbar}{2m} \vec{s}$


Note: $\eta = 2 \cdot 10^{-15}$ for $d = 10^{-29}$ ecm, $G \approx 1.79$ for protons

Spin Precession: Thomas-BMT Equation



$$\frac{d\vec{s}}{dt} = \vec{\Omega} \times \vec{s} = \frac{-q}{m} \left[\underbrace{G\vec{B} + \left(G - \frac{1}{\gamma^2 - 1}\right) \vec{v} \times \vec{E}}_{\vec{\Omega}_{\text{MDM}} = 0, \text{ frozen spin}} + \frac{\eta}{2}(\vec{E} + \vec{v} \times \vec{B}) \right] \times \vec{s}$$

achievable with pure electric field if $G = \frac{1}{\gamma^2 - 1}$, works only for $G > 0$, e.g. proton
or with special combination of E , B fields and γ , i.e. momentum

Momentum and ring radius for proton in frozen spin condition



Momentum and ring radius for proton in frozen spin condition



Different Options

		
1.) pure electric ring	no $\vec{B}$ field needed, $\odot, \ominus$ beams simultaneously	works only for particles with $G > 0$ (e.g. p)
2.) combined ring	works for $p, d, {}^3\text{He}$, smaller ring radius	both $\vec{E}$ and $\vec{B}$ B field reversal for $\odot, \ominus$ required
3.) pure magnetic ring	existing (upgraded) COSY ring can be used, shorter time scale	lower sensitivity, precession due to G , i.e. no frozen spin

Statistical Sensitivity

beam intensity	$N = 4 \cdot 10^{10}$ per fill
polarization	$P = 0.8$
spin coherence time	$\tau = 1000$ s
electric fields	$E = 8$ MV/m
polarimeter analyzing power	$A = 0.6$
polarimeter efficiency	$f = 0.005$

$$\sigma_{\text{stat}} \approx \frac{2\hbar}{\sqrt{Nf\tau PAE}} \Rightarrow \sigma_{\text{stat}}(1\text{year}) = 2.4 \cdot 10^{-29} \text{ e}\cdot\text{cm}$$

challenge: get σ_{sys} to the same level

Systematic Sensitivity

observable: $\Omega_{\text{EDM}} = \frac{dE}{s\hbar} = 2.4 \cdot 10^{-9} \text{ s}^{-1}$ for $d = 10^{-29} \text{ e cm}$

- radial B -field of $B_r = 10^{-17} \text{ T}$:

$$\Omega_{B_r} = \frac{eGB_r}{m} = 1.7 \cdot 10^{-9} \text{ s}^{-1}$$

- geometric Phases (non-commutation of rotations), $B_{\text{long}}, B_{\text{vert}} \approx 1 \text{ nT}$

$$\Omega_{\text{GP}} = \left(\frac{eGB}{16m} \right)^2 \frac{1}{f_{\text{rev}}} \approx 3.7 \cdot 10^{-9} \text{ s}^{-1}$$

- ...


Remedy:

$$\odot: \Omega_{\text{CW}} = \Omega_{\text{EDM}} + \Omega_{\text{GP}} + \Omega_{B_r},$$


$$\oslash: \Omega_{\text{CCW}} = \Omega_{\text{EDM}} - \Omega_{\text{GP}} + \Omega_{B_r}.$$

Ω_{GP} drops out in sum, $\Omega_{\text{CW}} + \Omega_{\text{CCW}}$, effect of B_r can be subtracted by observing displacement of the two beams.

Systematics



Systematics



Systematics

Gravity:

$$\Omega_{\text{grav}} = \frac{2\gamma + 1}{\gamma + 1} \frac{\beta g}{c} = 3 \cdot 10^{-8} \text{ s}^{-1}$$

$$g = 9.81 \text{ m/s}^2$$

second effect: vertical electric (E_V) and radial magnetic (B_r) field needed to counteract force due to gravity $\left(F_{\text{grav}} = \frac{2\gamma^2 - 1}{\gamma} mg \right)$

Conclusion:

Statistically one can reach sensitivity of $\approx 10^{-29} \text{ e cm}$, many systematic effects can be controlled using $\odot$ and $\ominus$ beams, needs further investigation

→ **staged approach**

Towards a storage ring EDM measurement

Staged approach

Stage 1

precursor experiment
at COSY (FZ Jülich)




- magnetic storage ring

now

Stage 2

prototype ring




- initially electrostatic storage ring
- simultaneous $\odot$ and $\ominus$ beams

5 years

Stage 3


dedicated storage ring



- magic momentum
(701 MeV/c)



10 years

$$\sigma_{EDM}/(e \cdot cm)$$



Stage 1: Precursor Experiment

- Ongoing at COSY/ Forschungszentrum Jülich
- Achievements:
 - Long Spin Coherence time > 1000 s ✓




- measurement and manipulation and polarisation vector ✓
- First deuteron EDM measurement underway $\rightarrow$ V. Shmakova



Step 2: Prototype Ring

- operate electrostatic ring
- store $10^9 - 10^{10}$ particles for 1000 s
- simultaneous $\odot$ and $\ominus$ beams
- frozen spin (only possible with additional magnetic bending)
- develop and benchmark simulation tools
- develop key technologies:
beam cooling, deflector, beam position monitors, shielding . . .
- perform EDM measurement

Ring Lattice & Bending Element




Ring Lattice & Bending Element



→ A. Saleev

Step 3: Dedicated Ring

- pure electric ring:
frozen spin ($p = 701 \text{ MeV}/c$ $E_{kin}=233 \text{ MeV}$):



Summary

- EDMs are unique probe to search for new CP-violating interactions (and contribute to axion searches)
- **charged** particle EDMs can be measured in storage rings
- staged approach:
precursor at COSY → prototype (100 m) → dedicated ring (500 m)

Document submitted to ESPP in Dec. 2018 (arXiv:1812.08535, CERN yellow report CERN-PBC-REPORT-2019-002 in preparation)



Spare


$\mathcal{T}$ and $\mathcal{P}$ violation of EDM

$\vec{d}$: EDM

$\vec{\mu}$: magnetic moment (MDM)

both $\parallel$ to spin $\vec{s}$

	$H = -\mu \frac{\vec{s}}{s} \cdot \vec{B} - d \frac{\vec{s}}{s} \cdot \vec{E}$
$\mathcal{T}$:	$H = -\mu \frac{\vec{s}}{s} \cdot \vec{B} + d \frac{\vec{s}}{s} \cdot \vec{E}$
$\mathcal{P}$:	$H = -\mu \frac{\vec{s}}{s} \cdot \vec{B} + d \frac{\vec{s}}{s} \cdot \vec{E}$



$\Rightarrow$ EDM measurement tests violation of fundamental symmetries $\mathcal{P}$ and $\mathcal{T}(\overset{\mathcal{CP}\mathcal{T}}{=}\mathcal{CP})$

EDM activities around the world

Neutrons: (~ 200 ppl.)

- Beam EDM @ Bern
- LANL nEDM @ LANL
- nEDM @ PSI
- nEDM @ SNS
- PanEDM @ ILL
- PNPI/FTI/ILL @ ILL
- TUCAN @ TRIUMF

Storage rings: (~ 400 ppl.)


- CPEDM/JEDI
- muEDM @ PSI
- g-2 @ FNAL
- g-2 @ JPARC

High Energy Physics: (~ 20 ppl.)

- Λ -baryon @ LHCb

Atoms: (~ 60 ppl.)


- Cs @ Penn State
- Fr @ Riken
- Hg @ Bonn
- Ra @ Argonne
- Xe @ Heidelberg
- Xe @ PTB
- Xe @ Riken



Molecules: (~ 55 ppl.)



- BaF (EDM³) @ Toronto
- BaF (NLeEDM) @ Groningen/Nikhef
- HfF⁺ @ JILA
- ThO (ACME) @ Yale
- YBF @ Imperial

Axion Searches




S. P. Chang, S. Hacıomeroglu, O. Kim, S. Lee, S. Park and Y. K. Semertzidis, PoS PSTP **2017** (2018) 036 [arXiv:1710.05271 [hep-ex]].

Momentum and ring radius for deuteron in frozen spin condition



Why Charged Particle EDMs?



J. de Vries