

Chiral constraints on the isoscalar electromagnetic spectral functions of the nucleon from leading order vector meson couplings



Y. Ünal^{a,b,*}, Ulf-G. Meißner^{a,c,d}

^a Helmholtz-Institut für Strahlen- und Kernphysik and Bethe Center for Theoretical Physics, Universität Bonn, D-53115 Bonn, Germany

^b Physics Department, Çanakkale Onsekiz Mart University, 17100 Çanakkale, Turkey

^c Institute for Advanced Simulation, Institut für Kernphysik, and Jülich Center for Hadron Physics, Forschungszentrum Jülich, D-52425 Jülich, Germany

^d Tbilisi State University, 0186 Tbilisi, Georgia

ARTICLE INFO

Article history:

Received 13 February 2019

Received in revised form 16 April 2019

Accepted 20 April 2019

Available online 22 May 2019

Editor: W. Haxton

Keywords:

Electromagnetic form factors

Baryon chiral perturbation theory

ABSTRACT

Using baryon chiral perturbation theory including vector mesons, we analyze various continuum contributions to the isoscalar electromagnetic spectral functions of the nucleon induced by the leading order couplings.

© 2019 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

A premier tool to analyse the electromagnetic structure of the nucleon are dispersion relations. The physics is encoded in the so-called spectral functions, which feature vector meson resonances and continuum contributions. Much is known about the continuum contribution to the isovector spectral function of the electric and magnetic nucleon form factors, see e.g. Ref. [1] for the most recent and most precise investigation. The situation for the corresponding continuum contributions to the isoscalar spectral functions is different, as discussed in detail in Ref. [2]. Chiral perturbation theory has been used to analyze the three-pion continuum [3,4] within the threshold region starting at $t_0 = 9M_\pi^2$. These two-loop contributions, however, turn out to be rather small, because the potential enhancement from the anomalous threshold at $t \simeq 8.9M_\pi^2$ is efficiently masked by phase space factors. On the other hand, earlier modelling suggests an important contribution from the $\pi\rho$ continuum [5] and dispersion relations have been used to calculate the $K\bar{K}$ contribution [6,7]. In the sector with strange quarks, there have been suggestions of cancellation between contributions from $K\bar{K}$, KK^* and K^*K^* loops, although these have very different thresholds [8]. It might be worth to put our work in proper

context: We work with a chiral Lagrangian with explicit vector mesons, whereas in Ref. [5] a meson-exchange model was used. In contrast to that, the approach used here can be systematically improved. As concerns the dispersive approach used in Refs. [6,7], it in principle contains the vector meson contributions. However, as noted in these papers, the required data for the analytic continuation of the kaon-nucleon scattering amplitudes are not precise enough to uniquely fix the isoscalar spectral functions. Thus, the approach used here gives complementary information.

In this paper, we want to re-analyze some of these continua as well as some vector meson contributions to the isoscalar electromagnetic spectral functions based on chiral perturbation theory with explicit vector mesons. This extends the earlier tree-level analyses of vector meson contributions to the nucleon electromagnetic (em) form factors as given e.g. in Refs. [9,10]. Here, we work at leading one-loop order and consider the vector coupling induced contributions. The paper is organized as follows: In Sec. 2, we briefly recall the pertinent definitions for the electromagnetic form factors and their spectral functions. The $\pi\rho$ contribution is worked out and discussed in Sec. 3. The contributions involving K and K^* mesons are given in Sec. 4. We end with a summary and outlook. Some technicalities are given in the Appendix.

2. Isoscalar electromagnetic spectral functions

The electromagnetic form factors of the nucleon are defined via the matrix element of the electromagnetic current,

* Corresponding author at: Helmholtz-Institut für Strahlen- und Kernphysik and Bethe Center for Theoretical Physics, Universität Bonn, D-53115 Bonn, Germany.

E-mail addresses: unal@hiskp.uni-bonn.de (Y. Ünal), meissner@hiskp.uni-bonn.de (U.-G. Meißner).

$$\begin{aligned} & \langle N(p_f) | \bar{q} \gamma^\mu Q q | N(p_i) \rangle \\ &= \bar{u}(p_f) \left[\gamma^\mu F_1(t) + \frac{i}{2m_N} \sigma^{\mu\nu} (p_f - p_i)_\nu F_2(t) \right] u(p_i), \quad (1) \end{aligned}$$

with $t = (p_f - p_i)^2$ the invariant momentum transfer squared, m_N the nucleon mass and Q the quark charge matrix. Throughout, we work in the isospin limit $m_N = (m_p + m_n)/2$, with m_p (m_n) the proton (neutron) mass. The functions $F_1(t)$ and $F_2(t)$ are the Dirac and Pauli form factors of the nucleon, respectively. In the isospin basis, they can be decomposed into isoscalar and isovector components, following the notation of Ref. [11]

$$F_i(t) = F_i(t)^S + \tau^3 F_i^V(t), \quad i = 1, 2.$$

$F_1(t)$ and $F_2(t)$ are related to the commonly used electric and magnetic Sachs form factors in the following fashion

$$\begin{aligned} G_E^{S,V}(t) &= F_1^{S,V}(t) + \frac{t}{4m_N^2} F_2^{S,V}(t), \\ G_M^{S,V}(t) &= F_1^{S,V}(t) + F_2^{S,V}(t). \end{aligned} \quad (2)$$

An unsubtracted dispersion relation can be written down for each form factor defined by

$$F(t) = \frac{1}{\pi} \int_{t_0}^{\infty} \frac{\text{Im}F(t')}{t' - t - i\epsilon} dt' \quad (3)$$

where t_0 is the threshold energy for hadronic intermediate states. Here, we focus entirely on the isoscalar spectral function with $t_0 = 9M_\pi^2$, as the three-pion state is the lowest mass intermediate state possible. For a more general discussion of the spectral functions, see e.g. Refs. [2,12,13].

3. The $\pi\rho$ contribution

In this section we focus on the calculation of the imaginary parts of the isoscalar nucleon electromagnetic form factors generated from the $\pi\rho$ continuum based on relativistic two-flavor baryon chiral perturbation theory. At lowest order in the quark mass and momentum expansion, the relevant interaction Lagrangians are given by [14–16]

$$\begin{aligned} \mathcal{L}_{\pi N} &= \frac{g_A}{2} \bar{\Psi} \gamma^\mu \gamma_5 u_\mu \Psi, \\ \mathcal{L}_{\rho N} &= g_{\rho N} \bar{\Psi} \rho_\mu \gamma^\mu \Psi, \\ \mathcal{L}_{\omega\rho\pi} &= \frac{g_{\omega\rho\pi}}{2} \epsilon^{\mu\nu\alpha\beta} \omega_\nu \langle \rho_{\alpha\beta} u_\mu \rangle, \\ \mathcal{L}_{\omega\gamma} &= \frac{f_\omega}{2} e (\partial_\mu \omega_\nu - \partial_\nu \omega_\mu) (\partial^\mu A^\nu - \partial^\nu A^\mu). \end{aligned} \quad (4)$$

Here, Ψ denotes the nucleon doublet, ρ_μ the isovector-vector ρ -meson, ω_μ the isoscalar-vector ω -meson, A^μ the photon field, $\langle \dots \rangle$ denotes the trace in flavor space and $\epsilon^{\mu\nu\alpha\beta}$ is the totally antisymmetric Levi-Civita tensor with $\epsilon^{0123} = 1$. Further, g_A is the nucleon axial-vector coupling, $g_A \simeq 1.27$, and the various vector meson couplings are taken as $g_{\rho N} = 5.92$, $g_{\omega\rho\pi} = 11.6$ [17] and $f_\omega = 0.1$ [18]. As f_ω is not well determined in the analysis of Ref. [18], we also use $f_\omega = 0.4$ corresponding to the larger coupling g_ω from Ref. [2], keeping the product $g_\omega f_\omega$ fixed. This defines a reasonable range for this coupling constant. In the SU(2) sector, due to the universality of the ρ -meson coupling the equality of the ρ -meson self-coupling and the coupling to the nucleons $g = g_{\rho N}$, holds [19]. As can be seen from the Lagrangian in Eq. (4), we work in the vector meson dominance approximation, i.e. the




Fig. 1. Continuum $\pi\rho$ contribution to the imaginary parts of the isoscalar electromagnetic form factors of the nucleon through the anomalous coupling to the ω . Dashed and solid lines denote pions and nucleons, respectively. The dashed lines represent the ω - and ρ -mesons.

photon couples only via vector mesons to the hadrons. This is e.g. realized in the hidden symmetry approach of Ref. [20] for the parameter $a = 2$. However, in general, the gauged Wess-Zumino-Witten term also leads to a direct $\gamma\rho\pi$ coupling e.g. in the massive Yang-Mills approach. It can, however, be shown that all these different approaches are equivalent, see e.g. the detailed discussion in Ref. [21]. Our choice is driven by simplicity. For more thorough discussion of the Wess-Zumino-Witten term in the presence of vector mesons and photons, see the reviews [20,21]. Note also that we neglect the small OZI-violating π - ρ contribution due to the ϕ -meson coupling.

The following power counting rules for the loop diagrams are used: vertices from $\mathcal{L}^{(n)}$ count as $\mathcal{O}(q^n)$, so we count the vector meson, nucleon and pion propagators as $\mathcal{O}(q^0)$, $\mathcal{O}(q^{-1})$ and $\mathcal{O}(q^{-2})$, in order. Thus the order of the diagrams in Fig. 1 is $\mathcal{O}(q^7)$ at low energies, i.e. for small t .

We work in the centre of momentum frame of the nucleon pair with $q = (-2E_p, \vec{0})$. The initial and the final momentum of the nucleons are, respectively, $p_i = (E_p, \vec{p})$, $p_f = (E_p, \vec{p})$, with $|\vec{p}| = (t/4 - m_N^2)^{1/2}$ and $E_p = (m_N^2 + |\vec{p}|^2)^{1/2}$. To calculate the imaginary part of the amplitude for the diagrams which are shown in Fig. 1, the Cutkosky rules are applied. The imaginary part of the loop diagram corresponds to a cut diagram for the momentum transfer squared $t \geq (M_\rho + M_\pi)^2$. For this calculation, we perform a reduction to scalar loop integrals and thus require the basic scalar loop integrals of two- and three- point functions, called H and F , respectively,

$$\begin{aligned} H &= i \int \frac{d^4k}{(2\pi)^4} \frac{1}{[k^2 - M_\pi^2 + i\epsilon^+][(k+q)^2 - M_\rho^2 + i\epsilon^+]}, \\ F &= i \int \frac{d^4k}{(2\pi)^4} \frac{1}{[k^2 - M_\pi^2 + i\epsilon^+][(p_i - k)^2 - m_N^2 + i\epsilon^+][(k+q)^2 - M_\rho^2 + i\epsilon^+]}, \end{aligned} \quad (5)$$

with $q^2 = t = (p_f - p_i)$ and ϵ^+ stands for $\epsilon \rightarrow 0^+$. The corresponding imaginary parts can be readily determined

$$\begin{aligned} \text{Im} H : \mathcal{H}(t) &= \frac{M_\pi}{8\pi\sqrt{t}} \sqrt{\frac{(M_\rho^2 - M_\pi^2 - t)^2}{4M_\pi^2 t} - 1}, \\ \text{Im} F : \mathcal{F}(t) &= \frac{1}{16\pi\sqrt{t}} \sqrt{t - 4m_N^2} \\ &\quad \times \log \left[\frac{\frac{t - M_\pi^2 - M_\rho^2}{\sqrt{t - 4m_N^2}} - \sqrt{\frac{M_\pi^4 + (M_\rho^2 - t)^2 - 2M_\pi^2(M_\rho^2 + t)}{t}}}{\frac{t - M_\pi^2 - M_\rho^2}{\sqrt{t - 4m_N^2}} + \sqrt{\frac{M_\pi^4 + (M_\rho^2 - t)^2 - 2M_\pi^2(M_\rho^2 + t)}{t}}} \right]. \end{aligned} \quad (6)$$

From these, the expressions for the imaginary part of the isoscalar electric and magnetic form factor can be given as follows




Fig. 2. Continuum $\pi\rho$ contribution through the anomalous coupling to the ω to the isoscalar spectral functions of the electric (upper panel) and magnetic nucleon form factors (lower panel) weighted with and without $1/t^2$. The bands represent the variation of f_ω as discussed in the text.

$$\begin{aligned} \text{Im } G_E^S(t) &= \frac{3f_\omega g_{\omega\rho\pi} g_{\rho N} g_A}{32\pi^2 F_\pi^2} \frac{t}{(t - m_\omega^2)(t - 4m_N^2)} \\ &\times \left[2(t^2 m_N^2 + tM_\pi^2 M_\rho^2 - 2tM_\pi^2 m_N^2 - 2tM_\rho^2 m_N^2 \right. \\ &+ M_\pi^4 m_N^2 + M_\rho^4 m_N^2 - 2M_\pi^2 M_\rho^2 m_N^2) \mathcal{F}(t) \\ &\left. + 2t(t - M_\pi^2 - M_\rho^2) \mathcal{H}(t) \right], \end{aligned} \quad (7)$$

$$\begin{aligned} \text{Im } G_M^S(t) &= \frac{3f_\omega g_{\omega\rho\pi} g_{\rho N} g_A}{16\pi^2 F_\pi^2} \frac{m_N^2}{(t - m_\omega^2)(t - 4m_N^2)^2} \\ &\times \left[(2tm_N^2(3M_\pi^4 + 3M_\rho^4 - 2M_\pi^2 M_\rho^2 + 8M_\pi^2 m_N^2 \right. \\ &+ 8M_\rho^2 m_N^2) \\ &- t^2(M_\pi^4 + M_\rho^4 + 8m_N^4 - 4M_\pi^2 m_N^2 + 12M_\rho^2 m_N^2) \\ &+ 2t^3(M_\pi^2 + M_\rho^2 + 3m_N^2) - t^4 \\ &- 8m_N^4(M_\pi^4 + M_\rho^4 - 2M_\pi^2 M_\rho^2)) \mathcal{F}(t) \\ &\left. + (t^2 + tM_\pi^2 + tM_\rho^2)(t - 4m_N^2) \mathcal{H}(t) \right]. \end{aligned} \quad (8)$$

The resulting electric and magnetic spectral functions $\text{Im } G_{E,M}^S(t)$ and the weighted spectral functions $\text{Im } G_{E,M}^S(t)/t^2$ are shown in Fig. 2. The magnetic one shows a peak at $t \simeq 1.1 \text{ GeV}^2$, which is consistent with the modelling in Ref. [5], where the calculated $\pi\rho$ distribution was approximated by a sharp vector meson poles with a mass of about 1.1 GeV. For a more detailed comparison, we would also have to include the ρ -meson tensor couplings that appear at next-to-leading order.

4. The contribution from strange loops

Next, we consider the effect of loops with $K\bar{K}$, KK^* and K^*K^* loops. We note, however, that the latter two only start to have an imaginary part at about 2 GeV^2 and 3 GeV^2 , which is already far inside the resonance dominated region. The relevant effective Lagrangians in the SU(3) basis to obtain the contributions from $K\bar{K}$, KK^* and K^*K^* loops read [20–25]

$$\begin{aligned} \mathcal{L}_{V\Phi\Phi} &= -ig\langle V^\mu[\Phi, \partial_\mu\Phi] \rangle, \\ \mathcal{L}_{VV\Phi} &= \frac{1}{\sqrt{2}} G \epsilon^{\mu\nu\alpha\beta} \langle \partial_\mu V_\nu \partial_\alpha V_\beta \Phi \rangle, \\ \mathcal{L}_{VVV} &= ig\langle (\partial_\mu V_\nu - \partial_\nu V_\mu) V^\mu V^\nu \rangle, \\ \mathcal{L}_{BB\Phi} &= -\frac{F}{2} \langle \bar{B} \gamma^\mu \gamma_5 \{u_\mu, B\} \rangle - \frac{D}{2} \langle \bar{B} \gamma^\mu \gamma_5 \{u_\mu, B\} \rangle, \\ \mathcal{L}_{BBV} &= G_F \langle \bar{B} \gamma^\mu \{V_\mu, B\} \rangle + \langle \bar{B} \gamma^\mu B \rangle \langle V_\mu \rangle \\ &\quad + G_D \langle \bar{B} \gamma^\mu \{V_\mu, B\} \rangle + \langle \bar{B} \gamma^\mu B \rangle \langle V_\mu \rangle, \\ \mathcal{L}_{\phi\gamma} &= \frac{e}{3\sqrt{2}} f_V (\partial_\mu \phi_\nu - \partial_\nu \phi_\mu) (\partial^\mu A^\nu - \partial^\nu A^\mu) \end{aligned} \quad (9)$$

where V_μ and V_μ represent the vector meson octet and the vector meson nonet particles, respectively. We take the couplings as $G_F = g$ and $G_D = 0$ which results from the constraint analysis and the perturbative renormalizability condition on the vector meson octet and baryon octet interaction of Ref. [26]. The low-energy constants D and F can be determined by fitting to the semi-leptonic baryon decays at tree level. We use the values $F = 0.5$ and $D = 0.8$ from Ref. [27]. Throughout, we use the pion decay constant F_π for the leading order meson decay constant.

In addition to the scalar loop integrals in Eq. (6), we also need the following functions corresponding to the $\bar{K}K$, K^*K^* , $\Lambda\bar{K}K$, $\Sigma\bar{K}K$, ΛK^*K^* and ΣK^*K^* loops with the equal masses




Fig. 3. Contributions from $K\bar{K}$, KK^* and K^*K^* loops to the imaginary parts of the isoscalar electromagnetic form factors of the nucleon through the coupling to the ϕ . The double wiggly, double dashed and dashed lines denote ϕ -, K^* - and K -mesons, respectively. The dashed solid lines represent either Λ - or Σ -baryons. There is an additional permutation of the diagrams in group (b).

$$P = i \int \frac{d^4k}{(2\pi)^4} \frac{1}{[k^2 - M_i^2 + i\epsilon^+][(k+q)^2 - M_i^2 + i\epsilon^+]},$$

$$J = i \int \frac{d^4k}{(2\pi)^4} \frac{1}{[k^2 - M_i^2 + i\epsilon^+][(k+q)^2 - M_i^2 + i\epsilon^+][(p_i-k)^2 - m_j^2 + i\epsilon^+]}. \quad (10)$$

The corresponding imaginary parts are determined as

$$\text{Im } P : \mathcal{P}_i(t) = -\frac{1}{16\pi} \sqrt{1 - \frac{4M_i^2}{t}},$$

$$\text{Im } J : \mathcal{J}_j(t) = \frac{1}{16\pi \sqrt{t} \sqrt{t - 4m_j^2}} \times \log \left[\frac{M_i^2 - \frac{t}{2} + \frac{1}{2} \sqrt{-4m_j^2 + t} \sqrt{-4M_i^2 + t}}{M_i^2 - \frac{t}{2} - \frac{1}{2} \sqrt{-4m_j^2 + t} \sqrt{-4M_i^2 + t}} \right] \quad (11)$$

where M_i and m_j denote the masses of the K , K^* mesons and of the Λ , Σ baryons, respectively.

We work in the particle basis using the SU(3) symmetric Lagrangian densities given above. An example of such a calculation can be found in the appendix. The imaginary parts of the Sachs form factors for the KK loop are obtained as

$$\text{Im } G_E^S(t) = \frac{f_V g}{432 \sqrt{2} F_\pi^2 \pi^2 m_N (t - 4m_N^2) (t - M_\phi^2)} \times \left[\begin{aligned} &3(D + 3F)^2 t (m_\Lambda + m_N)^2 (4m_N^5 - 4M_K^4 m_N \\ &- 4m_\Lambda^4 m_N - 2m_\Lambda^2 m_N t - 2m_N^3 t - m_\Lambda^3 (2t - 8m_N^2)) \\ &- m_\Lambda (8m_N^4 - 6m_N^2 t + t^2) \\ &+ 2M_K^2 (4m_\Lambda^2 m_N + m_N t + m_\Lambda (t - 4m_N^2)) \mathcal{J}_\Lambda(t) \\ &- 2t (D^2 (6m_\Lambda^4 m_N + 60m_N^5 - 108m_N^3 m_\Sigma^2 + 54m_N m_\Sigma^4 \\ &+ 3m_\Lambda^3 t + 3m_\Lambda m_N^2 t - 20m_N^3 t + 27m_N^2 m_\Sigma t \\ &+ 54m_N m_\Sigma^2 t + 27m_\Sigma^3 t + 5m_N t^2 + 6m_\Lambda^2 m_N (t - 2m_N^2)) \\ &- 2M_K^2 m_N (3m_\Lambda^2 + 6m_\Lambda m_N - 10m_N^2 + 54m_N m_\Sigma \\ &+ 27m_\Sigma + 10t) - 6DF (-6m_\Lambda^4 m_N + 12m_N^5 \\ &- 36m_N^3 m_\Sigma^2 + 18m_N m_\Sigma^4 - 3m_\Lambda^3 t - 3m_\Lambda m_N^2 t \\ &- 4m_N^3 t + 9m_N^2 m_\Sigma t + 18m_N m_\Sigma^2 t + 9m_\Sigma^3 t + m_N t^2 \\ &+ 6m_\Lambda^2 (2m_N^3 - m_N t) + 2M_K^2 m_N (3m_\Lambda^2 + 6m_\Lambda m_N \\ &+ 2m_N^2 - 18m_N m_\Sigma - 9m_\Sigma^2 - 2t)) \\ &- 9F^2 (-6m_\Lambda^4 m_N - 12m_N^5 + 12m_N^3 m_\Sigma^2 - 6m_N m_\Sigma^4 \end{aligned} \right]$$

$$\begin{aligned} &- 3m_\Lambda^3 t - 3m_\Lambda m_N^2 t + 4m_N^3 t - 3m_N^2 m_\Sigma t - 6m_N m_\Sigma^2 t \\ &- 3m_\Sigma^3 t - m_N t^2 + 6m_\Lambda^2 (2m_N^3 - m_N t) \\ &+ 2M_K^2 m_N (3m_\Lambda^2 + 6m_\Lambda m_N - 2m_N^2 + 6m_N m_\Sigma \\ &+ 3m_\Sigma^2 + 2t)) \mathcal{P}_K(t) \\ &- 27(D - F)^2 t (m_N + m_\Sigma)^2 (4M_K^4 m_N - 4m_N^5 + 8m_N^4 m_\Sigma \\ &+ 2m_N^3 t + 2m_N m_\Sigma^2 (2m_\Sigma^2 + t) + m_\Sigma t (2m_\Sigma^2 + t) \\ &- 2m_N^2 (4m_\Sigma^3 + 3m_\Sigma t) \\ &+ M_K^2 (8m_N^2 m_\Sigma - 2m_\Sigma t - 2m_N (4m_\Sigma^2 + t))) \mathcal{J}_\Sigma(t) \end{aligned} \quad (12)$$

and

$$\text{Im } G_M^S(t) = \frac{f_V g}{432 \sqrt{2} F_\pi^2 \pi^2 (t - 4m_N^2) (t - M_\phi^2)} \times \left[\begin{aligned} &6(D + 3F)^2 t (m_\Lambda + m_N)^2 (M_K^4 + m_\Lambda^4 + m_N^4 \\ &- 2M_K^2 (m_\Lambda^2 + m_N^2) + m_\Lambda^2 (t - 2m_N^2)) \mathcal{J}_\Lambda(t) \\ &- t (D^2 (-6m_\Lambda^4 - 12m_\Lambda^3 m_N + 12m_\Lambda m_N^3 \\ &+ 60m_N^4 + 108m_N^3 m_\Sigma - 108m_N m_\Sigma^3 - 54m_\Sigma^4 - 3m_\Sigma^2 t \\ &- 6m_\Lambda m_N t - 70m_N^2 t - 54m_N m_\Sigma t - 27m_\Sigma^2 t + 10t^2 \\ &+ 2M_K^2 (3m_\Lambda^2 + 6m_\Lambda m_N + 110m_N^2 + 54m_N m_\Sigma \\ &+ 27m_\Sigma^2 - 20t) + 6DF (-6m_\Lambda^4 - 12m_\Lambda^3 m_N \\ &+ 12m_\Lambda m_N^3 - 12m_N^4 - 36m_N^3 m_\Sigma + 36m_N m_\Sigma^3 + 18m_\Sigma^4 \\ &- 3m_\Lambda^2 t - 6m_\Lambda m_N t + 14m_N^2 t + 18m_N m_\Sigma t + 9m_\Sigma^2 t \\ &- 2t^2 + 2M_K^2 (3m_\Lambda^2 + 6m_\Lambda m_N - 22m_N^2 - 18m_N m_\Sigma \\ &- 9m_\Sigma^2 + 4t)) + 9F^2 (-6m_\Lambda^4 - 12m_\Lambda^3 m_N \\ &+ 12m_\Lambda m_N^3 + 12m_N^4 + 12m_N^3 m_\Sigma - 12m_N m_\Sigma^3 \\ &- 6m_\Sigma^4 - 3m_\Lambda^2 t - 6m_\Lambda m_N t - 14m_N^2 t - 6m_N m_\Sigma t \\ &- 3m_\Sigma^2 t + 2t^2 + 2M_K^2 (3m_\Lambda^2 \\ &+ 6m_\Lambda m_N + 22m_N^2 + 6m_N m_\Sigma + 3m_\Sigma^2 - 4t)) \mathcal{P}_K(t) \\ &+ 54(D - F)^2 t (m_N + m_\Sigma)^2 (M_K^4 + m_\Lambda^4 - 2m_N^2 m_\Sigma^2 + m_\Lambda^4 \\ &- 2M_K^2 (m_N^2 + m_\Sigma^2) + m_\Sigma^2 t) \mathcal{J}_\Sigma(t) \end{aligned} \right] \quad (13)$$

The definitions of $\mathcal{J}_\Lambda(t)$, $\mathcal{J}_\Sigma(t)$ and $\mathcal{P}_K(t)$ can be found in Eq. (11). In Fig. 4, we show the $K\bar{K}$ loop contribution including the one from the ϕ -meson. To account for its finite width, we have substituted M_ϕ by $M_\phi - i\Gamma_\phi/2$, where $\Gamma_\phi \simeq 4$ MeV is the width of




Fig. 4. $K\bar{K}$ contribution through the anomalous coupling to the ϕ to the isoscalar spectral functions weighted with and without $1/t^2$. The red (solid) and blue (dashed) lines refer to the electric and magnetic form factors, $\text{Im } G_E^S(t)$ and $\text{Im } G_M^S(t)$, respectively.




Fig. 5. K^*K contribution through the anomalous coupling to the ϕ to the isoscalar spectral functions weighted with and without $1/t^2$. For notations, see Fig. 4.




Fig. 6. K^*K^* contribution through the anomalous coupling to the ϕ to the isoscalar spectral functions weighted with and without $1/t^2$. For notations, see Fig. 4.

the ϕ . We find similar spectral functions for the electric and magnetic form factors except for the sign. Similar to Ref. [6], there is no enhancement on the left wing of the ϕ which sits directly at the $K\bar{K}$ threshold. Our results with vector mesons at one loop can, however, not directly be compared to dispersion-theoretical result of Ref. [7] as their calculation does not include all of the ϕ -meson strength and ours would also require the inclusion of the NLO tensor coupling. Still, it is comforting to see that in the region of the ϕ there is little continuum strength. As noted earlier, the spectral functions generated by K^*K and K^*K^* start at much larger t and turn out to be rather small, see Fig. 5 and Fig. 6, respectively.

5. Summary

In this paper, we have used baryon chiral perturbation theory including vector mesons to derive new chiral constraints on the isoscalar electromagnetic spectral functions based on the leading order vector coupling. As noted earlier, the π - ρ loop contribution is the most substantial non-resonant effect and needs to be included. The non-resonant $K\bar{K}$ contribution that sits under the ϕ -meson is less pronounced, which is consistent with earlier findings, used e.g. in Ref. [2]. The inclusion of the tensor coupling effects should be considered next.

Acknowledgements

We are grateful to Jambul Gegelia for helpful discussions. We acknowledge partial financial support from the Deutsche Forschungsgemeinschaft (SFB/TRR 110, ‘‘Symmetries and the Emergence of Structure in QCD’’), by the Chinese Academy of Sciences (CAS) President’s International Fellowship Initiative (PIFI) (grant no. 2018DM0034) and by VolkswagenStiftung (grant no. 93562).

Appendix A. A sample SU(3) calculation

As an example for the pertinent SU(3) calculations, we consider the $K\bar{K}\Lambda$ contribution depicted in Fig. 7. The Feynman rules for the $\phi\gamma$, ϕK^+K^- , ΛK^-p and $\Lambda K^+\bar{p}$ vertices appearing in this diagram are obtained as follows

$$\begin{aligned} \phi\gamma &: \frac{2ief_V}{3\sqrt{2}}(g^{\mu\nu}q^2 - q^\mu q^\nu) \\ \phi K^+K^- &: 2(\partial^\mu K^+)K^- \phi_\mu - 2(\partial^\mu K^-)K^+ \phi_\mu \\ i\mathcal{L} &\rightarrow 2ig(k_1^\mu - k_2^\mu) \end{aligned}$$




Fig. 7. A Feynman diagram for the $\bar{K}K\Lambda$ loop contribution.

$$pK^-\bar{\Lambda} : -\frac{D}{2\sqrt{3}}\bar{\Lambda}(\partial^\mu K^-)p - \frac{F\sqrt{3}}{2}\bar{\Lambda}(\partial^\mu K^-)p$$

$$i\mathcal{L} \rightarrow \frac{(D+3F)}{2\sqrt{3}F_\pi}\gamma_5 k_2$$

$$\Lambda K^+\bar{p} : -\frac{D}{2\sqrt{3}}\bar{p}(\partial^\mu K^+)\Lambda - \frac{F\sqrt{3}}{2}\bar{p}(\partial^\mu K^+)\Lambda$$

$$i\mathcal{L} \rightarrow \frac{(D+3F)}{2\sqrt{3}F_\pi}\gamma_5 k_1$$

Here, k_1 and k_2 denote the momenta for the K^+ and the K^- , in order. The ϕ vector meson has momentum q and Lorentz index μ . The amplitude for the diagram in Fig. 7 reads

$$\Gamma^\mu(p_i, p_f, q) = \frac{egf_V(D+3F)^2}{24\sqrt{2}F_\pi^2\pi^4}$$

$$\times \mu^{4-n} \int d^n k \frac{(q^\mu q^\mu - q^2 g^{\mu\nu})(2k^\delta - q^\delta)(g_{\delta\eta}M_\phi^2 - q_\delta q_\eta)\gamma^5(g-k)_{(\rho_i+k+m_\Lambda)}\gamma^5 k}{(q^2 - M_\phi^2)(k^2 - M_K^2)((k+p_i)^2 - m_\Lambda^2)((q-k)^2 - M_K^2)},$$

with n the number of dimensions and μ the scale of dimensional regularization.

References

- [1] M. Hoferichter, B. Kubis, J. Ruiz de Elvira, H.-W. Hammer, U.-G. Meißner, On the $\pi\pi$ continuum in the nucleon form factors and the proton radius puzzle, *Eur. Phys. J. A* 52 (11) (2016) 331, <https://doi.org/10.1140/epja/i2016-16331-7>, arXiv:1609.06722 [hep-ph].
- [2] M.A. Belushkin, H.-W. Hammer, U.-G. Meißner, Dispersion analysis of the nucleon form-factors including meson continua, *Phys. Rev. C* 75 (2007) 035202, <https://doi.org/10.1103/PhysRevC.75.035202>, arXiv:hep-ph/0608337.
- [3] V. Bernard, N. Kaiser, U.-G. Meißner, Nucleon electroweak form-factors: analysis of their spectral functions, *Nucl. Phys. A* 611 (1996) 429–441, [https://doi.org/10.1016/S0375-9474\(96\)00291-6](https://doi.org/10.1016/S0375-9474(96)00291-6), arXiv:hep-ph/9607428.
- [4] N. Kaiser, E. Passemar, Spectral functions of nucleon form factors: three-pion continua at low energies, *Eur. Phys. J. A* 55 (2) (2019) 16, arXiv:1901.02865 [nucl-th].
- [5] U.-G. Meißner, V. Mull, J. Speth, J.W. van Orden, Strange vector currents and the OZI rule, *Phys. Lett. B* 408 (1997) 381–386, [https://doi.org/10.1016/S0370-2693\(97\)00828-9](https://doi.org/10.1016/S0370-2693(97)00828-9), arXiv:hep-ph/9701296.
- [6] H.W. Hammer, M.J. Ramsey-Musolf, K anti-K continuum and isoscalar nucleon form-factors, *Phys. Rev. C* 60 (1999) 045204, <https://doi.org/10.1103/PhysRevC.60.045204>, Erratum: *Phys. Rev. C* 62 (2000) 049902, arXiv:hep-ph/9903367.
- [7] H.W. Hammer, M.J. Ramsey-Musolf, Spectral content of isoscalar nucleon form-factors, *Phys. Rev. C* 60 (1999) 045205, <https://doi.org/10.1103/PhysRevC.60.045205>, Erratum: *Phys. Rev. C* 62 (2000) 049903, arXiv:hep-ph/9812261.
- [8] L.L. Barz, H. Forkel, H.W. Hammer, F.S. Navarra, Marina Nielsen, M.J. Ramsey-Musolf, K^* mesons and nucleon strangeness, *Nucl. Phys. A* 640 (1998) 259–280, [https://doi.org/10.1016/S0375-9474\(98\)00438-2](https://doi.org/10.1016/S0375-9474(98)00438-2), arXiv:hep-ph/9803221.
- [9] Bastian Kubis, Ulf-G. Meißner, Low-energy analysis of the nucleon electromagnetic form-factors, *Nucl. Phys. A* 679 (2001) 698–734, [https://doi.org/10.1016/S0375-9474\(00\)00378-X](https://doi.org/10.1016/S0375-9474(00)00378-X), arXiv:hep-ph/0007056.
- [10] Matthias R. Schindler, J. Gegelia, Stefan Scherer, Electromagnetic form-factors of the nucleon in chiral perturbation theory including vector mesons, *Eur. Phys. J. A* 26 (2005) 1–5, <https://doi.org/10.1140/epja/i2005-10145-8>, arXiv:nucl-th/0509005.
- [11] P. Mergell, Ulf-G. Meißner, D. Drechsel, Dispersion theoretical analysis of the nucleon electromagnetic form-factors, *Nucl. Phys. A* 596 (1996) 367–396, [https://doi.org/10.1016/0375-9474\(95\)00339-8](https://doi.org/10.1016/0375-9474(95)00339-8), arXiv:hep-ph/9506375.
- [12] I.T. Lorenz, Ulf-G. Meißner, H.-W. Hammer, Y.-B. Dong, Theoretical constraints and systematic effects in the determination of the proton form factors, *Phys. Rev. D* 91 (1) (2015) 014023, <https://doi.org/10.1103/PhysRevD.91.014023>, arXiv:1411.1704 [hep-ph].
- [13] Stefan Leupold, The nucleon as a test case to calculate vector-isovector form factors at low energies, *Eur. Phys. J. A* 54 (1) (2018) 1, <https://doi.org/10.1140/epja/i2018-12447-0>, arXiv:1707.09210 [hep-ph].
- [14] G. Ecker, J. Gasser, H. Leutwyler, A. Pich, E. de Rafael, Chiral lagrangians for massive spin 1 fields, *Phys. Lett. B* 223 (1989) 425–432, [https://doi.org/10.1016/0370-2693\(89\)91627-4](https://doi.org/10.1016/0370-2693(89)91627-4).
- [15] G. Ecker, J. Gasser, A. Pich, E. de Rafael, The role of resonances in chiral perturbation theory, *Nucl. Phys. B* 321 (1989) 311–342, [https://doi.org/10.1016/0550-3213\(89\)90346-5](https://doi.org/10.1016/0550-3213(89)90346-5).
- [16] J. Gasser, M.E. Sainio, A. Svarc, Nucleons with chiral loops, *Nucl. Phys. B* 307 (1988) 779–853, [https://doi.org/10.1016/0550-3213\(88\)90108-3](https://doi.org/10.1016/0550-3213(88)90108-3).
- [17] K. Nakayama, Yongseok Oh, J. Haidenbauer, T.-S.H. Lee, On the sign of the $\pi-\rho-\omega$ coupling constant, *Phys. Lett. B* 648 (2007) 351–356, <https://doi.org/10.1016/j.physletb.2007.03.034>, arXiv:nucl-th/0611101.
- [18] T. Bauer, J.C. Bernauer, S. Scherer, Electromagnetic form factors of the nucleon in effective field theory, *Phys. Rev. C* 86 (2012) 065206, <https://doi.org/10.1103/PhysRevC.86.065206>, arXiv:1209.3872 [nucl-th].
- [19] D. Djukanovic, M.R. Schindler, J. Gegelia, G. Japaridze, S. Scherer, Universality of the ρ -meson coupling in effective field theory, *Phys. Rev. Lett.* 93 (2004) 122002, <https://doi.org/10.1103/PhysRevLett.93.122002>, arXiv:hep-ph/0407239.
- [20] Masako Bando, Taichiro Kugo, Koichi Yamawaki, Nonlinear realization and hidden local symmetries, *Phys. Rep.* 164 (1988) 217–314, [https://doi.org/10.1016/0370-1573\(88\)90019-1](https://doi.org/10.1016/0370-1573(88)90019-1).
- [21] Ulf-G. Meißner, Low-energy hadron physics from effective chiral lagrangians with vector mesons, *Phys. Rep.* 161 (1988) 213, [https://doi.org/10.1016/0370-1573\(88\)90090-7](https://doi.org/10.1016/0370-1573(88)90090-7).
- [22] Michael C. Birse, Effective chiral Lagrangians for spin 1 mesons, *Z. Phys. A* 355 (1996) 231–246, <https://doi.org/10.1007/s002180050105>, arXiv:hep-ph/9603251.
- [23] B. Borasoy, Ulf-G. Meißner, Chiral Lagrangians for baryons coupled to massive spin 1 fields, *Int. J. Mod. Phys. A* 11 (1996) 5183–5202, <https://doi.org/10.1142/S0217751X96002376>, arXiv:hep-ph/9511320.
- [24] A. Krause, Baryon matrix elements of the vector current in chiral perturbation theory, *Helv. Phys. Acta* 63 (1990) 3–70, <https://doi.org/10.5169/seals-116214>.
- [25] A. Bramon, A. Grau, G. Pancheri, Chiral perturbation theory and radiative $V^0 \rightarrow P^0 P^0 \gamma$ decays, *Phys. Lett. B* 289 (1992) 97–102, [https://doi.org/10.1016/0370-2693\(92\)91368-J](https://doi.org/10.1016/0370-2693(92)91368-J).
- [26] Y. Unal, A. Kucukarslan, S. Scherer, Interaction of the vector-meson octet with the baryon octet in effective field theory, *Phys. Rev. C* 92 (5) (2015) 055208, <https://doi.org/10.1103/PhysRevC.92.055208>, arXiv:1510.02205 [nucl-th].
- [27] B. Borasoy, Baryon axial currents, *Phys. Rev. D* 59 (1999) 054021, <https://doi.org/10.1103/PhysRevD.59.054021>, arXiv:hep-ph/9811411.