

Aircraft-based observations of ozone-depleting substances in the upper troposphere and lower stratosphere in and above the Asian summer monsoon

Karina E. Adcock¹, Paul J. Fraser², Brad D. Hall³, Ray L. Langenfelds², Geoffrey Lee¹, Stephen A. Montzka³, David E. Oram^{1,4}, Thomas Röckmann⁵, Fred Stroh⁶, William T. Sturges¹, Bärbel Vogel⁶ and Johannes C. Laube^{1,6}

¹Centre for Ocean and Atmospheric Sciences, School of Environmental Sciences, University of East Anglia, Norwich, United Kingdom

²Oceans and Atmosphere, Climate Science Centre, Commonwealth Scientific and Industrial Research Organisation, Aspendale, Victoria, Australia

³Global Monitoring Laboratory, National Oceanic and Atmospheric Administration, Boulder, CO, USA

⁴National Centre for Atmospheric Science, School of Environmental Sciences, University of East Anglia, Norwich, United Kingdom

⁵Institute for Marine and Atmospheric Research, Utrecht University, Utrecht, the Netherlands

⁶Institute of Energy and Climate Research – Stratosphere (IEK-7), Forschungszentrum Jülich GmbH, Jülich, Germany

Key points of research:

- First study to measure a comprehensive set of ozone-depleting substances in air entering the stratosphere above the Asian Summer Monsoon
- Higher than expected mixing ratios found for many compounds, particularly chlorinated very short-lived substances
- Regional estimate but the extra equivalent chlorine could significantly enhance the chlorine and bromine loading of the entire stratosphere

Abstract

Recent studies show that the Asian Summer Monsoon Anticyclone (ASMA) transports emissions from the rapidly industrializing nations in Asia into the tropical upper troposphere. Here we present a unique set of measurements on over 100 air samples collected on multiple flights of the M55 Geophysika high altitude research aircraft over the Mediterranean, Nepal and northern India during the summers of 2016 and 2017 as part of the European Union project StratoClim. These air samples were measured for 27 ozone-depleting substances (ODSs), many of which were enhanced above expected levels, including the chlorinated very short-lived substances, dichloromethane (CH_2Cl_2), 1,2-dichloroethane ($\text{CH}_2\text{ClCH}_2\text{Cl}$) and chloroform (CHCl_3). CH_2Cl_2 mixing ratios in the tropopause region were 65-136 ppt in comparison to previous estimates of mixing ratios in the tropical tropopause layer of 30-44 ppt in 2013-2014. Backward trajectories, calculated with the trajectory module of the chemistry-transport model CLaMS and driven by the ERA5 reanalysis, indicate possible source regions of CH_2Cl_2 in South Asia. We derived total Equivalent Chlorine (ECI), and Equivalent Effective Stratospheric Chlorine (EESC) and found that these quantities were substantially higher than previous estimates in the literature. EESC at mean age-of-air of 3 years based on the 2016 measurements was 1861-1872 ppt in comparison to a previously estimated EESC of 1646 ppt. Our findings show that the ASMA transports larger than expected mixing ratios of long-lived and very short-lived ODSs into the upper troposphere and lower stratosphere, likely leading to an impact on the stratospheric ozone layer.

45 1. Introduction

The Asian Summer Monsoon Anticyclone (ASMA) occurs during the boreal summer (July, August & September) over East and South Asia. It is a major meteorological system characterised by deep convection and anticyclonic flow in the upper troposphere and lower stratosphere (UTLS) and is subject to strong dynamical variability (e.g. Li et al., 2005; 50 Randel and Park, 2006; Park et al., 2007, 2008, 2009; Garny and Randel, 2013; Vogel et al., 2015). Within the ASMA, air masses are rapidly uplifted from the boundary layer into the UTLS (e.g. Park et al., 2009; Randel et al., 2010; Brunamonti et al., 2018).

In this study, we investigate whether the Asian summer monsoon convection transports elevated levels of ozone-depleting substances (ODSs) including very short-lived substances (VSLs) into the lower stratosphere. VSLs are defined here as having atmospheric lifetimes 55 of 6 months or less (Engel and Rigby et al., 2018). VSLs are not regulated under the Montreal Protocol on Substances that Deplete the Ozone Layer and have large sinks in the troposphere, which limits their contribution to stratospheric ozone depletion. However, the ASMA provides an effective pathway to transport air containing tropospheric trace gases 60 from the surface into the lower stratosphere on the time scale of a few days to a few weeks. Thus, VSLs emitted from surface regions in the ASMA should be able to reach the stratosphere at significantly elevated mixing ratios (e.g. Brioude et al., 2010; Vogel et al., 2014, 2019; Orbe et al., 2015; Hossaini et al., 2016; Tissier and Legras, 2016).

The tropics are the main entrance region to the stratosphere associated with the large-scale 65 Brewer-Dobson circulation. Transport via the ASMA provides an additional seasonal source of air into the global stratosphere, especially into the northern hemispheric lower stratosphere (Orbe et al., 2016; Vogel et al., 2016; Ploeger et al. 2017). Convection uplifts air over polluted regions, such as the Indian subcontinent and South-East Asia and this air is then horizontally confined in the UTLS by the anticyclonic winds (e.g. Park et al., 2007; Ploeger 70 et al., 2015; Vogel et al., 2015). Then air in the ASMA is transported either vertically towards the tropical stratosphere or quasi-horizontally into the Northern Hemisphere extra-tropical lower stratosphere (e.g. Orbe et al., 2015; Garny and Randel, 2016; Vogel et al., 2016, 2019; Ploeger et al., 2017).

Using the model-based results of Ploeger et al. (2017) it can be estimated that, on an annual 75 average basis, air from the ASMA contributes about 5 % of the air in the Northern Hemisphere extra-tropical stratosphere (at a potential temperature level of 380 K), whereas in the tropical stratosphere the contribution is about 2 % in the tropical pipe (at 460 K) and in the Southern Hemisphere extra-tropical stratosphere less than 0.5 % (at 380 K, Ploeger et al., 2017). The maximum contribution of air from the Asian monsoon anticyclone in the Northern 80 Hemispheric lower stratosphere is found during autumn and is around 15 % (Ploeger et al., 2017). The transport of ODSs from the ASMA region into the Northern Hemisphere extra-tropical lower stratosphere therefore has the potential to change the chemical composition of this part of the atmosphere. In addition, it has recently been found that ozone at mid-latitudes in the lower stratosphere is still decreasing, and although this is an ongoing discussion, it has 85 been suggested that VSLs may be one of the factors contributing to this (Hossaini et al., 2015; Ball et al., 2018; Chipperfield et al., 2018).

The contribution of chlorinated VSLs to total chlorine at the surface increased in recent years, from an average rate of around ~ 2.5 ppt Cl yr⁻¹ (2004-2012) to $\sim 4.3 \pm 4.9$ ppt Cl yr⁻¹ (2012-2016) (Engel and Rigby et al., 2018) and could continue to increase in the future (Leedham Elvidge et al., 2015; Oram et al., 2017; Feng et al., 2018; Fang et al., 2019). Total tropospheric chlorine has been decreasing in recent years due to the Montreal Protocol. However, this rate of decrease is slowing down: from 11.8 ± 6.9 ppt Cl yr⁻¹ during 2008-2012 to 4.4 ± 4.1 ppt Cl yr⁻¹ during 2012-2016 (Engel and Rigby et al., 2018). This is due, in part, to increasing mixing ratios of chlorinated VSLs - in particular in East and South-East Asia (Engel and Rigby et al., 2018; Fang et al., 2019; Claxton et al., 2020). This could undermine some of the progress made by the Montreal Protocol and its amendments and further offset the reduction in emissions and mixing ratios of long-lived ODSs.

The three major chlorinated VSLs are dichloromethane (CH₂Cl₂), 1,2-dichloroethane (CH₂ClCH₂Cl) and chloroform (trichloromethane, CHCl₃) (Engel and Rigby et al., 2018). CHCl₃ and CH₂Cl₂ are usually co-produced industrially (McCulloch, 2017). CH₂Cl₂ has an atmospheric lifetime of about 6 months and its global atmospheric abundances are believed to be at least 90 % anthropogenic in origin in recent years (Engel and Rigby et al., 2018; Claxton et al., 2020). It has a wide range of industrial applications in chemical and pharmaceutical processes and in the production of HFC-32 (CH₂F₂) (McCulloch, 2017). CHCl₃ has an atmospheric lifetime of about 6 months and is estimated to be about 50 % anthropogenic in origin (Engel and Rigby et al., 2018). The principal use for CHCl₃ is as a chemical feedstock for the production of HCFC-22 (CHClF₂) (Oram et al., 2017). CH₂ClCH₂Cl has an atmospheric lifetime of about 3 months and it is likely fully anthropogenic in origin (Engel and Rigby et al., 2018). Its primary use is in the manufacture of vinyl chloride, the precursor to polyvinyl chloride (PVC), and other chlorinated solvents (Oram et al., 2017). It is likely that there are major source regions of these chlorinated VSLs in China and India (Oram et al., 2017; Fang et al., 2019; Say et al., 2019; Claxton et al., 2020).

Recent studies, using air samples collected at ground-based measurement sites and on board aircraft at altitudes of 0.02-12 km, found enhancements of chlorinated VSLs in the South and East Asian region, both at the surface and in the lower and upper troposphere (Leedham Elvidge et al., 2015; Oram et al., 2017; Say et al., 2019). Upper tropospheric levels of these chlorinated VSLs are likely to ascend into the lower stratosphere. This is supported by satellite observations of unusually high levels of phosgene (COCl₂) in the stratosphere, a product gas of the photochemical decomposition of CH₂Cl₂, CHCl₃ and other gases (Harrison et al., 2019). Our study investigates this hypothesis with air samples collected via a high-altitude research aircraft in the ASMA region but at higher altitudes (up to 20 km), i.e. within the lower stratosphere where no *in situ* data exists for many ODSs.

2. Methods


2.1 Aircraft campaigns

Air samples were collected during two campaigns of the M55 Geophysika high-altitude research aircraft that were part of the StratoClim EU project (www.stratoclim.org). Sample

collection and measurement details such as coordinates, altitudes, pressures, potential temperatures, and mixing ratios of various trace gases can be found in the Supplement.

130 The first campaign took place over the Mediterranean in September 2016. The aim of this campaign was to measure the composition of the outflow from the ASMA. The monsoon circulation system has a large variability in its spatial extent and can reach from the Mediterranean and North-East Africa to East Asia (Annamalai and Slingo, 2001; Garny and Randel, 2013; Vogel et al., 2015; Pan et al., 2016). This campaign is referred to as AMO-16
135 (the Asian Monsoon Outflow 2016 campaign). During AMO-16, 24 air samples were collected during two flights (1st and 6th September 2016). The aircraft operated from Kalamata, Greece (37.1°, 22.0°) and samples were collected in the region of 33° - 41° N, 23° - 31° E, 10 - 20 km altitude (Figure 1).

140 The second campaign took place over the Indian subcontinent in July-August 2017. The aim of this campaign was to measure the composition of the ASMA. This campaign is referred to as AMA-17 (the Asian Monsoon Anticyclone 2017 campaign). During AMA-17, 94 samples were collected during 6 flights (27-Jul-2017, 29-Jul-2017, 31-Jul-17, 02-Aug-2017, 04-Aug-2017, and 06-Aug-2017). The campaign base was Tribhuvan International Airport at Kathmandu, Nepal (27.7°, 85.4°) and samples were collected in the region of 21° - 29° N, 79°
145 - 91° E, 10 - 20 km altitude (Figure 1). In addition to the flight samples, 9 air samples were collected at ground level, 2 samples at Kathmandu University and the rest at Kathmandu airport. Kathmandu is situated at an elevation of about 1400 m.



150 Figure 1. The latitude and longitude locations where air samples were collected on board the Geophysika research aircraft during the AMO-16 (red) and AMA-17 (blue) campaigns.

2.2 Analytical technique

155 Air samples were collected with the whole air sampler (WAS) of Utrecht University operated on board the Geophysika research aircraft (Kaiser et al., 2006; Cairo et al., 2010; Laube et al., 2010b). Ambient air was compressed into evacuated stainless-steel canisters (2 litre) using a metal bellows pump that has been previously shown to not impact trace gas mixing ratios (Laube et al., 2010b). In addition, for AMA-17 the internal surfaces of 30 canisters were passivated using a common passivation technique ('SilcoTM-treatment') to minimise the

160 breakdown of more reactive gases in the canisters between collection and analysis (25 filled on board the aircraft and 5 at ground level).

The samples were transported to the University of East Anglia (UEA) for analysis on a high sensitivity gas chromatograph - tri-sector mass spectrometer system (Waters AutoSpec GC-MS). The trace gases were cryogenically extracted and pre-concentrated. A full description of this system can be found in Laube et al. (2010a). The different compounds were separated on a GS-GasPro column (length ~ 50 m; ID: 0.32 mm). Additionally, all of the AMO-16 and some of the AMA-17 samples were reanalysed on a KCl-passivated CP-PLOT Al₂O₃ column (length: 50 m; ID: 0.32 mm) (Laube et al., 2016). During analysis on the Al₂O₃ column, an Ascarite (NaOH-coated silica) trap was used to remove carbon dioxide, which can distort or reduce the signal of a number of compounds. A full list of measured species and their calibration scales can be found in Table S1.

All samples were bracketed by measurements of a working standard (in AMO-16: AAL-071170, and in AMA-17: SX-3591). Calibrations of our target compounds in these working standards were in part provided by the National Oceanic and Atmospheric Administration's Global Monitoring Laboratory in Boulder, USA and complemented by UEA internal calibrations for some gases. UEA intercomparisons of these working standards with four other working standards, two of which had internal surfaces of passivated aluminium, were used to ensure that results were consistent over time and in relation to long-term tropospheric trends. It was confirmed that the mixing ratios of all compounds presented here remained constant in the two primary working standards within 3 standard deviations during the 2008-2017 period, except for methyl bromide (CH₃Br) in the SX-3591 standard. This is likely to be due to loss (absorption and/or breakdown) of CH₃Br on the internal walls of the cylinders and CH₃Br mixing ratios for SX-3591 were drift-corrected accordingly (see Supplement). The dry-air mole fractions (mixing ratios) were measured for 27 ODSs (Table S1), and the unit, parts per trillion (ppt), is used in this study as an equivalent to picomole per mole. Some additional non-ODSs were measured that are good tracers for stratospheric mean age-of-air calculations and can be used to derive tropospheric emissions, including sulphur hexafluoride (SF₆), perfluoroethane (C₂F₆), HFC-125 (CHF₂CF₃) and HFC-32 (CH₂F₂).

Due to potential loss during storage for some of the compounds of interest, the time between flask collection and measurement was kept as short as possible. During AMO-16, the time between collection and measurement was 14-19 days. During AMA-17, the time between collection and measurement was more variable between 2-8 weeks/16-54 days. Only CCl₄ (carbon tetrachloride, tetrachloromethane) and CH₃Cl (methyl chloride, chloromethane) were affected by the longer time delay in AMA-17 and for these compounds, the samples that were measured later and were not stored in SilcoTM-treated canisters have been removed from the analysis. CCl₄ had previously been found to be unstable in stainless steel canisters containing dry air over long periods (Laube et al., 2008, 2013). For more information on this see the supplementary material.

2.3 CLaMS backward trajectory calculations

200 There are multiple factors that influence the mixing ratios of VSLs in the UTLS: the emissions near the surface, the time taken for air to be transported from the surface to the UTLS (i.e. convection), and the mixing processes in the troposphere and the stratosphere. In

order to investigate the possible source regions of enhanced CH₂Cl₂ mixing ratios, 10-day backward trajectories were calculated for each sample collected during AMA-17 using the trajectory module of the Chemical Lagrangian Model of the Stratosphere (CLaMS; McKenna, 2002; Konopka et al., 2012; Pommrich et al., 2014 and references therein). Only trajectories for AMA-17 were calculated, as AMO-16 samples that were influenced by the ASMA have much longer transport times from CH₂Cl₂ sources in Asia to the location of the measurements at UTLS altitudes in the Mediterranean area. Therefore, they will have experienced more mixing and chemical break down and so will have a more diluted source signal than AMA-17 samples. This is in agreement with the lower mixing ratios of most compounds in AMO-16 in comparison to AMA-17. Additionally, only AMA-17 samples measured at potential temperatures of less than 390 K were used for backward trajectory calculations, i.e. 53 out of 94 samples, because we focus here on enhanced mixing ratios of CH₂Cl₂.

The CLaMS backward trajectory calculations are driven by horizontal winds from the high-resolution ERA5 reanalysis (Hersbach et al., 2020) recently released by the European Centre for Medium-Range Weather Forecasts (ECMWF). ERA5 provides data on a 0.3° × 0.3° horizontal grid every hour on 137 hybrid levels from the surface to 0.01 hPa. In general, this results in a much better representation of convective updrafts and tropical cyclones in ERA5 (e.g. Hoffmann et al., 2019, Li et al., 2020, Legras and Bucci, 2020) compared with the earlier ERA-Interim re-analysis (Dee et al., 2011). To derive the vertical velocities, we used diabatic heating rates including latent heat release (for details, see Ploeger et al., 2010).

The CLaMS model trajectory calculations were used to find the last location the air parcel was in the model boundary layer (trajectory end point). The model boundary layer was set to ≈2-3km above the surface following orography (for more details see Vogel et al., 2015). The location of trajectory end points provides an indication of the regions where surface sources last influenced the mixing ratios measured in the air samples. Each air sample typically took 2-3 minutes to be collected. To investigate the variability of the trajectories within the whole time interval of the measurements, ERA5-based CLaMS backward trajectories were calculated every second for the entire time interval over which an air sample was collected during the flight.

2.4 Equivalent chlorine (ECI)

Equivalent chlorine (ECI) is the sum of the mixing ratios of chlorine and bromine atoms from all halogen source gases; the bromine mixing ratios are multiplied by a weighting factor of 60 as bromine is about 60-65 times more effective at depleting ozone than chlorine (Daniel and Velders et al., 2006; Sinnhuber et al., 2009). The ECI calculation was used to investigate the eventual impact on stratospheric ozone of ODSs measured in the tropopause region of the ASMA in comparison to estimates of ECI based on measurements in other atmospheric regions.

2.5 Equivalent Effective Stratospheric Chlorine (EESC)

The Equivalent Effective Stratospheric Chlorine (EESC) is defined as the “chlorine-equivalent sum of chlorine and bromine derived from ODS tropospheric abundances,

245 weighted to reflect their expected depletion of stratospheric ozone.” (Engel and Rigby et al.,
2018). EESC – like the ECl – takes into account the sum of the mixing ratios of chlorine and
bromine atoms from all halogen source gases with bromine multiplied by 60. However,
EESC – in contrast to ECl – takes into account the effects of stratospheric transport and
250 chemistry on the amount of chlorine and bromine released from long-lived ODSs at a given
location and time.

As stratospheric circulation is slow, air sampled in the stratosphere may have entered it
several years ago. The ‘mean age-of-air’ is defined as the average amount of time an air
parcel has spent in the stratosphere. Inert compounds can be used as ‘age-of-air tracers’ to
calculate the ‘mean age-of-air’ of the air sampled in the stratosphere provided that there are
255 a) long-term measurements of their global tropospheric mixing ratios and b) these mixing
ratios have been monotonically increasing over time at sufficient rates. Our mean age
calculation also takes into account the underlying transit time distribution (the “age
spectrum”) using the parameterisation introduced by Engel et al. (2002).

The mean ages-of-air were calculated using ground-level background mixing ratio trends of
260 selected gases from 1978 to January 2018 from air samples collected at the Cape Grim,
Tasmania (40.7°S, 144.7°E) baseline station and analysed at the UEA (Laube et al., 2010a,
2013). These mixing ratio trends were shifted backwards in time by 6 months, which has
been proven to be a good proxy for air entering the stratosphere via the upper troposphere in
the tropics, provided the gas is inert enough in the troposphere, (i.e. no significant
265 decomposition on tropospheric transport time scales, Leedham Elvidge et al., 2018).
Commonly used age-of-air tracers include sulphur hexafluoride (SF₆) and carbon dioxide
(CO₂) (e.g. Volk et al., 1997; Andrews et al., 2001; Engel et al., 2002). However, recent
research has introduced other potential age tracers (Leedham Elvidge et al., 2018). We
compared three different age-of-air tracers: SF₆, C₂F₆, and HFC-125 in air samples collected
270 during AMO-16 and AMA-17. The mean ages of air were then calculated using the same
methods described in Leedham Elvidge et al. (2018). For more information see the
supplementary material.

For trace gases with stratospheric sinks such as ODSs, at mid-latitudes a mean age of 3 years
is generally used as a reference to estimate the EESC in that region (Engel and Rigby et al.,
275 2018). The mixing ratios of ODSs measured in the stratosphere at a mean age-of-air of 3
years, for example, would be roughly similar to their mixing ratios in the upper troposphere 3
years earlier, assuming no decomposition. As ODSs are at least partly broken down by strong
UV radiation and/or reaction with OH radicals and O(¹D) in the stratosphere, the mixing
ratios of these compounds are however impacted not just by the age of the air but also by
280 different reactions and reaction rates.

For long-lived ODSs, similarly time-shifted tropospheric trends from Cape Grim were
propagated into the stratosphere and mixing ratios assuming no decomposition were
calculated for the mean ages of air, with the latter based on the measured age-of-air tracers.
However, the actual mixing ratios measured in the samples were lower, indicating that there
285 was decomposition of these ODSs and mixing with “older” air masses. We used the
difference between the mixing ratios assuming no decomposition (given a particular mean
age-of-air) and the measured mixing ratios in AMO-16 and AMA-17 to calculate the fraction

of the ECI that had already been released for each compound. These measures are known as Fractional Release Factors (FRFs). The more long-lived a compound is, the less decomposition takes place and the smaller their FRF is at the same mean age-of-air. Importantly, we used an improved method of FRF calculation (Ostermöller et al., 2017), which takes into account the dependency of the FRFs on the stratospheric lifetime of an ODS. The FRF uncertainties were calculated using the stratospheric measurement precisions, a proxy of the interhemispheric gradient in the troposphere, and the uncertainty in the tropospheric time series based on ground-based measurements.

For some of the FRF-mean-age correlations, the FRF was negative around mean ages of air of zero years. This was because our samples were collected above a polluted continental region so in some cases the mixing ratios of the remote tropospheric monitoring stations were lower than the mixing ratios observed in the ASMA tropopause region. This would cause our FRFs to have a low bias. Therefore, FRFs at a mean age-of-air of zero years were calculated from the respective FRF-mean age correlations (using a second order polynomial fit function) and when significantly different from zero, they were subtracted from the FRFs to calculate corrected FRFs. The FRFs of CFC-115, CFC-114a, HCFC-133a, CH₃CCl₃, Halon-1202, CH₃Cl, and CH₃Br were shifted in this way. This correction method relies on the assumption that the trace gases measured in 3-year old air originated from a similarly polluted air mass below the tropopause, whereas it could in principal have come from somewhere else. The fact that we do observe continuity throughout the profile gives some confidence, but we note that this introduces an additional uncertainty to these FRFs, which we therefore refer to as regional FRFs. The FRFs in general agree with other FRFs calculated in the literature. For more information see the supplementary material.

For the VSLs, it was not possible to use tropospheric trends at background stations to calculate FRFs because of significant loss and variability in the troposphere and the wide range of mixing ratios observed near the tropopause. Therefore, simplified FRFs were calculated by comparing the highest and lowest measured mixing ratios in the tropopause region (355 K – 375 K) to the measured mixing ratios above 375 K in the campaigns.

For AMA-17, measurements of CFC-114, CFC-114a, CFC-13, CH₃Cl, Halon-1202 and CCl₄ mixing ratios were only available for some of the samples. Therefore, the correlations of the available mixing ratios and FRFs with those of CFC-11 were used to estimate the values for the missing samples. Due to the set-up of the instrument some compounds were not measured or exhibited poor quality and were therefore excluded from the EESC estimate: CHClCCl₂, CCl₂CCl₂, CHBr₃, CH₂BrCl, CHBr₂Cl, CHBrCl₂, CFC-112, CFC-112a, HCFC-124, and Halon-2402.

To summarise, EESC depends on three factors: the mixing ratios of ODSs in the troposphere, the transport from the troposphere to the stratosphere (mean age-of-air) and breakdown of ODSs in the stratosphere (FRFs). The EESC was calculated and compared to other estimates in the literature in order to assess the overall impact on stratospheric ozone from ODSs in both campaigns.

3. Results and discussion

3.1 Long-lived ozone-depleting substances




Figure 2. Mixing ratios of CFC-12, Halon-1211, HCFC-22 and methyl bromide (CH_3Br) as a function of potential temperature (a pseudo-vertical coordinate) for AMO-16, AMA-17 and the ground samples collected during AMA-17. The dotted vertical line indicates the globally averaged background mixing ratios of CFC-12 and Halon-1211 for July-August 2017 from NOAA ground-based data (Table S1). Two background levels are indicated for HCFC-22 and CH_3Br , the lower one is from measurements at American Samoa (14.2°S , 170.6°W) and the higher one from Mauna Loa (19.5°N , 155.6°W) (Table S1).

CFC-12 and Halon-1211 in Figure 2 illustrate the observed distributions of long-lived ODSs that have been phased out under the Montreal Protocol. These two gases have only relatively small emission sources to the atmosphere in the monsoon input region and therefore their mixing ratios in the tropopause are not very variable. The ground-based AMA-17 samples and the lower aircraft measurements generally agree with the expected low variability as well as the NOAA background mixing ratios (Figure 2, top). This pattern is similar to what is found for many other long-lived ODSs in this study (see supplement). Conversely, HCFC-22 and CH_3Br are significantly enhanced above expected background mixing ratios, on average by 5 % for HCFC-22 and 25 % for CH_3Br (Figure 2, bottom). Enhancements are also observed for HCFC-141b (6 %) and CH_3Cl (14 %) (see supplement). These enhancements indicate continued large emissions of these compounds in the monsoon input region.

Among long-lived gases, Halon-1211 has a relatively short atmospheric lifetime of about 16 years. Halon-1211 mixing ratios in the tropopause region are similar to mixing ratios at the surface and NOAA ground-based mixing ratios indicating relatively little photolytic

destruction of Halon-1211 in the troposphere in the ASMA region, as expected (Figure 2). At higher potential temperatures and therefore altitudes, in the stratosphere, the mixing ratios decrease due to photochemical degradation and mixing with other stratospheric air masses (Figure 2). Halon-1211 has a similar vertical profile to CFC-11, which also begins to decrease around the same potential temperature (see supplement). The beginning of the decrease of Halon-1211 and CFC-11 mixing ratios therefore indicates that air has entered the lower stratosphere and constrains the location of the chemical tropopause. Other data (Brunamonti et al., 2018; Vogel et al., 2019) indicate that there is a transition region between the troposphere and the stratosphere at the top of the anticyclone. For our purposes using the ODS-based one is most appropriate as we are looking at similar gases. In this study we define the location of the chemical transition layer between the troposphere and stratosphere to be the region of 355 – 375 K (i.e. just below the level at which Halon-1211 starts to decrease) which is represented by the horizontal grey bar in Figure 2. The location of the tropopause region is important because of the slow ascent rates representing a transport barrier that limits particularly the contribution of VSLs (due to their quicker chemical decomposition) to ozone depletion in the stratosphere.

3.2 Very short-lived substances




Figure 3. Mixing ratios for dichloromethane, 1,2-dichloroethane, chloroform and dibromomethane (CH₂Br₂) as a function of potential temperature for AMO-16, AMA-17 and the AMA-17 ground samples. The red shaded region indicates the estimated mixing ratios at the Level of Zero Radiative Heating (LZRH) for 2013-2014 from the WMO 2018 report (Engel and Rigby et al., 2018; Table S1).

The mixing ratios of the three major chlorinated VSLs, CH₂Cl₂, CH₂ClCH₂Cl and CHCl₃, and one brominated VSL, dibromomethane (CH₂Br₂) are shown in Figure 3 as a function of

potential temperature. The chlorinated VSLs show a large range of mixing ratios both in the ground-based samples (similar to many regions in Asia, e.g., Oram et al., 2017; Fang et al., 2019) and in the tropopause region indicating continued large emissions, which reach the tropopause (Figure 3). At higher potential temperatures, their mixing ratios decrease rapidly as in the stratosphere they are broken down, predominantly by hydroxyl radical oxidation (Figure 3).

From WMO 2018 (Engel and Rigby et al., 2018), estimated mixing ratios of these compounds in the tropical tropopause region (as derived from multiple aircraft campaigns in 2013-2014) are represented by the red-shaded areas indicated at the tropopause levels (Table S1, Figure 3). The chlorinated VSLs in the tropopause region during AMA-17 are enhanced above the results from measurements in the tropical tropopause layer in 2013-2014 (Engel and Rigby et al., 2018) on average by 124 % for CH_2Cl_2 , by 76 % for $\text{CH}_2\text{ClCH}_2\text{Cl}$, and by 136 % for CHCl_3 . Chlorine loading in the tropopause region from these three VSLs is 169-393 ppt in AMA-17 compared to the 89-132 ppt stated in the WMO 2018 report (Table 1). The global mean CH_2Cl_2 mixing ratio measured at NOAA increased by about 10 % from 2013-2014 to 2017, but this increase is not large enough to explain the difference. The mixing ratios are still enhanced above these data presented in WMO 2018 until about 400 K potential temperature. These estimates are for two different stratospheric entry points. The estimates in this study are for the ASMA tropopause region where there is a co-location of efficient vertical lifting from Earth's boundary layer to UTLS altitudes, with surface industrial emissions of chlorinated VSLs in Asian countries. Whereas, the WMO 2018 estimates are largely based on measurements from the tropical tropopause region, which is also influenced by strong convection but likely represents an average of regions with smaller underlying surface emissions of chlorinated VSLs than the ASMA region. CH_2Br_2 , in contrast to the three chlorinated VSLs, has mostly natural oceanic sources, and its mixing ratios measured near the tropopause agree within the range of these data presented in WMO 2018 (Figure 3).

Samples collected during AMO-16 often have lower mixing ratios for the chlorinated VSLs than the samples from AMA-17 in the aircraft measurements at lower levels of potential temperature. This is possibly because during AMO-16 in general two different types of air masses were sampled: mid-latitude extra-tropical air with lower mixing ratios and monsoon outflow-influenced air with higher mixing ratios, but not necessarily Asian sources.

Mixing ratios of these three chlorinated VSLs are well correlated, even in the more source-influenced upper tropospheric region. For AMA-17, the three aircraft samples with the highest CH_2Cl_2 mixing ratios are also the samples with the three highest $\text{CH}_2\text{ClCH}_2\text{Cl}$ and CHCl_3 mixing ratios. For AMO-16, the sample with the highest CH_2Cl_2 mixing ratio also has the highest $\text{CH}_2\text{ClCH}_2\text{Cl}$ and CHCl_3 mixing ratios. There is one outlier in AMA-17 which has high CH_2Cl_2 mixing ratios but is not particularly enhanced for the other compounds (CH_2Cl_2 107 ppt, 305 K, see Supplement). This suggests that the enhanced CH_2Cl_2 in this sample originates from a different source. In addition, an influence from the tropospheric trends could play a role in these correlations as both CHCl_3 and CH_2Cl_2 have been increasing in recent years (Engel and Rigby et al., 2018). Given that the samples with high mixing ratios of these three chlorinated VSLs are all at low altitudes and contain relatively young air this influence can however be neglected here.

In addition to having strong correlations with each other, the chlorinated VSLs are also correlated with other compounds. When investigating only the samples in the tropopause region using Spearman correlations (i.e. assuming a monotonic but not necessarily linear relationship) and excluding the outlier sample, CH_2Cl_2 has the strongest Spearman's correlation coefficients with CHCl_3 ($R = 0.87$), HFC-32 ($R = 0.87$), HCFC-22 ($R = 0.87$), $\text{CH}_2\text{ClCH}_2\text{Cl}$ ($R = 0.75$) and HFC-125 ($R = 0.74$). $\text{CH}_2\text{ClCH}_2\text{Cl}$ has the strongest positive spearman's correlations with CH_2Cl_2 ($R = 0.75$), HCFC-133a ($R = 0.71$), CHCl_3 ($R = 0.70$), CCl_4 ($R = 0.66$) and HFC-23 ($R = 0.66$). The compounds that are well correlated with CHCl_3 are CH_2Cl_2 ($R = 0.87$), HFC-32 ($R = 0.82$), HFC-125 ($R = 0.72$), SF_6 ($R = 0.71$) and $\text{CH}_2\text{ClCH}_2\text{Cl}$ ($R = 0.70$) (see supplement). These correlations are all significant ($p < 0.01$) and all of the compounds have known strong industrial emissions in East and South-East Asia (Kim et al., 2010; Vollmer et al., 2015; Fang et al., 2018, 2019; Lunt et al., 2018); suggesting the sources of the halogenated compounds in these air samples are from continental industrial areas and indicating that they are either emitted from co-located sources or co-produced. The strongest correlation is between CH_2Cl_2 and CHCl_3 and these two compounds are known to be co-produced in large quantities in East Asia (Oram et al., 2017).

3.3 CLaMS backward trajectory calculations

The possible source regions in the Earth's model boundary layer of CH_2Cl_2 in the AMA-17 air samples were investigated using 10-day CLaMS backward trajectories driven by ERA5. Figure 4 shows the end points of the 10-day backward trajectories, the location where the air was last in the model boundary layer i.e. 2-3 km above the surface. It should be noted that of all the 10-day backward trajectories calculated, only 9 % of them have end points in the model boundary layer; the rest of the trajectories end in the free troposphere and stratosphere. Therefore, the trajectory end points reflect only the contributions of the very young air masses (<10 days) to the composition of the air samples.

Our analysis shows that clusters of trajectory end points in the model boundary layer exist that correspond to air samples with high CH_2Cl_2 mixing ratios (red colours) located in east Asia (mainly in China) and at the north-western flank of the Tibetan Plateau (northwest India and north Pakistan) and for a trajectory length of 15 days also in northern India (not shown here). Red points are only found if CH_2Cl_2 -polluted air masses in the lower troposphere experienced strong vertical uplift by convection. Therefore, the location of red points indicates possible source regions in South Asia with enhanced CH_2Cl_2 emissions. Many of the possible source regions of high CH_2Cl_2 mixing ratios found in our trajectory calculations are in China which is in agreement with sources of anthropogenic CH_2Cl_2 emissions in Asia (e.g. Feng et al., 2018).

Clusters of trajectory end points in the model boundary layer with low CH_2Cl_2 mixing ratios (blue colours) are mainly found in the region of the Tibetan Plateau, in the Bay of Bengal and in surrounding continental area, indicating the absence of significant anthropogenic CH_2Cl_2 sources in these regions. For a trajectory length of 15 days, sources of low CH_2Cl_2 mixing ratios are also found in Southeast Asia and over the western Pacific (not shown here). In general, the longer the trajectories the wider the trajectory endpoints are distributed over South Asia caused by the trajectory dispersion depending on the trajectory length.




Figure 4. Location of end points of the CLaMS 10-day backward trajectories, initialized for each air sample (with 1-sec resolution). Shown are only trajectory end points for air samples collected below 390 K potential temperature during AMA-17, where the trajectory reaches the model boundary layer (i.e. 2-3 km above the Earth's surface following orography), within 10 days. The colour of the data points indicates the CH_2Cl_2 mixing ratios in the air sample collected at UTLS altitudes.

3.4 Equivalent chlorine (ECI)

After analysing the individual VSLs, their contribution to the total equivalent chlorine (ECI) in the tropopause region was investigated. Table 1 shows two estimates of ECI, one estimate
 455 derived using a similar methodology to that used in the WMO 2018 report (Engel and Rigby et al., 2018) and the other estimate based on the air samples from AMA-17.

In the first method for calculating ECI, the tropospheric reference mixing ratios of the long-lived compounds were, as explained in Section 2.5, taken from NOAA
 (https://www.esrl.noaa.gov/gmd/dv/ftpdata.html). Similar to previous approaches (Laube et
 460 al., 2010b, 2013) either global mixing ratios were used, or in the case of compounds with significant tropospheric sinks such as HCFCs and methyl halides, data from the NOAA ground-based measurement sites at Mauna Loa (19.5°N, 155.6°W) and American Samoa (14.2°S, 170.6°W) in July-August 2017, the same time period as AMA-17, were used. As
 465 HCFCs and methyl halides have significant tropospheric sinks, using ground-based measurements can lead to a slight overestimation of their mixing ratios in the tropopause region. UEA measurements of air samples collected at Cape Grim, Tasmania, in early 2018 and shifted back in time by 6 months were used to complement this approach (see also Table S1). For the VSLs that are broken down rapidly in the troposphere and for which it is
 470 therefore much harder to estimate the proportion transported to the tropopause, the global estimates of tropospheric reference mixing ratios come from the WMO 2018 report for the

Level of Zero Radiative Heating (LZRH) (Engel and Rigby et al., 2018). These mixing ratios of VSLs in the WMO 2018 report were based on results from the CAST (Coordinated Airborne Studies in the Tropics), CONTRAST (Convective Transport of Active Species in the Tropics) and ATTREX (Airborne Tropical Tropopause Experiment) aircraft campaigns that took place in the tropical west Pacific in 2013 and 2014 (Harris et al., 2017; Jensen et al., 2017; Pan et al., 2017). They were used to investigate the differences between the tropics and the ASMA input of VSLs in the lower stratosphere. It should be noted that mixing ratios of some of these compounds have changed over time and some of the reported values were on different calibration scales to the measurements in this study which may cause small differences.

There are some halogenated gases that were considered in Engel and Rigby et al. (2018) but were not available from the AMA-17 samples: CHClCCl_2 , CCl_2CCl_2 , CHBr_3 , CH_2BrCl , CHBr_2Cl , CHBrCl_2 , CFC-112, CFC-112a, HCFC-124, and Halon-2402. For these compounds, we used the same mixing ratios from the Engel and Rigby et al. (2018) in both estimates to make them comparable. In total, these compounds contribute 79-253 ppt to ECl. See the supplement for more information.

For the global estimates of ECl, a potential temperature range of 355-365 K was used. This is the range of the tropical tropopause layer or the LZRH in the WMO 2018 report (Engel and Rigby et al., 2018). Air at this altitude is likely to continue to ascend into the lower tropical stratosphere by radiative heating. Some studies have suggested that the LZRH may be pushed upwards by the ASMA as air from the troposphere rises, creating an elevated tropopause (e.g. Dunkerton, 1995; Highwood and Hoskins, 1998; Dethof et al., 1999; Ploeger et al., 2017). To take this into account, we used an increased potential temperature range of 355-375 K for estimates based on the AMA-17 air samples, which also agrees with our own observations of gases with exclusively stratospheric sinks such as Halon-1211 (Figure 2). There were 27 samples collected in this range during AMA-17 and these were used to estimate the ECl. An ECl estimate for AMO-16 was not calculated as no air samples were collected in the tropopause region in this campaign (and much fewer samples were collected in general). Note that we did not consider ECl contributions from the breakdown products of VSLs in this study, so our ECl can be considered a lower limit of the total chlorine and bromine entering the stratosphere via the ASMA.

Table 1 shows that the total ECl from the AMA-17 aircraft campaign is higher than the total ECl from the global estimates based on Cape Grim, NOAA and WMO mixing ratios. However, the AMA-17-based ECl also has a wider range and the lower end of the range overlaps with the higher end of the range for the global estimate (Table 1). So there is not a significant difference between the overall ECl range estimates. This is due to many of our samples in the tropopause having higher mixing ratios of ODSs than in the previous global estimates while some samples also show mixing ratios in the range of the global estimates (Sections 3.1, 3.2).

The compounds contributing to the ECl were divided into four categories: chlorinated VSLs, brominated VSLs (CH_2Br_2), long-lived chlorine and long-lived bromine (Table 1). Both estimates for CH_2Br_2 have large ranges that overlap. The long-lived chlorine in the AMA-17 samples is slightly higher than the global estimate. This is because mixing ratios of

the CFCs and carbon tetrachloride (CCl_4) agree within our range, methyl chloroform (CH_3CCl_3) is slightly smaller in the AMA-17 estimate, and HCFCs and methyl chloride (CH_3Cl) are slightly larger. This is qualitatively in line with the findings of Umezawa et al. (2014) who found enhancements of CH_3Cl in the ASMA region at lower altitudes that may be related to biomass and biofuel emissions. For the long-lived bromine species, the AMA-17 estimate is also slightly higher than the global estimate. Here, the halon mixing ratios agree with the AMA-17 estimate, so the higher long-lived bromine estimate for AMA-17 is almost completely due to methyl bromide (CH_3Br) (6.5 - 6.8 ppt vs 7.4 - 9.1 ppt) indicating larger sources of methyl bromide in the monsoon input region.

Mixing ratios of chlorinated VSLs are higher in the AMA-17 estimate than in the estimate based on WMO 2018 mixing ratios at the LZRH. In the latter estimate chlorinated VSLs contribute about 2 to 3 % to the total ECl entering the stratosphere whereas in the AMA-17 estimate this is higher, 4 to 8 % of the total ECl. We note that the AMA-17 estimate is for a particular region and a particular time of year when there is likely to be a very high injection rate during the ASMA (Leedham Elvidge et al., 2015). The WMO 2018 values incorporate measurements from various campaigns and are more representative of the input from the tropics. In both estimates chlorinated VSLs make up a relatively small fraction of the total ECl in the tropopause region.

The influence of the ASMA on the stratosphere is largest in the Northern Hemisphere, there is a smaller influence on air in the tropical stratosphere and an even smaller influence on the stratosphere in the Southern Hemisphere (Ploeger et al., 2017; Yan et al., 2019). Using the results in Ploeger et al. (2017) we calculated that, averaged over the whole year, 5 % of the air in the Northern Hemispheric lower stratosphere comes from the ASMA. If we assume that 5 % of the additional equivalent chlorine from the AMA-17 estimate ends up in the lower stratosphere of the Northern Hemisphere, this translates to an additional 0.3-34.9 ppt of ECl from all measured compounds, of which 1.6-15.2 ppt are from chlorinated VSLs. Total tropospheric chlorine from controlled substances has been decreasing by 12.7 ± 0.92 ppt Cl yr^{-1} while that from uncontrolled substances (CH_3Cl and chlorinated VSLs) has been increasing by 8.3 ± 4.9 ppt Cl yr^{-1} , leading to an overall decrease of 4.4 ± 4.1 ppt Cl yr^{-1} between 2012 and 2016 (Engel and Rigby et al., 2018). The decrease in chlorine from controlled ODSs has been partially offset by the increase in CH_3Cl and chlorinated VSLs (Engel and Rigby et al., 2018). These current annual decreases are up to six times smaller than the additional 0.3-34.9 ppt we calculated from chlorinated VSLs. Although these elevated mixing ratios are only observed in one part of the atmosphere it indicates that the influence of the enhanced mixing ratios of the methyl halides as well as the chlorinated VSLs may be significant.

Table 1. Comparison of the global estimate of equivalent chlorine (ECl) based on Cape Grim, NOAA and WMO mixing ratios (see Table S1) and the regional estimate based on the air samples from AMA-17. ¹For the compounds that were not measured in this study the WMO 2018 reported values were used in both estimates. ²ECl excluding compounds that were not measured in this study.

	Global estimate ECl (ppt) 355 K – 365 K	AMA-17 ECl (ppt) 355 K – 375 K
Chlorinated VSLs	89-132 (2-3%)	163-393 (4-8%)

CH ₂ Cl ₂	59-89	130-272
CHCl ₃	19-24	24-74
CH ₂ ClCH ₂ Cl	10-19	9-47
Brominated VSLs	71-118	58-92
CH ₂ Br ₂	71-118	58-92
Long-lived chlorine	3159-3186	3188-3356
CFCs	1960	1939-1997
HCFCs	310	317-343
CH ₃ CCl ₃	6.5	4.8-6.0
CCl ₄	321	321-338
CH ₃ Cl	558-586	603-669
Halon-1211	3.4	3.4-3.5
Long-lived bromine	789-811	842-963
Halons	402	398-414
CH ₃ Br	387-409	445-549
Estimated (not measured)¹	79-253	79-253
Total Equivalent Chlorine (ECI)	4186-4499 (4107-4246)²	4331-5057 (4252-4804)²

3.5 Equivalent Effective Stratospheric Chlorine (EESC)

EESC was calculated for a mean age-of-air of 3 years for AMO-16 as this is the mean age-of-air usually used in the literature to approximate mid-latitudinal ozone depletion. EESC was also calculated for a mean age-of-air of 2.4 years for both campaigns in order to compare the campaigns. The EESC was calculated using tropospheric trends from the same NOAA and Cape Grim data sets mentioned above (Section 3.4), mixing ratios from the aircraft samples to calculate time-independent FRFs (see Supplement) and an adjusted mean age-of-air (see Supplement).

For both campaigns, EESC was calculated using the ‘relevant age EESC’ method from Engel et al. (2018). The ‘relevant age EESC’ is a refinement in the calculation method of EESC. A previous method (Newman et al., 2007) was based on the assumption assumes that the age spectrum for an inert species is representative of the age spectrum of a chemically reactive species. This is not the case as the average age-of-air for source gases that have been dissociated in the stratosphere is longer than the average age of inert tracers in the same stratospheric location (Plumb et al., 1999; Engel et al., 2018). Younger air contains more reactive species than older air so the organic fraction of a chemically active species is largely determined by the fraction of the air with shorter transit times (Plumb et al., 1999). To take into account the interaction between chemical loss and transit time, the relevant age EESC method uses new time-independent FRFs (Ostermüller et al., 2017) and an age spectrum weighted by chemical loss (Engel et al., 2018). The EESC was also calculated based on the

older mean-age based method (Newman et al., 2007). This gives similar results with a quantitative comparison being available in the supplement.

Additionally, each estimate has a range because of the relatively short lifetime of CH₃Br and CH₃Cl. The EESC contribution of these two compounds was calculated twice: using again tropospheric trends from Mauna Loa and American Samoa. The lower end of the range is based on Mauna Loa trends and the higher end of the range is based on American Samoa trends. Neither of these sites are ideal for estimates of the amount of shorter-lived ODSs reaching the tropical upper troposphere, but they are the closest ground-based approximations available and yield similar FRFs (see supplement).

The EESCs from this study were calculated using the same method used in Engel et al. (2018). However, there are some differences between our EESCs and the Engel et al. (2018) estimates:

1. The shifting of FRFs explained in section 2.5 means that for some species the mixing ratios observed near the tropopause were significantly lower (CFC-115, CH₃CCl₃) or higher (CFC-114a, HCFC-133a, Halon-1202, CH₃Cl, CH₃Br) than expected from surface-based trends. This is likely due to the special nature of this region and season. The adjusted FRFs compare well with those from other studies in line with the expectation that they are dominated by common sinks and global tropospheric trends. It does however not take into account that the actual amount found near the tropopause is different, which is important for deriving a regional EESC. The difference was therefore calculated and added to all our regional EESC estimates.
2. We included some minor compounds that Engel et al. (2018) did not include (CFC-113a, HCFC-133a and CFC-13). They also did not include CFC-114a but their CFC-114 mixing ratio is a combination of CFC-114 and CFC-114a.
3. They included Halon-2402 and due to small contamination problems, this was not possible here.

Therefore, to compare our EESC to Engel et al. (2018) we recalculated our EESC excluding CFC-113a, HCFC-133a and CFC-13. Including these species makes very little difference, it adds only ~1 ppt to the EESCs, well within the uncertainty of the estimate. We determined the contribution of Halon-2402 to our EESC estimates using the 2017 mean tropospheric mixing ratio and the FRF given in Engel et al. (2018), to arrive at an additional contribution of ~33 ppt from this molecule.

Table 2. Regional Equivalent Effective Stratospheric Chlorine (EESC) estimates from the AMO-16 and AMA-17 campaigns calculated using the relevant age. Also shown is the global EESC estimate from Engel et al. (2018).

Campaign	EESC ^a	EESC+VSLS ^b
AMA-17 (age-of-air 2.4 years)	1630 – 1650	1804 – 2087
AMO-16 (age-of-air 2.4 years)	1483 – 1495	1604 – 1692
AMO-16 (age-of-air 3 years)	1861 – 1872	1988 – 2075
Engel et al. (2018) (age-of-air 3 years)	1646 (in 2017)	–

^a EESC including CFC-13, CFC-113a, HCFC-133a and Halon-2402.

^b EESC^a with an additional contribution from very short-lived substances (VSLSs).

For AMO-16 the EESC is 12 % higher (1861-1872 ppt) than the Engel et al. (2018) estimate for an age-of-air of 3 years (1646 ppt). At an age-of-air of 2.4 years EESC is lower in AMO-16 than in AMA-17 by about 135 ppt. This is likely because during AMO-16 different types of air masses were sampled: outflow from the Asian monsoon and Northern Hemispheric extra-tropical air. It implies that, if it was measured, the EESC above the ASMA at an age-of-air of 3 years may be much higher than the Engel et al. (2018) EESC estimate.

To investigate the impact of VSLs on lower stratospheric ozone depletion, an EESC contribution for these substances was calculated. The EESC contribution at 2.4 years mean age-of-air from our VSLs is between 121-197 ppt based on the air samples collected during AMO-16 and 174-437 ppt based on the air samples collected during AMA-17 (Figure 5).

This is about 8-26% of the EESC from long-lived compounds (Figure 5). This inclusion increases both the EESC and its range (Table 2) and indicates that the contribution of VSLs is relatively larger in the northern hemispheric lower stratosphere where there is less fractional release of reactive chlorine from the longer-lived species. Moreover, the total EESC contribution from these VSLs may be higher than our estimate as some fraction of the halogenated ‘product gases’ from their tropospheric breakdown may also be injected in to the stratosphere (‘product gas injection’ is discussed in Engel and Rigby et al., 2018). There are also other VSLs not measured in this study that may contribute (e.g. CHBr_3 , $\text{CH}_3(\text{CH}_2)_2\text{Br}$ and other anthropogenic chlorocarbons).

Assuming a linear trend, EESC at mid-latitudes has decreased by 14-16 ppt per year from its peak values to 2017 (Engel and Rigby et al., 2018). This gives some context to the size of the contribution from VSLs observed in measurements from both campaigns compared to the size of the decreasing trend.




Figure 5. EESC at mean age-of-air of 2.4 years for the AMO-16 and AMA-17 campaigns and at mean age-of-air of 3 years for the AMO-16 campaign, calculated using relevant age and mean age, showing the contributions from long-lived and very short-lived chlorine and bromine. AoA = age-of-air.

4. Conclusions

Air samples collected in the upper troposphere and lower stratosphere on board the high-altitude Geophysika research aircraft in the vicinity of and within the ASMA were found to have substantially elevated mixing ratios of chlorinated VSLs compared to WMO 2018 estimates (Engel and Rigby et al., 2018). For example, CH₂Cl₂ mixing ratios were 30-44 ppt in the tropical tropopause region in 2013-2014, as stated in the WMO 2018 report, but 65-136 ppt in the ASMA tropopause region in 2017, based on the AMA-17 samples. This is likely largely due to the rapid transport of emissions of these substances from South Asia to the UTLS via the ASMA and higher-than-global emission rates in this region. We show that VSLs are transported irreversibly from the ASMA into the lower stratosphere where they will contribute to ozone depletion. The contribution of chlorinated VSLs is significantly higher in this region than that reported in the WMO Scientific Assessment of Ozone Depletion (Engel and Rigby et al., 2018) in terms of Equivalent Chlorine (ECI) in the tropopause region (89-132 ppt vs 169-393 ppt). These additional VSL contributions increase the estimate of Equivalent Effective Stratospheric Chlorine (EESC) in the Northern Hemisphere extra-tropical lower stratosphere by 8-26%.

Our estimates of ECI and EESC from long-lived species in the stratosphere in this region are generally larger than global average values based on Engel et al. (2018). For example, EESC

at 3 years in Engel et al. (2018) is 1646 ppt, whereas the AMO-16-based range is 1861-1872 ppt. ECl from long-lived species is 3947-3997 ppt in the global estimate and 4031-4319 ppt in the AMA-17 estimate. The ASMA region generally has large continental emissions and more input into the stratosphere than many other regions. This explains why the AMA-17 estimate is larger than the global average from WMO as it is impacted by regional emissions to a larger degree than global mean estimates.

A previous study found similar enhanced mixing ratios of CH_2Cl_2 , $\text{CH}_2\text{ClCH}_2\text{Cl}$ and CHCl_3 in the upper tropopause region during boreal winter over South-East Asia, indicating that rapid upward transport also occurs in the winter monsoon (Oram et al., 2017). The additional input of chlorine into the stratosphere from these sources could delay the recovery of the ozone layer if the growth rate of chlorinated VSLs persists in the future (Hossaini et al., 2017). Since our observations are both spatially and temporally limited the quantification of this possible future impact is beyond the scope of this study.

However, when combining the differences between the most recent WMO ECl estimates (Engel and Rigby, 2018) and our AMA-17-based ECl (Table 1) with the estimate from Ploeger et al. (2017) of the monsoon contributing an annual average of about 5 % to northern hemispheric lower stratospheric air, we derive a difference ranging from 0.3-34.9 ppt of ECl from all measured compounds, much of it in the form of chlorinated VSLs. Depending on the inter-annual monsoon variability as well as how much ECl enters this part of the stratosphere via the tropical west pacific and through extratropical isentropic transport, the available levels of chlorine and bromine might thus be substantially higher than the global average derived from global ground-based measurements. In summary, this work highlights the importance of both the ASMA as a fast transport mechanism in an important ODS emission region, and the role of chlorinated VSLs for stratospheric ozone, particularly in the northern extra-tropical lower stratosphere during the Northern Hemisphere summer and autumn.

Acknowledgements

We are grateful for the contributions of Michel Bolder, Carina van der Veen (both Utrecht University), and the Geophysika team (sample collection and campaign organisation), Guus Velders and Andreas Engel (EESC comparisons), Elinor Tuffnell (data processing) and the NOAA Global Monitoring Division (surface halocarbon data). We thank the staff at the Cape Grim station and at CSIRO GASLAB Aspendale for collecting and maintaining the Cape Grim air archive and preparing the UEA flasks and sub-samples. We also acknowledge CSIRO and the Bureau of Meteorology for funding these activities. The StratoClim flights were funded by the European Commission (FP7 project Stratoclim-603557, www.stratoclim.org). This work was supported by the Natural Environment Research Council through the EnvEast Doctoral Training Partnership (grant number NE/L002582/1) as well as the ERC project EXC³ITE (EXC3ITE-678904-ERC-2015-STG). Johannes C. Laube received funding from the UK Natural Environment Research Council (Research Fellowship NE/I021918/1). The data are available from the Zenodo repository, [doi:10.5281/zenodo.3836630](https://doi.org/10.5281/zenodo.3836630).

Author contributions

Karina Adcock wrote the article and did most of the data analysis. Bärbel Vogel produced the output from the CLaMS chemistry-transport model, which Karina Adcock then compared to the atmospheric observations. Geoffrey Lee, Johannes Laube, and Karina Adcock measured and analysed the air samples used in this study. Additionally, Johannes Laube and Karina Adcock worked together to calculate the Fractional Release Factors, Equivalent Chlorine and Equivalent Effective Stratospheric Chlorine. Johannes Laube and William Sturges coordinated activities for the University of East Anglia (UEA) related to the StratoClim aircraft campaigns. Paul Fraser, Ray Langenfelds and David Oram organised the collection of samples from the Cape Grim Monitoring Station. Bradley Hall and Stephen Montzka were involved in producing the NOAA atmospheric observations that were used in this study. Fred Stroh coordinated the aircraft campaigns while Thomas Röckmann coordinated the operation a whole air sampler on the research aircraft to collect the air samples used in this study. In addition, all of the co-authors contributed comments and suggestions for editing this work.

Conflict of interest declaration

The authors declare no conflict of interest.

References

- Andrews, A. E., Boering, K. A., Daube, B. C., Wofsy, S. C., Loewenstein, M., Jost, H., Podolske, J. R., Webster, C. R., Herman, R. L., Scott, D. C., Flesch, G. J., Moyer, E. J., Elkins, J. W., Dutton, G. S., Hurst, D. F., Moore, F. L., Ray, E. A., Romashkin, P. A. and Strahan, S. E.: Mean ages of stratospheric air derived from in situ observations of CO₂, CH₄, and N₂O, *J. Geophys. Res. Atmos.*, 106(D23), 32295–32314, doi:10.1029/2001JD000465, 2001.
- Annamalai, H. and Slingo, J. M.: Active / break cycles: diagnosis of the intraseasonal variability of the Asian Summer Monsoon, *Clim. Dyn.*, 18(1–2), 85–102, doi:10.1007/s003820100161, 2001.
- Ball, W. T., Alsing, J., Mortlock, D. J., Stachelin, J., Haigh, J. D., Peter, T., Tummon, F., Stübi, R., Stenke, A., Anderson, J., Bourassa, A., Davis, S. M., Degenstein, D., Frith, S., Froidevaux, L., Roth, C., Sofieva, V., Wang, R., Wild, J., Yu, P., Ziemke, J. R. and Rozanov, E. V.: Evidence for a continuous decline in lower stratospheric ozone offsetting ozone layer recovery, *Atmos. Chem. Phys.*, 18(2), 1379–1394, doi:10.5194/acp-18-1379-2018, 2018.
- Brioude, J., Portmann, R. W., Daniel, J. S., Cooper, O. R., Frost, G. J., Rosenlof, K. H., Granier, C., Ravishankara, A. R., Montzka, S. A. and Stohl, A.: Variations in ozone depletion potentials of very short-lived substances with season and emission region, *Geophys. Res. Lett.*, 37(19), 3–7, doi:10.1029/2010GL044856, 2010.
- Brunamonti, S., Jorge, T., Oelsner, P., Hanumanthu, S., Singh, B. B., Kumar, K. R., Sonbawne, S., Meier, S., Singh, D., Wienhold, F. G., Luo, B. P., Böttcher, M., Poltera, Y., Jauhiainen, H., Kayastha, R., Dirksen, R., Naja, M., Rex, M., Fadnavis, S. and Peter, T.: Balloon-borne measurements of temperature, water vapor, ozone and aerosol backscatter at the southern slopes of the Himalayas during StratoClim 2016-2017, *Atmos. Chem. Phys. Discuss.*, 1–38, doi:10.5194/acp-2018-222, 2018.

- 740 Cairo, F., Pommereau, J. P., Law, K. S., Schlager, H., Garnier, A., Fierli, F., Ern, M.,
Streibel, M., Arabas, S., Borrmann, S., Berthelier, J. J., Blom, C., Christensen, T., D'Amato,
F., Di Donfrancesco, G., Deshler, T., Diedhiou, A., Durr, G., Engels, O., Goutail, F.,
Harris, N. R. P., Kerstel, E. R. T., Khaykin, S., Konopka, P., Kylling, A., Larsen, N., Lebel,
T., Liu, X., MacKenzie, A. R., Nielsen, J., Oulanowski, A., Parker, D. J., Pelon, J., Polcher,
J., Pyle, J. A., Ravegnani, F., Rivi re, E. D., Robinson, A. D., R ckmann, T., Schiller, C.,
745 Sim es, F., Stefanutti, L., Stroh, F., Some, L., Siegmund, P., Sitnikov, N., Vernier, J. P.,
Volk, C. M., Voigt, C., Von Hobe, M., Viciani, S. and Yushkov, V.: An introduction to the
SCOUT-AMMA stratospheric aircraft, balloons and sondes campaign in West Africa, August
2006: Rationale and roadmap, *Atmos. Chem. Phys.*, 10(5), 2237–2256, doi:10.5194/acp-10-
2237-2010, 2010.
- 750 Chipperfield, M. P., Dhomse, S., Hossaini, R., Feng, W., Santee, M. L., Weber, M., Burrows,
J. P., Wild, J. D., Loyola, D. and Coldewey-Egbers, M.: On the Cause of Recent Variations in
Lower Stratospheric Ozone, *Geophys. Res. Lett.*, 45(11), 5718–5726,
doi:10.1029/2018GL078071, 2018.
- 755 Claxton, T., Hossaini, R., Wilson, C., Montzka, S. A., Chipperfield, M. P., Wild, O.,
Bednarz, E. M., Carpenter, L. J., Andrews, S. J., Hackenberg, S. C., M hle, J., Oram, D.,
Park, S., Park, M. K., Atlas, E., Navarro, M., Schauffler, S., Sherry, D., Vollmer, M., Schuck,
T., Engel, A., Krummel, P. B., Maione, M., Arduini, J., Saito, T., Yokouchi, Y., O'Doherty,
S., Young, D. and Lunder, C.: A Synthesis Inversion to Constrain Global Emissions of Two
Very Short Lived Chlorocarbons: Dichloromethane, and Perchloroethylene, *J. Geophys. Res.*
760 *Atmos.*, 125(12), doi:10.1029/2019JD031818, 2020.
- Daniel, J. S., Velders, G. J. M., (Lead Authors), Douglass, A. R., Forster, P. M. D.,
Hauglustaine, D. A., Isaksen, I. S. A., Kuijpers, L. J. M., McCulloch, A. and Wallington, T.
J.: Chapter 8 Halocarbon Scenarios, Ozone Depletion Potentials, and Global Warming
Potentials, in *Scientific Assessment of Ozone Depletion: 2006, Global Ozone Research and*
765 *Monitoring Project-Report No. 50*, World Meteorological Organization, Geneva,
Switzerland., 2007.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U.,
Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L.,
Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L.,
770 Healy, S. B., Hersbach, H., H lm, E. V., Isaksen, L., K llberg, P., K hler, M., Matricardi,
M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay,
P., Tavolato, C., Th paut, J.-N. and Vitart, F.: The ERA-Interim reanalysis: configuration and
performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137(656), 553–597,
doi:10.1002/qj.828, 2011.
- 775 Dethof, A., O'Neill, A., Slingo, J. M. and Smit, H. G. J.: A mechanism for moistening the
lower stratosphere involving the Asian summer monsoon. [online] Available from:
<https://rmets.onlinelibrary.wiley.com/doi/pdf/10.1002/qj.1999.49712555602> (Accessed 18
April 2019), 1999.
- 780 Dunkerton, T. J.: Evidence of meridional motion in the summer lower stratosphere adjacent
to monsoon regions. [online] Available from:
<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/95JD01263> (Accessed 18 April
2019), 1995.

- Engel, A., Strunk, M., Müller, M., Haase, H., Poss, C., Levin, I. and Schmidt, U.: Temporal development of total chlorine in the high-latitude stratosphere based on reference
785 distributions of mean age derived from CO₂ and SF₆, *J. Geophys. Res.*, 107(D12), ACH 1-1-ACH 1-11, doi:10.1029/2001JD000584, 2002.
- Engel, A., Bönisch, H., Ostermöller, J., Chipperfield, M. P., Dhomse, S. and Jöckel, P.: A refined method for calculating equivalent effective stratospheric chlorine, *Atmos. Chem. Phys.*, 18(2), 601–619, doi:10.5194/acp-18-601-2018, 2018a.
- 790 Engel, A., Rigby, M., (Lead Authors), Burkholder, J. B., Fernandez, R. P., Froidevaux, L., Hall, B. D., Hossaini, R., Saito, T., Vollmer, M. K. and Yao, B.: Chapter 1 Update on Ozone-Depleting Substances (ODSs) and Other Gases of Interest to the Montreal Protocol, in *Scientific Assessment of Ozone Depletion: 2018*, Global Ozone Research and Monitoring Project–Report No. 58, World Meteorological Organization, Geneva, Switzerland., 2018b.
- 795 Fang, X., Ravishankara, A. R., Velders, G. J. M., Molina, M. J., Su, S., Zhang, J., Hu, J. and Prinn, R. G.: Changes in Emissions of Ozone-Depleting Substances from China Due to Implementation of the Montreal Protocol, *Environ. Sci. Technol.*, 52(19), 11359–11366, doi:10.1021/acs.est.8b01280, 2018.
- 800 Fang, X., Park, S., Saito, T., Tunnicliffe, R., Ganesan, A. L., Rigby, M., Li, S., Yokouchi, Y., Fraser, P. J., Harth, C. M., Krummel, P. B., Mühle, J., O'Doherty, S., Salameh, P. K., Simmonds, P. G., Weiss, R. F., Young, D., Lunt, M. F., Manning, A. J., Gressent, A. and Prinn, R. G.: Rapid increase in ozone-depleting chloroform emissions from China, *Nat. Geosci.*, 12(2), 89–93, doi:10.1038/s41561-018-0278-2, 2019.
- 805 Feng, Y., Bie, P., Wang, Z., Wang, L. and Zhang, J.: Bottom-up anthropogenic dichloromethane emission estimates from China for the period 2005–2016 and predictions of future emissions, *Atmos. Environ.*, 186(May), 241–247, doi:10.1016/j.atmosenv.2018.05.039, 2018.
- 810 Garny, H. and Randel, W. J.: Dynamic variability of the Asian monsoon anticyclone observed in potential vorticity and correlations with tracer distributions, *J. Geophys. Res. Atmos.*, 118(24), 13421–13433, doi:10.1002/2013JD020908, 2013.
- Garny, H. and Randel, W. J.: Transport pathways from the Asian monsoon anticyclone to the stratosphere, *Atmos. Chem. Phys.*, 16, 2703–2718, doi:10.5194/acp-16-2703-2016, 2016.
- 815 Harris, N. R. P., Carpenter, L. J., Lee, J. D., Vaughan, G., Filus, M. T., Jones, R. L., Ouyang, B., Pyle, J. A., Robinson, A. D., Andrews, S. J., Lewis, A. C., Minaeian, J., Vaughan, A., Dorsey, J. R., Gallagher, M. W., Le Breton, M., Newton, R., Percival, C. J., Ricketts, H. M. A., Bauguutte, S. J. B., Nott, G. J., Wellpott, A., Ashfold, M. J., Flemming, J., Butler, R., Palmer, P. I., Kaye, P. H., Stopford, C., Chemel, C., Boesch, H., Humpage, N., Vick, A., MacKenzie, A. R., Hyde, R., Angelov, P., Meneguz, E. and Manning, A. J.: Coordinated airborne studies in the tropics (CAST), *Bull. Am. Meteorol. Soc.*, 98(1), 145–
820 162, doi:10.1175/BAMS-D-14-00290.1, 2017.
- Harrison, J. J., Chipperfield, M. P., Hossaini, R., Boone, C. D., Dhomse, S., Feng, W. and Bernath, P. F.: Phosgene in the Upper Troposphere and Lower Stratosphere: A Marker for Product Gas Injection Due to Chlorine-Containing Very Short Lived Substances, *Geophys. Res. Lett.*, 46(2), 1032–1039, doi:10.1029/2018GL079784, 2019.
- 825 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X.,

- Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146, 1999–2049, doi.org/10.1002/qj.3803, 2020.
- Highwood, E. J. and Hoskins, B. J.: The tropical tropopause, *Q. J. R. Meteorol. Soc.*, 124(549), 1579–1604, 1998.
- Hossaini, R., Chipperfield, M. P., Montzka, S. A., Rap, A., Dhomse, S. and Feng, W.: Efficiency of short-lived halogens at influencing climate through depletion of stratospheric ozone, *Nat. Geosci.*, 8(3), 186, doi:10.1038/NGEO2363, 2015.
- Hossaini, R., Patra, P. K., Leeson, A. A., Krysztofiak, G., Abraham, N. L., Andrews, S. J., Archibald, A. T., Aschmann, J., Lidster, T., Wilson, C., Ziska, F. and Hossaini, R.: A multi-model intercomparison of halogenated very short-lived substances (TransCom-VSLS): linking oceanic emissions and tropospheric transport for a reconciled estimate of the stratospheric source gas injection of bromine, *Atmos. Chem. Phys.*, 16, 9163–9187, doi:10.5194/acp-16-9163-2016, 2016.
- Hossaini, R., Chipperfield, M. P., Montzka, S. A., Leeson, A. A., Dhomse, S. S. and Pyle, J. A.: The increasing threat to stratospheric ozone from dichloromethane, *Nat. Commun.*, 8, 15962, doi:10.1038/ncomms15962, 2017.
- Jensen, E. J., Pfister, L., Jordan, D. E., Bui, T. V., Ueyama, R., Singh, H. B., Thornberry, T. D., Rollins, A. W., Gao, R.-S., Fahey, D. W., Rosenlof, K. H., Elkins, J. W., Diskin, G. S., DiGangi, J. P., Lawson, R. P., Woods, S., Atlas, E. L., Navarro Rodriguez, M. A., Wofsy, S. C., Pittman, J., Bardeen, C. G., Toon, O. B., Kindel, B. C., Newman, P. A., McGill, M. J., Hlavka, D. L., Lait, L. R., Schoeberl, M. R., Bergman, J. W., Selkirk, H. B., Alexander, M. J., Kim, J.-E., Lim, B. H., Stutz, J. and Pfeilsticker, K.: The NASA Airborne Tropical Tropopause Experiment: High-Altitude Aircraft Measurements in the Tropical Western Pacific, *Bull. Am. Meteorol. Soc.*, 98(1), 129–143, doi:10.1175/BAMS-D-14-00263.1, 2017.
- Kaiser, J., Engel, A., Borchers, R. and Röckmann, T.: Probing stratospheric transport and chemistry with new balloon and aircraft observations of the meridional and vertical N₂O isotope distribution, *Atmos. Chem. Phys.*, 6(11), 3535–3556, doi:10.5194/acp-6-3535-2006, 2006.
- Kim, J., Li, S., Kim, K. R., Stohl, A., Mühle, J., Kim, S. K., Park, M. K., Kang, D. J., Lee, G., Harth, C. M., Salameh, P. K. and Weiss, R. F.: Regional atmospheric emissions determined from measurements at Jeju Island, Korea: Halogenated compounds from China, *Geophys. Res. Lett.*, 37(12), 2007–2011, doi:10.1029/2010GL043263, 2010.
- Ko, M. K. W., Newman, P. A., Reimann, S., and Strahan, S. E. (Eds.): SPARC Report on Lifetimes of Stratospheric Ozone-Depleting Substances, Their Replacements, and Related Species, SPARC Report No. 6, WCRP-15/2013, SPARC Office. [online] Available from: <http://www.sparc-climate.org/publications/sparc-reports/sparc-report-no-6/>, 2013.
- Konopka, P., Ploeger, F. and Müller, R.: Entropy-based and static stability based Lagrangian model grids, in: *Lagrangian Modeling of the Atmosphere*, Geophysical Monograph Series, 200, AGU Chapman Conference on Advances in Lagrangian Modeling of the Atmosphere, Grindelwald, Switzerland, 9–14 Octo., 2012.

- Laube, J. C., Engel, A., Bönisch, H., Möbius, T., Worton, D. R., Sturges, W. T., Grunow, K. and Schmidt, U.: Contribution of very short-lived organic substances to stratospheric chlorine and bromine in the tropics – a case study, *Atmos. Chem. Phys.*, 8(23), 7325–7334, doi:10.5194/acp-8-7325-2008, 2008.
- 875 Laube, J. C., Martinerie, P., Witrant, E., Blunier, T., Schwander, J., Brenninkmeijer, C. A. M., Schuck, T. J., Bolder, M., Röckmann, T., Van Der Veen, C., Bönisch, H., Engel, A., Mills, G. P., Newland, M. J., Oram, D. E., Reeves, C. E. and Sturges, W. T.: Accelerating growth of HFC-227ea (1,1,1,2,3,3,3- heptafluoropropane) in the atmosphere, *Atmos. Chem. Phys.*, 10(13), 5903–5910, doi:10.5194/acp-10-5903-2010, 2010a.
- 880 Laube, J. C., Engel, A., Bönisch, H., Möbius, T., Sturges, W. T., Braß, M. and Röckmann, T.: Fractional release factors of long-lived halogenated organic compounds in the tropical stratosphere, *Atmos. Chem. Phys.*, 10(3), 1093–1103, doi:10.5194/acp-10-1093-2010, 2010b.
- Laube, J. C., Keil, A., Bönisch, H., Engel, A., Röckmann, T., Volk, C. M. and Sturges, W. T.: Observation-based assessment of stratospheric fractional release, lifetimes, and ozone
885 depletion potentials of ten important source gases, *Atmos. Chem. Phys.*, 13(5), 2779–2791, doi:10.5194/acp-13-2779-2013, 2013.
- Laube, J. C., Mohd Hanif, N., Martinerie, P., Gallacher, E., Fraser, P. J., Langenfelds, R., Brenninkmeijer, C. A. M., Schwander, J., Witrant, E., Wang, J.-L., Ou-Yang, C.-F., Gooch, L. J., Reeves, C. E., Sturges, W. T. and Oram, D. E.: Tropospheric observations of CFC-114
890 and CFC-114a with a focus on long-term trends and emissions, *Atmos. Chem. Phys.*, 16(23), 15347–15358, doi:10.5194/acp-16-15347-2016, 2016.
- Leedham Elvidge, E., Bönisch, H., Brenninkmeijer, C. A. M., Engel, A., Fraser, P. J., Gallacher, E., Langenfelds, R., Mühle, J., Oram, D. E., Ray, E. A., Ridley, A. R., Röckmann, T., Sturges, W. T., Weiss, R. F. and Laube, J. C.: Evaluation of stratospheric age of air from
895 CF₄, C₂F₆, C₃F₈ CHF₃, HFC-125, HFC-227ea and SF₆; implications for the calculations of halocarbon lifetimes, fractional release factor, *Atmos. Chem. Phys.*, 18(5), 3369–3385, doi:10.5194/acp-18-3369-2018, 2018.
- Leedham Elvidge, E. C., Oram, D. E., Laube, J. C., Baker, A. K., Montzka, S. A., Humphrey, S., O’Sullivan, D. A. and Brenninkmeijer, C. A. M.: Increasing concentrations of
900 dichloromethane, CH₂Cl₂, inferred from CARIBIC air samples collected 1998-2012, *Atmos. Chem. Phys.*, 15(4), 1939–1958, doi:10.5194/acp-15-1939-2015, 2015. Legras, B. and Bucci, S.: Confinement of air in the Asian monsoon anticyclone and pathways of convective air to the stratosphere during the summer season, *Atmos. Chem. Phys.*, 20, 11045–11064, doi:10.5194/acp-20-11045-2020, 2020.
- 905
- Li, D., Vogel, B., Müller, R., Bian, J., Günther, G., Ploeger, F., Li, Q., Zhang, J., Bai, Z., Vömel, H. and Riese, M.: Dehydration and low ozone in the tropopause layer over the Asian monsoon caused by tropical cyclones: Lagrangian transport calculations using ERA-Interim and ERA5 reanalysis data, *Atmos. Chem. Phys.*, 20(7), 4133–4152, doi:10.5194/acp-20-
910 4133-2020, 2020.
- Li, Q., Jiang, J. H., Wu, D. L., Read, W. G., Livesey, N. J., Waters, J. W., Zhang, Y., Wang, B., Filipiak, M. J., Davis, C. P., Turquety, S., Wu, S., Park, R. J., Yantosca, R. M. and Jacob, D. J.: Convective outflow of South Asian pollution: A global CTM simulation compared with EOS MLS observations, *Geophys. Res. Lett.*, 32(14), 1–4, doi:10.1029/2005GL022762,

915 2005.

Lunt, M. F., Park, S., Li, S., Henne, S., Manning, A. J., Ganesan, A. L., Simpson, I. J., Blake, D. R., Liang, Q., O'Doherty, S., Harth, C. M., Mühle, J., Salameh, P. K., Weiss, R. F., Krummel, P. B., Fraser, P. J., Prinn, R. G., Reimann, S. and Rigby, M.: Continued Emissions of the Ozone-Depleting Substance Carbon Tetrachloride From Eastern Asia, *Geophys. Res. Lett.*, 45(20), 11,423–11,430, doi:10.1029/2018GL079500, 2018.

920 McCulloch, A.: Dichloromethane in the environment, A note prepared for the European Chlorinated Solvents Association (ECSA) and the Halogenated Solvents Industry Alliance (HSIA), 9th November 2017. [online] Available from: [http://www.chlorinated-solvents.eu/images/Documents/Newsroom/Dichloromethane paper.pdf](http://www.chlorinated-solvents.eu/images/Documents/Newsroom/Dichloromethane%20paper.pdf) (Accessed 19 February 2018), 2017.

925 McKenna, D. S.: A new Chemical Lagrangian Model of the Stratosphere (CLaMS) 1. Formulation of advection and mixing, *J. Geophys. Res.*, 107(D16), Pages ACH 15-1-ACH 15-15, doi:10.1029/2000JD000114, 2002.

930 Newman, P. A., Daniel, J. S., Waugh, D. W. and Nash, E. R.: A new formulation of equivalent effective stratospheric chlorine (EESC), *Atmos. Chem. Phys.*, 7(17), 4537–4552, doi:10.5194/acp-7-4537-2007, 2007.

935 Oram, D. E., Ashfold, M. J., Laube, J. C., Gooch, L. J., Humphrey, S., Sturges, W. T., Leedham-Elvidge, E., Forster, G. L., Harris, N. R. P., Mead, M. I., Samah, A. A., Phang, S. M., Ou-Yang, C.-F., Lin, N.-H., Wang, J.-L., Baker, A. K., Brenninkmeijer, C. A. M. and Sherry, D.: A growing threat to the ozone layer from short-lived anthropogenic chlorocarbons, *Atmos. Chem. Phys.*, 17(19), 11929–11941, doi:10.5194/acp-17-11929-2017, 2017.

940 Orbe, C., Waugh, D. W. and Newman, P. A.: Air-mass origin in the tropical lower stratosphere: The influence of Asian boundary layer air, *Geophys. Res. Lett.*, 42(10), 4240–4248, doi:10.1002/2015GL063937, 2015.

Ostermöller, J., Bönisch, H., Jöckel, P. and Engel, A.: A new time-independent formulation of fractional release, *Atmos. Chem. Phys.*, 17(6), 3785–3797, doi:10.5194/acp-17-3785-2017, 2017.

945 Pan, L. L., Honomichl, S. B., Kinnison, D. E., Abalos, M., Randel, W. J., Bergman, J. W. and Bian, J.: Transport of chemical tracers from the boundary layer to stratosphere associated with the dynamics of the asian summer monsoon, *J. Geophys. Res. Atmos.*, 121(23), 14,159–14,174, doi:10.1002/2016JD025616, 2016.

950 Pan, L. L., Atlas, E. L., Salawitch, R. J., Honomichl, S. B., Bresch, J. F., Randel, W. J., Apel, E. C., Hornbrook, R. S., Weinheimer, A. J., Anderson, D. C., Andrews, S. J., Baidar, S., Beaton, S. P., Campos, T. L., Carpenter, L. J., Chen, D., Dix, B., Donets, V., Hall, S. R., Hanisco, T. F., Homeyer, C. R., Huey, L. G., Jensen, J. B., Kaser, L., Kinnison, D. E., Koenig, T. K., Lamarque, J. F., Liu, C., Luo, J., Luo, Z. J., Montzka, D. D., Nicely, J. M., Pierce, R. B., Riemer, D. D., Robinson, T., Romashkin, P., Saiz-Lopez, A., Schauffler, S., Shieh, O., Stell, M. H., Ullmann, K., Vaughan, G., Volkamer, R. and Wolfe, G.: The convective transport of active species in the tropics (Contrast) experiment, *Bull. Am. Meteorol. Soc.*, 98(1), 106–128, doi:10.1175/BAMS-D-14-00272.1, 2017.

955 Park, M., Randel, W. J., Gettelman, A., Massie, S. T. and Jiang, J. H.: Transport above the

- Asian summer monsoon anticyclone inferred from Aura Microwave Limb Sounder tracers, *J. Geophys. Res. Atmos.*, 112(16), 1–13, doi:10.1029/2006JD008294, 2007.
- 960 Park, M., Randel, W. J., Emmons, L. K., Bernath, P. F., Walker, K. A. and Boone, C. D.: Chemical isolation in the Asian monsoon anticyclone observed in Atmospheric Chemistry Experiment (ACE-FTS) data, *Atmos. Chem. Phys.*, 8(3), 757–764, doi:10.5194/acp-8-757-2008, 2008.
- 965 Park, M., Randel, W. J., Emmons, L. K. and Livesey, N. J.: Transport pathways of carbon monoxide in the Asian summer monsoon diagnosed from Model of Ozone and Related Tracers (MOZART), *J. Geophys. Res. Atmos.*, 114(8), 1–11, doi:10.1029/2008JD010621, 2009.
- 970 Ploeger, F., Konopka, P., Günther, G., Grooß, J.-U. and Müller, R.: Impact of the vertical velocity scheme on modeling transport in the tropical tropopause layer, *J. Geophys. Res.*, 115(D3), D03301, doi:10.1029/2009JD012023, 2010.
- 975 Ploeger, F., Gottschling, C., Griessbach, S., Grooß, J.-U., Guenther, G., Konopka, P., Müller, R., Riese, M., Stroh, F., Tao, M., Ungermann, J., Vogel, B. and von Hobe, M.: A potential vorticity-based determination of the transport barrier in the Asian summer monsoon anticyclone, *Atmos. Chem. Phys.*, 15(22), 13145–13159, doi:10.5194/acp-15-13145-2015, 2015.
- Ploeger, F., Konopka, P., Walker, K. and Riese, M.: Quantifying pollution transport from the Asian monsoon anticyclone into the lower stratosphere, *Atmos. Chem. Phys.*, 17(11), 7055–7066, doi:10.5194/acp-17-7055-2017, 2017.
- 980 Plumb, I. C., Vohralik, P. F. and Ryan, K. R.: Normalization of correlations for atmospheric species with chemical loss, *J. Geophys. Res. Atmos.*, 104(D9), 11723–11732, doi:10.1029/1999JD900014, 1999.
- 985 Pommrich, R., Müller, R., Grooß, J.-U., Konopka, P., Ploeger, F., Vogel, B., Tao, M., Hoppe, C. M., Günther, G., Spelten, N., Hoffmann, L., Pumphrey, H.-C., Viciani, S., D’amato, F., Volk, C. M., Hoor, P., Schlager, H. and Riese, M.: Tropical troposphere to stratosphere transport of carbon monoxide and long-lived trace species in the Chemical Lagrangian Model of the Stratosphere (CLaMS), *Geosci. Model Dev.*, 7, 2895–2916, doi:10.5194/gmd-7-2895-2014, 2014.
- 990 Randel, W. J. and Park, M.: Deep convective influence on the Asian summer monsoon anticyclone and associated tracer variability observed with Atmospheric Infrared Sounder (AIRS), *J. Geophys. Res. Atmos.*, 111(12), 1–13, doi:10.1029/2005JD006490, 2006.
- Randel, W. J., Park, M., Emmons, L., Kinnison, D., Bernath, P., Walker, K. A., Boone, C. and Pumphrey, H.: Asian monsoon transport of pollution to the stratosphere, *Science*, 328(5978), 611–613, doi:10.1126/science.1182274, 2010.
- 995 Say, D., Ganesan, A. L., Lunt, M. F., Rigby, M., O’Doherty, S., Harth, C., Manning, A. J., Krummel, P. B. and Bauguitte, S.: Emissions of halocarbons from India inferred through atmospheric measurements, *Atmos. Chem. Phys.*, 19(15), 9865–9885, doi:10.5194/acp-19-9865-2019, 2019.
- 1000 Sinnhuber, B. M., Sheode, N., Sinnhuber, M., Chipperfield, M. P. and Feng, W.: The contribution of anthropogenic bromine emissions to past stratospheric ozone trends: A modelling study, *Atmos. Chem. Phys.*, 9(8), 2863–2871, doi:10.5194/acp-9-2863-2009, 2009.

2009.

Tissier, A.-S. and Legras, B.: Convective sources of trajectories traversing the tropical tropopause layer, *Atmos. Chem. Phys.*, 16, 3383–3398, doi:10.5194/acp-16-3383-2016, 2016.

- 1005 Umezawa, T., Baker, A. K., Oram, D., Sauvage, C., O’Sullivan, D., Rauthe-Schöch, A., Montzka, S. A., Zahn, A. and Brenninkmeijer, C. A. M.: Methyl chloride in the upper troposphere observed by the CARIBIC passenger aircraft observatory: Large-scale distributions and Asian summer monsoon outflow, *J. Geophys. Res. Atmos.*, 119(9), 5542–5558, doi:10.1002/2013JD021396, 2014.
- 1010 Velders, G. J. M. and Daniel, J. S.: Uncertainty analysis of projections of ozone-depleting substances: Mixing ratios, EESC, ODPs, and GWPs, *Atmos. Chem. Phys.*, 14(6), 2757–2776, doi:10.5194/acp-14-2757-2014, 2014.
- 1015 Vogel, B., Günther, G., Müller, R., Grooß, J.-U., Hoor, P., Krämer, M., Müller, S., Zahn, A. and Riese, M.: Fast transport from Southeast Asia boundary layer sources to northern Europe: rapid uplift in typhoons and eastward eddy shedding of the Asian monsoon anticyclone, *Atmos. Chem. Phys.*, 14(23), 12745–12762, doi:10.5194/acp-14-12745-2014, 2014.
- Vogel, B., Günther, G., Müller, R., Grooß, J.-U. and Riese, M.: Impact of different Asian source regions on the composition of the Asian monsoon anticyclone and of the extratropical lowermost stratosphere, *Atmos. Chem. Phys.*, 15(23), 13699–13716, doi:10.5194/acp-15-13699-2015, 2015.
- 1020 Vogel, B., Günther, G., Müller, R., Grooß, J.-U., Afchine, A., Bozem, H., Hoor, P., Krämer, M., Müller, S., Riese, M., Rolf, C., Spelten, N., Stiller, G. P., Ungermann, J. and Zahn, A.: Long-range transport pathways of tropospheric source gases originating in Asia into the northern lower stratosphere during the Asian monsoon season 2012, *Atmos. Chem. Phys.*, 16(23), 15301–15325, doi:10.5194/acp-16-15301-2016, 2016.
- 1025 Vogel, B., Müller, R., Günther, G., Spang, R., Hanumanthu, S., Li, D., Riese, M. and Stiller, G. P.: Lagrangian simulations of the transport of young air masses to the top of the Asian monsoon anticyclone and into the tropical pipe, *Atmos. Chem. Phys.*, 19, 6007–6034, doi:10.5194/acp-19-6007-2019, 2019.
- 1030 Volk, C. M., Elkins, J. W., Fahey, D. W., Dutton, G. S., Gilligan, J. M., Loewenstein, M., Podolske, J. R., Chan, K. R. and Gunson, M. R.: Evaluation of source gas lifetimes from stratospheric observations, *J. Geophys. Res. Atmos.*, 102(D21), 25543–25564, doi:10.1029/97JD02215, 1997.
- 1035 Vollmer, M. K., Rigby, M., Laube, J. C., Henne, S., Siek, T., Gooch, L. J., Wenger, A., Young, D., Steele, L. P., Langenfelds, L., Brenninkmeijer, C. A. M., Wang, J., Wyss, S. A., Hill, M., Oram, D. E., Krummel, P. B., Schoenenberger, F., Zellweger, C., Fraser, P. J., William, T., Doherty, S. O. and Reimann, S.: Abrupt reversal in emissions and atmospheric abundance of HCFC-133a (CF₃CH₂Cl), *Geophys. Res. Lett.*, 42, 8702–8710, doi:10.1002/2015GL065846, 2015.
- 1040 Yan, X., Konopka, P., Ploeger, F., Wright, J. S., Wright, J. S., Müller, R. and Riese, M.: The efficiency of transport into the stratosphere via the Asian and North American summer monsoon circulations, *Atmos. Chem. Phys.*, 19(24), 15629–15649, doi:10.5194/acp-19-15629-2019, 2019.