

Hidden Spin-Isospin Exchange Symmetry

Dean Lee¹, Scott Bogner¹, B. Alex Brown¹, Serdar Elhatisari², Evgeny Epelbaum³, Heiko Hergert¹,
Morten Hjorth-Jensen^{1,4}, Hermann Krebs³, Ning Li⁵, Bing-Nan Lu⁶, and Ulf-G. Meißner^{7,8,9}

¹*Facility for Rare Isotope Beams and Department of Physics and Astronomy, Michigan State University,
East Lansing, Michigan 48824, USA*

²*Faculty of Engineering, Karamanoglu Mehmetbey University, Karaman 70100, Turkey*

³*Ruhr-Universität Bochum, Fakultät für Physik und Astronomie, Institut für Theoretische Physik II, D-44780 Bochum, Germany*

⁴*Department of Physics and Center for Computing in Science Education, University of Oslo, N-0316 Oslo, Norway*

⁵*Sun Yat-sen University, School of Physics, 135 Xingang Road, Haizhu District, Guangzhou, Guangdong Province, 510000, China*

⁶*China Academy of Engineering Physics, Graduate School, Building 8, No. 10 Xi'er Road, ZPark II,
Haidian District, Beijing, 100193, China*

⁷*Helmholtz-Institut für Strahlen- und Kernphysik and Bethe Center for Theoretical Physics, Universität Bonn,
D-53115 Bonn, Germany*

⁸*Institute for Advanced Simulation, Institut für Kernphysik, and Jülich Center for Hadron Physics,
Forschungszentrum Jülich, D-52425 Jülich, Germany*

⁹*Tbilisi State University, 0186 Tbilisi, Georgia*



(Received 9 November 2020; revised 24 April 2021; accepted 23 June 2021; published 2 August 2021)

The strong interactions among nucleons have an approximate spin-isospin exchange symmetry that arises from the properties of quantum chromodynamics in the limit of many colors, N_c . However this large- N_c symmetry is well hidden and reveals itself only when averaging over intrinsic spin orientations. Furthermore, the symmetry is obscured unless the momentum resolution scale is close to an optimal scale that we call $\Lambda_{\text{large-}N_c}$. We show that the large- N_c derivation requires a momentum resolution scale of $\Lambda_{\text{large-}N_c} \sim 500$ MeV. We derive a set of spin-isospin exchange sum rules and discuss implications for the spectrum of ^{30}P and applications to nuclear forces, nuclear structure calculations, and three-nucleon interactions.

DOI: [10.1103/PhysRevLett.127.062501](https://doi.org/10.1103/PhysRevLett.127.062501)

Quantum chromodynamics is the fundamental theory of the strong interactions. While the physical universe has $N_c = 3$ colors, it is useful to consider how nuclear physics might appear when N_c is large [1–7]. In the large- N_c picture, Kaplan, Savage, and Manohar [8,9] found that the leading nucleon-nucleon (NN) interaction between two nucleons can be written in the form

$$V_{\text{large-}N_c}^{2N} = V_C + \vec{\sigma}_1 \cdot \vec{\sigma}_2 \vec{\tau}_1 \cdot \vec{\tau}_2 W_S + S_{12} \vec{\tau}_1 \cdot \vec{\tau}_2 W_T + \dots, \quad (1)$$

where the ellipses refer to terms subleading in the large- N_c expansion, $\vec{\sigma}_{1,2}$ represent the Pauli matrices for intrinsic spins, $\vec{\tau}_{1,2}$ are the Pauli matrices for isospin, and S_{12} is the tensor operator $3\hat{r} \cdot \vec{\sigma}_1 \hat{r} \cdot \vec{\sigma}_2 - \vec{\sigma}_1 \cdot \vec{\sigma}_2$. Each of the scalar functions V_C , W_S , and W_T are local interactions, meaning that they depend only on the separation vector between the two nucleons and not their velocities. The strength of these leading interactions scales linearly with N_c , while all other terms scale as $1/N_c$ or smaller. The relative $1/N_c^2$ suppression of the subleading terms is fairly strong even for $N_c = 3$.

In the literature, the large- N_c limit is often linked with Wigner's approximate SU(4) spin-isospin symmetry [10]

where the four spin and isospin nucleon components transform as an SU(4) quartet. See for example, Refs. [8,11–16]. In the low-energy limit, nucleon-nucleon scattering is dominated by the two S -wave channels. Therefore, if one reduces Eq. (1) to only the S -wave channels, then the tensor interaction vanishes and we can replace $\vec{\sigma}_1 \cdot \vec{\sigma}_2 \vec{\tau}_1 \cdot \vec{\tau}_2$ by -3 . We therefore seem to derive the result that the low-energy S -wave interactions satisfy Wigner's SU(4) symmetry. On the other hand, the deuteron is bound by 2.22 MeV while the spin-singlet channel is unbound. The discrepancy seems larger than the predicted $1/N_c^2$ relative error of the large- N_c expansion.

As we will show in the following derivation, Eq. (1) is only valid when the momentum resolution scale is not too high or too low (where the precise meaning of high and low will become clear later). At the proper resolution scale we find that the large- N_c limit predicts a spin-isospin exchange symmetry that is satisfied at the $1/N_c^2$ level and provides a valuable guide for understanding nuclear interactions and nuclear structure.

The derivation of Eq. (1) can be sketched as follows. Let us assume that N_c is a large odd integer, $N_c = 2k + 1$. The nucleon wave function in its rest frame has a fairly simple

structure in the limit of nonrelativistic quarks [2,17]. The N_c quarks are completely antisymmetric with respect to color, and the spatial wave function is a simple Hartree product state. The spin and isospin content can be understood by dividing the quarks into two partitions of size $k+1$ and k . The partition with $k+1$ quarks has isospin $(k+1)/2$ and spin $(k+1)/2$. The partition with k quarks has isospin $k/2$ and spin $k/2$. These maximal values for isospin and spin mean that the wave functions for each partition are symmetric in both isospin and spin indices. The two partitions are combined together to have total isospin $1/2$ and total spin $1/2$, and we symmetrize over all ways to divide the quarks into the two partitions. The nucleon wave function has a spin-isospin exchange symmetry, which we can define at the quark level as the exchange of spin-up down quarks $d_\uparrow$ with spin-down up quarks $u_\downarrow$. At the nucleon level, the spin-isospin exchange symmetry corresponds with the exchange of spin-up neutrons $n_\uparrow$ with spin-down protons $p_\downarrow$.

If we now relax the condition that the quarks are nonrelativistic, the generalization to the relativistic mean-field picture becomes more complicated. However, the phenomenological success of the nonrelativistic quark model in predicting the large isovector part of the nucleon anomalous magnetic moment and nearly vanishing isoscalar part [18,19] suggests that the quantitative features of this description of the nucleon wave function remain valid. We note that there have been lattice QCD studies of baryon magnetic moments [20,21] and spin-flavor symmetries [22,23].

The expectation value of the quark bilinear operator $\bar{q}q$ for the nucleon will scale linearly with the number quarks in the nucleon and thus linearly with N_c . Similarly an operator such as $\bar{q}\sigma^a\tau^b q$ will also scale linearly with N_c . However, since the spin of the nucleon is $1/2$ and the isospin is $1/2$, the expectation values of $\bar{q}\vec{\sigma}q$ and $\bar{q}\vec{\tau}q$ do not grow with N_c . This argument no longer holds for $\bar{q}\vec{\sigma}q$ if the nucleon is moving with speed, $v > 0$. In that case the Pauli-Lubanski pseudovector W_μ will have a nonzero time component that scales as v times N_c .

Let us now consider two nucleons with relative velocities low enough so that we have an approximate Galilean invariance for the nucleon wave functions. Following Ref. [9], we can apply a multipole expansion to the effective nucleon-nucleon interactions that we can build from the leading quark operators $\bar{q}q$ and $\bar{q}\sigma^a\tau^b q$. That process leads to an effective Hamiltonian where the leading terms in the large- N_c limit are given in Eq. (1). This construction yields local interactions, and the connection with meson exchange interactions has been studied [24,25] as well as predictions for scattering observables [26].

We note that Eq. (1) is not renormalization group invariant, and we are somehow implicitly setting a preferred momentum resolution scale. How this happens can be understood as follows. Our simple description of the

nucleon wave function has corrections proportional to the square of the nucleon velocity, v^2 . We must control these high-energy modes in our effective theory in order for the derivation of Eq. (1) to hold with errors of size $1/N_c^2$. In order to restrict v^2 so that it is of size $1/N_c^2$ or smaller, we require that the momentum resolution scale Λ is proportional to the nucleon mass times a factor of $1/N_c$. Since the nucleon mass is proportional to N_c , the upper bound on Λ remains a constant independent of N_c . This is consistent with the discussion of nucleon momenta in Ref. [24]. We discuss the relevant energy and momentum scales further in the Supplemental Material[27].

While there are no problems with nucleon-nucleon scattering at low energies, the form of Eq. (1) will no longer hold if we also choose to lower the momentum resolution scale Λ so much that the distinction between the orbital angular momentum and intrinsic spin of the nucleon is not fully resolved. In that case the nucleon intrinsic spin should be viewed as an effective spin composed of both intrinsic spin and orbital angular momentum. In order to suppress these effects, it suffices that $1/\Lambda$ is comparable to the size of the nucleon or smaller. Since the size of the nucleon is independent of N_c , the resulting lower bound on Λ is also a constant independent of N_c .

We note that a similar phenomenon of hidden symmetry at low energies can also be seen in nucleon-nucleon scattering within the Kaplan-Savage-Wise (KSW) scheme introduced in Ref. [34,35]. In that case the difference between the two S -wave spin channels is magnified at low energies due to the different scattering lengths, one large and positive and the other large and negative.

The renormalization group dependence of the large- N_c approximation was first studied in Refs. [36–38]. The authors noticed that there was a momentum resolution scale at which the large- N_c constraints are satisfied far better than at other scales and speculated that it might be a numerical accident. The renormalization group analysis for pionless effective field theory was also considered in Ref. [39]. See also Ref. [40] for a discussion of emergent symmetries as arising from entanglement suppression. In this Letter we make the stronger statement that the leading large- N_c reduction of the nuclear interaction has error corrections of size $1/N_c^2$ only when the momentum resolution scale is near an optimal resolution scale that we call $\Lambda_{\text{large-}N_c}$. As we will see in the examples below, the optimal momentum resolution scale is $\Lambda_{\text{large-}N_c} \sim 500$ MeV, consistent with the momentum scale found in Ref. [36]. We posit that any effective field theory description of the nuclear interactions at momentum resolution scale $\Lambda_{\text{large-}N_c}$ with local interactions will satisfy Eq. (1) with corrections of size $1/N_c^2$. For interactions with a small amount of velocity dependence and therefore nonlocality, there will also be small corrections arising from the velocity dependence of the interactions. For some explicit examples demonstrating this behavior, see Fig. 1 of

TABLE I. Lattice chiral effective field theory coefficients of short-range interactions for the spin-singlet and spin-triplet S -wave and D -wave channels at lattice spacing $a = 1.32$ fm.

Channel, order	Coupling (l.u.)	Channel, order	Coupling (l.u.)
$^1S_0, Q^0$	1.45(5)	$^3S_1, Q^0$	1.56(3)
$^1S_0, Q^2$	-0.47(3)	$^3S_1, Q^2$	-0.53(1)
$^1S_0, Q^4$	0.129(13)	$^3S_1, Q^4$	0.115(4)
$^1D_2, Q^4$	-0.088(1)	$^3D_{\text{all}}, Q^4$	-0.070(2)

Ref. [41] for local chiral potentials at $N^2\text{LO}$ and Figs. 6,7 of Ref. [42] for the long-range part of chiral potentials up to $N^4\text{LO}$.

As an example, in Table I we show the S -wave short-range interaction coefficients in lattice units (l.u.) for lattice chiral effective field theory with lattice spacing $a = 1.32$ fm. This corresponds to a momentum cutoff of $\pi/a = 471$ MeV. Each of these terms are central interactions, meaning that the spin components form rotational invariants through contractions with themselves. We note the very good agreement between the $S = 0, T = 1$ and $S = 1, T = 0$ channels, where S is intrinsic spin and T is isospin. The error bars indicate uncertainties in the fit to empirical data. The full next-to-next-to-next-to-leading-order ($N^3\text{LO}$) chiral interaction includes these short-range interactions together with a leading-order (LO) interaction composed of an $\text{SU}(4)$ symmetric interaction and one-pion exchange potential that together have the form described in Eq. (1). Full details are given in the Supplemental Materials [27]. In the last row of Table I we show the D -wave short-range interaction coefficients in lattice units. In this case the interactions are not purely central, and so we average over all possible total angular momentum channels J for the spin-triplet channels to remove contributions from the tensor force and spin-orbit interactions, and this is denoted as $^3D_{\text{all}}$. This averaging over J was also used in Ref. [14]. We see again very good agreement between the $S = 0, T = 1$ channel and the average over the $S = 1, T = 0$ channels.

Let us now consider the matrix element between any two-nucleon states A and B , both with total intrinsic spin S and total isospin T . Let H be the isospin-invariant part of the nucleon-nucleon Hamiltonian. We define the matrix element $M(S, T)$ as

$$\frac{1}{2S+1} \sum_{S_z=-S}^S \langle A; S, S_z; T, T_z | H | B; S, S_z; T, T_z \rangle. \quad (2)$$

The statement of spin-isospin exchange symmetry is the constraint $M(S, T) = M(T, S)$.

As an example, let us consider the structure of ^{30}P . In a minimal shell model description, ^{30}P has one proton and one neutron in the $1s_{1/2}$ orbitals. The actual wave function is considerably more complicated than this, but we can make the rough approximation that the spatial wave

functions for the two lowest-lying states of ^{30}P are the same. If true, then the prediction of spin-isospin exchange symmetry is that the $S = 0, T = 1$ and $S = 1, T = 0$ states are degenerate. The actual data is that the 1^+ ground state is lower than the 0^+ state by about 0.677 MeV. This is a fairly good agreement, far better than the splitting between the deuteron and the spin-singlet proton-neutron pair in vacuum. In that case the tensor force significantly modifies the deuteron wave function. Our analysis here suggests that the interactions of the proton-neutron pair with the ^{28}Si core are suppressing spatial correlations of the 1^+ wave function caused by the tensor interaction.

We can also look at the two-body matrix elements for the proton-neutron pair in the $1s_{1/2}$ orbitals. For the following calculations we use a harmonic oscillator frequency of 13.92 MeV. Starting from the AV18 potential [43] and using a renormalization group flow to construct the corresponding effective interaction V_{lowk} [44] at momentum scale $\Lambda = 2.5 \text{ fm}^{-1} = 490$ MeV, we get two-body matrix elements of -2.54 MeV for the 1^+ state and -2.37 MeV for the 0^+ state, respectively, showing a relative error consistent with the expected $1/N_c^2$ size. Using the same V_{lowk} renormalization group flow for the chiral $N^3\text{LO}$ interaction [45] at $\Lambda = 2.5 \text{ fm}^{-1}$, we get similar results of -2.48 MeV for the 1^+ state and -2.36 MeV for the 0^+ state. However, the spin-isospin exchange symmetry is obscured at other resolution scales. For AV18 at $\Lambda = 6.0 \text{ fm}^{-1}$ we find 1.58 MeV for the 1^+ state and 0.19 MeV for the 0^+ state, which is a violation larger than the expected $1/N_c^2$ size. This lends support to our argument that the large- N_c derivation is only valid for momentum resolution scales in the range $\Lambda_{\text{large-}N_c} \sim 500$ MeV. We note that the chiral $N^3\text{LO}$ interaction has some nonlocality while the AV18 is a local interaction. In the Supplemental Materials [27] we consider several other interactions and check for spin-isospin exchange symmetry as measured by scattering phase shifts.

Our test of spin-isospin exchange symmetry is sensitive to the effective resolution scale of the nuclear interactions. We can therefore use spin-isospin exchange symmetry as a tool for estimating the effective resolution scale Λ of any nuclear interaction which might use one of many possible methods for regulating high-energy behavior. For $\Lambda < \Lambda_{\text{large-}N_c}$ the $(S, T) = (1, 0)$ central interaction is more attractive, and for $\Lambda > \Lambda_{\text{large-}N_c}$ the $(S, T) = (0, 1)$ central interaction is more attractive. We can understand this as follows. As we lower Λ , the effect of the tensor coupling between the S and D partial waves becomes weaker. In order to compensate for the loss of tensor attraction, the $(S, T) = (1, 0)$ central interaction must be more attractive than the $(S, T) = (0, 1)$ central interaction. As we increase Λ , the tensor attraction becomes stronger and the reverse effect occurs.

TABLE II. Selected two-body matrix elements for the $1s-0d$ shell.

Matrix element	L_1	L_2	L_3	L_4	L_{12}	L_{34}
1	2	2	2	2	0	0
2	2	2	2	2	2	2
3	2	2	2	2	4	4
4	2	2	2	0	2	2
5	2	2	0	0	0	0
6	2	0	2	0	2	2
7	0	0	0	0	0	0

We can now test spin-isospin exchange for general two-body matrix elements in the $1s-0d$ shell. For this analysis we use the spin-tensor analysis developed in Refs. [46–48]. We consider seven two-body matrix elements for $(S, T) = (1, 0)$ and $(S, T) = (0, 1)$. The seven matrix elements are listed in Table II. L_3 and L_4 are the orbital angular momenta of the incoming orbitals corresponding to state B in Eq. (2), and L_{34} is the total orbital angular momentum. L_1 and L_2 are the orbital angular momenta of the outgoing orbitals corresponding to state A in Eq. (2), and L_{12} is the total orbital angular momentum. Matrix element 7 is defined by the $1s_{1/2}$ orbitals that we have discussed above. In addition to performing the average over S_z in Eq. (2), we also set $L_z = (L_{12})_z = (L_{34})_z$ and average over L_z as well. The results for AV18 and the chiral $N^3\text{LO}$ interaction [45] at $\Lambda = 2.0, 2.5, 3.0, 3.5 \text{ fm}^{-1}$ are shown in Fig. 1. We see that the $(S, T) = (1, 0)$ and $(S, T) = (0, 1)$ results are nearly

equal at $\Lambda = 2.5 \text{ fm}^{-1}$ for both AV18 and the $N^3\text{LO}$ interaction, with a relative error of size $1/N_c^2$. We also confirm that for $\Lambda < \Lambda_{\text{large-}N_c}$ the $(S, T) = (1, 0)$ channel is more attractive, and for $\Lambda > \Lambda_{\text{large-}N_c}$ the $(S, T) = (0, 1)$ channel is more attractive. We should point out that the chiral $N^3\text{LO}$ interaction is defined with a momentum cutoff scale that is comparable to $\Lambda \sim 2.5 \text{ fm}^{-1}$, and so any renormalization scale dependence above this momentum value is relatively minor. A detailed summary of the methods used in these shell model calculations can be found in Ref. [48], and a description of the $V_{\text{low}k}$ effective shell model interactions can be found in Ref. [44].

The large- N_c analysis can also be applied to three-nucleon interactions [49,50]. In the large- N_c limit, the leading central three-nucleon interactions will have the form

$$V_{\text{large-}N_c}^{3N} = V_C^{3N} + [(\vec{\sigma}_1 \times \vec{\sigma}_2) \cdot \vec{\sigma}_3][(\vec{\tau}_1 \times \vec{\tau}_2) \cdot \vec{\tau}_3]W_{123}^{3N} + \vec{\sigma}_1 \cdot \vec{\sigma}_2 \vec{\tau}_1 \cdot \vec{\tau}_2 W_{12}^{3N} + \vec{\sigma}_2 \cdot \vec{\sigma}_3 \vec{\tau}_2 \cdot \vec{\tau}_3 W_{23}^{3N} + \vec{\sigma}_3 \cdot \vec{\sigma}_1 \vec{\tau}_3 \cdot \vec{\tau}_1 W_{31}^{3N} + \dots, \quad (3)$$

provided that the momentum resolution scale is close to $\Lambda_{\text{large-}N_c}$. In this case the subleading central three-nucleon interactions are of size $1/N_c$ due to terms such as $\vec{\sigma}_1 \cdot \vec{\sigma}_2 [(\vec{\tau}_1 \times \vec{\tau}_2) \cdot \vec{\tau}_3]$ and $[(\vec{\sigma}_1 \times \vec{\sigma}_2) \cdot \vec{\sigma}_3] \vec{\tau}_1 \cdot \vec{\tau}_2$ [49]. This simplification should be helpful in constraining the many short-range three-nucleon interactions that appear at higher orders in chiral effective field theory. The spin-isospin exchange symmetry of the leading interactions also




FIG. 1. Two-body matrix elements for the $1s-0d$ shell. Panels (a)–(d): results for AV18 for $(S, T) = (1, 0)$ with red dots and $(S, T) = (0, 1)$ with blue lines at $\Lambda = 2.0 \text{ fm}^{-1}$, 2.5 fm^{-1} , 3.0 fm^{-1} , and 3.5 fm^{-1} , respectively. Panels (e)–(h): results for the chiral $N^3\text{LO}$ interaction for $(S, T) = (1, 0)$ with red dots and $(S, T) = (0, 1)$ with blue lines at $\Lambda = 2.0 \text{ fm}^{-1}$, 2.5 fm^{-1} , 3.0 fm^{-1} , and 3.5 fm^{-1} , respectively.

severely limits the isospin-dependent contributions of the three-nucleon interactions to the nuclear equation of state. This fact is relevant for calculations of the nuclear symmetry energy and its density dependence in dense nuclear matter.

Spin-isospin exchange symmetry is also useful for constraining calculations that use local regularization to produce a nonzero interaction range for the three-nucleon contact interaction at N^2 LO. Local regularization produces local interactions with some strength in higher partial waves, and it has been observed that different isospin-dependent structures for the three-nucleon contact interaction produce different behaviors in many-body calculations [51,52]. The results of our analysis here suggest that such many-body calculations may be more reliable if one imposes spin-isospin exchange symmetry on the three-nucleon contact interaction for calculations with a momentum cutoff of $\Lambda_{\text{large-}N_c}$.

In this work we have confirmed that the strong interactions have an approximate spin-isospin exchange symmetry that can be derived in the limit of many colors, N_c . The derivation is valid with relative error $1/N_c^2$ only if the momentum resolution scale is near $\Lambda_{\text{large-}N_c} \sim 500$ MeV. While our work relies on the conceptual foundations provided by others, our findings may provide new insights into the empirical fact that chiral effective field theory is most effective for momentum cutoff scales near this value. We believe that our findings provide new motivation for investigating and applying the constraints of the large- N_c limit in future nuclear structure calculations.

We have derived a set of spin-isospin exchange sum rules and discussed applications to the spectrum of ^{30}P and as well as applications to nuclear forces, nuclear structure calculations, and constraining three-nucleon interactions. Our results using modern forces and methods are nevertheless entirely consistent with the results presented in Fig. 3 of Ref. [9], which shows the dominance of meson exchange coupling terms in the Nijmegen NN potential that respect spin-isospin exchange symmetry. In the Supplemental Material [27] we present evidence for spin-isospin exchange symmetry as seen from the nucleon-nucleon scattering phase shifts as well as further information about the lattice interactions.

We are grateful for illuminating discussions with Enrique Ruiz Arriola, David Kaplan, Witold Nazarewicz, Gautam Rupak, Martin Savage, and Robert Wiringa. We acknowledge funding by the U.S. National Science Foundation (Grants No. PHY-1404159, No. PHY-1811855, and No. PHY-2013047), U.S. Department of Energy (DE-SC0017887 and DE-SC0018638), the Nuclear Computational Low-Energy Initiative (NUCLEI) SciDAC-4 project (DE-SC0018083), Deutsche Forschungsgemeinschaft (Project-ID 196253076—TRR 110), VolkswagenStiftung (93562), the BMBF (Verbundprojekt 05P18PCFP1), Chinese

Academy of Sciences (CAS) through a President's International Fellowship Initiative (PIFI) (2018DM0034). The authors gratefully acknowledge computational resources provided by the Oak Ridge Leadership Computing Facility through the INCITE award “Ab-initio nuclear structure and nuclear reactions,” the Gauss Centre for Supercomputing e.V. [53] for computing time at Jülich Supercomputing Centre (JSC), and Michigan State University.

-
- [1] G. 't Hooft, *Nucl. Phys.* **B72**, 461 (1974).
 - [2] E. Witten, *Nucl. Phys.* **B160**, 57 (1979).
 - [3] H. Mütter, C. Engelbrecht, and G. Brown, *Nucl. Phys.* **A462**, 701 (1987).
 - [4] R. F. Dashen and A. V. Manohar, *Phys. Lett. B* **315**, 425 (1993).
 - [5] R. F. Dashen, E. E. Jenkins, and A. V. Manohar, *Phys. Rev. D* **51**, 3697 (1995).
 - [6] C. Carone, H. Georgi, and S. Osofsky, *Phys. Lett. B* **322**, 227 (1994).
 - [7] E. E. Jenkins and A. V. Manohar, *Phys. Lett. B* **335**, 452 (1994).
 - [8] D. B. Kaplan and M. J. Savage, *Phys. Lett. B* **365**, 244 (1996).
 - [9] D. B. Kaplan and A. V. Manohar, *Phys. Rev. C* **56**, 76 (1997).
 - [10] E. Wigner, *Phys. Rev.* **51**, 106 (1937).
 - [11] T. Mehen, I. W. Stewart, and M. B. Wise, *Phys. Rev. Lett.* **83**, 931 (1999).
 - [12] J.-W. Chen, D. Lee, and T. Schäfer, *Phys. Rev. Lett.* **93**, 242302 (2004).
 - [13] D. Lee, *Phys. Rev. Lett.* **98**, 182501 (2007).
 - [14] A. Calle Cordon and E. Ruiz Arriola, *Phys. Rev. C* **80**, 014002 (2009).
 - [15] S. König, H. W. Griebhammer, H. W. Hammer, and U. van Kolck, *Phys. Rev. Lett.* **118**, 202501 (2017).
 - [16] B.-N. Lu, N. Li, S. Elhatisari, D. Lee, E. Epelbaum, and U.-G. Meißner, *Phys. Lett. B* **797**, 134863 (2019).
 - [17] G. Karl and J. E. Paton, *Phys. Rev. D* **30**, 238 (1984).
 - [18] P. Mergell, U.-G. Meißner, and D. Drechsel, *Nucl. Phys.* **A596**, 367 (1996).
 - [19] M. A. Belushkin, H.-W. Hammer, and U.-G. Meißner, *Phys. Rev. C* **75**, 035202 (2007).
 - [20] P. E. Shanahan, R. Horsley, Y. Nakamura, D. Pleiter, P. E. L. Rakow, G. Schierholz, H. Stüben, A. W. Thomas, R. D. Young, and J. M. Zanotti (CSSM, QCDSF/UKQCD Collaborations), *Phys. Rev. D* **89**, 074511 (2014).
 - [21] A. Parreno, M. J. Savage, B. C. Tiburzi, J. Wilhelm, E. Chang, W. Detmold, and K. Orginos, *Phys. Rev. D* **95**, 114513 (2017).
 - [22] M. L. Wagman, F. Winter, E. Chang, Z. Davoudi, W. Detmold, K. Orginos, M. J. Savage, and P. E. Shanahan, *Phys. Rev. D* **96**, 114510 (2017).
 - [23] M. Illa *et al.*, *Phys. Rev. D* **103**, 054508 (2021).
 - [24] M. K. Banerjee, T. D. Cohen, and B. A. Gelman, *Phys. Rev. C* **65**, 034011 (2002).
 - [25] T. D. Cohen, *Phys. Rev. C* **66**, 064003 (2002).
 - [26] T. D. Cohen and B. A. Gelman, *Phys. Lett. B* **540**, 227 (2002).

- [27] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.127.062501> for details on phase shifts, lattice interactions, and momentum and energy scales. The following references correspond to the Supplemental Material, which includes Refs. [28–33].
- [28] V. G. J. Stoks, R. A. M. Klomp, M. C. M. Rentmeester, and J. J. de Swart, *Phys. Rev. C* **48**, 792 (1993).
- [29] R. Machleidt, *Phys. Rev. C* **63**, 024001 (2001).
- [30] D. R. Entem, R. Machleidt, and Y. Nosyk, *Phys. Rev. C* **96**, 024004 (2017).
- [31] N. Li, S. Elhatisari, E. Epelbaum, D. Lee, B.-N. Lu, and U.-G. Meißner, *Phys. Rev. C* **98**, 044002 (2018).
- [32] D. Lee, *Prog. Part. Nucl. Phys.* **63**, 117 (2009).
- [33] T. A. Lähde and U.-G. Meißner, *Nuclear Lattice Effective Field Theory: An Introduction* (Springer, New York, 2019), Vol. 957, ISBN 978-3-030-14187-5, 978-3-030-14189-9.
- [34] D. B. Kaplan, M. J. Savage, and M. B. Wise, *Phys. Lett. B* **424**, 390 (1998).
- [35] D. B. Kaplan, M. J. Savage, and M. B. Wise, *Nucl. Phys. B* **534**, 329 (1998).
- [36] V. S. Timoteo, S. Szpigel, and E. Ruiz Arriola, *Phys. Rev. C* **86**, 034002 (2012).
- [37] E. R. Arriola, V. Timoteo, and S. Szpigel, *Proc. Sci.*, CD12 (2013) 106 [[arXiv:1302.3978](https://arxiv.org/abs/1302.3978)].
- [38] E. Ruiz Arriola, *Symmetry* **8**, 42 (2016).
- [39] M. R. Schindler, H. Singh, and R. P. Springer, *Phys. Rev. C* **98**, 044001 (2018).
- [40] S. R. Beane, D. B. Kaplan, N. Klco, and M. J. Savage, *Phys. Rev. Lett.* **122**, 102001 (2019).
- [41] A. Gezerlis, I. Tews, E. Epelbaum, M. Freunek, S. Gandolfi, K. Hebeler, A. Nogga, and A. Schwenk, *Phys. Rev. C* **90**, 054323 (2014).
- [42] P. Reinert, H. Krebs, and E. Epelbaum, *Eur. Phys. J. A* **54**, 86 (2018).
- [43] R. B. Wiringa, V. G. J. Stoks, and R. Schiavilla, *Phys. Rev. C* **51**, 38 (1995).
- [44] S. Bogner, T. Kuo, L. Coraggio, A. Covello, and N. Itaco, *Phys. Rev. C* **65**, 051301(R) (2002).
- [45] D. Entem and R. Machleidt, *Phys. Rev. C* **68**, 041001(R) (2003).
- [46] M. Kirson, *Phys. Lett.* **47B**, 110 (1973).
- [47] B. A. Brown, W. A. Richter, and B. H. Wildenthal, *J. Phys. G* **11**, 1191 (1985).
- [48] B. A. Brown, W. A. Richter, R. E. Julies, and B. Wildenthal, *Ann. Phys. (N.Y.)* **182**, 191 (1988).
- [49] D. R. Phillips and C. Schat, *Phys. Rev. C* **88**, 034002 (2013).
- [50] E. Epelbaum, A. Gasparyan, H. Krebs, and C. Schat, *Eur. Phys. J. A* **51**, 26 (2015).
- [51] D. Lonardoni, J. Carlson, S. Gandolfi, J. E. Lynn, K. E. Schmidt, A. Schwenk, and X. B. Wang, *Phys. Rev. Lett.* **120**, 122502 (2018).
- [52] D. Lonardoni, S. Gandolfi, J. E. Lynn, C. Petrie, J. Carlson, K. E. Schmidt, and A. Schwenk, *Phys. Rev. C* **97**, 044318 (2018).
- [53] GCS Supercomputer JUWELS <https://www.gauss-centre.eu/>.