

**Physiological, biochemical, and structural bioinformatic analysis of the  
multiple inositol dehydrogenases from *Corynebacterium glutamicum***

Paul Ramp<sup>1,2</sup>, Christopher Pfleger<sup>3</sup>, Jonas Dittrich<sup>2,3</sup>, Christina Mack<sup>1</sup>, Holger Gohlke<sup>2,3,4,#</sup>, and  
Michael Bott<sup>1,2,#</sup>

<sup>1</sup>IBG-1: Biotechnology, Institute of Bio- and Geosciences, Forschungszentrum Jülich, Jülich,  
Germany

<sup>2</sup>The Bioeconomy Science Center (BioSC), Forschungszentrum Jülich, Jülich, Germany

<sup>3</sup>Institut für Pharmazeutische und Medizinische Chemie, Heinrich-Heine-Universität  
Düsseldorf, 40225 Düsseldorf, Germany

<sup>4</sup>John von Neumann Institute for Computing (NIC), Jülich Supercomputing Centre (JSC),  
Institute of Biological Information Processing (IBI-7: Structural Biochemistry) & Institute of  
Bio- and Geosciences (IBG-4: Bioinformatics), Forschungszentrum Jülich GmbH, 52425  
Jülich, Germany

<sup>#</sup>Corresponding authors: [m.bott@fz-juelich.de](mailto:m.bott@fz-juelich.de) and [h.gohlke@fz-juelich.de](mailto:h.gohlke@fz-juelich.de)

Running title: Inositol dehydrogenases of *Corynebacterium glutamicum*

## ABSTRACT

Inositols (cyclohexanehexols) comprise nine isomeric cyclic sugar alcohols, several of which occur in all domains of life with various functions. Many bacteria can utilize inositols as carbon and energy source via a specific pathway involving inositol dehydrogenases (IDHs) as the first step of catabolism. The microbial cell factory *Corynebacterium glutamicum* can grow with *myo*-inositol as sole carbon source. Interestingly, this species encodes seven potential IDHs, raising the question for this multiplicity. We therefore investigated the seven IDHs to determine their function, activity, and selectivity towards the biologically most important isomers *myo*-, *scyllo*-, and *D-chiro*-inositol. We created an  $\Delta$ IDH strain lacking all seven IDH genes, which could not grow on the three inositols. *Scyllo*- and *D-chiro*-inositol were identified as novel growth substrates of *C. glutamicum*. Complementation experiments showed that only four of the seven IDHs (IolG, OxiB, OxiD, OxiE) enabled growth of the  $\Delta$ IDH strain on two of the three inositols. The kinetics of the four purified enzymes agreed with the complementation results. IolG and OxiD are NAD<sup>+</sup>-dependent IDHs accepting *myo*- and *D-chiro*-inositol, but not *scyllo*-inositol. OxiB is an NAD<sup>+</sup>-dependent *myo*-IDH with a weak activity also for *scyllo*-inositol, but not for *D-chiro*-inositol. OxiE on the other hand is an NAD<sup>+</sup>-dependent *scyllo*-IDH showing also good activity for *myo*-inositol and a very weak activity for *D-chiro*-inositol. Structural models, molecular docking experiments, and sequence alignments enabled the identification of the substrate binding sites of the active IDHs and of residues allowing predictions on the substrate specificity.

## IMPORTANCE

*Myo*-, *scyllo*- and *D-chiro*-inositol are C<sub>6</sub> cyclic sugar alcohols with various biological functions, which also serve as carbon sources for microbes. Inositol catabolism starts with an oxidation to keto-inositols catalyzed by inositol dehydrogenases (IDHs). The soil bacterium

*Corynebacterium glutamicum* encodes seven potential IDHs. Using a combination of microbiological, biochemical, and modeling approaches we analyzed the function of these enzymes and identified four IDHs involved in the catabolism of inositols. They possess distinct substrate preferences for the three isomers and modeling and sequence alignments allowed the identification of residues important for substrate specificity. Our results expand the knowledge of bacterial inositol metabolism and provide an important basis for the rational development of producer strains for these valuable inositols, which show pharmacological activities against e.g., Alzheimer's disease, polycystic ovarian syndrome, or type II diabetes.

## **KEY WORDS**

inositol metabolism, *myo*-inositol, *scyllo*-inositol, *D-chiro*-inositol, enzyme kinetics, structural models, molecular docking

## INTRODUCTION

Inositols (cyclohexanehexols) comprise a group of nine isomeric forms of C<sub>6</sub>-sugar alcohols having a cyclic structure formed by the six carbons atoms, each linked to a hydroxyl group. Depending on the orientation of the hydroxyl groups, nine isomers are possible, termed *myo*-, *scyllo*-, *epi*-, *allo*-, *muco*-, *neo*-, *D-chiro*, *L-chiro*-, and *cis*-inositol, all of which except the latter occur in nature (1, 2). *Myo*-inositol (MI) is the predominant isomer used in biology and occurs in all kingdoms of life (3). It is synthesized from glucose 6-phosphate, which is converted by inositol 1-phosphate synthase (Ino1) to *myo*-inositol 1-phosphate followed by dephosphorylation to *myo*-inositol (MI) by an inositol monophosphatase (4-6). The other naturally occurring isomers are known or assumed to be derived from MI via epimerization (7, 8).

Numerous biological functions have been identified for inositols. For example, MI-containing phospholipids are constituents of the membranes of many archaea and all eukaryotes (3). Also, *scyllo*-inositol (SI) and *D-chiro*-inositol (DCI) were identified in lipids in some plant species (9, 10). Polyphosphorylated inositols (IP<sub>1-3</sub>) are key components of eukaryotic signaling pathways (3, 11), and MI hexakisphosphate (IP<sub>6</sub>), also known as phytic acid, is an abundant plant constituent serving as the main storage form of phosphate in seeds (12). In the bacterial kingdom, inositols play a prominent role, particularly in *Actinobacteria*. In this large phylum, MI is one of the precursors for the synthesis of mycothiol, a metabolite substituting glutathione (13), and a precursor of phosphatidylinositol, an abundant phospholipid in the cytoplasmic membrane and the precursor of more complex lipids of the cell envelope such as phosphatidylinositol mannosides, lipomannan, and lipoarabinomannan (14).

Many bacteria are able to utilize MI as carbon and energy source, such as *Klebsiella aerogenes* (15), *Rhizobium leguminosarum* (16), *Bacillus subtilis* (17, 18), *Sinorhizobium meliloti* (19, 20), *Paracoccus laeviglucoosivorans* (21, 22), *Legionella pneumophila* (23), or *Thermotoga maritima* (24). After uptake via specific inositol transporters,

MI is first oxidized by an inositol dehydrogenase (IDH) to yield the intermediate 2-keto-*myo*-inositol (2KMI), which is dehydrated to 3D-(3,5/4)-trihydroxycyclohexane-1,2-dione (THcHDO) by 2KMI dehydratase (25, 26). This intermediate is converted in subsequent steps to dihydroxyacetone phosphate, acetyl-CoA, and CO<sub>2</sub> (15). The genes encoding the responsible enzymes are organized in large operons (27-29), which are usually regulated by a repressor called IolR that dissociates from its operator and enables gene expression when it forms a complex with intermediates of MI catabolism (30, 31).

*Corynebacterium glutamicum* is a soil-dwelling Gram-positive actinobacterium that is used as industrial cell factory, in particular for large-scale production of L-glutamate and L-lysine (32-34). It can grow with MI as the sole carbon source (35). During growth on MI, more than 20 genes showed increased expression, most of which were located in two clusters on the genome. Cluster *iolI* (Fig. 1) contains 16 genes, which include a putative operon comprising 10 genes (cg0197-cg0207), including those for the seven enzymes assumed to be responsible for MI conversion to dihydroxyacetone, acetyl-CoA, and CO<sub>2</sub>. Whereas many genes of cluster *iolI* such as *iolD* are essential for growth on MI, the genes of cluster *iol2* are dispensable (35). Two secondary transporters for MI uptake were identified in *C. glutamicum*, called IolT1 and IolT2 (35). In the absence of MI, expression of the genes involved in MI transport and degradation was shown to be repressed by the GntR-type transcriptional regulator IolR (36). *C. glutamicum* is not only able to degrade MI but also has the intrinsic capability to synthesize MI via MI phosphate synthase (Ino1, Cg3323) and MI phosphate monophosphatase (ImpA, Cg2298) (37). Expression of the *inoI* gene is activated by the LacI-type transcriptional regulator IpsA in response to the cytoplasmic MI concentration. When sufficient MI is present, it binds to IpsA and abolishes activation of *inoI* expression (38).

In many inositol-degrading bacteria, multiple paralogous genes annotated or shown to encode IDHs were identified (39-41), which, in the case of *B. subtilis*, for example, enable growth not only on MI, but also on SI and DCI (42, 43). In *C. glutamicum*, seven potential

inositol oxidoreductases were annotated, three in cluster *iol1* (IolG, IolW, OxiB), three in cluster *iol2* (OxiC, OxiD, OxiE), and another one elsewhere in the genome (IdhA3) (Fig. 1). This work aimed at a detailed characterization of the IDHs of *C. glutamicum*. Using a  $\Delta$ IDH strain lacking the genes for all seven potential IDHs as a host for overexpression of the seven genes individually, we could show that four of the seven IDHs enable growth with MI, SI, or DCI (Fig. S1) as sole carbon and energy source. Biochemical characterization of the four purified enzymes revealed different activity profiles for the tested inositol isomers. We used structural modeling and molecular docking to elucidate the molecular basis responsible for the varying substrate specificity of the four enzymes that may be helpful in predicting the substrate specificity of yet uncharacterized inositol dehydrogenases in other organisms.

## RESULTS

**Growth on different inositols.** *C. glutamicum* can grow in minimal medium with MI as the sole carbon and energy source (35). The IDH IolG was shown to be important for growth on MI, as inactivation of *iolG* led to a reduced growth rate. Additional deletion of the gene cluster comprising *oxiC*-cg3390-*oxiD*-*oxiE* abolished the growth on MI, suggesting redundant MI dehydrogenase activities in *C. glutamicum* (35). To determine the potential of *C. glutamicum* to utilize further inositols besides MI for growth, we cultivated *C. glutamicum* MB001(DE3) in CGXII media with either glucose, MI, SI, or DCI as the sole carbon and energy source using a BioLector microcultivation system. Growth was monitored by measuring backscatter at 620 nm over a time period of 48 h. This experiment showed that *C. glutamicum* is not only able to grow with MI, but also with DCI and SI (Fig. 2A). The growth rate ( $\mu$ ) on MI ( $\sim 0.42 \text{ h}^{-1}$ ) and DCI ( $\sim 0.42 \text{ h}^{-1}$ ) was comparable to that on glucose ( $\sim 0.46 \text{ h}^{-1}$ ), while the growth rate on SI was slower ( $\sim 0.26 \text{ h}^{-1}$ ).

*C. glutamicum* possesses the two inositol transporters IolT1 and IolT2, both contributing to the uptake of MI (35). To test if DCI and SI enter the cells the same way, we analyzed the

growth of the transporter deletion mutant strains  $\Delta iolT1$ ,  $\Delta iolT2$  and  $\Delta iolT1\Delta iolT2$  on glucose, MI, SI, and DCI. Indeed, both transporters contributed to the uptake of all tested inositols (Fig. 2B, C, D). With MI as carbon source, the strains  $\Delta iolT1$  and  $\Delta iolT2$  showed a comparable growth rate ( $\sim 0.42 \text{ h}^{-1}$ ). In case of DCI and SI, strain  $\Delta iolT1$  grew slightly slower (DCI:  $\sim 0.38 \text{ h}^{-1}$ ; SI:  $\sim 0.24 \text{ h}^{-1}$ ) than strain  $\Delta iolT2$  (DCI:  $\sim 0.41 \text{ h}^{-1}$ ; SI:  $\sim 0.26 \text{ h}^{-1}$ ), suggesting that IolT1 has a higher activity for DCI and SI uptake than IolT2 (Fig. 2B, C). Deletion of both *iolT1* and *iolT2* abolished growth on each of the three inositols completely (Fig. 2D), indicating that *C. glutamicum* does not possess an additional transporter for the uptake of inositols.

As an efficient approach for investigating the role of the seven annotated IDHs of *C. glutamicum* for growth on MI, SI, and DCI, we constructed the *C. glutamicum* strain  $\Delta IDH$ , in which all seven IDH genes and the putative sugar phosphate isomerase gene *cg3390*, which is part of the *oxiC-cg3390-oxiD-oxiE* operon, were deleted (Table 1). *C. glutamicum*  $\Delta IDH$  was transformed with pMKEx2-based expression plasmids encoding one of the seven IDHs and tested for growth on the different inositols. The successful synthesis of the individual IDH proteins was confirmed by SDS-PAGE (Fig. S2). As controls, the parent strain *C. glutamicum* MB001(DE3) and the  $\Delta IDH$  strain were transformed with pMKEx2-*eyfp*. Target gene expression was induced by adding 20  $\mu\text{M}$  IPTG to the second, overnight preculture and the main culture to enable an immediate start of growth.

In contrast to strain MB001(DE3) (pMKEx2-*eyfp*), strain  $\Delta IDH$  (pMKEx2-*eyfp*) was unable to grow on MI, SI, and DCI, confirming that this mutant is suitable to test the functionality of the different IDHs. Growth of strain  $\Delta IDH$  on MI comparable to that of the positive control was obtained by expressing either *iolG* or *oxiD* (Fig. 3). Expression of *oxiB* and *oxiE* also enabled growth on MI, but at slower growth rates of  $0.24 \text{ h}^{-1}$  and  $0.11 \text{ h}^{-1}$ , respectively. Expression of *iolW*, *oxiC*, and *idhA* did not restore growth on MI and showed the same profile as the negative control expressing *eyfp*.

Similar to the growth with MI, the growth of strain  $\Delta$ IDH with DCI was made possible by the expression of either *iolG* or *oxiD* (Fig. 3) and enabled the same growth rate ( $0.45 \text{ h}^{-1}$ ) as the positive control strain. In contrast to growth with MI, no growth on DCI was observed for strain  $\Delta$ IDH expressing *oxiB* or *oxiE*. Growth of strain  $\Delta$ IDH on SI was only enabled by expressing *oxiE* or *oxiB* (Fig. 3). Plasmid-based expression of *oxiE*, even at low induction levels, enabled faster growth ( $0.40 \text{ h}^{-1}$ ) than the positive control strain ( $0.26 \text{ h}^{-1}$ ), indicating that native *oxiE* expression limited growth on SI. The expression of *oxiB* led to slower growth on SI ( $0.09 \text{ h}^{-1}$ ) and a lower final backscatter after 48 h of cultivation.

The results of the growth experiments suggest that IolG and OxiD function as efficient MI and DCI dehydrogenases. OxiB and OxiE also possess MI dehydrogenase activity, but apparently not DCI dehydrogenase activity. OxiE probably has a high SI dehydrogenase activity, whereas OxiB has a weak activity for SI.

**Kinetic properties of the enzymes IolG, OxiD, OxiE and OxiB.** To confirm the conclusions derived from the growth experiments, we biochemically characterized those IDHs that enabled growth on the tested inositols, i.e., IolG, OxiD, OxiB, and OxiE. The enzymes were overproduced with a C-terminal Strep-tag II in *C. glutamicum* MB001(DE3) using the newly constructed pPREx6 vector. It enables the direct fusion of the target protein to a C-terminal Strep-tag II and strong inducible overexpression under the control of the T7 promoter. Enzymes were purified via StrepTactin sepharose affinity chromatography followed by size exclusion chromatography. The purity of the proteins was confirmed by SDS-PAGE and Coomassie blue staining (Fig. S3).

The purified proteins were used for enzyme activity measurements via spectrophotometric assays measuring the decrease in absorbance of NADH at 340 nm with MI, SI, and DCI as substrates. The results of these experiments agreed with the conclusions derived from the growth experiments and revealed clear differences in substrate preferences and activities (Table

1 and Fig. S4). IolG and OxiD both accept MI and DCI as substrates with a preference for MI. OxiD showed a 2.5 times higher turnover number for MI than IolG. Also, the  $K_m$  of OxiD for MI was 3 times lower than the one of IolG. OxiB showed activity for MI and SI, but not for DCI. The specific activity of OxiB for MI was 4 times lower than the one of IolG and 10 times lower than the one of OxiD, corresponding to the slower growth of strain  $\Delta$ IDH (pMKEx2-oxiB) on MI (Fig. 3). The  $K_m$  values of OxiB for MI and SI were similar and comparable to the  $K_m$  values of IolG for MI. OxiE was the only IDH that showed activity for all three tested inositols with the highest activity and lowest  $K_m$  for SI. The activity for DCI was more than 1000-fold lower than for MI and SI. This low activity was apparently not sufficient to enable the growth of strain  $\Delta$ IDH (pMKEx2-OxiE) on DCI.

**Analysis of the *C. glutamicum* IDHs by sequence alignments.** Explanations for the substrate specificity of different IDHs are scarce. Previous studies dealt with the structure elucidation of IDHs in complex with inositols to understand the interactions between enzyme, substrate, and cofactor. For OxiD of *C. glutamicum*, a crystal structure (3EUW) with a resolution of 2.3 Å has been deposited in the Protein Data Bank (PDB) (44, 45) but without a bound cofactor or substrate. The structure of IolG of *B. subtilis* complexed with  $NAD^+$  and MI enabled the identification of important residues for cofactor and substrate binding. Structure-based sequence alignments led to the definition of six conserved sequence motifs (46). Motifs I and II contain amino acid residues that are important for cofactor binding, whereas motifs III-VI contain residues responsible for substrate binding and the catalytic triade consisting of Lys97, Asp172, and His176 (BsIolG numbering).

We compared the amino acid sequences of all seven annotated IDHs of *C. glutamicum* with the sequences of the biochemically characterized inositol dehydrogenases reported in literature (Table S1) and sorted them into four groups (Fig. S5): (i)  $NAD^+$ -dependent IDHs known to have activity for MI and DCI; (ii)  $NAD^+$ -dependent IDHs known to have activity for MI and

214 SI; (iii) NADP<sup>+</sup>-dependent IDHs catalyzing the reduction of 2KMI to SI; (iv) IDHs with no  
 215 activity for any tested inositol. In our comparison we focused on the previously reported motifs  
 216 to identify differences in functionally important residues within motifs I-VI. In group (i), which  
 217 includes CgIolG and CgOxiD, the sequence G<sub>124</sub>FM/NRRY/FD<sub>130</sub> in motif III and Y<sub>233</sub>GY<sub>235</sub>  
 218 in motif V (BsIolG numbering) seem to be more conserved in comparison to the other groups.  
 219 F<sub>125</sub>M/N<sub>126</sub>R<sub>127</sub> and Y<sub>235</sub> were reported as substrate binding sites for BsIolG. In group (ii)  
 220 IDHs, the sequence G<sub>124</sub>FM/NRRY/FD<sub>130</sub> can also be found in some cases, however, the  
 221 Y<sub>233</sub>GY<sub>235</sub> sequence does not occur in any representative. In most cases, the second Tyr residue  
 222 is replaced by a positively charged amino acid (H>R>K). As shown below, this residue is  
 223 involved in substrate binding and, therefore, can serve as a marker to discriminate between  
 224 IDHs specific for SI and DCI, although exceptions are possible (Fig. 4).

225 We previously identified IolW as an NADP<sup>+</sup>-dependent *scyllo*-IDH that catalyzes the  
 226 reduction of 2KMI to SI (47) and therefore assigned it to the group (iii). At position 35, IolW  
 227 contains an Ala residue, while most other IDHs contain an Asp or Glu residue. Asp- or Glu  
 228 residues at this position are conserved in NAD<sup>+</sup>-dependent IDHs, in which they form hydrogen  
 229 bonds with the ribose moiety of NAD<sup>+</sup>. NADP<sup>+</sup>-dependent enzymes typically replace Asp or  
 230 Glu with a small, neutral residue, as the negatively charged carboxylate of Asp or Glu would  
 231 effectively repel the phosphate group in this position. Often, a basic residue follows the small  
 232 neutral residue, like Arg36 in IolW, that can interact with the 2'-phosphate group of NADPH  
 233 (48-50). Also BsIolW and BsIolU, both of which have been characterized as NADPH-  
 234 dependent KMI reductases, possess a Ser or Thr residue rather than Asp or Glu at position 35  
 235 (BsIolG numbering) (51). This difference between NAD<sup>+</sup>- and NADP<sup>+</sup>-dependent IDHs  
 236 suggests that IolW is the only IDH of *C. glutamicum* favoring NADPH as cofactor.

237 Among all analyzed IDHs, the motifs of CgOxiC differ the most from the published ones  
 238 (Fig. S5). It is the only protein within the annotated IDHs of *C. glutamicum* that does not contain  
 239 a complete GxGxxG consensus sequence in motif I. Additionally, instead of Asp179, a residue

of the catalytic triad, OxiC contains an Ile residue. The lack of Asp179 suggests that OxiC is not active as IDH, which is supported by the fact that expression of *oxiC* did not enable the growth of the  $\Delta$ IDH strain on MI, DCI, or SI (Fig. 3). IdhA3 also differs at the corresponding position 172, as it contains a Glu residue instead of Asp, similar to the *myo*-IDH Gk1899, for which activity towards MI was reported (40). Despite being a conservative exchange, the difference in size might prevent IDH activity of IdhA3. Like in the case of OxiC, the expression of *idhA3* did not allow growth of the  $\Delta$ IDH strain on MI, DCI, or SI (Fig. 3). We constructed an IdhA3 variant in which we exchanged Glu172 by Asp. However, the expression of *idhA3*-E172D also did not allow growth of *C. glutamicum*  $\Delta$ IDH on MI, DCI, or SI (data not shown).

**Structural models of *C. glutamicum* IDHs and inositol docking.** Amino acid sequence comparisons of IDHs allow the prediction of cofactors and potential substrates when looking at highly conserved motifs. However, structural models and docking experiments are required to further understand inositol preference and binding mechanisms. To this end, we generated structural models with their corresponding cofactor of the IDHs IolG, OxiB, OxiC, OxiD, OxiE, and IdhA3 (Table S2). All structures show an intermediate to good global model quality (Table S2) and good local model quality near the inositol binding sites (Fig. S6), with regions of lower quality located mainly in the loops and the central tetrameric interface.

The models served as input for docking experiments using AutoDock3 (52) in combination with DrugScore<sup>2018</sup> (53) to probe the potential interaction between MI, SI, or DCI and the catalytic site of each IDH. To test if a docked solution likely adopts a favorable position in the catalytic site, we measured the distance  $d$  between the C2 atom in MI, the C1 or C6 atom in DCI, or any C atom in SI and the C4 atom of the nicotinamide group from the NAD<sup>+</sup> cofactor. As the orientation of the reactive carbon atoms towards the cofactor and the distance between these atoms is crucial for the reaction to take place, we only considered a binding pose valid if  $d < 5$  Å. For validation of the docking approach, we performed redocking experiments using

the X-ray structures of *L. casei* IDH1 (PDB id 4MIO) and *L. casei* IDH2 (PDB id 4N54) in complex with MI and SI, respectively. All dockings converged perfectly, and the poses show RMSD values  $< 2.0 \text{ \AA}$  to the respective bound MI and SI in the X-ray structures (Fig. S7A, B). Furthermore, the docked solutions of the inositols showed  $d < 5 \text{ \AA}$  for both IDHs (Fig. S7C, D), even if the docked MI poses are slightly rotated in a counter-clockwise manner.

Of the systems investigated here, the two dehydrogenases, OxiC and IdhA3, served as negative controls as both show no activity for either MI or SI (47). The two IDHs differ markedly in the structure of the entrance region to the catalytic site. In OxiC, the presence of the  $\alpha$ -helix<sub>V159-Q169</sub> narrows the catalytic site's accessibility, thus, hampering the interaction of the inositols and the enzyme (Fig. 5A). This helical element is missing in IolG, OxiD, OxiE, and IdhA3. The latter shows an open catalytic site (Fig. 5B). In OxiB, an extended loop region is located at the same position as the helix in OxiC. However, the loop does not narrow the entrance to the catalytic site.

We obtained converged docking results for all combinations of these IDHs and the investigated inositols, indicating that the docking method finds a single most favorable binding pose for each inositol, except for the combinations of IdhA3 with MI and of OxiE with DCI, where two binding poses were found (Table S3). All docked solutions of the inositols in OxiC showed distances of  $d > 5 \text{ \AA}$  (Fig. 5C), since helix<sub>V159-Q169</sub> blocks the catalytic site. Despite the accessible catalytic site in IdhA3, we also observed no valid docking pose ( $d > 5 \text{ \AA}$ ). This finding is remarkable as both IDHs showed no activity for MI or SI in previous experiments (47) and also could not recover growth of *C. glutamicum*  $\Delta$ IDH on any tested inositol

In the case of OxiE, valid docking poses were found for MI ( $d = 4.4 \text{ \AA}$ ) and SI ( $d = 5.0 \text{ \AA}$ ), which agrees with the activity data from experiments (Fig. 5C). Furthermore, the hydroxyl groups of docked inositol poses of MI and SI form interactions with the charged amino acids R13, K101, R130, D160, D178, H182, and H243 (Fig. 5D, E). Interestingly, we observed a slight incline of the docked MI compared to the SI orientation, which, though less pronounced,

agrees with results reported previously for *L. casei* IDH1 (39). The docking with OxiE failed to generate valid docking poses for DCI, although purified OxiE showed very weak activity for DCI (Table 1), which was insufficient to enable growth on DCI.

For OxiD, we observed valid docking poses for MI ( $d = 3.9 \text{ \AA}$ ), but not for SI ( $d = 9.7 \text{ \AA}$ ), which agrees with the enzymatic activity data (Fig. 5C and Fig. S8). Here, the computed pose of MI interacts with the charged amino acids K94, D153, and H175 and the hydroxyl groups Y235 and Y280 (Fig. 5F,G). Compared to the orientation of the docked MI in OxiE, the incline is more pronounced in OxiD and like the orientation reported before for *L. casei* IDH1 (39). Surprisingly, for OxiD no valid docking pose was found for DCI (Fig. 5C), even though purified OxiD shows high activity for DCI as substrate (Table 1).

For IolG, our docking results agree with the enzyme activity data (Fig. 5C). Here, MI and DCI produced valid docking poses ( $d = 4.5 \text{ \AA}$  for both inositols), whereas for SI no valid docking pose was found ( $d = 5.7 \text{ \AA}$ ) (Fig. S8). Nevertheless, MI, SI, and DCI show the same interactions in our docking experiments with the amino acids H155, H176, S173, and Y235 (Fig. S9). The larger distance between SI and the cofactor is due to a wrong orientation of the inositol that shows a similar incline as observed for MI. Finally, SI is located further away from the cofactor than MI and DCI.

In the case of OxiB, the predicted binding poses agree only for SI with the enzyme activity data (Fig. 5C). For SI, we observed a valid docking pose with a distance  $d = 3.8 \text{ \AA}$  (Fig. S8). The position of all docked inositols is strongly overlapping, thus, interacting with the same amino acids Y138, Y166, D193, H197, and N278 (Fig. S9). The larger distance between MI and the cofactor results from the misplaced C2 atom of the inositol. Here, the C2 atom points away from the cofactor and reveals a parallel orientation of the carbocyclic ring and the nicotinamide group of the cofactor. In the case of DCI, the C1/C6 atoms are oriented toward the cofactor and show a slight incline, suggesting an optimal interaction between DCI and the cofactor. Here, the docking result deviates from the enzyme activity data, as OxiB showed no

activity with DCI as substrate (Table 1). Overall, we were able to identify valid binding poses for MI, SI, and DCI in six IDHs in 15 out of 18 docking experiments.

## DISCUSSION

The genome of *C. glutamicum* harbors seven genes that potentially encode IDHs. In this study, we characterized the physiological functions and biochemical properties of these IDHs and employed bioinformatics and molecular modeling to obtain more detailed information on their structural differences and substrate preferences. Our initial growth experiments with the strain MB001(DE3) derived from *C. glutamicum* ATCC13032 revealed that it can grow not only with MI, but also with SI and DCI as sole carbon and energy source. In Fig. 6, we present an overview of our current knowledge of the inositol metabolism in *C. glutamicum*. We showed that both inositol transporters, IolT1 and IolT2, catalyze not only the uptake of MI, but also of DCI and SI, which further underlines the fact that these transporters have a broad substrate specificity including not only inositols, but also glucose, fructose, and xylose (54-57). According to the observed growth rates, IolT1 seems to have a higher activity for DCI and SI uptake than IolT2, while both transporters are comparably effective with respect to MI uptake (35). Studies on the three inositol transporters of *B. subtilis* showed that they exhibit different preferences for different inositols (58).

By creating the  $\Delta$ IDH strain of *C. glutamicum*, which lacks all seven known or putative IDH genes, we were able to test the role of each of the IDHs individually for their ability to enable growth on MI, SI, and DCI. The experiments showed that, besides IolG, also OxiD allows for fast growth on MI. OxiB and OxiE also enabled growth on MI, but at a much slower rate. These results agree with the kinetic properties of the purified enzymes with MI as substrate: IolG and OxiD showed about 5- to 10-fold higher  $V_{\max}$  values than OxiB and OxiE (Table 1). Growth on DCI was only possible with IolG and OxiD and the kinetic properties confirmed a high activity of these enzymes with DCI as substrate. OxiE showed superior growth

and faster kinetics for SI compared to MI, suggesting that this enzyme primarily functions as *scyllo*-IDH. OxiB also allowed growth on SI, but at a much lower rate. The kinetic properties of OxiE and OxiB for SI were in agreement with the growth data (Table 1). The observation that the IDHs possess activity for more than one inositol isomer has also been reported for IDHs of other bacteria. They are often classified either as *myo*-IDH with the highest activity for MI and lower activity for DCI or as *scyllo*-IDH with a preference for SI and lower activity for MI (21, 22, 39). OxiB is unusual in that it shows activity for SI and MI, but has a strong preference for MI.

Expression of *iolW*, *oxiC*, and *idhA3* did not allow growth of the  $\Delta$ IDH strain on MI, DCI, or SI, indicating that these proteins do not possess the required enzymatic activities. For *IolW*, this result was expected as our previous studies showed that this enzyme catalyzes the NADPH-dependent reduction of 2KMI to SI (47). In the case of *OxiC*, several reasons for the lack of enzymatic activity were identified. *OxiC* lacks the Asp-179 residue, which is part of the catalytic triad, and contains an incomplete GxGxxG motif. Furthermore, the structural model shows that the substrate binding site of *OxiC* is blocked by an  $\alpha$ -helix (Fig. 5A). Also the docking experiments revealed no valid binding poses for the tested inositols. Therefore, all evidence argues against an enzymatic activity of *OxiC* as IDH and the function of this protein remains unknown. In the case of *IdhA3*, the Asp-172 residue of the catalytic triad is replaced by a Glu residue and the exchange of the Glu residue by Asp did not enable growth on MI, SI or DCI (data not shown). The structural model of *IdhA3* shows that the catalytic site is more exposed than the binding pockets of active IDHs (Figure 5B) and the docking experiments revealed no valid binding poses. As in the case of *OxiC*, the function of *IdhA3* is currently unclear.

The structural models generated for the *C. glutamicum* IDHs in this study were used in blind docking experiments, which in 15 out of 18 cases were in good agreement with the experimental growth and kinetic data when the distance between the reactive carbon atom from

each inositol and the C4 atom of the cofactor's nicotinamide group was evaluated as a criterion for activity. Only for the pairs OxiD/DCI and OxiB/MI, we obtained false-negative and for OxiB/DCI false-positive docking solutions, which might be due to treating the protein and the cofactor as rigid. Using computationally more demanding investigations with molecular dynamics simulations may overcome these limitations.

Amino acid sequence alignments and the predicted interactions of the inositols with the residues of the binding pocket suggested that motif V plays a role in the selectivity of IDHs. In group (i) IDHs using MI and DCI as substrates, a YGY<sub>245</sub> motif (BsIolG numbering) with Y245 serving as substrate binding site is highly conserved, whereas this motif is absent in group (ii) IDHs using MI or SI as substrates and in group (iii) IDHs serving as 2KMI reductases. In group (ii) Y245 is exchanged mainly for a positively charged residue (H>R>K) and the model proposes H243 as a substrate interaction site in OxiE (Fig. 5D, E). Therefore, this position can be used to estimate the substrate preferences of IDHs (Fig. 4).

As several inositols were reported to show pharmacological activities against e.g., Alzheimer's disease, polycystic ovarian syndrome, or type II diabetes (2), the biotechnological production of these sugar alcohols is of high interest and was shown e.g., for SI production with *B. subtilis* (59) and *C. glutamicum* (47). These processes require the activity of IDHs for epimerization of MI to the desired inositol and the knowledge of biochemical properties of the IDHs is a prerequisite for the design of appropriate synthesis pathways and chassis strains preventing for example the reoxidation of the target inositol. The structural models enable rational engineering of the IDHs to change the substrate or cofactor selectivity, which can provide new synthetic routes for the interconversion of inositol isomers. Besides MI, DCI, and SI, also other inositols were reported to have pharmacological activities (2) and our strategy for analyzing the properties of IDHs can be employed to identify novel IDHs suitable for production of rare inositols.

Our study revealed the functions of four of the seven putative IDHs present in *C. glutamicum* (IolG, OxiD, OxiB, and OxiE), not including the 2KMI reductase IolW reported previously (47). The functions of OxiC and IdhA3 remain unknown and especially OxiC is unlikely to be an active IDH. The overlapping substrate specificities of several of the four active NAD<sup>+</sup>-dependent IDHs might provide an advantage for scavenging inositols in the natural habitat. The oxidation product of MI and SI is 2KMI (or *scyllo*-inosose), which is subsequently converted by 2KMI dehydratase (IolE) to 3D-(3,5/4)-trihydroxy-cyclohexane-1,2-dione (Fig. 6). The oxidation product of DCI, however, is 1-keto-D-*chiro*-inositol, which in *B. subtilis* is converted by the isomerase IolI to 2KMI (43). Current studies aim at identifying the *C. glutamicum* isomerase involved in growth on DCI.

## MATERIAL AND METHODS

**Bacterial strains, plasmids, and growth conditions.** All bacterial strains and plasmids used in this work are listed in Table 2. All cloning steps were performed with *Escherichia coli* DH5 $\alpha$  as host. *E. coli* strains were cultivated at 37 °C on LB agar plates or in lysogeny broth (LB) (60) with 50  $\mu$ g/mL kanamycin. For growth characterization, *C. glutamicum* was cultivated in a BioLector microcultivation system (m2p-labs, Baesweiler, Germany). Single colonies were transferred in BHI medium and cultivated for 8 h at 30°C as a first preculture. The second preculture containing defined CGXII medium (61) with 0.03 g/L protocatechuic acid and 2% (wt/vol) glucose was inoculated with 10% (vol/vol) of the first preculture and cultivated for 16 h at 30°C. Before inoculation of the main cultures, cells were washed once with CGXII medium without a carbon source. BioLector microcultivation was performed in 800  $\mu$ L CGXII medium, which was supplemented with 1% (wt/vol) of the indicated carbon source in 48-well FlowerPlates (m2p-labs, Baesweiler, Germany) at 1200 rpm at 30 °C. Growth in this system was measured online as scattered light at 620 nm (62). For protein production, *C. glutamicum* was cultivated in 200 mL BHI medium supplemented with 2% (wt/vol) glucose

in 2 L baffled shake flasks at 100 rpm and 30°C. When appropriate, 25 µg/mL kanamycin was added to the medium. Gene expression was induced via the addition of isopropyl-β-D-thiogalactoside (IPTG) at the indicated concentrations. Bacterial growth was followed by measuring the optical density at 600 nm (OD<sub>600</sub>).

**Recombinant DNA work and construction of deletion mutants.** Plasmids and oligonucleotides used in this study are listed in Table 2 and Table S4, respectively. PCRs, DNA restrictions, and plasmid constructions were performed according to established protocols (63, 64). DNA sequencing and oligonucleotide synthesis were performed by Eurofins Genomics (Ebersberg, Germany). Chemically competent *E. coli* cells were transformed according to an established protocol (65). *C. glutamicum* was transformed via electroporation as described previously (66). The deletion mutant *C. glutamicum* MB001(DE3)ΔIDH was constructed via consecutive double homologous recombination as described previously (67) using the plasmids pK19mobsacBΔiol2, pK19mobsacBΔiolG, pK19mobsacBΔiolW, pK19mobsacBΔidhA3, and pK19mobsacBΔoxiB. The chromosomal deletions were confirmed via colony-PCR using oligonucleotides annealing outside the deleted region.

For the construction of the pMKEx2-based expression plasmids, the corresponding target genes were cloned downstream of the *C. glutamicum* consensus ribosome binding site (RBS) via Gibson assembly. For protein overproduction and purification, the inositol dehydrogenase genes were cloned into the newly constructed pPREx6 plasmid, which is a derivative of pPREx2 (68) in which the promoter *P<sub>tac</sub>* was replaced by the T7 promoter. For promoter exchange, the plasmid backbone was amplified using oligonucleotides P027 and P028, and the T7 promoter was amplified from pMKEx2 with oligonucleotides P029 and P030. DNA fragments were joined via Gibson assembly yielding pPREx6.

**Protein overproduction and purification.** *C. glutamicum* MB001(DE3) was transformed with pPREx6-based expression plasmids for inositol dehydrogenase production and cultivated in 200 mL BHI medium supplemented with 20 g/L glucose. Gene overexpression was induced with 250  $\mu$ M IPTG after 3 h, and cells were harvested after 24 h of cultivation via centrifugation at 5,000 g for 20 min at 4 °C. Cell pellets were washed and resuspended in 4 mL lysis buffer (100 mM KPO<sub>4</sub> pH 7.5, 150 mM NaCl, 1 mM MgSO<sub>4</sub>) per g cell wet weight and lysed by five passages through a French Press at 124 MPa. The resulting cell extract was first centrifuged at 5,000 g and 4°C for 20 min, and the supernatant was then subjected to ultracentrifugation at 45,000 g and 4°C for 1 h. The resulting supernatant was incubated with avidin (25  $\mu$ g/mg protein) for 30 min on ice before performing purification on an Äkta pure protein purification system (Cytiva) via StrepTactin Sepharose affinity chromatography and subsequent size exclusion chromatography.

A StrepTrap HP 1 mL column was equilibrated with binding buffer (100 mM KPO<sub>4</sub> pH 7.5, 150 mM NaCl, 1 mM MgSO<sub>4</sub>) before loading the protein extract. The column was washed with 10 column volumes (CV) of binding buffer and the remaining proteins were then eluted in six 0.5 ml fractions with elution buffer I (100 mM KPO<sub>4</sub> pH 7.5, 150mM NaCl, 1 mM MgSO<sub>4</sub>, 2.5 mM desthiobiotin). The protein-containing elution fractions were combined and concentrated by using a 10 kDa AMICON filter and centrifuging at 3,500 g and 4°C to a final volume of 500  $\mu$ L. The concentrated protein was then applied to a Superdex 200 Increase size exclusion chromatography column that had been equilibrated with 2 CV of elution buffer II (100 mM KPO<sub>4</sub> pH 7.5, 1 mM MgSO<sub>4</sub>). Protein was eluted with 1.5 CV of elution buffer II and collected in 2 mL fractions. The purity and apparent molecular mass of the proteins after both purification steps were determined by 12% (wt/vol) SDS-PAGE according to standard procedures (64). Protein concentrations were determined using the Bradford assay (69).

**Inositol dehydrogenase activity assays.** Inositol dehydrogenase activity was determined as described before with some adjustments (68). Measurements were performed in 600  $\mu$ L reaction volume using 1 mL cuvettes containing 0.25-600  $\mu$ g purified enzyme in elution buffer II at 30°C. A reaction mixture without substrate was used as a blank, and the reaction was initiated by the addition of substrate. Kinetic assays were performed with varying concentrations of MI, DCI, and SI (0.5-50 mM) at a constant concentration of 5 mM NAD<sup>+</sup>. Kinetic constants were determined via a non-linear regression fit based on the Michaelis-Menten equation with the GraphPad Prism software.

**Structural bioinformatics methods.** Homology models of the IDHs were generated with the protein structure homology modeling server of SWISS-MODEL (70, 71). The template search against the SWISS-MODEL template library (SMTL, last update: 2021-02-10, last included PDB release: 2021-02-05) was performed with BLAST (72) and HHblits (73): Initially, the target sequence was searched with BLAST against the primary amino acid sequences contained in the SMTL. A total of 23 (OxiC), 79 (IdhA3), 42 (OxiE), 61 (OxiD), 28 (IolG), and 19 (OxiB) templates were found. An initial HHblits profile was built using the procedure outlined in ref. (73), followed by one iteration of HHblits against Uniclust30 (74). Next, the obtained profile was searched against all profiles of the SMTL. A total of 1846 (OxiC), 3608 (IdhA3), 2552 (OxiE), 3393 (OxiD), 3946 (IolG), and 2505 (OxiB) templates were found. Based on the found template structures, we chose the ones that included a bound cofactor and showed the highest sequence identity (Table S2). Models are built based on the target-template alignment using ProMod3 (75). Coordinates conserved between the target and the template are copied from the template to the model. Insertions and deletions are remodeled using a fragment library. Side chains are then rebuilt. Finally, the resulting model's geometry is regularized using a force field. The global and per-residue model quality was assessed using the QMEANDisCo scoring function (76) (Fig. S6). The cofactors' position was determined

from the template structures and carried over to the structural models using PyMOL Molecular Graphics System, version 2.3.0 Schrödinger, LLC, New York (77).

For the molecular docking, the 3D structure of the substrates MI, SI, and DCI were generated based on their corresponding SMILES codes using RDKit: Open-source chemoinformatics (DOI: 10.5281/zenodo.3732262). The substrates were subsequently docked into the catalytic sites of the respective IDH utilizing a combination of AutoDock3 (52) as a docking engine and DrugScore<sup>2018</sup> (53) as an objective function. Docking grids were generated with DrugScore<sup>2018</sup> using converged pair-potentials for all atom pairs. The position and dimension of the grids were calculated using the positions of inositols in crystal structures as reference point. Accounting for a margin of 8 Å in every direction, the final docking grid shows box dimensions of approximately 23 Å x 23 Å x 20 Å and is centered in the pocket of the IDHs (Fig. S7A, B). Following an established procedure (53), the docking protocol considered 100 independent runs for each ligand using an initial population size of 100 individuals, a maximum number of  $27.0 \times 10^3$  generations, a maximum number of  $5.0 \times 10^6$  energy evaluations, a mutation rate of 0.02, a crossover rate of 0.8, and an elitism value of 1. The Lamarckian genetic algorithm was chosen for sampling in all approaches. The distance between the reactive carbon of the docked substrates and the cofactor was measured using the PyMOL Molecular Graphics System.

**Data availability.** The strains and plasmids used in this work will be made available by the corresponding author (MB) upon request.

## ACKNOWLEDGEMENTS

This project was financially supported by the CLIB-Competence Center Biotechnology (CKB) funded by the European Regional Development Fund ERDF [grant number 34.EFRE-0300097] and by the German Federal Ministry of Education and Research (BMBF) [grant number

031B0918A] as part of the innovation lab "AutoBiotech" within the project "BioökonomieREVIER". The study is part of the scientific activities of the Bioeconomy Science Center (BioSC), which is financially supported by the Ministry of Innovation, Science and Research of the German Federal State of North Rhine-Westphalia (project IDH). We are grateful for computational support and infrastructure provided by the "Zentrum für Informations- und Medientechnologie" (ZIM) at the Heinrich Heine University Düsseldorf and the computing time provided by the John von Neumann Institute for Computing (NIC) to HG on the supercomputer JUWELS at the Jülich Supercomputing Centre (JSC) (user ID: HKF7, VSK33).

## REFERENCES

1. Thomas MP, Mills SJ, Potter BV. 2016. The "other" inositols and their phosphates: synthesis, biology, and medicine. *Angew Chem Int Ed* 55:1614-1650.
2. Lopez-Gamero AJ, Sanjuan C, Serrano-Castro PJ, Suarez J, Rodriguez de Fonseca F. 2020. The biomedical uses of inositols: a nutraceutical approach to metabolic dysfunction in aging and neurodegenerative diseases. *Biomedicines* 8:295.
3. Michell RH. 2008. Inositol derivatives: evolution and functions. *Nat Rev Mol Cell Biol* 9:151-161.
4. Donahue TF, Henry SA. 1981. *myo*-Inositol-1-phosphate synthase. Characteristics of the enzyme and identification of its structural gene in yeast. *J Biol Chem* 256:7077-7085.
5. Bachhawat N, Mande SC. 1999. Identification of the INO1 gene of *Mycobacterium tuberculosis* H37Rv reveals a novel class of inositol-1-phosphate synthase enzyme. *J Mol Biol* 291:531-536.
6. Nigou J, Besra GS. 2002. Characterization and regulation of inositol monophosphatase activity in *Mycobacterium smegmatis*. *Biochem J* 361:385-390.
7. Pak Y, Huang LC, Lilley KJ, Lerner J. 1992. *In vivo* conversion of [3H]-*myo*-inositol to [3H]-*chiro*-inositol in rat tissues. *J Biol Chem* 267:16904-16910.
8. Hipps PP, Ackermann KE, Sherman WR. 1982. Inositol epimerase - inosose reductase from bovine brain. *Meth Enzymol* 89 Pt D:593-598.
9. Kinnard RL, Narasimhan B, Pliska-Matyshak G, Murthy PP. 1995. Characterization of *scyllo*-inositol-containing phosphatidylinositol in plant cells. *Biochem Biophys Res Commun* 210:549-555.
10. Majumder AL, Biswas BB. 2006. *Biology of inositols and phosphoinositides*. Springer, New York.
11. Mikoshiba K. 2007. IP3 receptor/Ca<sup>2+</sup> channel: from discovery to new signaling concepts. *J Neurochem* 102:1426-1446.
12. Lott JNA, Ockenden I, Raboy V, Batten GD. 2000. Phytic acid and phosphorus in crop seeds and fruits: a global estimate. *Seed Sci Res* 10:11-33.
13. Fahey RC. 2013. Glutathione analogs in prokaryotes. *Biochim Biophys Acta* 1830:3182-3198.

14. Morita YS, Fukuda T, Sena CB, Yamaryo-Botte Y, McConville MJ, Kinoshita T. 2011. Inositol lipid metabolism in mycobacteria: biosynthesis and regulatory mechanisms. *Biochim Biophys Acta* 1810:630-641.
15. Anderson WA, Magasanik B. 1971. The pathway of *myo*-inositol degradation in *Aerobacter aerogenes*. Conversion of 2-deoxy-5-keto-D-gluconic acid to glycolytic intermediates. *J Biol Chem* 246:5662-5675.
16. Fry J, Wood M, Poole PS. 2001. Investigation of *myo*-inositol catabolism in *Rhizobium leguminosarum* bv. *viciae* and its effect on nodulation competitiveness. *Mol Plant Microbe Interact* 14:1016-1025.
17. Yoshida KI, Yamaguchi M, Morinaga T, Kinehara M, Ikeuchi M, Ashida H, Fujita Y. 2008. *myo*-Inositol catabolism in *Bacillus subtilis*. *J Biol Chem* 283:10415-10424.
18. Ramaley R, Fujita Y, Freese E. 1979. Purification and properties of *Bacillus subtilis* inositol dehydrogenase. *J Biol Chem* 254:7684-7690.
19. Kohler PR, Zheng JY, Schoffers E, Rossbach S. 2010. Inositol catabolism, a key pathway in *Sinorhizobium meliloti* for competitive host nodulation. *Appl Environ Microbiol* 76:7972-7980.
20. Galbraith MP, Feng SF, Borneman J, Triplett EW, de Bruijn FJ, Rossbach S. 1998. A functional *myo*-inositol catabolism pathway is essential for rhizopine utilization by *Sinorhizobium meliloti*. *Microbiology* 144 2915-2924.
21. Shimizu T, Takaya N, Nakamura A. 2012. An L-glucose catabolic pathway in *Paracoccus* species 43P. *J Biol Chem* 287:40448-40456.
22. Fukano K, Ozawa K, Kokubu M, Shimizu T, Ito S, Sasaki Y, Nakamura A, Yajima S. 2018. Structural basis of L-glucose oxidation by *scyllo*-inositol dehydrogenase: Implications for a novel enzyme subfamily classification. *PLoS One* 13:e0198010.
23. Manske C, Schell U, Hilbi H. 2016. Metabolism of *myo*-inositol by *Legionella pneumophila* promotes infection of amoebae and macrophages. *Appl Environ Microbiol* 82:5000-5014.
24. Rodionova IA, Leyn SA, Burkart MD, Boucher N, Noll KM, Osterman AL, Rodionov DA. 2013. Novel inositol catabolic pathway in *Thermotoga maritima*. *Environ Microbiol* 15:2254-2266.
25. Berman T, Magasanik B. 1966. The pathway of *myo*-inositol degradation in *Aerobacter aerogenes*. Dehydrogenation and dehydration. *J Biol Chem* 241:800-806.
26. Yoshida KI, Yamaguchi M, Ikeda H, Omae K, Tsurusaki KI, Fujita Y. 2004. The fifth gene of the *iol* operon of *Bacillus subtilis*, *iolE*, encodes 2-keto-*myo*-inositol dehydratase. *Microbiology* 150:571-580.
27. Kawsar HI, Ohtani K, Okumura K, Hayashi H, Shimizu T. 2004. Organization and transcriptional regulation of *myo*-inositol operon in *Clostridium perfringens*. *FEMS Microbiol Lett* 235:289-295.
28. Yoshida KI, Aoyama D, Ishio I, Shibayama T, Fujita Y. 1997. Organization and transcription of the *myo*-inositol operon, *iol*, of *Bacillus subtilis*. *J Bacteriol* 179:4591-4598.
29. Yebra MJ, Zuniga M, Beaufils S, Perez-Martinez G, Deutscher J, Monedero V. 2007. Identification of a gene cluster enabling *Lactobacillus casei* BL23 to utilize *myo*-inositol. *Appl Environ Microbiol* 73:3850-3858.
30. Yoshida KI, Shibayama T, Aoyama D, Fujita Y. 1999. Interaction of a repressor and its binding sites for regulation of the *Bacillus subtilis* *iol* divergon. *J Mol Biol* 285:917-929.
31. Kohler PR, Choong EL, Rossbach S. 2011. The RpiR-like repressor IolR regulates inositol catabolism in *Sinorhizobium meliloti*. *J Bacteriol* 193:5155-5163.
32. Becker J, Rohles CM, Wittmann C. 2018. Metabolically engineered *Corynebacterium glutamicum* for bio-based production of chemicals, fuels, materials, and healthcare products. *Metab Eng* 50:122-141.

33. Eggeling L, Bott M. 2015. A giant market and a powerful metabolism: L-lysine provided by *Corynebacterium glutamicum*. Appl Microbiol Biotechnol 99:3387-3394.
34. Wendisch VF. 2020. Metabolic engineering advances and prospects for amino acid production. Metab Eng 58:17-34.
35. Krings E, Krumbach K, Bathe B, Kelle R, Wendisch VF, Sahm H, Eggeling L. 2006. Characterization of *myo*-inositol utilization by *Corynebacterium glutamicum*: the stimulon, identification of transporters, and influence on L-lysine formation. J Bacteriol 188:8054-8061.
36. Klaffl S, Brocker M, Kalinowski J, Eikmanns BJ, Bott M. 2013. Complex regulation of the phosphoenolpyruvate carboxykinase gene *pck* and characterization of its GntR-Type regulator IolR as a repressor of *myo*-inositol utilization genes in *Corynebacterium glutamicum*. J Bacteriol 195:4283-4296.
37. Chen C, Chen K, Su T, Zhang B, Li G, Pan J, Si M. 2019. *Myo*-inositol-1-phosphate synthase (Ino-1) functions as a protection mechanism in *Corynebacterium glutamicum* under oxidative stress. MicrobiologyOpen 8:e00721.
38. Baumgart M, Luder K, Grover S, Gätgens C, Besra GS, Frunzke J. 2013. IpsA, a novel LacI-type regulator, is required for inositol-derived lipid formation in *Corynebacteria* and *Mycobacteria*. BMC Biol 11:122.
39. Aamudalapalli HB, Bertwistle D, Palmer DRJ, Sanders DAR. 2018. *myo*-Inositol dehydrogenase and *scyllo*-inositol dehydrogenase from *Lactobacillus casei* BL23 bind their substrates in very different orientations. Biochim Biophys Acta 1866:1115-1124.
40. Yoshida K, Sanbongi A, Murakami A, Suzuki H, Takenaka S, Takami H. 2012. Three inositol dehydrogenases involved in utilization and interconversion of inositol stereoisomers in a thermophile, *Geobacillus kaustophilus* HTA426. Microbiology 158:1942-1952.
41. Zhang WY, Sun ZH, Yu DL, Airideng C, Chen W, Meng H, Zhang HP. 2010. Comparative analysis of *iol* clusters in *Lactobacillus casei* strains. World J Microbiol Biotechnol 26:1949-1955.
42. Morinaga T, Ashida H, Yoshida K. 2010. Identification of two *scyllo*-inositol dehydrogenases in *Bacillus subtilis*. Microbiology 156:1538-1546.
43. Yoshida K, Yamaguchi M, Morinaga T, Ikeuchi M, Kinehara M, Ashida H. 2006. Genetic modification of *Bacillus subtilis* for production of D-*chiro*-inositol, an investigational drug candidate for treatment of type 2 diabetes and polycystic ovary syndrome. Appl Environ Microbiol 72:1310-1315.
44. Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, Weissig H, Shindyalov IN, Bourne PE. 2000. The protein data bank. Nucleic Acids Res 28:235-42.
45. Kumaran D, Mahmood A, Burley SK, Swaminathan S. 2021. 3EUW - Crystal structure of a *myo*-inositol dehydrogenase from *Corynebacterium glutamicum* ATCC 13032 doi:10.2210/pdb3euw/pdb.
46. van Straaten KE, Zheng H, Palmer DR, Sanders DA. 2010. Structural investigation of *myo*-inositol dehydrogenase from *Bacillus subtilis*: implications for catalytic mechanism and inositol dehydrogenase subfamily classification. Biochem J 432:237-47.
47. Ramp P, Lehnert A, Matamouros S, Wirtz A, Baumgart M, Bott M. 2021. Metabolic engineering of *Corynebacterium glutamicum* for production of *scyllo*-inositol, a drug candidate against Alzheimer's disease. Metab Eng 67:173-185.
48. Carugo O, Argos P. 1997. NADP-dependent enzymes. II: Evolution of the mono- and dinucleotide binding domains. Proteins 28:29-40.
49. Carugo O, Argos P. 1997. NADP-dependent enzymes. I: Conserved stereochemistry of cofactor binding. Proteins 28:10-28.

50. Zheng H, Bertwistle D, Sanders DA, Palmer DR. 2013. Converting NAD-specific inositol dehydrogenase to an efficient NADP-selective catalyst, with a surprising twist. *Biochemistry* 52:5876-5883.
51. Kang DM, Tanaka K, Takenaka S, Ishikawa S, Yoshida K. 2017. *Bacillus subtilis* *iolU* encodes an additional NADP<sup>+</sup>-dependent *scyllo*-inositol dehydrogenase. *Biosci Biotechnol Biochem* 81:1026-1032.
52. Morris GM, Goodsell DS, Halliday RS, Huey R, Hart WE, Belew RK, Olson AJ. 1998. Automated docking using a Lamarckian genetic algorithm and an empirical binding free energy function. *J Comp Chem* 19:1639-1662.
53. Dittrich J, Schmidt D, Pflieger C, Gohlke H. 2019. Converging a knowledge-based scoring function: DrugScore<sup>2018</sup>. *J Chem Inf Model* 59:509-521.
54. Bäumchen C, Krings E, Bringer S, Eggeling L, Sahm H. 2009. *Myo*-inositol facilitators IolT1 and IolT2 enhance D-mannitol formation from D-fructose in *Corynebacterium glutamicum*. *FEMS Microbiol Lett* 290:227-235.
55. Ikeda M, Mizuno Y, Awane S, Hayashi M, Mitsuhashi S, Takeno S. 2011. Identification and application of a different glucose uptake system that functions as an alternative to the phosphotransferase system in *Corynebacterium glutamicum*. *Appl Microbiol Biotechnol* 90:1443-1451.
56. Lindner SN, Seibold GM, Henrich A, Krämer R, Wendisch VF. 2011. Phosphotransferase system-independent glucose utilization in *Corynebacterium glutamicum* by inositol permeases and glucokinases. *Appl Environ Microbiol* 77:3571-3581.
57. Brüsseler C, Radek A, Tenhaef N, Krumbach K, Noack S, Marienhagen J. 2018. The *myo*-inositol/proton symporter IolT1 contributes to D-xylose uptake in *Corynebacterium glutamicum*. *Bioresour Technol* 249:953-961.
58. Bettaney KE, Sukumar P, Hussain R, Siligardi G, Henderson PJ, Patching SG. 2013. A systematic approach to the amplified expression, functional characterization and purification of inositol transporters from *Bacillus subtilis*. *Mol Membr Biol* 30:3-14.
59. Michon C, Kang CM, Karpenko S, Tanaka K, Ishikawa S, Yoshida K. 2020. A bacterial cell factory converting glucose into *scyllo*-inositol, a therapeutic agent for Alzheimer's disease. *Comm Biol* 3:93.
60. Bertani G. 1951. Studies on lysogenesis. The mode of phage liberation by lysogenic *Escherichia coli*. *J Bacteriol* 62:293-300.
61. Keilhauer C, Eggeling L, Sahm H. 1993. Isoleucine synthesis in *Corynebacterium glutamicum*: molecular analysis of the *ilvB-ilvN-ilvC* operon. *J Bacteriol* 175:5595-5603.
62. Kensy F, Zang E, Faulhammer C, Tan RK, Büchs J. 2009. Validation of a high-throughput fermentation system based on online monitoring of biomass and fluorescence in continuously shaken microtiter plates. *Microb Cell Fact* 8:31.
63. Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, Smith HO. 2009. Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Meth* 6:343-345.
64. Green MR, Sambrook J. 2012. *Molecular Cloning. A Laboratory Manual*, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
65. Hanahan D. 1983. Studies on transformation of *Escherichia coli* with plasmids. *J Mol Biol* 166:557-580.
66. van der Rest ME, Lange C, Molenaar D. 1999. A heat shock following electroporation induces highly efficient transformation of *Corynebacterium glutamicum* with xenogeneic plasmid DNA. *Appl Microbiol Biotechnol* 52:541-545.
67. Niebisch A, Bott M. 2001. Molecular analysis of the cytochrome *bc<sub>1</sub>-aa<sub>3</sub>* branch of the *Corynebacterium glutamicum* respiratory chain containing an unusual diheme cytochrome *c<sub>1</sub>*. *Arch Microbiol* 175:282-294.

68. Bakkes PJ, Ramp P, Bida A, Dohmen-Olma D, Bott M, Freudl R. 2020. Improved pEKEx2-derived expression vectors for tightly controlled production of recombinant proteins in *Corynebacterium glutamicum*. Plasmid 112.
69. Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Anal Biochem 72:248-254.
70. Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, Heer FT, de Beer TAP, Rempfer C, Bordoli L, Lepore R, Schwede T. 2018. SWISS-MODEL: homology modelling of protein structures and complexes. Nucleic Acids Res 46:W296-W303.
71. Bienert S, Waterhouse A, de Beer TA, Tauriello G, Studer G, Bordoli L, Schwede T. 2017. The SWISS-MODEL repository - new features and functionality. Nucleic Acids Res 45:D313-D319.
72. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. BMC Bioinformatics 10:421.
73. Steinegger M, Meier M, Mirdita M, Vohringer H, Haunsberger SJ, Soding J. 2019. HH-suite3 for fast remote homology detection and deep protein annotation. BMC Bioinformatics 20:473.
74. Mirdita M, von den Driesch L, Galiez C, Martin MJ, Soding J, Steinegger M. 2017. Uniclust databases of clustered and deeply annotated protein sequences and alignments. Nucleic Acids Res 45:D170-D176.
75. Studer G, Tauriello G, Bienert S, Biasini M, Johnner N, Schwede T. 2021. ProMod3 - A versatile homology modelling toolbox. PLoS Comput Biol 17:e1008667.
76. Studer G, Rempfer C, Waterhouse AM, Gumienny R, Haas J, Schwede T. 2020. QMEANDisCo-distance constraints applied on model quality estimation. Bioinformatics 36:1765-1771.
77. DeLano WL. 2002. The PyMOL molecular graphics system. [www.pymol.org](http://www.pymol.org). Accessed 2022.
78. Kortmann M, Kuhl V, Klaffl S, Bott M. 2015. A chromosomally encoded T7 RNA polymerase-dependent gene expression system for *Corynebacterium glutamicum*: construction and comparative evaluation at the single-cell level. Microb Biotechnol 8:253-265.
79. Schäfer A, Tauch A, Jäger W, Kalinowski J, Thierbach G, Pühler A. 1994. Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. Gene 145:69-73.
80. Tenhaef N, Brüsseler C, Radek A, Hilmes R, Unrean P, Marienhagen J, Noack S. 2018. Production of D-xylonic acid using a non-recombinant *Corynebacterium glutamicum* strain. Bioresour Technol 268:332-339.

**Table 1.** Role of the indicated IDHs for growth on MI, DCI, and SI, and kinetic constants for oxidation of MI, DCI, and SI by purified IolG, OxiB, OxiD, and OxiE.

Enzyme	Substrate	Growth <sup>1</sup>	V <sub>max</sub> (μmol min <sup>-1</sup> mg <sup>-1</sup> )	K <sub>m</sub> (mM)	k <sub>cat</sub> (s <sup>-1</sup> )	k <sub>cat</sub> /K <sub>m</sub> (M <sup>-1</sup> s <sup>-1</sup> )
IolG	<i>myo</i> -Inositol	+++	23.1 ± 3.2	60.9 ± 13.7	14.0 ± 1.9	235.1 ± 23.4
	<i>D-chiro</i> -Inositol	+++	14.3 ± 0.9	61.93 ± 5.61	8.66 ± 0.54	140.39 ± 7.4
	<i>scyllo</i> -Inositol	-	n.d. <sup>2</sup>			
OxiD	<i>myo</i> -Inositol	+++	59.0 ± 2.3	19.6 ± 1.8	35.6 ± 1.4	1831.8 ± 107.4
	<i>D-chiro</i> -Inositol	+++	25.5 ± 2.7	50.6 ± 8.7	15.4 ± 1.7	307.6 ± 20.0
	<i>scyllo</i> -Inositol	-	n.d.			
OxiB	<i>myo</i> -Inositol	++	5.8 ± 0.5	62.1 ± 12.5	4.2 ± 0.4	69.1 ± 7.3
	<i>D-chiro</i> -Inositol	-	n.d.			
	<i>scyllo</i> -Inositol	+	0.05 ± 0.01	28.8 ± 7.1	0.03 ± 0.00	1.03 ± 0.11
OxiE	<i>myo</i> -Inositol	+	3.1 ± 0.2	51.6 ± 3.6	3.9 ± 0.3	76.7 ± 5.2
	<i>D-chiro</i> -Inositol	-	0.005 ± 0.001	54.1 ± 8.0	0.003 ± 0.00	0.06 ± 0.00
	<i>scyllo</i> -Inositol	+++	13.4 ± 0.1	12.4 ± 0.8	8.5 ± 0.1	688.6 ± 41.5

<sup>1</sup>Growth of ΔIDH strain expressing the genes encoding the indicated IDHs, with "+++" indicating very good growth and "-" indicating no growth

<sup>2</sup>n.d. = no detectable activity


758 **Table 2.** Bacterial strains and plasmids used in this study

Strain or plasmid	Relevant characteristics	Source or reference
<i>E. coli</i>		
DH5 $\alpha$	F <sup>-</sup> $\Phi$ 80 <i>dlac</i> $\Delta$ ( <i>lacZ</i> )M15 $\Delta$ ( <i>lacZYA-argF</i> ) U169 <i>endA1 recA1 hsdR17</i> (r $\kappa^-$ , m $\kappa^+$ ) <i>deoR thi-1 phoA supE44 <math>\lambda^-</math> gyrA96 relA1</i> ; strain used for cloning procedures	(65)
<i>C. glutamicum</i>		
MB001(DE3)	Derivative of the prophage-free strain MB001 with a chromosomally encoded <i>E. coli lacI</i> gene under control of its native promoter followed by the T7 RNA polymerase gene under control of the <i>lacUV5</i> promoter.	(78)
MB001(DE3) $\Delta$ IDH	MB001(DE3) derivative with deletion of the genes <i>oxiC</i> -cg3390- <i>oxiD</i> - <i>oxiE</i> (cg3389-cg3392), <i>iolG</i> (cg0204), <i>iolW</i> (cg0207), <i>idhA3</i> (cg2313) and <i>oxiB</i> (cg0211)	This work
ATCC13032 $\Delta$ <i>iolT1</i>	Derivative of the wild type ATCC13032 in which the inositol transporter <i>iolT1</i> was deleted	(35)
ATCC13032 $\Delta$ <i>iolT2</i>	Derivative of the wild type ATCC13032 in which the inositol transporter <i>iolT2</i> was deleted	(35)
ATCC13032 $\Delta$ <i>iolT1</i> $\Delta$ <i>iolT2</i>	Derivative of the wild type ATCC13032 in which the inositol transporters <i>iolT1</i> and <i>iolT2</i> were deleted	(35)
Plasmids		
pK19mobsacB	Kan <sup>R</sup> ; plasmid for allelic exchange in <i>C. glutamicum</i> ; (pK18 <i>oriV<sub>E.c.</sub></i> , <i>sacB</i> , <i>lacZ<math>\alpha</math></i> )	(79)
pK19mobsacB $\Delta$ <i>iol2</i>	Kan <sup>R</sup> ; plasmid for deletion of the genes cg3389-cg3392 containing two 1-kb PCR products which cover the upstream flanking region of <i>oxiC</i> (cg3389) and the downstream flanking region of <i>oxiE</i> (cg3392)	(47)
pK19mobsacB $\Delta$ <i>iolG</i>	Kan <sup>R</sup> ; plasmid for deletion of <i>iolG</i> (cg0204)	(80)
pK19mobsacB $\Delta$ <i>iolW</i>	Kan <sup>R</sup> ; plasmid for deletion of <i>iolW</i> (cg0207)	(57)
pK19mobsacB $\Delta$ <i>oxiB</i>	Kan <sup>R</sup> ; plasmid for deletion of <i>oxiB</i> (cg0211)	This work
pK19mobsacB $\Delta$ <i>idhA3</i>	Kan <sup>R</sup> ; plasmid for deletion of <i>idhA3</i> (cg2313)	This work
pMKEx2	Kan <sup>R</sup> ; <i>E. coli</i> - <i>C. glutamicum</i> shuttle vector ( <i>lacI</i> , P <sub>T7</sub> , <i>lacO1</i> , pHM1519 <i>ori<sub>Cg</sub></i> ; pACYC177 <i>ori<sub>Ec</sub></i> ) for expression of target genes under control of the T7 promoter	(78)
pMKEx2- <i>eyfp</i>	Kan <sup>R</sup> ; pMKEx2 derivative containing the <i>eyfp</i> gene under control of P <sub>T7</sub>	(78)
pMKEx2- <i>IolG</i>	Kan <sup>R</sup> ; pMKEx2 derivative containing the <i>iolG</i> gene under control of P <sub>T7</sub>	This work
pMKEx2- <i>IolW</i>	Kan <sup>R</sup> ; pMKEx2 derivative containing the <i>iolW</i> gene under control of P <sub>T7</sub>	This work
pMKEx2- <i>OxiB</i>	Kan <sup>R</sup> ; pMKEx2 derivative containing the <i>oxiB</i> gene under control of P <sub>T7</sub>	This work


pMKEx2-IdhA3	Kan <sup>R</sup> ; pMKEx2 derivative containing the <i>idhA3</i> gene under control of P <sub>T7</sub>	(47)
pMKEx2-OxiC	Kan <sup>R</sup> ; pMKEx2 derivative containing the <i>oxiC</i> gene under control of P <sub>T7</sub>	(47)
pMKEx2-OxiD	Kan <sup>R</sup> ; pMKEx2 derivative containing the <i>oxiD</i> gene under control of P <sub>T7</sub>	(47)
pMKEx2-OxiE	Kan <sup>R</sup> ; pMKEx2 derivative containing the <i>oxiE</i> gene under control of P <sub>T7</sub>	(47)
pPREx2	Kan <sup>R</sup> ; <i>E. coli</i> - <i>C. glutamicum</i> shuttle vector (P <sub>tac</sub> , <i>lacI</i> <sup>q</sup> , pBL1 <i>ori</i> <sub>Cg</sub> ; ColE1 <i>ori</i> <sub>Ec</sub> with a Strep-tag II encoding sequence	(68)
pPREx6	Kan <sup>R</sup> ; pPREx2 derivative with with P <sub>tac</sub> exchanged for P <sub>T7</sub> promoter	This work
pPREx6-IolG	Kan <sup>R</sup> ; pPREx6 derivative containing the <i>iolG</i> gene under control of P <sub>T7</sub> and fused to Strep-tag II sequence	This work
pPREx6-OxiB	Kan <sup>R</sup> ; pPREx6 derivative containing the <i>oxiB</i> gene under control of P <sub>T7</sub> and fused to Strep-tag II sequence	This work
pPREx6-OxiD	Kan <sup>R</sup> ; pPREx6 derivative containing the <i>oxiD</i> gene under control of P <sub>T7</sub> and fused to Strep-tag II sequence	This work
pPREx6-OxiE	Kan <sup>R</sup> ; pPREx6 derivative containing the <i>oxiE</i> gene under control of P <sub>T7</sub> and fused to Strep-tag II sequence	This work

759

760



**Fig. 1.** Organization of *C. glutamicum* genes involved inositol transport and metabolism with the seven different IDH genes highlighted in color. The gene clusters *iol1* and *iol2* comprising genes involved in inositol metabolism are indicated.



769

770

771

772

773

774


775

776

777


778

**Fig. 2.** Growth of the *C. glutamicum* strain MB001(DE3) (A), ATCC13032  $\Delta iolT1$  (B), ATCC13032  $\Delta iolT2$ , and ATCC13032  $\Delta iolT1\Delta iolT2$  on different inositols compared to glucose. The strains were cultivated in a BioLector system using CGXII minimal medium supplemented with glucose, MI, SI, or DCI at 10 g/L. The cultures were incubated for 48 h at 30°C, 1200 rpm, and 85% humidity. Mean values and standard deviations of three biological replicates are shown.



**Fig. 3.** Growth on MI, SI, and DCI of *C. glutamicum*  $\Delta$ IDH expressing one of the seven IDH genes or as negative control *eyfp* using the corresponding pMKEx2-based plasmids. *C. glutamicum* MB001(DE3) transformed with pMKEx2-eyfp was used as a positive control. The strains were cultivated in a BioLector cultivation system for 48 h at 30°C, 1200 rpm, and 85% humidity in CGXII minimal medium supplemented with 10 g/L of MI, DCI, SI, or glucose. Mean values and standard deviations of three biological replicates are shown.


787



788

789


790 **Fig. 4.** Consensus sequences for motif V of the four IDH subgroups with the substrate binding  
791 site highlighted. Letter height is proportional to the relative abundance of that residue at each  
792 position and letter width is proportional to the fraction of valid symbols at that position. Letter  
793 color corresponds to the chemical properties of the amino acid (black, hydrophobic; red, acidic;  
794 blue, basic; green, polar; pink, carboxamides). The figure was generated using WebLogo 3  
795 ([WebLogo 3 - About \(threeplusone.com\)](http://threeplusone.com/weblogo)).



**Fig 5.** Docking results for *myo*-, *scyllo*-, and *D-chiro*-inositol into structural models of *C. glutamicum* IDHs. Structural comparison between IdhA3 (A) and OxiC (B) shows an  $\alpha$ -helix (magenta) blocking the entrance to the catalytic site of OxiC. Compared to active IDHs, the catalytic site in IdhA3 is more exposed. (C) Distances were measured for the docked inositol poses between the reactive carbon atom from each inositol and the C4 atom of the cofactors

803 nicotinamide group; black diamonds depict those inositols that result in growth. The horizontal  
804 dashed line indicates the threshold for considering a valid docking pose. The bar color depicts  
805 if the distance is within the threshold (green) or outside (red). (D-G) Comparison of the docked  
806 solutions for MI and SI into OxiE and OxiD.  
807

808



809

810

811 **Fig. 6.** Schematic overview of *myo*-, *scyllo*-, and *D-chiro*-inositol catabolism in *C. glutamicum*.

812 Reactions leading to cell constituents requiring L-*myo*-inositol-1-phosphate or *myo*-inositol for

813 synthesis are indicated with dashed arrows.