



RESEARCH ARTICLE

A model-based estimate of nitrate leaching in Germany for GHG reporting

Max Eysholdt¹ | Ralf Kunkel² | Claus Rösemann³ | Frank Wendland² |
Tim Wolters² | Maximilian Zinnbauer¹ | Roland Fuß³

¹Thünen Institute of Rural Studies,
Braunschweig, Germany

²Forschungszentrum Jülich GmbH, Institute of
Bio- and Geosciences (IBG-3) - Agrosphere,
Jülich, Germany

³Thünen Institute of Climate-Smart
Agriculture, Braunschweig, Germany

Correspondence

Maximilian Zinnbauer, Institute of Rural
Studies of the Johann Heinrich von
Thünen-Institute, Bundesallee 64, 38116
Braunschweig, Germany.
Email: maximilian.zinnbauer@thuenen.de

This article has been edited by Horst Gerke.

Abstract

Background: Conversion of leached and runoff nitrate (NO_3^-) from agricultural land into emissions of the greenhouse gas (GHG) nitrous oxide (N_2O) by denitrification in water bodies has to be reported in national GHG inventories. The global IPCC default methodology for estimating these indirect N_2O emissions assumes that a fixed fraction of nitrogen (N) inputs ($\text{Frac}_{\text{LEACH}}$) is lost through leaching and runoff. However, this method does not consider all relevant country-specific conditions that may influence NO_3^- leaching.

Aims: The aim of this study was to apply a model-based approach for estimating indirect N_2O emissions through NO_3^- leaching and runoff from agricultural soils for use in Germany's national GHG inventory.

Methods: High-resolution spatial data and a comprehensive model system (RAUMIS-mGROWA-DENUZ) were used to derive regionally differentiated and temporarily dynamic $\text{Frac}_{\text{LEACH}}$ values from N surplus and hydrogeological conditions. These were then used to estimate indirect N_2O emissions in accordance with the IPCC methodology.

Results: The nationwide average of the new implied $\text{Frac}_{\text{LEACH}}$ values was $0.099 \text{ kg N (kg N input)}^{-1}$ in 2019. The new estimate of indirect N_2O emissions was $10.4 \text{ Gg N}_2\text{O}$ in 1990 and $5.7 \text{ Gg N}_2\text{O}$ in 2019, which are 27 and 52% less than the calculation based on the 2006 IPCC Tier 1 methodology.

Conclusions: The model-based method for estimating $\text{Frac}_{\text{LEACH}}$ incorporates relevant factors that influence NO_3^- leaching and runoff and considers site-specific, spatially varying conditions and differences in the agrarian structure. The use of N surplus as the model driver allows annual changes in cropping conditions and the effects of N-regulating policies and mitigation measures to be represented.

KEYWORDS

emission inventory, indirect N_2O emissions, nitrate leaching model, nitrogen surplus

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Journal of Plant Nutrition and Soil Science* published by Wiley-VCH GmbH

1 | INTRODUCTION

Nitrogen (N) is an essential input to crop production. Since the invention of the Haber-Bosch process, there has been an overabundant supply of N. N flows are considered to exceed the safe operating space for the planet (Steffen et al., 2015). Modern intensive agriculture is the major source of reactive N losses to the environment (Galloway et al., 2003), presenting a wide range of challenges for ecosystems and human health (Brink et al., 2011). Reactive N is lost via different pathways, for example, through nitrate (NO_3^-) leaching (N leaching), gaseous ammonia (NH_3) losses or nitrous oxide (N_2O), nitric oxide (NO), and dinitrogen (N_2) emissions from denitrification processes.

N leaching is a hydrological pathway from topsoil into groundwater, through which mineral N is dissolved and transported in seepage water. Leaching occurs when precipitation and/or irrigation exceeds field capacity and evapotranspiration, and soil NO_3^- has accumulated because N inputs have exceeded current crop needs. N leaching has numerous environmental impacts and potential implications for human health. These include negative effects on surface water and groundwater quality, eutrophication of ecosystems, acidification, and reduced biodiversity (Di & Cameron, 2002; Galloway et al., 2003; Hashemi et al., 2018; Molina-Navarro et al., 2014; Wang et al., 2019; Zhang & Yu, 2020).

The natural process of N conversion via nitrification and denitrification that occur in soil and in groundwater and surface water bodies can contribute to indirect emissions of the long-lived greenhouse gas (GHG) N_2O , a contributor to global warming and ozone depletion (IPCC, 2014; Ravishankara et al., 2009). Its 100-year global warming potential is 265 times greater than that of an equal mass of carbon dioxide (Butterbach-Bahl et al., 2013; IPCC, 2014). The increasing use of N to fertilize agricultural crops has altered the natural N cycle, and it is estimated that 22% of current global N_2O emissions result from N inputs in agriculture (Tian et al., 2020).

Progress in meeting the GHG reduction goals set in the Kyoto Protocol and subsequent agreements within the United Nations Framework Convention of Climate Change (UNFCCC) is evaluated based on national GNG inventory reports, which have to be compiled annually by countries listed in its Annex I. The Intergovernmental Panel on Climate Change (IPCC) provides internationally agreed guidelines for the estimation of GHG emissions for national GNG inventories (IPCC, 2006, 2019). The guidelines distinguish between two pathways that lead to N_2O emissions from N inputs in agriculture. In the direct pathway, N_2O is emitted directly from the soil, where N is added through fertilization, harvest residues, or mineralization of soil organic matter. A smaller but still important source are two indirect pathways: (1) volatilization of NH_3 and oxides of N (NO_x) and their subsequent redeposition and (2) N_2O emissions originating from N losses from agricultural land through leaching and runoff ($\text{N}_2\text{O}_{(\text{L})}$). Both pathways result in N_2O emissions in other locations. In the IPCC Tier 1 approach, total $\text{N}_2\text{O}_{(\text{L})}$ emissions are calculated using the fraction of N that is lost through leaching and runoff ($\text{Frac}_{\text{LEACH}}$), all N inputs to managed soils in regions where leaching and runoff occur, and an emissions factor (EF) (IPCC, 2019,

Equation 11.10):

$$\text{N}_2\text{O}_{(\text{L})} - \text{N} = \text{N}_{\text{input}} * \text{Frac}_{\text{LEACH}} * \text{EF}_5. \quad (1)$$

These total N_2O emissions are the sum of emissions that “may take place in the groundwater below the land to which the N was applied, or in riparian zones receiving drain or runoff water, or in the ditches, streams, rivers and estuaries (and their sediments) into which the land drainage water eventually flows” (IPCC, 2019, p. 11.21). EF_5 functions as a combined EF for these locations (IPCC, 1996, 2006, 2019). Distinct EFs for each location are not available in the IPCC methodology (IPCC, 2019) and therefore only the combined emission of N_2O from denitrification and other processes at these locations can be estimated.

Germany's National Greenhouse Gas Inventory currently calculates indirect N_2O emissions from N leaching and surface runoff with this methodology using the default values for $\text{Frac}_{\text{LEACH}}$ ($0.3 \text{ kg N } [\text{kg N input}]^{-1}$) and EF_5 ($0.0075 \text{ N}_2\text{O-N } [\text{kg N leaching/runoff}]^{-1}$), as stated in the 2006 IPCC guidelines (Federal Environment Agency, 2021; IPCC, 2006; Rösemann et al., 2021). The IPCC has recently updated estimates of $\text{Frac}_{\text{LEACH}}$ to a value of $0.24 \text{ kg N } (\text{kg N input})^{-1}$ and EF_5 to a value of $0.011 \text{ N}_2\text{O-N } (\text{kg N leaching/runoff})^{-1}$ for wet climates (IPCC, 2019). However, this global estimate of $\text{Frac}_{\text{LEACH}}$ remains very uncertain, with a confidence interval of between 0.1 and 0.8 $\text{kg N } (\text{kg N input})^{-1}$ (IPCC, 2019). This updated $\text{Frac}_{\text{LEACH}}$ value was derived from a global meta-analysis and does not consider different soil properties and varying hydrological conditions (Wang et al., 2019). As N leaching is dependent on soil properties, different climatic factors, agricultural management practices, and agricultural N surplus (Blicher-Mathiesen et al., 2014; Di & Cameron, 2002; De Notaris et al., 2018; Wang et al., 2019), a global default value may not be a true reflection of respective national conditions (Zhou & Butterbach-Bahl, 2014). N_2O emissions from agricultural soils are a key category in Germany's GHG inventory (Federal Environment Agency, 2021), for which IPCC guidelines require the use of a Tier 2 (country-specific) or Tier 3 (modeling) method, which should consider regional information and thereby reduce uncertainty and improve accuracy.

To facilitate fulfillment of this requirement, the aim of this study was to estimate regional and dynamic $\text{Frac}_{\text{LEACH}}$ values utilizing a model system. It considered high-resolution data on animal and crop production together with site-specific climate and hydrological factors and soil properties to reflect the heterogeneity of climatic and hydrogeological conditions, as well as farm types and agricultural production structures in Germany.

Western Germany is influenced by an oceanic climate with moderate temperatures and high precipitation, whereas in the east of the country, a more continental climate with particularly low precipitation rates in the north-east is prevalent. Due to intensive animal production, N surpluses are highest in north-west Germany, and even though some soils display high denitrification potentials (Ackermann et al., 2015), N leaching rates are generally high in this region. The north-east is dominated by intensive arable farming. N surpluses are lower, but NO_3^- concentrations in the leachate are elevated due to

predominantly sandy soils and low precipitation (Kunkel et al., 2017). In the southern part of Germany, dairy farming on grassland is widespread, especially in the Alpine foothills. The heterogeneous natural conditions and agricultural structure necessitate a spatially differentiated modeling approach to derive a new estimate of $\text{Frac}_{\text{LEACH}}$ values for use in Germany's GHG inventory.

In addition to including detailed spatial characterization of soil properties in the model, a particular focus was put on agricultural N surpluses as a driver of N leaching (Di & Cameron, 2002; Wang et al., 2019). The agricultural N surplus of an area is defined as the difference between N application and removal through crop harvest products. Studies have found a positive correlation between N surplus and N leaching (Blicher-Mathiesen et al., 2014; Huang et al., 2017; Zhou & Butterbach-Bahl, 2014), and a greater influence of N surplus than N input on N leaching (De Notaris et al., 2018).

Simulation models have been used for more than 30 years to calculate NO_3^- concentrations in the leachate (Groenendijk et al., 2014). Each of these models has been developed against the backdrop of both a specific research question and a certain spatial scale of application. Physically based models, such as HYDRUS-1D (Šimůnek et al., 2008) and the Daisy model (Manevski et al., 2016), may be suitable for simulating site-specific pore water fluxes of NO_3^- at field scale (Colombani et al., 2020). However, their applicability on the scale of larger regions or entire countries is limited due to the fact that numerous input data are not available on this scale (Kunkel et al., 2017). For application at the spatial resolution of German federal states, models such as SWAT (Arnold et al., 1998), HYPE (Arheimer et al., 2012), and MONERIS (Fuchs et al., 2010) are more suitable. However, their maximum spatial resolution is limited to the level of subcatchment areas. Thus, the identification of site-specific hotspot areas of N leaching below this level is impeded. The model system RAUMIS-mGROWA-DENUZ (Heidecke et al., 2015; Herrmann et al., 2015; Kunkel et al., 2017; Wendland et al., 2009) is not only suitable for applications on a state scale, but also allows the NO_3^- concentration in the leachate to be determined with high spatial resolution (i.e., on a 100×100 m grid).

The model system has been applied at regional scales, for instance, in the Weser catchment (Heidecke et al., 2015; Hirt et al., 2012; Kreins et al., 2011; Kuhn et al., 2016; Wendland et al., 2009, 2010) and in the federal state of Lower Saxony (Ackermann et al., 2015; Heidecke et al., 2016). The models have also been applied to several other German states (Kuhr et al., 2014; Kunkel et al., 2017; Tetzlaff et al., 2009, 2017, 2021; Wendland et al., 2014, 2015, 2021). Most recently, the model system has been deployed nationwide in the AGRUM-DE project (Schmidt et al., 2020). Model results from this project have been used as a basis of information to help German authorities identify nitrate polluted areas, in accordance with the German Fertilizer Ordinance (Fertilizer Ordinance, 2020), and prepare river basin management plans (2021–2027) under the EU Water Framework Directive (The European Parliament and The Council of the European Union, 2000).

The model system has been proven to be a capable instrument for evaluating N-related issues at a high spatial resolution. The main aim of this study was to apply the RAUMIS-mGROWA-DENUZ model system




FIGURE 1 GHG inventory time series of national N surplus (Gg N) for the inventory time period and calibration data. Points depict annual inventory N surpluses. In order to highlight trends, a curve was fitted using local polynomial regression (LOESS) with a span equal to 0.3.

to derive regionally differentiated $\text{Frac}_{\text{LEACH}}$ values thereby enabling quantification of indirect agricultural N_2O emissions for GHG reporting in a more suitable way than the estimation based on the global IPCC default approach.

2 | MATERIALS AND METHODS

The aforementioned models were used and linked to model high-resolution N surplus and resulting N losses through leaching and runoff for the years 2014–2016, calculated as an average over the 3 years to reduce the impact of outliers. Estimations of N surplus excluded atmospheric N deposition. All units indicating hectare refer to hectare agricultural area, excluding all other land uses. A depiction of the workflow of the applied methodology is supplied in the supporting information (Figure 1 and Supporting Information 1).

2.1 | Data

The model input database was compiled for the years 2014–2016, primarily from official statistical information. For RAUMIS land use and livestock production, data at municipality level were taken from the Thünen-Agraratlas (Processed data. Original data from statistical offices of the federal states [district data from agricultural census 2016]; research data center of Germany and the federal states, agricultural census 2010, and und AFiD panel “Agrarstruktur” 1999, 2003, 2007, 2016 [own calculations: 1999–2016. Cluster estimator]; © GeoBasis-DE/BKG [2016]. Derived using the method of Gocht & Röder [2014]. Version 2020.). Crop yield data were obtained from the Federal Statistical Office (www.destatis.de). Data on agricultural use of compost and sewage sludge were available at NUTS-1 level (Landsamt für Statistik Niedersachsen, 2019; StBA, vy.). Data on biogas plants and biogas production were assembled from publicly available

TABLE 1 German NUTS-1 regions and their abbreviations

Region (NUTS-1)	Abbreviation
Northrhine-Westphalia	NW
Lower Saxony North-West and Bremen	NI-W + HB
Lower Saxony South-East	NI-E
Bavaria	BY
Mecklenburg Western Pomerania	MV
Saxony-Anhalt	ST
Schleswig Holstein and Hamburg	SH + HH
Brandenburg and Berlin	BB + BE
Saxony	SN
Baden-Württemberg	BW
Thuringia	TH
Hesse	HE
Rhineland-Palatinate and Saarland	RP + SL

databases of the plant register (www.marktstammdatenregister.de) and national network operators. Substrate use data for biogas production were derived from Daniel-Gromke et al. (2017) and federal state authority data. Manure transportation data were available for some states from federal state authorities. National data on the use of mineral fertilizer were taken from the mineral fertilizer sales statistics (StBA, 2019).

For mGROWA-DENUZ, the German soil map 1:200,000 (BUEK200) (BGR, 2018) with horizon-wise data on soil type, humus content, field capacity, depth, waterlogging and groundwater influence was used. Long-term data on precipitation and the grass reference evapotranspiration were obtained from the German Meteorological Service. Land use and topography data were taken from the Federal Agency for Cartography and Geodesy. A detailed list of all data used for RAUMIS-mGROWA-DENUZ is supplied in the supporting information (Table 1 and Supporting Information 1).

For calculation of indirect N_2O emissions from leaching and runoff, N inputs as reported in the GHG inventory (Rösemann et al., 2021) were used. N removal by harvest and grazing was estimated from harvest data (StBA, FS3 R3.1.3, FS3 R3.2.1). N inputs and harvest data are a mandatory part of the annual GHG reporting for countries listed in the UNFCCC's Annex 1.

2.2 | N surplus model

N surpluses were modeled using the Regional Agricultural and Environmental Information System (RAUMIS). RAUMIS is a regional agricultural supply model based on "regional farms" at NUTS-3 level (Eurostat, 2020), developed in the 1990s as a tool for spatial agricultural policy analysis (Henrichsmeyer et al., 1996). It represents agricultural production, factor input and income on a regional level based on statistical information and a number of environmental indicators such as N balances. For the latter, an extension at municipality level is available,

allowing the high-resolution estimation of N surpluses, while also being able to include a variety of heterogeneous data sources. A number of studies conducting agricultural policy analysis of agricultural N and GHG emissions in Germany have been carried out using the model system (e.g., Ackermann et al., 2016; Gömann et al., 2002; Henseler & Dechow, 2014; Kreins et al., 2007). N surpluses have been modeled in the form of net area balances for the period 2014–2016 as part of the AGRUM-DE project (Schmidt et al., 2020). The RAUMIS database also comprises regional yields, spatial information on biogas plants, and manure transportation data.

Let k represent German municipalities, A_{ck} the area of a specific crop c in a municipality k . M_k is the total amount of mineral N, and N_{jk} the total amount of organic N applied from source j (animal manure, biogas digestates, sewage sludge and compost). F_{ck} represents crop-specific N fixation rates from legumes, Y_{ck} is regional per hectare crop yields, xn_c is the respective crop's N content and G_{jk} are total gaseous emissions from organic fertilizer application. N field surpluses NS_k are thus calculated as:

$$NS_k = \sum_j (N_{jk} - G_{jk}) + M_k + \sum_c F_{ck} * A_{ck} - \sum_c Y_{ck} * A_{ck} * xn_c, \quad (2)$$

with $G_{jk} = 0$ for $j \in$ (compost, sewage sludge).

Data on regional mineral fertilizer use are usually not available. Therefore, mineral fertilizer application in municipality k is modeled by means of yield-dependent and crop-dependent linear N requirement functions. The total N requirement per municipality TR_k is calculated as:

$$TR_k = xs_k * \sum_c [(a_c * Y_{ck} + b_c) * A_{ck}], \quad (3)$$

where xs_k reflects local site specifics such as soil and climate conditions, affecting TR_k (Krüll, 1988; as cited in Henrichsmeyer et al., 1996), and a_c and b_c are individual parameters of the model's N requirement function for every crop (Kreins et al., 2009). These functions are derived from national fertilization recommendations and reflect regional yield levels as well as site conditions (Henrichsmeyer et al., 1996). Finally, total mineral fertilizer application at municipality level is represented by:

$$M_k = \beta * TR_k - \alpha * \sum_j (N_{jk} - G_{jk}), \quad (4)$$

taking into consideration the share of total organic N being available for the crops (α) and a calibration factor (β). These factors were chosen in a way that $\sum_{k=1}^n M_k$ equals the amount of mineral fertilizer reported in the national mineral fertilizer sales statistics (StBA, 2019). For this study, aggregate N surpluses at NUTS-3 level were used.

2.3 | Model of nitrate leaching

The N surplus of agriculturally used soils determined with the RAUMIS model does not generally correspond with the N output from these

soils. This is due to microbial conversion processes by which the surplus is reduced into gaseous N compounds that can leave the soil and enter the atmosphere. An area-differentiated simulation of these N conversion processes in the soil has been carried out using the DENUZ model (Kunkel & Wendland, 2006; Wendland, 1994).

For grassland, it is assumed that 30% of the sum of N balance surpluses from agriculture and atmospheric deposition is stored in the soil and contributes to the formation of soil organic matter (Wendland, Herrmann et al., 2020). For arable land, it is assumed that no immobilization occurs as these soils are N-saturated after decades of fertilization (Wendland et al., 2009). Accordingly, the N output from arable soils corresponds to the leachable NO_3^- -N in soil (N surpluses from agriculture plus atmospheric deposition minus N immobilization) minus the denitrification losses in the root zone of the soil. For the purposes of this study, N input from atmospheric deposition was excluded to avoid double counting of indirect emissions from volatilization in the GHG inventory. The process of denitrification is simulated in the DENUZ model for both grassland and arable land based on Michaelis–Menten kinetics:

$$\frac{dN(t)}{dt} + D_{\max} \times \frac{N(t)}{k + N(t)} = 0, \quad (5)$$

where $N(t)$ = N content in soil after time t (kg N [ha y]^{-1}); t = residence time of the leachate in soil (y); D_{\max} = maximum denitrification rate (kg N [ha y]^{-1}); k = Michaelis constant (kg N [ha y]^{-1}).

The first step in modeling NO_3^- degradation in soil with the DENUZ model was to classify the denitrification conditions of soils throughout the study area. This then allowed the assignment of regional soil type-specific maximum denitrification rates (D_{\max}) per year. The classification of NO_3^- degradation conditions in the soil depends on various soil properties. For example, high water storage capacities and humus contents are favorable for achieving high denitrification rates in the soil, whereas low denitrification rates can occur in soils with limited water storage capacities and reduced humus content (Köhne & Wendland, 1992; Kunkel et al., 2010; Wendland et al., 1998, 2005; Wendland, Herrmann et al., 2020; Wienhaus et al., 2008).

The D_{\max} values applied in this study originated from relevant research work in the federal states of Mecklenburg-Western Pomerania (Kunkel et al., 2017), Schleswig-Holstein (Wendland et al., 2014), Saxony-Anhalt (Kühr et al., 2014), Lower Saxony (Heidecke et al., 2016), North Rhine-Westphalia (Kühr et al., 2013; Wendland, Bergmann et al., 2020), Rhineland-Palatinate (Wendland et al., 2021), and Thuringia (Tetzlaff et al., 2016), as well as the Weser River basin (Kuhn et al., 2016). These studies distinguished between five classes of denitrification conditions, depending on the initial substrate of the soil stratum, the humus content and the influence of groundwater and waterlogging. They found that the denitrification rates assigned to the denitrification conditions in the soil during model parameterization can differ significantly from region to region, therefore regionally differentiated D_{\max} values have been allocated to the soil types indicated in the German soil map 1:200,000 (BUEK200) (Federal Institute for Geosciences and Natural Resources, 2018). Value ranges according to Wienhaus et al. (2008) were used as starting points for the regionally differentiated parameterization of D_{\max} for different soil types.

The residence time of the leachate in the soil was derived from the field capacity of the soil and the leachate rate (Hennings, 2000; Müller & Raissi, 2002), in which the index i is run over all the denitrifying layers in the soil profile:

$$t = \frac{1}{Q_{SW}} \sum_i nFK_i \times d_i, \quad (6)$$

where t = residence time of the leachate in soil (y); Q_{SW} = leachate rate (mm year^{-1}); nFK = effective field capacity (mm dm^{-1}); d = thickness of soil layer (dm).

Using the kinetic parameters of denitrification, the initial displaceable N surpluses in the soil and the residence times of leachate in the soil, the Michaelis–Menten differential equation can be solved numerically and the remaining N outputs from the soil can be calculated. It is then possible to combine the N emissions from the soil with the leachate rate (Q_{SW}) in order to calculate the potential NO_3^- concentration in the leachate ($C_{\text{NO}_3^-}$):

$$C_{\text{NO}_3^-} = \frac{443 \times N(t)}{Q_{SW}}, \quad (7)$$

where $C_{\text{NO}_3^-}$ = potential NO_3^- concentration in the leachate (mg L^{-1}); $N(t)$ = N output from soil after residence NO_3^- time t in soil (kg N [ha y]^{-1}); Q_{SW} = leachate rate (mm year^{-1}).

The leachate rate (Q_{SW}) is an important parameter for calculating both, residence time in soil according to (6) and the NO_3^- concentration in the leachate (7). In this study, leachate rate was determined on the basis of the mGROWA model (Herrmann et al., 2015), which simulates the hydrologic processes at the earth's surface and in the root zone of soils. In particular, soil moisture dynamics including the movement of the leachate in the soil, capillary rise from groundwater to the root zone, actual evapotranspiration, and total runoff generation were calculated in daily time steps on the basis of grass reference evapotranspiration, land use-specific crop coefficients, and a topography correction function (Wendland, Bergmann et al., 2020). Leachate rates, that is, the amount of water that leaves the root zone of the soil vertically downward, are aggregated to mean long-term averages in order to exclude short-term climate-induced blurring of leachate rates and to guarantee that the modeled NO_3^- concentrations in the leachate are representative for the regional long-term hydrologic conditions (Wendland, Bergmann et al., 2020).

Against this background, the NO_3^- concentration in the leachate is calculated based on the DENUZ model. In this model, the mean long-term leachate rates from the mGROWA model are combined with the N output from the root zone of soil after residence time in soil. For the latter, agricultural N balance surpluses are considered as N input and denitrification (5) and immobilization processes in soil as N losses.

2.4 | Creating a time series of nitrate leaching for the national inventory

To enable estimation of indirect N_2O emissions from N leaching for national GHG reporting, data for the whole inventory time series are

needed. Since high-quality input data for the leaching model were only available as the mean of the years 2014 to 2016, a regional transfer coefficient was calculated:

$$F_{\text{Leach,Surplus}} = \frac{N_{\text{Leach,ref}}}{N_{\text{Surplus,ref}}}, \quad (8)$$

where $N_{\text{Leach,ref}}$ is the average annual amount of N leached in 2014–2016 modeled with RAUMIS-mGROWA-DENUZ and $N_{\text{Surplus,ref}}$ is the average N surplus in the same years derived from data available in the German national inventory (Federal Environment Agency, 2021). The transfer coefficient was then multiplied with the inventory N surplus for each region and year in order to derive the time series of leached and runoff N. Since the inventory data do not currently include data on manure transports and also contain substantial uncertainties regarding spatial distributions, the transfer coefficients were calculated at NUTS-1 level. Only Lower Saxony was divided into two regions: the north-west of the state (NI-W), which has the highest livestock density in Germany, and the south-east of the state (NI-E), where cash-crop farming is more prevalent. City states were merged with neighboring federal states, that is, Hamburg with Schleswig-Holstein, Bremen with Lower Saxony (north-west), and Berlin with Brandenburg (Table 1).

For the calculation of time series, N surplus was derived from N inputs reported in the inventory (Rösemann et al., 2021), that is, synthetic fertilizers, manure application, crop residues, grazing, sewage sludge, biogas digestates, and mineralization of soil organic matter. N input from N deposition was excluded to avoid double counting of indirect emissions from volatilization, which are already estimated by the inventory and include, at least in theory, the pathway N deposition–N leaching–denitrification. N removal by harvest and grazing was estimated from harvest data (StBA, FS3 R3.1.3, FS3 R3.2.1).

As the model calculates leached N out of the root zone, the combined Tier 1 EF ($EF_5 = 0.011 \text{ kg N}_2\text{O-N} [\text{kg N leaching/runoff}]^{-1}$) for indirect N_2O emissions from leaching and runoff in downstream water bodies can be applied. A multiplication of the leached/runoff N with the EF gave the indirect N_2O emissions. For comparison with the IPCC Tier 1 $\text{Frac}_{\text{LEACH}}$ value (IPCC, 2019), implied $\text{Frac}_{\text{LEACH}}$ values were calculated as N leached/runoff divided by the inventory's N input for the respective year and region.

2.5 | Validation of N leaching model and uncertainty estimates

Modeled N outputs from soil can be validated using measured values from soil depth profiles, suction probes, and lysimeters. However, such measurements are only available in a few cases and do not allow a plausibility check to be undertaken on a national scale (Wolters et al., 2021). Instead, modeled concentrations in leachate can be compared with measured NO_3^- values from the upper aquifer, for which more data are available.

However, monitoring sites need to meet certain preselection criteria (Wolters et al., 2021). These include the measuring points being either springs or groundwater observation wells filtered near the

surface. Furthermore, the redox indicators should show an oxidized groundwater milieu so that denitrification processes in groundwater can be excluded as much as possible. In addition, the catchment area of the measuring point must be identified to calculate an average modeled NO_3^- concentration in the leachate for each area. If these criteria are met, observed NO_3^- concentrations in groundwater can be used to check the plausibility of the simulated NO_3^- concentrations in the leachate and thus indirectly the plausibility of the simulated N emissions from the soil.

Due to the different spatial and temporal reference of random point measurements in groundwater and model results for mean long-term NO_3^- concentration in leachate, but also due to the limited site-specific accuracy of the input data for modeling, an exact site-specific agreement of model values and measured values can hardly be expected. In order to assess systematically and comprehensively if the spatial patterns and magnitudes of modeled NO_3^- concentrations in the leachate are confirmed by measured NO_3^- values from the upper aquifer, the observed and measured values were first categorized. Subsequently, the compliance of the measured and observed values to the resulting classes was assessed (Wolters et al., 2021). The class width was determined by considering the number and range of the measured values. Since there was a very large range in the measured values and the modeled values, a uniform class width of $25 \text{ mg NO}_3^-/\text{L}$ was defined to ensure comparability. The comparison was based on seven classes (0–25, 25–50, ..., $>150 \text{ mg NO}_3^-/\text{L}$). The agreement between modeled and observed NO_3^- levels in each class was then evaluated. The difference between the classes was used as a measure of the assessment. In this context, agreement is considered good if the observed and measured concentrations are in the same class. Acceptable agreement is when there is a deviation of one concentration class. If the deviation is two concentration classes or more, the agreement is moderate or poor.

Since the uncertainties and covariances included in many model input parameters were not available for this study, it was not possible to perform a traditional uncertainty analysis based on error propagation. Nevertheless, as modeled NO_3^- concentrations in the leachate were validated using measured values from the upper aquifer (Wolters et al., 2021), an approximate uncertainty estimate could be derived.

3 | RESULTS

The resulting 3-year average N surplus for Germany's agricultural area for the calibration data was calculated to be $951.2 \text{ Gg N year}^{-1}$ (corresponds to $57.2 \text{ kg N ha}^{-1} \text{ year}^{-1}$). For the same period, the inventory calculates a mean N surplus of $1074.6 \text{ Gg N year}^{-1}$ (corresponds to $64.3 \text{ kg N ha}^{-1} \text{ year}^{-1}$) ($N_{\text{Surplus,ref}}$). Overall, since 1990, the nationwide inventory N surplus has decreased by approximately 41% from $1637 \text{ Gg N year}^{-1}$ to 966 Gg N a^{-1} (Figure 1). In recent years, there has been a slight increase in the N surplus due to consecutive years with very dry conditions, leading to lower crop yields and hence reduced N removal through N uptake by plants and through harvest (Klages et al., 2020). However, the development of the N surplus in Germany's NUTS-1 regions is heterogeneous (Figure 2). All regions experienced a strong




FIGURE 2 GHG inventory time series of N surplus per hectare agricultural area in the study regions. Points depict modeled annual values for each study region. In order to highlight trends, a curve was fitted using local polynomial regression (LOESS) with a span equal to 0.3.




FIGURE 3 GHG inventory time series of lost N per hectare agricultural area to water bodies through leaching and runoff in the study regions. Points depict modeled annual values for each study region. In order to highlight trends, a curve was fitted and smoothed using local polynomial regression fitting (LOESS) with a span equal to 0.3.

decline until 1995, with a moderate increase in the early 2000s due to the introduction of anaerobic digestion plants for biogas production. In regions with a high and increasing livestock density (NI-W + HB), since around 2000, the N surplus has grown significantly.

In the reference period, an estimated $N_{\text{Leach,ref}}$ of 385 Gg N year⁻¹ (corresponds to 23.1 kg N ha⁻¹ year⁻¹) left the root zone and was lost to water bodies through leaching and runoff. The average national $F_{\text{Leach,ref}}$ (Equation 8) is 0.36. Leached N was calculated as the product of inventory N surplus and $F_{\text{Leach,ref}}$ for each year and region. Figure 3 shows the N leached per hectare since 1990. Brandenburg and Saxony-Anhalt had the lowest and most stable leaching rates because they are generally dryer regions with moderate surpluses. The manure-abundant western part of Lower Saxony and North Rhine-Westphalia,

TABLE 2 Calculated annual amounts of leached N (Gg N) from agricultural soils in German NUTS-1 regions for the GHG inventory time series

Region	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
SH + HH	59.2	50.3	56.0	41.9	38.9	38.3	42.6	35.7	39.3	40.7	42.3	37.3	41.4	47.4	41.3	39.6	42.9	43.1	37.3	36.8	37.6	38.9	34.3	40.4	32.6	36.5	35.6	33.4	48.9	25.9
NI-W + HB	32.5	29.0	32.6	27.3	26.1	25.7	27.9	24.5	27.6	27.9	26.8	27.9	28.6	34.2	26.4	28.2	31.5	29.6	30.4	35.0	34.8	33.0	35.8	41.1	37.1	40.1	41.3	42.4	49.3	43.5
NI-E	55.8	48.0	51.4	43.3	39.2	37.1	39.5	34.1	38.2	40.7	38.2	37.2	39.5	45.6	34.0	35.5	37.2	36.2	28.9	29.5	36.6	34.6	37.0	37.8	33.8	36.6	39.6	39.9	48.8	38.8
NW	77.6	65.4	66.9	60.3	61.6	59.0	58.6	56.0	60.7	64.0	60.5	57.0	55.5	62.0	52.9	54.3	54.9	56.4	47.7	51.1	55.6	58.3	57.7	61.2	60.6	66.1	68.4	62.5	68.8	57.4
HE	35.7	35.0	29.8	27.5	23.7	22.9	22.5	23.8	18.6	21.4	18.2	20.0	18.6	25.5	19.1	19.8	17.9	17.0	15.8	12.5	15.7	17.7	15.3	10.4	11.2	17.9	15.7	11.8	19.8	10.9
RP + SL	48.2	45.6	38.1	34.7	29.3	27.1	30.0	27.7	27.0	32.2	28.3	29.5	25.6	34.0	22.9	27.3	23.4	22.5	16.6	15.6	15.6	23.6	19.0	13.9	14.8	21.8	19.5	11.3	16.8	10.0
BW	48.2	44.6	41.2	32.9	32.9	34.1	31.5	30.8	32.1	35.7	34.7	33.9	31.4	44.9	30.0	31.3	30.0	27.6	24.6	23.2	26.7	28.2	26.1	28.0	26.0	37.0	30.0	22.4	29.6	17.7
BY	89.1	79.5	75.1	65.5	62.4	61.9	61.0	59.4	57.0	68.3	62.2	64.7	54.5	79.1	45.1	49.4	48.0	45.0	41.5	40.4	51.0	50.7	55.1	63.4	46.7	71.3	53.4	42.2	55.7	42.7
BB + BE	17.4	15.3	18.2	12.1	12.3	11.4	11.5	11.4	11.5	13.1	14.3	11.5	11.4	16.2	10.6	10.5	11.6	11.0	10.0	9.1	8.9	9.9	9.0	9.4	7.7	10.4	10.8	8.3	12.2	9.5
MV	42.9	36.4	40.7	30.3	30.9	25.7	29.6	25.1	23.5	27.6	28.7	25.8	25.5	29.3	22.2	23.4	23.4	24.9	20.3	19.8	23.9	25.3	23.3	23.1	19.2	22.7	30.7	24.5	31.2	22.3
SN	36.6	32.2	34.4	25.1	26.9	24.9	26.4	24.7	25.7	28.1	30.1	27.5	27.5	34.4	23.3	24.2	26.4	23.8	21.2	20.9	21.8	21.6	22.8	24.6	18.6	24.7	20.4	21.2	26.5	21.3
ST	23.2	19.3	23.1	17.9	16.1	14.6	15.2	14.8	15.2	15.9	16.9	15.0	15.9	17.8	13.0	14.0	14.0	14.1	11.3	11.6	11.4	13.2	12.5	13.5	11.5	14.9	14.4	12.8	16.5	14.6
TH	34.5	31.4	31.0	25.1	24.6	22.7	23.9	23.6	23.6	24.6	25.9	23.4	24.5	29.0	21.0	22.7	21.5	21.2	20.1	17.4	17.8	19.3	19.4	18.8	15.5	21.3	17.2	15.8	21.6	15.8
Germany	600.9	532.0	538.4	443.7	425.0	405.5	420.0	391.6	400.0	440.3	427.2	410.6	400.0	499.4	361.7	380.2	382.8	372.5	325.8	323.0	357.3	374.1	367.2	385.8	335.3	421.2	397.0	348.5	445.7	330.4

TABLE 3 Annual indirect N_2O emissions ($Gg\ N_2O$) based on new implied $Frac_{LEACH}$ values in German NUTS-1 regions using the 2019 IPCC guidelines emissions factor ($EF_5 = 0.011\ kg\ N_2O-N$ [$kg\ N\ leaching/runoff$] $^{-1}$)

Region	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
SH + HH	1.02	0.87	0.97	0.72	0.67	0.66	0.74	0.62	0.68	0.7	0.73	0.64	0.71	0.82	0.71	0.68	0.74	0.75	0.64	0.64	0.65	0.67	0.59	0.7	0.56	0.63	0.62	0.58	0.85	0.45
NI-W + HB	0.56	0.5	0.56	0.47	0.45	0.44	0.48	0.42	0.48	0.48	0.46	0.48	0.49	0.59	0.46	0.49	0.55	0.51	0.53	0.6	0.6	0.57	0.62	0.71	0.64	0.69	0.71	0.73	0.85	0.75
NI-E	0.97	0.83	0.89	0.75	0.68	0.64	0.68	0.59	0.66	0.7	0.66	0.64	0.68	0.79	0.59	0.61	0.64	0.63	0.5	0.51	0.63	0.6	0.64	0.65	0.58	0.63	0.68	0.69	0.84	0.67
NW	1.34	1.13	1.16	1.04	1.07	1.02	1.01	0.97	1.05	1.11	1.05	0.99	0.96	1.07	0.91	0.94	0.95	0.98	0.83	0.88	0.96	1.01	1	1.06	1.05	1.14	1.18	1.08	1.19	0.99
HE	0.62	0.6	0.52	0.47	0.41	0.4	0.39	0.41	0.32	0.37	0.31	0.35	0.32	0.44	0.33	0.34	0.31	0.29	0.27	0.22	0.27	0.31	0.26	0.18	0.19	0.31	0.27	0.2	0.34	0.19
RP + SL	0.83	0.79	0.66	0.6	0.51	0.47	0.52	0.48	0.47	0.56	0.49	0.51	0.44	0.59	0.4	0.47	0.4	0.39	0.29	0.27	0.27	0.41	0.33	0.24	0.26	0.38	0.34	0.2	0.29	0.17
BW	0.83	0.77	0.71	0.57	0.57	0.59	0.54	0.53	0.55	0.62	0.6	0.59	0.54	0.78	0.52	0.54	0.52	0.48	0.42	0.4	0.46	0.49	0.45	0.48	0.45	0.64	0.52	0.39	0.51	0.31
BY	1.54	1.37	1.3	1.13	1.08	1.07	1.05	1.03	0.98	1.18	1.08	1.12	0.94	1.37	0.78	0.85	0.83	0.78	0.72	0.7	0.88	0.88	0.95	1.1	0.81	1.23	0.92	0.73	0.96	0.74
BB + BE	0.3	0.26	0.32	0.21	0.21	0.2	0.2	0.2	0.2	0.23	0.25	0.2	0.2	0.28	0.18	0.18	0.2	0.19	0.17	0.16	0.15	0.17	0.16	0.16	0.13	0.18	0.19	0.14	0.21	0.16
MV	0.74	0.63	0.7	0.52	0.53	0.45	0.51	0.43	0.41	0.48	0.5	0.45	0.44	0.51	0.38	0.4	0.41	0.43	0.35	0.34	0.41	0.44	0.4	0.4	0.33	0.39	0.53	0.42	0.54	0.39
SN	0.63	0.56	0.59	0.43	0.46	0.43	0.46	0.43	0.44	0.49	0.52	0.47	0.47	0.59	0.4	0.42	0.46	0.41	0.37	0.36	0.38	0.37	0.39	0.42	0.32	0.43	0.35	0.37	0.46	0.37
ST	0.4	0.33	0.4	0.31	0.28	0.25	0.26	0.26	0.26	0.28	0.29	0.26	0.28	0.31	0.22	0.24	0.24	0.24	0.2	0.2	0.2	0.23	0.22	0.23	0.2	0.26	0.25	0.22	0.28	0.25
TH	0.6	0.54	0.54	0.43	0.43	0.39	0.41	0.41	0.41	0.43	0.45	0.4	0.42	0.5	0.36	0.39	0.37	0.37	0.35	0.3	0.31	0.33	0.34	0.33	0.27	0.37	0.3	0.27	0.37	0.27
Germany	10.39	9.2	9.31	7.67	7.35	7.01	7.26	6.77	6.91	7.61	7.38	7.1	6.91	8.63	6.25	6.57	6.62	6.44	5.63	5.58	6.18	6.47	6.35	6.67	5.8	7.28	6.86	6.02	7.7	5.71




FIGURE 4 GHG inventory time series of $Frac_{LEACH}$ in the study regions compared with IPCC default values (IPCC, 2006, 2019). Points depict modeled annual values for each study region. In order to highlight trends, a curve was fitted using local polynomial regression (LOESS) with a span equal to 0.3.

both regions with the highest stocking densities in Germany, had the highest leaching rates per hectare. Most other regions showed moderate leaching rates of around $20\ kg\ N\ ha^{-1}$. Tables 2 and 3 shows the calculated absolute amounts of leached N leaving the root zone. Since 1990, the total amount of leached N decreased by 45% from 600.9 to 330.4 $Gg\ N$ in 2019. Whereas all other regions experienced a decrease, in NI-W + HB the amount of leached N increased since 1990.

3.1 | $Frac_{leach}$

To derive a time series of implied $Frac_{LEACH}$ values, leached N was divided by the N input from the inventory. The resulting $Frac_{LEACH}$ for each year is shown in Figure 4. In 2019, $Frac_{LEACH}$ in all regions was below the IPCC default value of $0.24\ kg\ N\ (kg\ N\ input)^{-1}$ (IPCC, 2019), with a nationwide average of $0.099\ kg\ N\ (kg\ N\ input)^{-1}$ and a range between $0.051\ kg\ N\ (kg\ N\ input)^{-1}$ (BB + BE) and $0.159\ kg\ N\ (kg\ N\ input)^{-1}$ (NW) in 2019. Since 1990, most regions have shown a general downwards trend, with a slight increase recently mainly due to drought conditions during the vegetation period.

3.2 | N_2O emissions

Indirect N_2O emissions were calculated for the whole time series in accordance with the 2019 IPCC guidelines (IPCC, 2019). Based on the new estimation, emissions decreased from 10.4 $Gg\ N_2O$ in 1990 to 5.7 $Gg\ N_2O$ in 2019 (Figure 5, Tables 2 and 3). The largest reduction in indirect N_2O emissions from leaching and runoff occurred until 1995, with a decrease in emissions of 3.38 $Gg\ N_2O$. While lower on average, indirect N_2O emissions have experienced only a slight downwards trend since then. Due to higher N surpluses in recent years, emissions have increased slightly since 2015. In 2019, emissions were again




FIGURE 5 Estimated annual national indirect N_2O emissions based on the 2006 IPCC guidelines Tier 1 method and based on new implied $\text{Frac}_{\text{LEACH}}$ value in inventory time series together with national average wheat yields (StBA, FS3 R3.2.1).

below 6 Gg N_2O . Compared with the current methodology (IPCC, 2006; Tier 1) used in the German inventory (Federal Environment Agency, 2021), the new Tier 3 method leads to a nationwide reduction of indirect N_2O emission estimates of 27% in 1990 and 52.1% in 2019. Overall, indirect N_2O emissions calculated with the new method showed a greater variance, since N surplus and N leaching varied depending on the environmental conditions in the respective years, compared with the calculations with the constant IPCC default Tier 1 approach.

3.3 | Validation of nitrate leaching model and uncertainty estimates

Figure 6 shows the comparison of the class widths of modeled NO_3^- concentrations in leachate and measured NO_3^- concentrations in groundwater as a frequency distribution for 1496 preselected groundwater quality monitoring wells in the land use categories cropland (left) and grassland (right).

For the land-use category cropland, 1068 groundwater monitoring wells remained for comparison after preselection. Here, the modeled values show a good agreement at 30% of the monitoring sites, while a deviation of one class occurred at 40% of the monitoring sites. Overall, there was no tendency to systematically overestimate or underestimate the measured values. The corresponding deviations exhibit a symmetric distribution and thereby indicate that the overall system, which determines the NO_3^- concentrations in the leachate, is well represented.

For the land use category grassland 428 monitoring sites were available for comparison. At these monitoring sites, 62% show good agreement in NO_3^- concentration classes, while 29% overestimate or underestimate concentrations by one class. Although the frequency distribution shows a slightly higher number of groundwater monitoring sites where modeled NO_3^- concentrations in leachate were lower

than measured NO_3^- concentrations in groundwater, the overall agreement of modeled and observed NO_3^- concentrations for the grassland land-use category can be considered very good.

For the approximate uncertainty estimate the percentage mean deviation of the modeled NO_3^- concentrations from measured concentrations in groundwater wells (-100% , $+200\%$) was transferred to modeled $\text{Frac}_{\text{LEACH}}$ values, resulting in a 95% confidence interval of (0, 0.198) kg N (kg N input) $^{-1}$ for 2019, which is considerably narrower than that of the IPCC Tier 1 value.

4 | DISCUSSIONS

The aim of this study was to apply a model-based approach to estimate indirect N_2O emissions through N leaching and runoff from agricultural soils for Germany's national GHG inventory. A comprehensive model system was used to estimate regionally differentiated, country-specific $\text{Frac}_{\text{LEACH}}$ values to supersede the IPCC Tier 1 default value. The link of an agricultural economic model with hydrological models and the use of high-resolution spatial data allowed the representation of a wide range of different hydrological and hydrogeological conditions, as well as the drivers of the agrarian structure in Germany. Similar to the Tier 2 and Tier 3 approaches of other countries (UNFCCC, 2021), the newly estimated $\text{Frac}_{\text{LEACH}}$ was lower than the IPCC default EF, indicating an overestimation by the IPCC Tier 1 value. Germany's implied nationwide $\text{Frac}_{\text{LEACH}}$ of 0.099 kg N (kg N input) $^{-1}$ in 2019 is similar to other countries that use Tier 3 approaches. The Netherlands reports a $\text{Frac}_{\text{LEACH}}$ value of 0.13 kg N (kg N input) $^{-1}$ for 2019 based on STONE model results (Groenendijk et al., 2008; van der Zee et al., 2021; Velthof & Mosquera, 2011), Ireland uses a national average value of 0.1 kg N (kg N input) $^{-1}$ (Environmental Protection Agency, 2021) and New Zealand reports a value of 0.07 kg N (kg N input) $^{-1}$ (Ministry for the Environment, 2021).

Räbiger et al. (2020) modeled N leaching and indirect N_2O emissions for oilseed rape from five sites with different growing conditions (e.g., soil, temperature, precipitation) in the main rapeseed-growing areas in Germany. Even though oil seed rape is a crop with low N efficiency and high leaching potential (Räbiger et al., 2020), they modeled $\text{Frac}_{\text{LEACH}}$ values between 0.05 and 0.176 kg N (kg N input) $^{-1}$ depending on the site and fertilizer used. Indirect N_2O emissions were calculated to be 60–90% lower than those calculated using the IPCC default values. Fu et al. (2017) used lysimeters to measure N leaching on intensively and extensively managed montane cut grassland in southern Germany. They observed values of between 0.008 and 0.069 kg N (kg N input) $^{-1}$. Since N leaching rates from grassland are generally lower (Di & Cameron, 2002), the new estimate in the present study is in agreement with their findings.

Räbiger et al. (2020) emphasizes the need for site-specific EFs. This is confirmed by the comparatively wide range of estimated $\text{Frac}_{\text{LEACH}}$ values of the respective regions (2019: 0.051–0.159 kg N [kg N input] $^{-1}$) in the present study. As N leaching depends on multiple factors, such as soil properties of the topsoil, N denitrification potential and N input, a country-specific calculation of $\text{Frac}_{\text{LEACH}}$ based only on




FIGURE 6 Frequency distribution of the deviation classes of simulated NO_3^- concentrations in the leachate and observed NO_3^- concentrations in the groundwater for the main land use types “arable land” (left) and “grassland” (right).

water balances, as used by some countries in their inventories as an IPCC Tier 2 approach, falls short of including all relevant controlling factors. $\text{Frac}_{\text{LEACH}}$ is highly dependent on denitrification conditions in the soil. Figure 7 shows the heterogeneity of denitrification losses in Germany. As outlined above, they depend not only on the denitrification conditions in the soil, but also on the residence time of the leachate in the soil. Thus, high denitrification rates were calculated for regions exhibiting high denitrification capacities in soil (lowland areas, soils with high organic carbon content), and regions exhibiting long residence times (loess regions), whereas low denitrification rates were calculated for the midland region, where low denitrification capacities in soil coincide with low residence times in soil. Along with differences in N surpluses in every region, this explains regionally different trends in the dynamics of $\text{Frac}_{\text{LEACH}}$. For instance, the regions of Rhineland-Palatinate and Saarland showed a strong decline in $\text{Frac}_{\text{LEACH}}$ between 1990 and 1995. In this five-year period, N surplus and N losses through leaching decreased by nearly 60%, whereas N inputs decreased by just 15%. And even though the N leaching rate in this area is the highest in Germany, $\text{Frac}_{\text{LEACH}}$ decreased greatly as a result. In contrast, denitrification capacities in the soil are generally high in north-west Germany (NW, NI-W + HB), yet high N surpluses led to the region having the highest $\text{Frac}_{\text{LEACH}}$ values in the country.

Substantial annual changes in indirect N_2O emissions could be observed because the use of N surplus rather than N inputs as the source of N leaching allowed the effects of annual changes in growing conditions to be reflected. High N inputs only result in high leaching when N uptake of the crops is low. In contrast to N input as




FIGURE 7 Denitrification loss in soil based on displaceable nitrogen input into soil, soil residence time, and maximum denitrification degradation according to the DENUZ model (Wendland et al., 2009).

an indicator, N surplus also considers N removal via harvest. A main cause of higher N surpluses is unfavorable growing conditions. Farmers plan crop N inputs at the beginning of the growing season, as required by the German Fertilizer Ordinance (Fertilizer Ordinance, 2020). Poor growing conditions lead to impeded crop growth and reduced N requirements. Consequently, N inputs are likely to exceed N uptake, and the increased N surpluses lead to higher N leaching risks, resulting in elevated indirect N_2O emissions. To illustrate this, national average wheat yields can be used (Figure 5). The years 2003 and 2018 were exceptionally dry and resulted in lower wheat yields. The N surplus increased N leaching and there was a rise in indirect N_2O emissions. Indirect N_2O emissions calculated based on the new $\text{Frac}_{\text{LEACH}}$ showed a highly significant, strongly negative correlation ($r = -0.71$, $p < 0.001$) with the national average wheat yield, whereas indirect N_2O emissions estimated based on the default $\text{Frac}_{\text{LEACH}}$ showed no significant correlation ($r = 0.19$, $p = 0.29$). Furthermore, in contrast to an input-based methodology, the N surplus-based approach was able to reflect altered management practices and technological progress regarding N-use efficiency.

In the inventory period since 1990, an overall reduction in N surplus and hence in indirect N_2O emissions can be observed. The magnitude and development of the N surpluses (Figures 1 and 2) estimated in this study agree with findings of Häußermann et al. (2020), based on a similar database as used in this study. They attribute the initial reduction of N surplus to the European Union's policy aimed at reducing N surplus and N leaching to protect water bodies from N pollution, formulated in the Nitrate Directive (The Council of the European Communities, 1991) and in its national implementation, the German Fertilizer Ordinance (Fertilizer Ordinance, 2020), which regulates N inputs and N application times. They state that increased biogas production in recent years has impeded mitigation efforts. However, the development of N surplus and thus indirect N_2O emissions is heterogeneous (Figure 2 and Table 2). Most regions showed a steady decrease in N surplus, whereas a few exhibited much higher or similar N surpluses compared with 1995 (e.g., NW, NI-E, NI-W + HB) (see Häußermann et al. [2020] for further discussion). The new, spatially differentiated method to estimate indirect N_2O emissions from N leaching allows that these developments are taken into consideration in GHG reporting.

Although the model system used in this study calculated $\text{Frac}_{\text{LEACH}}$ at a higher spatial resolution, the considerable uncertainty around input data from the 1990s necessitated aggregation of the results at NUTS-1 level for the inventory. Differences in the N surplus of the calibration data and the inventory data mainly arose as a result of different data sources being used. More accurate data from the 2016 agricultural census and additional data sources (e.g., on manure transportation) could be used in RAUMIS, whereas such detailed data are not available for the whole reporting period of the GHG inventory.

5 | CONCLUSIONS

The model system RAUMIS-mGROWA-DENUZ was designed to quantify and monitor regional water body pollution with nutrients from

agricultural sources and to assess corresponding environmental policies. This study aimed to illustrate the model's capability to be also used to quantify indirect N_2O emissions for GHG reporting by deriving regionally differentiated $\text{Frac}_{\text{LEACH}}$ values. The application of the model system for GHG reporting allowed the inclusion of site-specific and spatially varying soil properties, hydrological conditions, climatic factors, and agricultural structure in the estimation of $\text{Frac}_{\text{LEACH}}$, which are not considered in the IPCC Tier 1 method. The use of N surplus rather than N input as the model driver allows the national GHG inventory to represent annual changes in cropping conditions, changes in N-use efficiency and the effects of N regulating policies and mitigation measures (The Council of the European Communities, 1991; Fertilizer Ordinance, 2020) aimed at reducing N surpluses. A combination of agricultural supply models with a water transport and a denitrification model and the use of high-resolution input data allow derivation of more accurate and less uncertain estimates of indirect N_2O emissions originating from NO_3^- leaching and runoff for national GHG inventories than the IPCC Tier 1 approach.

ACKNOWLEDGMENTS

The project AGRUM-DE was an important basis for this study. It is jointly funded by the Bund/Länder-Arbeitsgemeinschaft Wasser (LAWA) (<https://www.lawa.de/>) and the Thünen-Institut. We appreciate the contribution of Claire Tarring in editing the style and language of the manuscript. The remaining language mistakes, if any, were made by the authors.

Open access funding enabled and organized by Projekt DEAL.

DATA AVAILABILITY STATEMENT

Research data are not shared.

ORCID

Max Eysholdt  <https://orcid.org/0000-0001-9761-2991>

Maximilian Zinnbauer  <https://orcid.org/0000-0002-8278-2768>

REFERENCES

- Ackermann, A., Heidecke, C., Hirt, U., Kreins, P., Kuhr, P., Kunkel, R., Mahnkopf, J., Schott, M., Tetzlaff, B., Venohr, M., & Wendland, F. (2015). *Der Modellverbund AGRUM als Instrument zum landesweiten Nährstoffmanagement in Niedersachsen*. Johann Heinrich von Thünen-Institut. <https://doi.org/10.3220/REP1450256145000>
- Ackermann, A., Mahnkopf, J., Heidecke, C., & Venohr, M. (2016). Reducing agricultural nitrogen inputs in the German Baltic Sea catchment—trends and policy options. *Water Science and Technology*, 74(5), 1060–1068.
- Arheimer, B., Dahné, J., Donnelly, C., Lindström, G., & Strömqvist, J. (2012). Water and nutrient simulations using the HYPE model for Sweden vs. the Baltic Sea basin—influence of input-data quality and scale. *Hydrology Research*, 43(4), 315–329.
- Arnold, J. G., Srinivasan, R., Muttiah, R. S., & Williams, J. R. (1998). Large area hydrologic modeling and assessment part I: Model development. *Journal of the American Water Resources Association*, 34(1), 73–89.
- Blicher-Mathiesen, G., Andersen, H. E., & Larsen, S. E. (2014). Nitrogen field balances and suction cup-measured N leaching in Danish catchments. *Agriculture, Ecosystems & Environment*, 196, 69–75.
- Brink, C., van Grinsven, H., Jacobsen, B. H., Rabl, A., Gren, I.-M., Holland, M., Klimont, Z., Hicks, K., Brouwer, R., Dickens, R., Willems, J., Termansen, M., Velthof, G., Alkemade, R., van Oorschoot, M., & Webb, J. (2011).

- The European nitrogen assessment: Sources, effects and policy perspectives.* Cambridge University Press.
- Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R., & Zechmeister-Boltenstern, S. (2013). Nitrous oxide emissions from soils: How well do we understand the processes and their controls? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1621), 20130122. <https://doi.org/10.1098/rstb.2013.0122>
- Colombani, N., Gervasio, M. P., Castaldelli, G., & Mastrocicco, M. (2020). Soil conditioners effects on hydraulic properties, leaching processes and denitrification on a silty-clay soil. *Science of The Total Environment*, 733, 139342.
- Daniel-Gromke, J., Rensberg, N., Denysenko, V., Trommler, M., Reinholz, T., Völler, K., Beil, M., & Beyrich, W. (2017). Anlagenbestand Biogas und Biomethan - Biogaserzeugung und -nutzung in Deutschland. Deutsches Biomasseforschungszentrum. https://www.dbfz.de/fileadmin/user_upload/Referenzen/DBFZ_Reports/DBFZ_Report_30.pdf
- De Notaris, C., Rasmussen, J., Sørensen, P., & Olesen, J. E. (2018). Nitrogen leaching: A crop rotation perspective on the effect of N surplus, field management and use of catch crops. *Agriculture, Ecosystems & Environment*, 255, 1–11.
- Di, H. J., & Cameron, K. C. (2002). Nitrate leaching in temperate agroecosystems: Sources, factors and mitigating strategies. *Nutrient Cycling in Agroecosystems*, 64(3), 237–256.
- Environmental Protection Agency. (2021). *National inventory report 2021 - greenhouse gas emissions 1990 - 2019 reported to the United Nations framework convention on climate change.* <https://unfccc.int/documents/272856>
- Eurostat (2020). *Statistical regions in the European union and partner countries: NUTS and statistical regions 2021: 2020 edition.* European Union, Publications Office.
- Federal Environment Agency. (2021). *National inventory report for the German greenhouse gas inventory 1990 - 2019.* <https://unfccc.int/documents/273433>
- Federal Institute for Geosciences and Natural Resources (2018). Soil map of Germany 1:200,000 (BUEK200). BGR.
- Fertilizer Ordinance. (2020). Verordnung über die Anwendung von Düngemitteln, Bodenhilfsstoffen, Kultursubstraten und Pflanzenhilfsmitteln nach den Grundsätzen der guten fachlichen Praxis beim Düngen. https://www.gesetze-im-internet.de/d_v_2017/
- Fu, J., Gasche, R., Wang, N., Lu, H., Butterbach-Bahl, K., & Kiese, R. (2017). Impacts of climate and management on water balance and nitrogen leaching from montane grassland soils of S-Germany. *Environmental Pollution*, 229, 119–131.
- Fuchs, S., Scherer, U., Wander, R., Behrendt, H., Venohr, M., Opitz, D., Hillenbrand, T., Marscheider-Weidemann, F., & Götz, T. (2010). *Berechnung von Stoffeinträgen in die Fließgewässer Deutschlands mit dem Modell MONERIS: Nährstoffe, Schwermetalle und Polyzyklische aromatische Kohlenwasserstoffe.* <https://www.umweltbundesamt.de/publikationen/berechnung-von-stoffeintragen-in-flie遝gewaesser>
- Galloway, J. N., Aber, J. D., Erisman, J. W., Seitzinger, S. P., Howarth, R. W., Cowling, E. B., & Cosby, B. J. (2003). The nitrogen cascade. *Bioscience*, 53(4), 341–356.
- Gocht, A., & Röder, N. (2014). Using a Bayesian estimator to combine information from a cluster analysis and remote sensing data to estimate high-resolution data for agricultural production in Germany. *International Journal of Geographical Information Science*, 28(9), 1744–1764.
- Gömann, H., Julius, C., & Kreins, P. (2002). Quantifying impacts of different agri-environmental policies on the environment using the regional agri-environmental information system RAUMIS. In Pillmann, W. (Ed.), *Environmental communication in the information society.* (pp. 209–216). ISEP International Society for Environmental Protection.
- Groenendijk, P., Heinen, M., Klammler, G., Fank, J., Kupfersberger, H., Pisinaras, V., Gemitzi, A., Peña-Haro, S., García-Prats, A., Pulido-Velazquez, M., Perego, A., Acutis, M., & Trevisan, M. (2014). Performance assessment of nitrate leaching models for highly vulnerable soils used in low-input farming based on lysimeter data. *The Science of the Total Environment*, 499, 463–480.
- Groenendijk, P., Renaud, L. V., Roelmsa, J., Janssen, G. M. C. M., Janssen, S., Heerdink, R., Griffioen, J., & van der Grift, B. (2008). A new compliance checking level for nitrate in groundwater. *Modelling nitrate leaching and the fate of nitrogen in the upper five meter of the groundwater system.* RIVM rapport 680100006.
- Hashemi, F., Olesen, J. E., Jabloun, M., & Hansen, A. L. (2018). Reducing uncertainty of estimated nitrogen load reductions to aquatic systems through spatially targeting agricultural mitigation measures using groundwater nitrogen reduction. *Journal of Environmental Management*, 218, 451–464.
- Häußermann, U., Klement, L., Breuer, L., Ullrich, A., Wechsung, G., & Bach, M. (2020). Nitrogen soil surface budgets for districts in Germany 1995 to 2017. *Environmental Sciences Europe*, 32(1), 1–14.
- Heidecke, C., Ackermann, A., Kreins, P., Kuhr, P., Kunkel, R., Tetzlaff, B., Wendland, F., Mahnkopf, J., & Venohr, M. (2016). Erreichbarkeit der Nährstoffreduzierungsziele in Niedersachsen fraglich. *Wasser und Abfall: Boden, Altlasten, Umweltschutz; Fachmagazin für Wasserwirtschaft, Abfallwirtschaft, Bodenschutz, Altlasten und Umweltrecht*, 18(7-8), 21–25.
- Heidecke, C., Hirt, U., Kreins, P., Kuhr, P., Kunkel, R., Mahnkopf, J., Schott, M., Tetzlaff, B., Venohr, M., Wagner, A., & Wendland, F. (2015). *Entwicklung eines Instrumentes für ein flussgebietsweites Nährstoffmanagement in der Flussgebietseinheit Weser AGRUM+-Weser; Endbericht zum Forschungsprojekt.* (project report). Johann Heinrich von Thünen-Institut.
- Hennings, V. (2000). *Methodendokumentation Bodenkunde; Auswertungsmethoden zur Beurteilung der Empfindlichkeit und Belastbarkeit von Böden; mit 112 Tabellen* (2., überarb. Aufl., Vol. 1). Ad-Hoc-Arbeitsgruppe Boden der Geologischen Landesämter und der Bundesanstalt für Geowissenschaften und Rohstoffe der Bundesrepublik Deutschland; Bundesanstalt für Geowissenschaften und Rohstoffe; Schweizerbart.
- Henrichsmeyer, W., Cypris, C., Löhe, W., Meudt, M., Sander, R., & v Sothen, F. (1996). *Entwicklung des gesamtdeutschen Agrarsektormodells RAUMIS96; Endbericht Zum Kooperationsprojekt.* (project report). <https://dl.idcst.org/pdfs/files/1/b0c4f65f0c8f06bd3e8fe6e93103c3ed.pdf>
- Henseler, M., & Dechow, R. (2014). Simulation of regional nitrous oxide emissions from German agricultural mineral soils: A linkage between an agro-economic model and an empirical emission model. *Agricultural Systems*, 124, 70–82.
- Herrmann, F., Keller, L., Kunkel, R., Vereecken, H., & Wendland, F. (2015). Determination of spatially differentiated water balance components including groundwater recharge on the Federal State level-A case study using the mGROWA model in North Rhine-Westphalia (Germany). *Journal of Hydrology: Regional Studies*, 4, 294–312.
- Hirt, U., Kreins, P., Kuhn, U., Mahnkopf, J., Venohr, M., & Wendland, F. (2012). Management options to reduce future nitrogen emissions into rivers: A case study of the Weser river basin, Germany. *Agricultural Water Management*, 115, 118–131.
- Huang, T., Ju, X., & Yang, H. (2017). Nitrate leaching in a winter wheat-summer maize rotation on a calcareous soil as affected by nitrogen and straw management. *Scientific Reports*, 7(1), 1–11.
- IPCC. (1996). *Revised 1996 IPCC guidelines for national greenhouse gas inventories; reference manual (Reference volume 3).* IPCC/OECD/IEA. UK Meteorological Office.
- IPCC. (2006). *2006 IPCC guidelines for national greenhouse gas inventories program, volume 4, Agriculture, forestry and other land use.* Institute for Global Environmental Strategies, Japan. <https://www.ipcc-nggip.iges.or.jp/public/2006gl/vol4.html>
- IPCC. (2014). *Climate change 2014: Synthesis report. Contribution of working groups I, II and III to the fifth assessment report of the intergovernmental panel on climate change.* (R. K. Pachauri & L. A. Meyer, Eds.). Intergovernmental Panel on Climate Change.

- IPCC. (2019). *2019 refinement to the 2006 IPCC guidelines for national greenhouse gas inventories*. Intergovernmental Panel on Climate Change.
- Klages, S., Heidecke, C., & Osterburg, B. (2020). The impact of agricultural production and policy on water quality during the dry year 2018, a case study from Germany. *Water*, 12(6), 1519. <https://doi.org/10.3390/w12061519>
- Köhne, C., & Wendland, F. (1992). Modellgestützte Berechnung des mikrobiellen Nitratabbaus im Boden. Interner Bericht KFA-STE-IB-1/92. Forschungszentrum Jülich GmbH, Programmgruppe STE Systemforschung und Technologische Entwicklung.
- Kreins, P., Behrendt, H., Gömann, H., Heidecke, C., Hirt, U., Kunkel, R., Seidel, K., Tetzlaff, B., & Wendland, F. (2009). Analyse von Agrar- und Umweltmaßnahmen im Bereich des landwirtschaftlichen Gewässerschutzes vor dem Hintergrund der EG-Wasserrahmenrichtlinie in der Flussgebietseinheit Weser; AGRUM-Weser. *Landbauforschung Sonderheft* (No. 336). Johann Heinrich von Thünen-Institut.
- Kreins, P., Gömann, H., Herrmann, S., Kunkel, R., & Wendland, F. (2007). Integrated agricultural and hydrological modeling within an intensive livestock region. In J. D. Erickson, F. Messner, & I. Ring (Eds.), *Ecological economics of sustainable watershed management*. (pp. 113–142). Emerald Group Publishing.
- Kreins, P., Heidecke, C., Gömann, H., Hirt, U., & Wendland, F. (2011). Möglichkeiten und Grenzen der wissenschaftlichen Politikanalyse zur Umsetzung der Wasserrahmenrichtlinie - Anwendung eines hydroökonomischen Modellverbundes für das Weser Einzugsgebiet. *Gesellschaft für Wirtschafts- und Sozialwissenschaften des Landbaus*, 46, 325–335.
- Krüll, H. (1988). Erstellung einer Stickstoffbilanz in den Kreisen der BRD. *Endbericht Zum Forschungsvorhaben Wirkungen Agrarpolitischer Maßnahmen Auf Ziele von Umwelt-, Natur- Und Landschaftsschutz* (pp. 8–39).
- Kuhn, U., Schmidt, B., Heidecke, C., Kreins, P., Ackermann, A., Wendland, F., Tetzlaff, B., Kunkel, R., Venohr, M., & Mahnkopf, J. (2016). Nährstoffmanagement in der Flussgebietseinheit Weser im Spannungsfeld zwischen Wasserwirtschaft und Landwirtschaft - Ergebnisse des Projektes AGRUM (project report). *Korrespondenz Wasserwirtschaft: KW; Organ der DWA; Wasser, Boden, Natur*, 9(4), 218–225.
- Kuhr, P., Haider, J., Kreins, P., Kunkel, R., Tetzlaff, B., Vereecken, H., & Wendland, F. (2013). Model based assessment of nitrate pollution of water resources on a federal state level for the dimensioning of agro-environmental reduction strategies. *Water resources management*, 27(3), 885–909.
- Kuhr, P., Kunkel, R., Tetzlaff, B., & Wendland, F. (2014). *Räumlich differenzierte Quantifizierung der Nährstoffeinträge in Grundwasser und Oberflächengewässer in Sachsen-Anhalt unter Anwendung der Modellkombination GROWA-WEKU-MEPHos; Endbericht eines Forschungsvorhabens im Auftrag des Landesbetriebs für Hochwasserschutz und Wasserwirtschaft Sachsen-Anhalt*. (project report). Forschungszentrum Jülich GmbH.
- Kunkel, R., Herrmann, F., Kape, H.-E., Keller, L., Koch, F., Tetzlaff, B., & Wendland, F. (2017). Simulation of terrestrial nitrogen fluxes in Mecklenburg-Vorpommern and scenario analyses how to reach n-quality targets for groundwater and the coastal waters. *Environmental Earth Sciences*, 76(4), 1–19.
- Kunkel, R., Kreins, P., Tetzlaff, B., & Wendland, F. (2010). Forecasting the effects of EU policy measures on the nitrate pollution of groundwater and surface waters. *Journal of Environmental Sciences*, 22(6), 872–877.
- Kunkel, R., & Wendland, F. (2006). Diffuse Nitratreinträge in die Grund- und Oberflächengewässer von Rhein und Ems, Ist-Zustands- und Massnahmenanalysen (project report). *Schriften des Forschungszentrums Jülich - Reihe Umwelt* (Vol. 62). Forschungszentrum Jülich.
- Landesamt für Statistik Niedersachsen. (2019). Request for agricultural compost application data of the federal states: Information via mail on 13.03.2019.
- Manevski, K., Børgesen, C. D., Li, X., Andersen, M. N., Abrahamsen, P., Hu, C., & Hansen, S. (2016). Integrated modelling of crop production and nitrate leaching with the Daisy model. *MethodsX*, 3, 350–363.
- Ministry for the Environment. (2021). *New Zealand's greenhouse gas inventory 1990–2019*. Ministry for the Environment New Zealand.
- Molina-Navarro, E., Trolle, D., Martínez-Pérez, S., Sastre-Merlín, A., & Jeppesen, E. (2014). Hydrological and water quality impact assessment of a Mediterranean limno-reservoir under climate change and land use management scenarios. *Journal of Hydrology*, 509, 354–366.
- Müller, U., & Raissi, F. (2002). *Arbeitshilfe für bodenkundliche Stellungnahmen und Gutachten im Rahmen der Grundwassernutzung*. Schweizerbart Science Publishers.
- Räbiger, T., Andres, M., Hegewald, H., Kesenheimer, K., Köbke, S., Quinones, T. S., Böttcher, U., & Kage, H. (2020). Indirect nitrous oxide emissions from oilseed rape cropping systems by NH₃ volatilization and nitrate leaching as affected by nitrogen source, N rate and site conditions. *European Journal of Agronomy*, 116, 126039. <https://doi.org/10.1016/j.eja.2020.126039>
- Ravishankara, A. R., Daniel, J. S., & Portmann, R. W. (2009). Nitrous oxide (N₂O): The dominant ozone-depleting substance emitted in the 21st century. *Science*, 326(5949), 123–125.
- Rösemann, C., Haenel, H.-D., Cora, V., Dämmgen, U., Döring, U., Wulf, S., Eurich-Menden, B., Freibauer, A., Döhler, H., Schreiner, C., Osterburg, B., & Fuß, R. (2021). Calculations of gaseous and particulate emissions from German agriculture 1990–2019; report on methods and data (RMD) submission 2021. Thünen Report (No. 84). Johann Heinrich von Thünen-Institut.
- Schmidt, B., Kuhn, U., Trepel, M., Kreins, P., Zinnbauer, M., Eysholdt, M., Osterburg, B., Löw, P., Wendland, F., Herrmann, F., Kunkel, R., Tetzlaff, B., Wolters, T., Venohr, M., & Nguyen, H. (2020). Modellansatz zur Bestimmung der Nährstoffbelastung und ihrer Reduktion in allen deutschen Flussgebieten. *Wasser und Abfall*, 22(1–2), 33–38.
- Šimůnek, J., van Genuchten, M. T., & Šejna, M. (2008). Development and applications of the HYDRUS and STANMOD software packages and related codes. *Vadose Zone Journal*, 7(2), 587–600.
- Statistisches Bundesamt (StBA FS3 R3.1.3). Gemüseerhebung – Anbau und Ernte von Gemüse und Erdbeeren. StBA, FS3 (No. R3.1.3). https://www.destatis.de/DE/Themen/Branchen-Unternehmen/Landwirtschaft-Forstwirtschaft-Fischerei/Feldfruechte-Gruenland/_inhalt.html#sprg239470
- Statistisches Bundesamt (StBA FS3 R3.2.1). Wachstum und Ernte – Feldfrüchte. In StBA, FS3 (No. R3.2.1). https://www.destatis.de/DE/Themen/Branchen-Unternehmen/Landwirtschaft-Forstwirtschaft-Fischerei/Obst-Gemuese-Gartenbau/Publikationen/_publikationen-innen-gemuese.html
- Statistisches Bundesamt (StBA) (2019). Produzierendes Gewerbe – Düngemittelversorgung; Fachserie 4 Reihe 8.2. <https://www.destatis.de/DE/Themen/Branchen-Unternehmen/Industrie-Verarbeitendes-Gewerbe/Publikationen/Downloads-Fachstatistiken/duengemittelversorgung-vierteljahr-2040820193234.pdf>
- Statistisches Bundesamt (StBA) (StBA various years). *Wasserwirtschaft: Klärschlamm Entsorgung aus der öffentlichen Abwasserbehandlung*. Statistisches Bundesamt.
- Steffen, W., Richardson, K., Rockström, J., Cornell, S. E., Fetzer, I., Bennett, E. M., Biggs, R., Carpenter, S. R., de Vries, W., de Wit, C. A., Folke, C., Gerten, D., Heinke, J., Mace, G. M., Persson, L. M., Ramanathan, V., Reyers, B., & Sörlin, S. (2015). Sustainability. Planetary boundaries: Guiding human development on a changing planet. *Science*, 347(6223), 1259855. <https://doi.org/10.1126/science.1259855>
- Tetzlaff, B., Keller, L., Kuhr, P., Kreins, P., Kunkel, R., & Wendland, F. (2017). *Nährstoffeinträge ins Grundwasser und die Oberflächengewässer Schleswig-Holsteins unter Anwendung der Modellkombination RAUMIS-GROWA-WEKU-MEPHos; Endbericht zum Forschungsprojekt*. (project report). Forschungszentrum Jülich GmbH.
- Tetzlaff, B., Kreins, P., Kuhr, P., Kunkel, R., & Wendland, F. (2016). *Quantifizierung der Stickstoff- und Phosphoreinträge ins Grundwasser und die Oberflächengewässer Thüringens mit eintragspfadbezogener und*

- regionaler Differenzierung; Endbericht zum Forschungsprojekt. (project report). Forschungszentrum Jülich GmbH.
- Tetzlaff, B., Kunkel, R., Ta, P., Wendland, F., & Wolters, T. (2021). Fortführung der Nährstoffmodellierung Mecklenburg-Vorpommern; Endbericht zum Forschungsprojekt (project report). Forschungszentrum Jülich GmbH.
- Tetzlaff, B., Schreiner, H., Vereecken, H., & Wendland, F. (2009). Modellgestützte Analyse signifikanter Phosphorbelastungen in hessischen Oberflächengewässern aus diffusen und punktuellen Quellen. *Reihe Energie & Umwelt /Energy & Environment*. (No. 29). Forschungszentrum Jülich GmbH.
- The Council of the European Communities (1991). Council directive 91/676/EEC of 12 December 1991 concerning the protection of waters against pollution caused by nitrates from agricultural sources; council directive 91/676/EEC. <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex%3A31991L0676>
- The European Parliament and The Council of the European Union. (2000). Directive 2000/60/EC of the European parliament and of the council of 23 October 2000 establishing a framework for community action in the field of water policy; directive 2000/60/EC. <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:32000L0060>
- Tian, H., Xu, R., Canadell, J. G., Thompson, R. L., Winiwarter, W., Suntharalingam, P., Davidson, E. A., Ciais, P., Jackson, R. B., Janssens, Maenhout, G., Prather, M. J., Regnier, P., Pan, N., Pan, S., Peters, G. P., Shi, H., Tubiello, F. N., Zaehle, S., Zhou, F., ... Yao, Y. (2020). A comprehensive quantification of global nitrous oxide sources and sinks. *Nature*, 586(7828), 248–256.
- UNFCCC (2021). United Nations Framework Convention on Climate Change national inventory submissions. <https://unfccc.int/ghg-inventories-annex-i-parties/2021>
- van der Zee, T., Bannink, A., van Bruggen, C., Groenestein, K., Huijsmans, J., van der Kolk, J., Lagerwerf, L., Luesink, H., Velthof, G., & Vonk, J. (2021). Methodology for estimating emissions from agriculture in the Netherlands. *Calculations for CH₄, NH₃, N₂O, NO_x, NMVOC, PM₁₀, PM_{2.5} and CO₂ using the national emission model for agriculture (NEMA)—update 2021*. Rijksinstituut voor Volksgezondheid en Milieu. <https://doi.org/10.21945/RIVM-2021-0008>
- Velthof, G., & Mosquera, J. (2011). Calculation of nitrous oxide emission from agriculture in the Netherlands; update of emission factors and leaching fraction. *Alterra* (No. 2151).
- Wang, Y., Ying, H., Yin, Y., Zheng, H., & Cui, Z. (2019). Estimating soil nitrate leaching of nitrogen fertilizer from global meta-analysis. *Science of the Total Environment*, 657, 96–102.
- Wendland, F. (1994). Modelling the nitrate flow in the ground-water provinces of the “old” federal states of the Federal Republic of Germany. *Ecological modelling*, 75, 385–397.
- Wendland, F., Bach, M., & Kunkel, R. (1998). The influence of nitrate reduction strategies on the temporal development of the nitrate pollution of soil and groundwater throughout Germany - a regionally differentiated case study. *Nutrient Cycling in Agroecosystems*, 50(1/3), 169–181.
- Wendland, F., Behrendt, H., Gömann, H., Hirt, U., Kreins, P., Kuhn, U., Kunkel, R., & Tetzlaff, B. (2009). Determination of nitrogen reduction levels necessary to reach groundwater quality targets in large river basins: The Weser basin case study, Germany. *Nutrient Cycling in Agroecosystems*, 85(1), 63–78.
- Wendland, F., Behrendt, H., Hirt, U., Kreins, P., Kuhn, U., Kuhr, P., Kunkel, R., & Tetzlaff, B. (2010). Analyse von Agrar- und Umweltmaßnahmen zur Reduktion der Stickstoffbelastung von Grundwasser und Oberflächengewässer in der Flussgebietseinheit Weser. *Hydrologie und Wasserbewirtschaftung*, 54(4), 231–244.
- Wendland, F., Bergmann, S., Eisele, M., Gömann, H., Herrmann, F., Kreins, P., & Kunkel, R. (2020). Model-based analysis of nitrate concentration in the leachate—The North Rhine-Westfalia case study, Germany. *Water*, 12(2), 550.
- Wendland, F., Bogen, H., Goemann, H., Hake, J. F., Kreins, P., & Kunkel, R. (2005). Impact of nitrogen reduction measures on the nitrogen loads of the river Ems and Rhine (Germany). *Physics and Chemistry of the Earth, Parts A/B/C*, 30(8–10), 527–541.
- Wendland, F., Herrmann, F., Kunkel, R., Ta, P., Tetzlaff, B., & Wolters, T. (2021). Quantifizierung der Stickstoff- und Phosphoreinträge ins Grundwasser und in die Oberflächengewässer in Rheinland-Pfalz mit eintragspfadbezogener und regionaler Differenzierung; Projektabschlussbericht. (project report). Forschungszentrum Jülich GmbH.
- Wendland, F., Herrmann, F., Kunkel, R., Tetzlaff, B., & Wolters, T. (2020). AGRUM-DE - Stickstoff- und Phosphoreinträge ins Grundwasser und die Oberflächengewässer Deutschlands mit eintragspfadbezogener und regionaler Differenzierung; 3. Zwischenbericht AGRUM-DE. (project report). Forschungszentrum Jülich GmbH.
- Wendland, F., Keller, L., Kuhr, P., & Tetzlaff, B. (2015). Regional differenzierte Quantifizierung der Nährstoffeinträge in das Grundwasser und in die Oberflächengewässer Mecklenburg-Vorpommerns unter Anwendung der Modellkombination GROWA-DENUZ-WEKU-MEPHOS; Endbericht zum Forschungsprojekt im Auftrag des Landesamts für Umwelt, Naturschutz und Geologie Mecklenburg-Vorpommern. (project report). Landesamt für Umwelt, Naturschutz und Geologie Mecklenburg-Vorpommern.
- Wendland, F., Keller, L., Kuhr, P., Tetzlaff, B., Heidecke, C., Kreins, P., Wagner, A., & Trepel, M. (2014). Räumlich differenzierte Quantifizierung der Stickstoffeinträge ins Grundwasser und die Oberflächengewässer Schleswig-Holsteins. *Korrespondenz Wasserwirtschaft*, 6(14), 327–332.
- Wienhaus, S., Höper, H., Eisele, M., Meesenburg, H., & Schäfer, W. (2008). Nutzung bodenkundlich-hydrogeologischer Informationen zur Ausweisung von Zielgebieten für den Grundwasserschutz. In Landesamt für Bergbau, Energie und Geologie (Ed.), *GeoBerichte* (1. Auflage). Landesamt für Bergbau, Energie und Geologie. https://doi.org/10.48476/geober_9_2008
- Wolters, T., Cremer, N., Eisele, M., Herrmann, F., Kreins, P., Kunkel, R., & Wendland, F. (2021). Checking the plausibility of modelled nitrate concentrations in the leachate on federal state scale in Germany. *Water*, 13(2), 226. <https://doi.org/10.3390/w13020226>
- Zhang, Y., & Yu, Q. (2020). Identification of current research intensity and influence factors of agricultural nitrogen loss from cropping systems. *Journal of Cleaner Production*, 276, 123308. <https://doi.org/10.1016/j.jclepro.2020.123308>
- Zhou, M., & Butterbach-Bahl, K. (2014). Assessment of nitrate leaching loss on a yield-scaled basis from maize and wheat cropping systems. *Plant and Soil*, 374(1), 977–991.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Eysholdt, M., Kunkel, R., Rösemann, C., Wendland, F., Wolters, T., Zinnbauer, M., & Fuß, R. (2022). A model-based estimate of nitrate leaching in Germany for GHG reporting. *Journal of Plant Nutrition and Soil Science*, 185, 850–863. <https://doi.org/10.1002/jpln.202200119>